Title

Prognosis of stroke subtypes in whole population health systems data: a matched cohort

study

Authors

Alice Hosking MBBS[1], Matthew H Iveson PhD[1], Laura Sherlock PhD[1], Mome Mukherjee PhD

[2], Claire Grover PhD[3], Beatrice Alex PhD[5,6,7], Sai Parepalli[1], Grant Mair MD[1], Fergus Doubal

PhD[1], Prof. Heather Whalley PhD[1], Richard Tobin MA[3], Prof Joanna Wardlaw PhD[1,8], Prof

Rustam Al-Shahi Salman PhD[1], Prof William N Whiteley PhD[1,2,4]

1. Institute for Neuroscience and Cardiovascular Research, The University of Edinburgh

2. Usher Institute, The University of Edinburgh

3. School of Informatics, The University of Edinburgh

4. British Heart Foundation Data Science Centre, HDRUK, London

5. School of Literatures, Languages and Cultures, The University of Edinburgh

6. Edinburgh Futures Institute, The University of Edinburgh

7. Advanced Care Research Centre, The University of Edinburgh

8. UK Dementia Research Institute Centre and BHF-UKDRI Centre for Vascular Dementia

Research, The University of Edinburgh, Edinburgh, UK

Corresponding author: Dr Alice Hosking. E: alice.hosking@ed.ac.uk. T: +4407846676318.

A: Chancellor's Building, 49 Little France Crescent, Edinburgh EH16 4SB.

Short title: Stroke subtypes in health systems data

Word count: 3270 words (main text); 5431 words (whole document).

Abstract

Background

Outcome after stroke varies according to stroke subtype by location, but healthcare

systems data studies do not include subtyping information. We linked natural language

processing (NLP) of brain imaging reports to routinely collected data to estimate risk of

death and other outcomes after stroke subtypes in a nationwide dataset.

Methods

We applied a previously validated NLP algorithm to all CT and MRI head scan reports in

Scotland between 2010 and 2018. We linked the reports to hospital readmissions,

prescriptions and death data to identify and characterize people with stroke, and to

categorize into deep and cortical ischemic stroke, deep and lobar intracerebral

hemorrhage (ICH), subarachnoid hemorrhage, and subdural hemorrhage. We used a

matched cohort design, and age- and sex-matched four controls per case who never had a

stroke. By subtype, we estimated rehospitalization with stroke, myocardial infarction (MI),

cancer, dementia, epilepsy and death, accounting for confounders and competing risk of

death.

Results

From 785,331 people with a head scan, we identified 64,219 with clinical stroke

phenotypes (mean age 73.4yrs, 49.5% male), and subtyped 12,616 with deep ischemic

stroke; 14,103 with cortical ischemic stroke; 1,814 with deep ICH; and 1,456 with lobar

ICH. There was higher absolute rate of 1-year hospital readmission for lobar compared

with deep ICH (4.9% [95%CI 3.9% - 6.1%] vs 3.4% [2.6% - 4.3%]), higher risk of dementia

beyond 6 months after lobar ICH compared to controls than for other stroke subtypes (aHR

3.5 [2.3-5.3]); and higher risk of MI within 6 months of cortical ischemic stroke than for

other stroke subtypes (aHR 4.6 [3.4-6.3]).

Conclusions

NLP of free-text reports linked to coded data successfully subtyped stroke at scale, and we

estimated risk of clinically relevant outcomes. Future work should use free text to enable

large-scale audit and epidemiology of people with stroke.

Key words: Routinely Collected Health Data, Natural Language Processing, Ischemic

Stroke, Intracerebral Hemorrhage

53
54

Introduction

Outcome after stroke varies by stroke type and stroke location, but existing studies of stroke subtypes by location are limited by coding systems ("ontologies") in healthcare data; and in cohort studies by cohort size, consent bias, and limited follow up. [1–3]

Healthcare systems data (HSD) provide the opportunity to examine outcomes after stroke in a large number of unselected people, reducing uncertainty and increasing relevance to clinical practice. [4–7] However, phenotyping in HSD is constrained by coding ontologies. Health systems typically record ischemic stroke, intracerebral hemorrhage (ICH) and subarachnoid hemorrhage (SAH) with ontologies such as the 10th revision of the International Classification of Diseases (ICD-10), [8,9] but ICD-10 codes give no information about ischemic stroke location, most people with intracerebral hemorrhage do not have a stroke location recorded, and many patients are not coded as ischemic or hemorrhagic stroke. [10] Although ICD-11 expands cerebrovascular granularity and SNOMED-CT can represent subtypes, they are not adopted widely for use in hospital coding systems.

Uncoded data (e.g., text or images) could improve phenotyping. Imaging reports are concise, clinically focused, and can be read with natural language processing (NLP). We previously developed and validated a rules-based NLP system for brain imaging reports, Edinburgh Information Extraction for Radiology reports (EdIE-R), which accurately tags MRI and CT head reports with stroke and subdural hemorrhage (SDH) types and locations. [11–13] Demonstrating that uncoded data can improve phenotyping of stroke would be useful to healthcare and research.

In this study, we identified cortical and deep ischemic strokes, and lobar and deep ICH

with NLP from free-text CT and MRI head scan reports in Scotland and linked coded HSD,

and estimated risks of death and other outcomes after stroke subtypes by location. We

age- and sex-matched people with stroke to controls without stroke. We examined

outcomes relevant to stroke survivors: hospital readmission, cancer, epilepsy, dementia,

myocardial infarction (MI) and death.

Methods

Data Sources

We accessed CT and MRI head scan reports in adults (≥18 years) from Scottish Medical

Imaging from 1st January 2010 to 31st August 2018.[14] Public Health Scotland

deterministically linked imaging data with a unique patient identifier (Community Health

Index number) to: general and psychiatric hospital admissions (Scottish Morbidity Record

(SMR)01 and SMR04); the Scottish Cancer Registry (SMR06); community prescriptions

(Prescribing Information System, from 1st January 2007 to 31st December 2020); and death

records from NRS Scotland (from 1st January 2010 to 31st December 2020). Prescription

data included drugs for vascular diseases, dementia and epilepsy (Supplement 2).

We tagged scan reports with an NLP algorithm developed and validated with a subset of

the Scottish data used in this study. Our previous work showed that the algorithm had

good precision and reasonable recall for each phenotype.[11–13] The algorithm generated the

following tags from text in the radiology report: ischemic stroke, ICH, SAH, SDH and stroke

of unknown type (when a stroke was identified but not the type). Ischemic stroke and ICH

were subclassified as recent or old, and cortical (for ischemic stroke), lobar (for ICH) or

deep (i.e. in deep brain structures). Additional tags included atrophy and white matter

hypoattenuation (on CT) or hyperintensities (on MRI) (WMH).

Population

Almost all healthcare in Scotland is delivered by the National Health Service (NHS). People

with a stroke can present either to primary care, where they may be referred on to the

emergency department or for outpatient assessment; or directly to the emergency

department; or be identified only at death, via autopsy. People in the emergency

department may either be discharged, be admitted to hospital, or die in the emergency

department. Primary data is not currently available in Scotland, and outpatient and

emergency department data does not code stroke types. We defined cases as people with

all the following: a stroke/SDH ICD-10 code in hospital admission and death data in the

main condition/primary cause of death field (code lists in Supplement 2; a head scan

within 30 days of the ICD-10 code; and no history of recorded stroke in hospital admission

between Scotland from 1st January 2010 to 31st August 2018. We included spontaneous

SDH as a spontaneous vascular event that leads to focal or global neurological symptoms.

Follow-up for outcomes continued to 31st December 2020.

Stroke Cohort and Matched Cohort

We defined six stroke (or SDH) categories: ischemic stroke, ICH, SAH, SDH, stroke

unspecified and non-traumatic hemorrhage unspecified; and four location subtypes:

119 cortical and deep ischemic stroke, and lobar and deep ICH. For each case identified in

120 hospital or death records, we compared the stroke-related ICD-10 code and NLP tag and

121 used an algorithm to assign diagnosis where the two differed (Supplement 1, eTable 1). The

122 algorithm prioritized common stroke types (e.g. ischemic stroke over intracerebral

123 hemorrhage), and recent strokes over old. We then used NLP to assign location subtypes

124 for ischemic stroke and ICH. We removed people with head trauma ICD-10 codes within

125 seven days of ICH, or brain cancer codes within one year of ICH, as these indicate bleeding

126 due to non-vascular pathology (brain tumor).

127 A matched cohort without recorded stroke/SDH was drawn from people with a record in

128 hospital admission data, community prescription data or death data. For each case, four

129 controls were matched on year of birth (+/- 2 years) and sex. Controls had to be alive at the

130 event date of their matched case.

131 Follow-up began at admission date for the index event (for controls, the date of their

132 matched case's index event) and stopped at date of death or each outcome. We did not

133 have reliable emigration data, but less than 2% of the Scottish population emigrate

134 annually.[15]

135 **Outcomes**

136 We defined death, readmission with stroke, dementia, epilepsy, dementia, cancer and MI

137 (Supplement 2). We ascertained outcome events for stroke/SDH types and subtypes which

138 correspond to defined pathological phenotypes: ischemic stroke, ICH, SAH and SDH. We

139 tabulated the most common underlying causes of death for each stroke type.

Readmission with stroke was defined as hospital admission with a stroke/SDH ICD-10

code in the primary position, starting from date of discharge after the index event. For

cancer, epilepsy and MI, we included relevant ICD-10 codes in the primary position in

hospital admission data or underlying cause of death from the stroke date onwards. We

included dementia in any position, but excluded people with events coded on the same

admission as the index stroke, which likely represented pre-existing dementia. We also

identified prevalent and incident epilepsy and dementia through prescriptions of anti-

epileptic or dementia drugs before or after the index date; and prevalent and incident

cancer cases in cancer registry data.

Covariates

Atrial fibrillation, diabetes and hypertension were identified in hospital records with ICD-10

codes in any position, before the date of defining stroke. Additionally, hypertension and

diabetes mellitus were identified with antihypertensive or diabetic medication

prescriptions in the year prior to the index event (Supplement 2). We identified statin,

anticoagulant and antiplatelet prescriptions in the year before the index event. We

recorded the number of hospital admissions and prescriptions dispensed in the year

before the index event. Additional phenotypes identified with NLP were atrophy and WMH.

Statistical Methods

We calculated cumulative incidence of each event type, accounting for competing risk of

death and censoring on death or 31st December 2020, and report by sex and age group (≤70

years and >70 years). We used cause-specific Cox proportional hazard models to estimate

adjusted hazard ratios (aHRs) with 95% CIs, adjusted for potential confounders (age, sex,

diabetes, hypertensive, atrial fibrillation, statins, anticoagulants, antiplatelets, number of

prescribed medicines in previous year, number of hospital admissions in previous year).

We compared each stroke type and subtype with their control group for each outcome,

except for readmission with stroke, where we compared stroke types to ischemic stroke,

as those without stroke could not be readmitted. For stroke subtypes, we compared deep

strokes to cortical/lobar.

As events were much more frequent in the first 6 months after stroke, the proportional

hazards assumption was violated, so we split our analysis into two periods: the first 6

months after stroke, and 6 months onwards. For each outcome, we confirmed the

proportional hazards assumption with visual inspection of plots of Schoenfeld residuals,

and confirmed the linearity of continuous input covariates or their logarithm by inspection

of plots of Martingale residuals.

Analyses were performed within the NHS Scotland Trusted Research Environment in R,

version 4.4.2, using the packages survival and survminer.

Standard Protocol Approvals, Registrations, and Patient Consents

The study was approved by the NRES Committee Northwest - Greater Manchester East

ethics committee (15/NW/0719), and the Public Benefit and Privacy Panel for Health and

Social Care (1516-0219) of NHS Scotland. Participant consent was not required.

Data Availability

Data is restricted as per Public Health Scotland guidelines, as it relates to individual health

records. Data can be made available within the National Safe Haven upon application and

approval by Public Health Scotland and the Public Benefit and Privacy panel.

Results

Of 5,667,954 people with any record in prescription, imaging, hospital or death data,

785,331 had a CT or MRI head scan. Of these, 79,045 had an ICD-10 stroke code recorded

at any time, and 66,917 had brain imaging within 30 days of a recorded stroke (Supplement

1, eFigure 1). The median follow-up time of this population was 3.3 years (IQR 0.46 – 5.82

years) and of their matched controls was 4.9 years (IQR 3.1 years to 7.3 years).

In hospital admission and death data, 34,739 people had an ICD-10 diagnosis of ischemic

stroke, 7,075 had ICH and 17,472 had unspecified stroke within 30 days of a head scan. No

further subtyping was possible. We selected the brain scan closest in time to the stroke

diagnosis, of which 64,178 were CT head scans, and 2,739 were MRI head scans. After

using NLP, 64,219 people had recognized clinical stroke phenotypes (mean age 73.4yrs,

49.5% male). 51,339 people had ischemic stroke [12,616 (24.6%) deep, 14,103 (27.5%)

cortical and 24,620 (48.0%) uncertain or multiple locations], 7,182 had ICH [1,814 (25.2%)

deep, 1,456 (20.3%) lobar and 3,912 (54.5%) uncertain or multiple locations], 2,984 had

SAH, 2,714 had SDH, and 2,618 unspecified stroke. Using NLP reduced the proportion of

people with unspecified stroke from 26.1% to 3.4% (Supplement 1, eFigure 2).

Compared with cortical ischemic stroke, those with deep ischemic stroke were more often

male (52.2% vs 49.9%), had lower prevalence of hypertension (66.6% vs 71.6%) and atrial

fibrillation (12.1% vs 18.9%), and higher rates of atrophy (52.8% vs 48.6%) and WMH

(68.5% vs 59.3%). Compared with lobar ICH, more of those with deep ICH were male

(50.3% vs 45.1%) and fewer had taken antiplatelets (33.9% vs 40.6%) (Table 1).

Readmission with stroke Over follow-up, 5,621 (8.71%) of people with stroke/SDH were

subsequently admitted with stroke/SDH. Readmission with stroke within 6 months

occurred in 2.7% to 2.9% of people with ischemic stroke, and 2.3% to 4.6% with ICH

(Supplement 1, eFigure 4), varying by age and sex. Those with lobar ICH were more likely to

be readmitted than those with deep ICH (aHR 1.71 (1.15-2.54), Figure 1) but there was no

difference between those with cortical ischemic stroke compared with deep (aHR 1.14

(0.99-1.31)). Beyond 6 months, there were no significant differences between stroke

subtypes.

MI During follow-up, 2680 of people with stroke/SDH had MI (number rounded to nearest

10 for data suppression purposes). The aHR for MI early after stroke (<6 months) was

highest for those with ischemic stroke (aHR 3.35 (2.79-4.03)) and SDH (aHR 3.97 (1.73-

9.1)). Among subtypes, the highest aHR was for those with cortical ischemic stroke (aHR

4.6 (3.35-6.31), Figure 2).

Dementia 5139/59720 (8.6%) of people with stroke, without pre-existing dementia,

developed dementia. Early after stroke (<6 months), the aHR for dementia was similar for

stroke types and subtypes. Late after stroke (>6 months), compared with matched

controls, the aHR for lobar ICH (3.49 (2.3-5.29)) was higher than for deep ICH (2.27 (1.47-

3.49)), cortical ischemic stroke (2.16 (1.9-2.46)) and deep ischemic stroke (1.94 (1.71-

2.21)) (Figure 3). Accounting for competing risk of death, cumulative incidence of dementia

by 5 years in men over 70 years was 17% for lobar ICH, 6.5% for deep ICH, 11% for cortical

ischemic stroke, and 10% for deep ischemic stroke (Figure 4).

Cancer During follow-up, 5424/56610 (9.6%) of people with stroke without pre-existing

cancer, developed cancer compared with 13.1% of matched controls. Early after stroke

(<6 months), there was elevated cancer risk for people with cortical ischemic stroke (aHR

1.54 (1.14-2.08); deep ischemic stroke (aHR 1.41 (1.03-1.93), and deep ICH (aHR 2.48

(1.01-6.05)). The aHR for lobar ICH was consistent with no increase (2.32 (0.85-6.37),

Figure 2). Beyond 6 months, we detected no difference in cancer incidence between

people without stroke and people with any stroke subtypes (Figure 3).

Epilepsy During follow-up, 4515/62548 (7.2%) of people with stroke, without preexisting

epilepsy, developed epilepsy. Cumulative incidence, accounting for competing risk of

death, was higher for those with cortical and lobar strokes (eFigure 4). Early after stroke the

aHR for epilepsy was higher for cortical than deep ischemic stroke (3.99 (3.18-5) vs 2.94

(2.26-3.84)), and for lobar than for deep ICH (10.98 (6.82-19.08) vs 3.42 (1.49-7.86), Figure

2). Late after stroke (>6 months), the same pattern was seen among subtypes, although

the aHRs were generally greater (Figure 3).

Death During follow-up, 36,914 (55.9%) of people with stroke died (Supplement 1, eFigure

5). Almost half (45.8%) of those with ICH died in the first 6 months, compared to 21.7%

with ischemic stroke. Both short- and long-term survival declined with increasing age

(Figure 5).

Early after stroke (<6 months), compared with matched controls, the aHR for death was

higher for ICH than ischemic stroke (23.06 (20-79-25.58) vs 8.63 (8.22-9.05)), and the aHR

for death was higher for cortical than deep ischemic stroke (10.9 (10.02-11.83) vs 6.91

(6.24-7,66)). Lobar and deep ICH had similar aHR (21.3 (16.8-27.0) vs 19.2 (15.4-23.9)),

Figure 2). Beyond 6 months, cortical ischemic stroke continued to have higher aHR for

death than deep ischemic stroke (2.04 (1.92-2.17) vs 1.86 (1.75-1.97)) (Figure 3).

The most common cause of death among people with ischemic stroke was unspecified

stroke; for ICH, SAH and SDH, it was the recorded stroke type. Among matched controls,

the most common cause of death was unspecified dementia (Supplement 1, eTable 4).

11,774 people with stroke/SDH died had stroke/SDH as underlying cause of death, of

whom 7,609 (65.6%) died in the first month after stroke.

Discussion

In linked healthcare systems data from NHS Scotland, we studied stroke subtypes at

unprecedented scale with bespoke NLP of radiology reports. We reduced the number of

people with unknown stroke types from 17,472 to 2,618. We confirmed the high rate of

early mortality for ICH compared to ischemic stroke; higher long-term risk of term

dementia after lobar ICH than for other stroke subtypes; higher short- and long-term risk of

epilepsy for cortical and lobar subtypes than for deep subtypes; and higher short-term risk

of MI for cortical ischemic stroke than for other subtypes. These findings validate our NLP

method for use in research and monitoring of care pathways.

We defined readmission with stroke only when a stroke code was in the primary position,

as a proxy for stroke recurrence. We did not include death records of stroke or SDH without

a corresponding hospital admission, as these likely represent death from the original

stroke. Readmission was increased for those with lobar ICH compared to deep ICH. Our

results are consistent with previous findings: in 674 people from two pooled British

cohorts, recurrence rates were higher in those with lobar ICH (RR 2.8 (1.5-5.5)).[1]

We found an association between cortical ischemic stroke and early MI (aHR 4.6 (3.35-

6.31). A previous systematic review found no difference in cardiac death in those with

lacunar and non-lacunar stroke, but based on relatively few events.[16] Cortical ischemic

stroke and MI may share pathological origins of large artery artherosclerosis.

People with stroke did not have long-term increased risk of cancer in this study and only a

modest early elevated risk. In the Canadian Longitudinal Health study, there was similarly

increased risk of cancer in the first year after stroke but not thereafter, compared with age-

matched controls without stroke;[17] and the Women's Health Initiative found stroke was

associated with a lower risk of subsequent cancer in post-menopausal women.[18]

Long-term dementia risk was increased after stroke, particularly lobar ICH (aHR 3.48 (2.3-

5.29)). The same pattern was found in single center cohorts of ICH in the US and

France.[19,20] Our detection rates of dementia are lower than cohort studies, presumably

because most dementia is diagnosed in outpatient settings (10.5% of a population-based

cohort of people with stroke in England developed dementia after one year, compared to

2.5% of our cohort).[21] Epilepsy risk was elevated for the first 6 months after stroke and

remained elevated beyond 6 months, particularly for cortical and lobar locations,

consistent with previous data .[22–24]

Strengths of our study were large size, representativeness of clinical practice, and

inclusion of a whole-country health-system including academic and non-academic

centers. We used an NLP algorithm with good positive predictive value for ischemic and

hemorrhagic stroke, and stroke location, allowing us to reclassify most people with non-

specific stroke codes, and to create large cohorts of stroke subtypes by location.[13] Our

innovative NLP approach can, with appropriate algorithms, be applied to other

conditions.[25]

Limitations were that we could not identify subtypes for around half of cases of ICH and

ischemic stroke, although the aHR for those with no location were similar to those with

identified locations (Supplement 1, eFigure 6). With direct access to images, an algorithm

could identify stroke subtypes without radiology reports, potentially reducing the number

with unknown location subtypes.[26] Our NLP method was rules-based, whereas most

modern systems employ large language models. We did not use these systems as the

governance requirements in secure data environments are onerous, and they may have

similar performance to smaller, expert developed models.[27] NLP identified atrophy and

WMH in scan reports, but these are not adjusted for as they are more likely to be recorded

in a person with stroke (who therefore had at least one head scan), compared to the

control population who may not have had a scan.

We have previously validated EdiE-R's categorization of stroke subtypes, but did not

validate our process of comparing ICD-10 codes and NLP tags for this study. However,

baseline characteristics aligned with our understanding of underlying pathology of

subtypes.[28,29] Those with cortical ischemic stroke had highest rates of atrial fibrillation

(18.9% vs 12.1% for deep ischemic stroke), and those with deep ischemic stroke had

higher rates of atrophy and WMH than those with cortical. An alternative study design

would use time-varying covariates; however, we chose a matched cohort study as for a

cohort of our size with variables updated daily, time-varying covariates are

computationally unfeasible. We chose cause-specific models rather than competing risk

models as they allow estimation of effects of covariates on the rate of occurrence of the

outcome in those subjects who are currently event free, and are suited to addressing

etiological questions, although we accounted for competing risk when estimating

cumulative incidence.[30]

We used real-world, nationwide, healthcare-data enhanced with NLP of imaging reports to

produce estimates for risks of outcomes directly relevant to people with stroke subtypes.

Our approach is one example of how NLP and other artificial intelligence technologies

allow analysis of large volumes of data without requiring expert input for each individual

case, with enormous possibilities for audit and epidemiology. Artificial intelligence tools

already assist in imaging analysis and clinical decision making in stroke medicine.[31]

Beyond routine clinical care, automated imaging analysis could identify phenotypes in

large-scale national cohort studies, such as UK Biobank and Our Future Health,[32,33] and

identify participants for trials of interventions for stroke and other conditions.

Acknowledgements

For the purpose of open access, the author(s) has applied a Creative Commons attribution (CC BY) licence to any Author Accepted Manuscript version arising.

Author Contributions: Alice Hosking had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

WNW and AH conceived the study. AH conducted the statistical analysis. BA, CG, and RT implemented the natural language processing algorithm and extracted phenotype data. WNW, MI, and LS provided methodological and statistical advice. AH and WNW drafted the manuscript. All authors reviewed the manuscript and provided feedback.

Sources of funding and support: AH is funded by a Medical Research Council (UK)/The Stroke Association fellowship (MR/Z504051/1). This project was funded by the Chief Scientist's Office (CSO-SCAF/17/01), the Medical Research Council (G0902303/1), the Alzheimer's Society (486). WNW is supported by CSO and Health Data Research UK. MHI is supported by the Wellcome Trust (220857/Z/20/Z; 226770/Z/22/Z, 104036/Z/14/Z; 216767/Z/19/Z) and by a Research Data Scotland Accelerator Award (RAS-24-2). MM is funded by Health Data Research UK. BA is supported by the Turing Fellowship and Turing project (EP/N510129/1) from The Alan Turing Institute, by Legal and General PLC as part of the Advanced Care Research Centre, and by the National Institute for Health Research (NIHR202639). JMW is part funded by the UK DRI which is funded by UK MRC, Als Soc and ARUK; JMW is also part funded by NIHR.

348 **Conflict of Interest Disclosures** GM has received consultancy fees from Canon Medical

349 Research, Europe, Ltd. The other authors declare no other conflicts of interest

References

1. Li L, Poon MTC, Samarasekera NE, et al. Risks of recurrent stroke and all serious vascular events after spontaneous intracerebral haemorrhage: pooled analyses of two population-based studies. *Lancet Neurol*. 2021;20(6):437-447. doi:10.1016/S1474-4422(21)00075-2

2. Verburgt E, Fellah L, Ekker MS, et al. Risk of Poststroke Epilepsy Among Young Adults With Ischemic Stroke or Intracerebral Hemorrhage. *JAMA Neurol*. 2025;82(6):597-604. doi:10.1001/JAMANEUROL.2025.0465

3. Wang Y, Jing J, Meng X, et al. The Third China National Stroke Registry (CNSR-III) for patients with acute ischaemic stroke or transient ischaemic attack: design, rationale and baseline patient characteristics. *Stroke Vasc Neurol*. 2019;4(3):158-164. doi:10.1136/SVN-2019-000242

4. Skajaa N, Adelborg K, Horváth-Puhó E, et al. Nationwide Trends in Incidence and Mortality of Stroke Among Younger and Older Adults in Denmark. *Neurology*. 2021;96(13):E1711-E1723. doi:10.1212/WNL.0000000000011636/ASSET/C2ACC646-606D-4FEC-95B2-62DA91FD81BD/ASSETS/GRAPHIC/6TTU1.JPEG

5.   Skajaa N, Adelborg K, Horváth-Puhó E, et al. Risks of Stroke Recurrence and Mortality After First and Recurrent Strokes in Denmark: A Nationwide Registry Study. *Neurology*. 2022;98(4):E329-E342. doi:10.1212/WNL.0000000000013118

6.   Bray BD, Paley L, Hoffman A, et al. Socioeconomic disparities in first stroke incidence, quality of care, and survival: a nationwide registry-based cohort study of 44 million adults in England. *Lancet Public Health*. 2018;3(4):e185-e193. doi:10.1016/S2468-2667(18)30030-6

7.   Jarman JW, Hunter TD, Hussain W, March JL, Wong T, Markides V. Stroke rates before and after ablation of atrial fibrillation and in propensity-matched controls in the UK. *Pragmat Obs Res*. 2017;8:107-118. doi:10.2147/POR.S134781

8.   Woodfield R, Grant I, Group UBSO, Group UBFU and OW, Sudlow CLM. Accuracy of Electronic Health Record Data for Identifying Stroke Cases in Large-Scale Epidemiological Studies: A Systematic Review from the UK Biobank Stroke Outcomes Group. *PLoS One*. 2015;10(10):e0140533. doi:10.1371/JOURNAL.PONE.0140533

9.   Roark C, Wilson MP, Kubes S, Mayer D, Wiley LK. Assessing the utility and accuracy of ICD10-CM non-traumatic subarachnoid hemorrhage codes for intracranial aneurysm research. *Learn Health Syst*. 2021;5(4):e10257. doi:10.1002/LRH2.10257

10.   Rannikmäe K, Ngoh K, Bush K, et al. Accuracy of identifying incident stroke cases from linked health care data in UK Biobank. *Neurology*. 2020;95(6):E697-E707. doi:10.1212/WNL.0000000000009924

11.   Wheater E, Mair G, Sudlow C, Alex B, Grover C, Whiteley W. A validated natural language processing algorithm for brain imaging phenotypes from radiology reports in UK electronic health records. *BMC Med Inform Decis Mak*. 2019;19(1):1-11. doi:10.1186/S12911-019-0908-7/TABLES/5

12.   Alex B, Grover C, Tobin R, Sudlow C, Mair G, Whiteley W. Text mining brain imaging reports. *J Biomed Semantics*. 2019;10(1):1-11. doi:10.1186/S13326-019-0211-7/TABLES/8

13.   Casey A, Davidson E, Grover C, et al. Understanding the performance and reliability of NLP tools: a comparison of four NLP tools predicting stroke phenotypes in radiology reports. *Frontiers in Digital Health , 5 , Article 1184919 (2023)* . 2023;5. doi:10.3389/FDGTH.2023.1184919

14.   Baxter R, Nind T, Sutherland J, et al. The Scottish Medical Imaging Archive: 57.3 Million Radiology Studies Linked to Their Medical Records. *Radiol Artif Intell*. 2023;6(1):e220266. doi:10.1148/RYAI.220266

15.   Migration flows - National Records of Scotland (NRS). Accessed December 19, 2024. https://www.nrscotland.gov.uk/publications/migration-flows/

16.   Jackson C, Sudlow C. Comparing risks of death and recurrent vascular events between lacunar and non-lacunar infarction. *Brain*. 2005;128(Pt 11):2507. doi:10.1093/BRAIN/AWH636

17.   Rioux B, Gioia LC, Keezer MR. Risk of Cancer Following an Ischemic Stroke in the Canadian Longitudinal Study on Aging. *Can J Neurol Sci*. 2022;49(2):225-230. doi:10.1017/CJN.2021.55

18.   Sealy-Jefferson S, Cote ML, Chlebowski RT, Rexrode KM, Simon MS. Post-Stroke Cancer Risk among Postmenopausal Women: The Women's Health Initiative. *Womens Health Issues*. 2018;28(1):29-34. doi:10.1016/J.WHI.2017.10.012

19.   Biffi A, Bailey D, Anderson CD, et al. Risk Factors Associated With Early vs Delayed Dementia After Intracerebral Hemorrhage. *JAMA Neurol*. 2016;73(8):969-976. doi:10.1001/JAMANEUROL.2016.0955

20.   Moulin S, Labreuche J, Bombois S, et al. Dementia risk after spontaneous intracerebral haemorrhage: a prospective cohort study. *Lancet Neurol*. 2016;15(8):820-829. doi:10.1016/S1474-4422(16)00130-7

21.   Pendlebury ST, Rothwell PM. Incidence and prevalence of dementia associated with transient ischaemic attack and stroke: analysis of the population-based Oxford Vascular Study. *Lancet Neurol*. 2019;18(3):248. doi:10.1016/S1474-4422(18)30442-3

22.   Biffi A, Rattani A, Anderson CD, et al. Delayed seizures after intracerebral haemorrhage. *Brain*. 2016;139(Pt 10):2694-2705. doi:10.1093/BRAIN/AWW199

23.   Galovic M, Döhler N, Erdélyi-Canavese B, et al. Prediction of late seizures after ischaemic stroke with a novel prognostic model (the SeLECT score): a multivariable

prediction model development and validation study. *Lancet Neurol*. 2018;17(2):143. doi:10.1016/S1474-4422(17)30404-0

24.   Graham NSN, Crichton S, Koutroumanidis M, Wolfe CDA, Rudd AG. Incidence and associations of poststroke epilepsy the prospective South London stroke register. *Stroke*. 2013;44(3):605-611. doi:10.1161/STROKEAHA.111.000220/SUPPL_FILE/STR202675_SUPPLEMENT.PDF

25.   Iveson MH, Mukerjee M, Davidson EM, et al. Clinically reported covert cerebrovascular disease and risk of neurological disease: a whole-population cohort of 395,273 people using natural language processing. *medRxiv*. Published online June 23, 2025:2025.06.12.25329472. doi:10.1101/2025.06.12.25329472

26.   Camilleri MPJ, Gouzou D, Al-Wasity S, et al. A large dataset of brain imaging linked to health systems data: the curation and access to a whole system national cohort from NHS Scotland. *medRxiv*. Published online October 24, 2025:2025.10.21.25338469. doi:10.1101/2025.10.21.25338469

27.   Peng Y, Wang X, Lu L, Bagheri M, Summers R, Lu Z. NegBio: a high-performance tool for negation and uncertainty detection in radiology reports. *AMIA Jt Summits Transl Sci Proc*. 2018;2017(2):188-196. Accessed January 20, 2026. http://www.ncbi.nlm.nih.gov/pubmed/29888070

28.   Dupré N, Drieu A, Joutel A. Pathophysiology of cerebral small vessel disease: a journey through recent discoveries. *J Clin Invest*. 2024;134(10). doi:10.1172/JCI172841

29.   Kamel H, Healey JS. Cardioembolic Stroke. *Circ Res*. 2017;120(3):514. doi:10.1161/CIRCRESAHA.116.308407

30.   Austin PC, Lee DS, Fine JP. Introduction to the Analysis of Survival Data in the Presence of Competing Risks. *Circulation*. 2016;133(6):601-609. doi:10.1161/CIRCULATIONAHA.115.017719/-/DC1

31.   Al-Janabi OM, El Refaei A, Elgazzar T, et al. Current Stroke Solutions Using Artificial Intelligence: A Review of the Literature. *Brain Sci*. 2024;14(12):1182. doi:10.3390/BRAINSCI14121182/S1

32.   Cook MB, Adams N, Adjetey A, et al. Cohort Profile: Our Future Health. *Int J Epidemiol*. 2025;54(6):dyaf171. doi:10.1093/IJE/DYAF171

33.   Sudlow C, Gallacher J, Allen N, et al. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med*. 2015;12(3). doi:10.1371/JOURNAL.PMED.1001779

Tables and figures

Table 1: Baseline data of people with ischemic stroke (IS), intracerebral hemorrhage (ICH), and subtypes by location. WMH= white matter hypointesities (CT)/hyperintensities(MRI).

	Ischemic stroke (51,339)				Intracerebral hemorrhage (7182)			
	Cortical (14,103)	Deep (12,616)	Cortical and deep (768)	No location (23,852)	Lobar (1456)	Deep (1814)	Lobar and deep (145)	No location (3767)
Sex (male)	7038 (49.9%)	6587 (52.2%)	361 (47.0%)	11695 (49.0%)	656 (45.1%)	912 (50.3%)	61 (42.1%)	1785 (47.4%)
Age (yrs, mean (SD))	75.8 (12.6)	74.5 (12.9)	74.2 (13.0)	72.7 (13.9)	75.2 (13.0)	73.3 (14.3)	75.3 (11.6)	73.1 (14.0)
Atrophy (n,%)	6854 (48.6%)	6661 (52.8%)	346 (45.1%)	10,805 (45.3%)	644(44.2%)	804(44.3%)	61 (42.1%)	1386 (36.8%)
WMH (n,%)	8363 (59.3%)	8642 (68.5%)	467 (60.8%)	13,095 (54.9%)	823 (56.5%)	1050 (57.9%)	72 (49.7%)	1793 (47.6%)
Diabetes (n,%)	1942 (13.8%)	1680 (13.3%)	111 (14.5%)	2903 (12.2%)	150 (10.3%)	168 (9.3%)	12 (8.3%)	361 (9.6%)
Hypertension (n,%)	10101 (71.6%)	8402 (66.6%)	504 (65.6%)	15,161 (63.6%)	902 (62.0%)	1154 (63.6%)	84 (57.9%)	2298 (61.0%)
Atrial fibrillation (n,%)	2671 (18.9%)	1525 (12.1%)	140 (18.2%)	3108 (13.0%)	195 (13.4%)	227 (12.5%)	14 (9.7%)	439 (11.7%)
On statins (n,%)	7239 (51.3%)	5812 (46.1%)	343 (44.7%)	10,502 (44.0%)	666 (45.7%)	773 (42.6%)	79 (48.3%)	1621 (43.0%)
On anticoagulants (n,%)	1710 (12.1%)	1100 (8.7%)	88 (11.5%)	2263 (9.5%)	217 (14.9%)	310 (17.1%)	21 (14.5%)	568 (15.1%)
On antiplatelets (n,%)	6729 (47.7%)	5383 (42.7%)	309 (40.2%)	9196 (38.6%)	591 (40.6%)	615 (33.9%)	66 (45.5%)	1393 (37.0%)
Hospital admissions prior year (mean, SD)	1.51 (2.90)	1.26 (2.46)	1.32 (2.46)	1.31 (2.90)	1.72 (4.70)	1.37 (3.70)	1.65 (3.29)	1.32 (2.87)
Prescriptions prior year (mean, SD)	3.86 (3.10)	3.46 (3.10)	3.40 (2.85)	3.33 (3.10)	3.35 (2.93)	3.20 (2.98)	3.12 (2.65)	3.29 (3.14)

Figure 1: Fully adjusted hazard ratios for readmission with stroke subtypes. Deep ischemic stroke is compared with cortical ischemic stroke; deep intracerebral hemorrhage is compared with lobar intracerebral hemorrhage.

Outcome	Stroke type		Hazard ratio	N with outcome
Readmission within 6 months	Ischaemic stroke, Deep		Reference	358
	Ischaemic stroke, Cortical		1.14 (0.99–1.31)	441
	Intracerebral haemorrhage, Deep		Reference	45
	Intracerebral haemorrhage, Lobar		1.71 (1.15–2.54)	55
Readmission beyond 6 months	Ischaemic stroke, Deep		Reference	826
	Ischaemic stroke, Cortical		1.04 (0.94–1.15)	811
	Intracerebral haemorrhage, Deep		Reference	100
	Intracerebral haemorrhage, Lobar		1.28 (0.96–1.7)	95

0.50 0.71 1.0 1.41 2.0 2.83

Figure 2: Fully adjusted hazard ratios for death, myocardial infarction, dementia, epilepsy and cancer early after stroke (<6 months), compared to age- and sex-matched controls without stroke. Abbreviations: MI=myocardial infarction, IS=ischemic stroke, ICH=intracerebral hemorrhage, SAH=subarachnoid hemorrhage, SDH=subdural hemorrhage.

Outcome	Stroke type	Hazard ratio	N with outcome
Death	Ischaemic stroke (all)	8.63 (8.22–9.05)	11138
	Ischaemic stroke, Cortical	10.89 (10.02–11.83)	4134
	Ischaemic stroke, Deep	6.91 (6.24–7.66)	2202
	Intracerebral haemorrhage (all)	23.06 (20.79–25.58)	3290
	Intracerebral haemorrhage, Lobar	21.3 (16.8–27.01)	626
	Intracerebral haemorrhage, Deep	19.18 (15.41–23.88)	706
	Subarachnoid haemorrhage	39.43 (31.37–49.57)	876
	Subdural haemorrhage	10.43 (8.71–12.49)	897
MI	Ischaemic stroke (all)	3.35 (2.79–4.03)	568
	Ischaemic stroke, Cortical	4.6 (3.35–6.31)	189
	Ischaemic stroke, Deep	2.45 (1.66–3.61)	135
	Intracerebral haemorrhage (all)	1.91 (0.92–3.96)	35
	Intracerebral haemorrhage, Lobar	0.85 (0.11–6.57)	<10
	Intracerebral haemorrhage, Deep	0.73 (0.11–4.83)	<10
	Subarachnoid haemorrhage	1.16 (0.15–8.73)	<10
	Subdural haemorrhage	3.97 (1.73–9.1)	28
Dementia	Ischaemic stroke (all)	2.97 (2.55–3.46)	750
	Ischaemic stroke, Cortical	3.57 (2.7–4.71)	234
	Ischaemic stroke, Deep	3.16 (2.38–4.19)	208
	Intracerebral haemorrhage (all)	2.95 (1.87–4.66)	99
	Intracerebral haemorrhage, Lobar	2.86 (1.15–7.07)	29
	Intracerebral haemorrhage, Deep	2.24 (0.82–6.13)	18
	Subarachnoid haemorrhage	4.51 (1.33–15.32)	13
	Subdural haemorrhage	2.94 (1.56–5.57)	46
Epilepsy	Ischaemic stroke (all)	3.37 (2.97–3.82)	1385
	Ischaemic stroke, Cortical	3.99 (3.18–5)	436
	Ischaemic stroke, Deep	2.94 (2.26–3.84)	286
	Intracerebral haemorrhage (all)	6.94 (5.23–9.22)	310
	Intracerebral haemorrhage, Lobar	10.98 (6.32–19.08)	87
	Intracerebral haemorrhage, Deep	3.43 (1.49–7.86)	41
	Subarachnoid haemorrhage	8.1 (5.47–12)	174
	Subdural haemorrhage	9.76 (6.74–14.11)	212
Cancer	Ischaemic stroke (all)	1.58 (1.35–1.85)	716
	Ischaemic stroke, Cortical	1.54 (1.14–2.08)	191
	Ischaemic stroke, Deep	1.41 (1.03–1.93)	184
	Intracerebral haemorrhage (all)	1.68 (1.03–2.75)	70
	Intracerebral haemorrhage, Lobar	2.32 (0.85–6.37)	16
	Intracerebral haemorrhage, Deep	2.48 (1.01–6.05)	23
	Subarachnoid haemorrhage	1.04 (0.37–2.95)	18
	Subdural haemorrhage	0.83 (0.35–1.97)	27

0.50 2.0 8.0 32.0

Figure 3: Fully adjusted hazard ratios for death, myocardial infarction, dementia, epilepsy and cancer, late after stroke (beyond 6 months), compared with age- and sex-matched controls without stroke. Abbreviations: MI=myocardial infarction, IS=ischemic stroke, ICH=intracerebral hemorrhage, SAH=subarachnoid hemorrhage, SDH=subdural hemorrhage.

Outcome	Stroke type		Hazard ratio	N with outcome
Death	Ischaemic stroke (all)		1.88 (1.82–1.94)	16814
	Ischaemic stroke, Cortical		2.04 (1.92–2.17)	4879
	Ischaemic stroke, Deep		1.86 (1.75–1.97)	4616
	Intracerebral haemorrhage (all)		2.36 (2.13–2.61)	1724
	Intracerebral haemorrhage, Lobar		2.58 (2.07–3.22)	407
	Intracerebral haemorrhage, Deep		2.32 (1.9–2.83)	467
	Subarachnoid haemorrhage		1.77 (1.4–2.24)	305
	Subdural haemorrhage		1.56 (1.35–1.8)	870
MI	Ischaemic stroke (all)		1.34 (1.22–1.48)	1777
	Ischaemic stroke, Cortical		1.16 (0.96–1.41)	476
	Ischaemic stroke, Deep		1.65 (1.39–1.96)	510
	Intracerebral haemorrhage (all)		1.49 (1.07–2.07)	138
	Intracerebral haemorrhage, Lobar		1.51 (0.72–3.17)	29
	Intracerebral haemorrhage, Deep		1.77 (0.97–3.24)	42
	Subarachnoid haemorrhage		1.35 (0.77–2.35)	51
	Subdural haemorrhage		1.15 (0.74–1.79)	78
Dementia	Ischaemic stroke (all)		1.97 (1.84–2.11)	3578
	Ischaemic stroke, Cortical		2.16 (1.9–2.46)	1057
	Ischaemic stroke, Deep		1.94 (1.71–2.21)	971
	Intracerebral haemorrhage (all)		2.54 (2.05–3.15)	407
	Intracerebral haemorrhage, Lobar		3.49 (2.3–5.29)	124
	Intracerebral haemorrhage, Deep		2.27 (1.47–3.49)	90
	Subarachnoid haemorrhage		2.16 (1.23–3.81)	56
	Subdural haemorrhage		1.81 (1.34–2.44)	190
Epilepsy	Ischaemic stroke (all)		3.63 (3.27–4.02)	1946
	Ischaemic stroke, Cortical		5.62 (4.68–6.75)	709
	Ischaemic stroke, Deep		2.35 (1.87–2.95)	354
	Intracerebral haemorrhage (all)		8.24 (6.2–10.96)	279
	Intracerebral haemorrhage, Lobar		9.33 (5.29–16.48)	77
	Intracerebral haemorrhage, Deep		4.7 (2.33–9.48)	43
	Subarachnoid haemorrhage		5 (3.32–7.52)	136
	Subdural haemorrhage		4.33 (2.54–7.36)	73
Cancer	Ischaemic stroke (all)		1.07 (1–1.14)	3948
	Ischaemic stroke, Cortical		1.03 (0.91–1.18)	887
	Ischaemic stroke, Deep		1.11 (0.98–1.24)	1068
	Intracerebral haemorrhage (all)		0.88 (0.7–1.1)	318
	Intracerebral haemorrhage, Lobar		0.67 (0.39–1.17)	61
	Intracerebral haemorrhage, Deep		1.26 (0.83–1.89)	88
	Subarachnoid haemorrhage		1.15 (0.85–1.57)	154
	Subdural haemorrhage		1.01 (0.75–1.37)	173

0.50   2.0   8.0

Figure 4: Cumulative incidence rates dementia accounting for competing risk of death (B) for ischemic stroke and intracerebral hemorrhage by location, stratified by age and sex.

Figure 5: Cumulative incidence rates of death for ischemic stroke and intracerebral hemorrhage by location, stratified by age and sex.