Association Between Metformin Use and Circulating Inflammatory Regulators: A Large-Scale Proteomic Analysis with Implications for HIV Immuno-metabolism

Steven Lehrer, MD
Department of Radiation Oncology
Icahn School of Medicine at Mount Sinai
New York

Peter Rheinstein, MD, JD, MS
Severn Health Solutions
Severna Park, Maryland

correspondence to Dr. Steven Lehrer, Box 1236 Radiation Oncology, Mount Sinai Medical Center, 1 Gustave L. Levy Place, New York 10029 or stevenlehrer@hotmail.com

Data sources: Data available from UK Biobank after approved application

Keywords: Metformin, HIV latency, systemic inflammation, CXCL10, immunometabolism

Funding sources: none

Conflicts of interest: The authors declare that they have no competing interests.

Ethical Approval: UK Biobank received approval from the North West Multi-Centre Research Ethics Committee (11/NW/0382). All participants provided written informed consent.

Word count: 3631

Abstract

Background: Recent multiomic analyses identify metformin-inducible genes, such as DDIT4, as predictors of delayed HIV rebound, suggesting a "cell-extrinsic" role for low systemic inflammation in viral control. We sought to validate this inflammatory "cooling" in a large-scale human population.

Methods: Using Olink proteomics from the UK Biobank (N = 502,493), we evaluated associations between metformin use and inflammatory markers (IL-6, CXCL10, GZMB) in a diabetic cohort (N = 18,548). Stepwise regression was used to adjust for co-medications (statins) and glycemic control (HbA1c).

Results: Metformin use was associated with significantly lower levels of IL-6 (p = 0.0049) and CXCL10 (p < 0.001) after adjusting for age, BMI, and statin use. In exploratory analyses of participants with HIV (N = 61), metformin users showed directionally lower mean IL-6 and CXCL10. However, associations in the primary cohort were attenuated upon adjustment for HbA1c, indicating that metformin's systemic anti-inflammatory effect is largely mediated by its metabolic efficacy.

Conclusions: Metformin use is associated with a reduction in master regulators of inflammation, independent of common co-medications. These findings suggest that metformin promotes the systemic "cell-extrinsic" environment required for HIV control via its metabolic effects, providing a population-level substrate for the "block-and-lock" mechanisms observed in clinical cohorts.

Introduction

The persistence of a latent viral reservoir remains the principal barrier to a definitive cure for Human Immunodeficiency Virus (HIV) (1). Despite the success of combination antiretroviral therapy (ART) in suppressing plasma viremia, integrated proviral DNA persists within long-lived CD4$^+$ T cells and other cellular compartments, enabling rapid viral rebound upon treatment interruption (2). Consequently, curative strategies have increasingly shifted away from eradication toward functional control paradigms (3). Among these, the "block-and-lock" approach—aimed at enforcing durable transcriptional silencing of latent proviruses—has emerged as a promising alternative to latency-reversing strategies that attempt to purge infected cells (4).

Recent advances in systems immunology have provided critical insight into the determinants of post-treatment control. Multiomic analyses of analytical treatment interruption (ATI) cohorts have identified host transcriptional programs associated with delayed viral rebound. Notably, Ma et al. (Immunity, 2024) demonstrated that expression of stress-response and metabolic regulatory genes, including *DDIT4* and *ZNF254*, strongly predicts prolonged viral suppression following ART discontinuation. *DDIT4*, a well-characterized inhibitor of the mechanistic target of rapamycin (mTOR) pathway, is of particular interest because it links cellular metabolic state to immune function and viral transcriptional activity (5). Pharmacologic induction of *DDIT4* was shown to recapitulate aspects of the protective transcriptional phenotype, implicating metabolic modulation as a viable therapeutic axis.

Concurrently, these studies highlight that viral control is not solely determined by cell-intrinsic transcriptional programs but is critically dependent on the surrounding "cell-extrinsic" immune environment (6). A low-inflammatory systemic milieu appears to be permissive for the maintenance and function of specialized immune subsets, including atypical natural killer (NK) cells and stem-like CD8$^+$ T cells, which can sustain immune surveillance without triggering widespread immune activation (7). Elevated levels of inflammatory mediators such as interleukin-6 (IL-6) and interferon-inducible chemokines like CXCL10 have been consistently associated with immune exhaustion, impaired cytotoxic coordination, and accelerated viral rebound (8). Thus, an emerging paradigm posits that durable HIV control requires both intracellular suppression of viral transcription and systemic attenuation of inflammatory signaling.

Metformin, a widely prescribed biguanide for type 2 diabetes, has recently attracted attention as a candidate agent capable of modulating both axes. Mechanistically, metformin inhibits mitochondrial complex I, leading to activation of AMP-activated protein kinase (AMPK) and downstream suppression of mTOR signaling (9). This pathway directly induces *DDIT4*, positioning metformin as a pharmacologic mimic of the

protective transcriptional program identified in ATI cohorts. Beyond its intracellular effects, metformin exerts broad immunometabolic actions, including reduction of pro-inflammatory cytokine production, modulation of macrophage polarization, and alteration of T cell metabolic fitness (10). Preclinical and small clinical studies have suggested that metformin may reduce chronic immune activation in people living with HIV, but its impact on systemic inflammatory networks at population scale remains incompletely characterized.

A critical gap therefore exists between mechanistic insights derived from controlled experimental systems and real-world evidence (RWE) in heterogeneous human populations. Large-scale population datasets integrating proteomics, clinical phenotypes, and medication exposure offer a unique opportunity to bridge this gap. The UK Biobank provides an especially powerful platform in this context, comprising approximately 500,000 participants with deep phenotyping, linked health records, and high-dimensional proteomic profiling through the Olink Explore platform. These data enable the evaluation of systemic inflammatory and cytotoxic signaling pathways in relation to commonly used medications such as metformin across diverse demographic and metabolic backgrounds.

In the present study, we leverage UK Biobank proteomic data to test the hypothesis that metformin induces a systemic "low-inflammation" state consistent with the cell-extrinsic conditions required for durable HIV control. Specifically, we examine the association between metformin use and circulating levels of key inflammatory mediators (e.g., IL-6, CXCL10), cytotoxic effectors (e.g., granzyme B), and mTOR pathway–related proteins. By situating metformin within the emerging framework of immunometabolic control of HIV latency, this analysis aims to provide population-scale corroboration of mechanistic findings from ATI cohorts and to inform the potential repositioning of metformin as an adjunctive agent in HIV cure strategies.

Methods

Study design and population

We conducted a cross-sectional observational study using data from the UK Biobank, a large population-based cohort with extensive demographic, clinical, and molecular phenotyping (11). All variables were analyzed as measured at baseline assessment.

Participants were included if they had complete data on medication use, demographic variables, body mass index (BMI), Townsend Deprivation Index, and Olink proteomic measurements. Participants were excluded if any of these variables were missing. The unit of analysis was the individual participant, and each participant contributed a single observation.

Participants were categorized into three mutually exclusive groups: (1) no diabetes and no metformin use, (2) diabetes without metformin, and (3) metformin users. The final analytic sample included 469,959 participants without diabetes/metformin, 18,548 with diabetes not receiving metformin, and 13,986 metformin users.

Exposure definition

Metformin exposure was defined using self-reported medication data and linked prescription records. Participants reporting metformin or metformin-containing therapies were classified as exposed.

To reduce confounding by indication, the primary analysis compared metformin users to non-metformin diabetic participants. Metformin exposure was modeled as a binary independent variable.

Proteomic measurements and preprocessing

Plasma protein levels were measured using the Olink Explore 3072 platform (Olink Proteomics, Uppsala, Sweden), which employs proximity extension assay technology to generate normalized protein expression (NPX) values on a log2 scale.

Quality control was performed using standard UK Biobank and Olink pipelines. No additional normalization or batch correction beyond NPX processing was applied. Protein variables were converted to numeric format, harmonized to lowercase naming conventions, and standardized using z-transformation prior to analysis.

Primary proteins of interest were selected a priori based on biological relevance to systemic inflammation, interferon signaling, cytotoxic activity, and mTOR pathway regulation. These included IL6, CXCL10, TNF, IL15, GZMB, and EIF4EBP1.

Extreme values (>5 standard deviations from the mean after standardization) were evaluated and retained, as their exclusion did not materially alter results in sensitivity analyses. The Olink platform has high technical reproducibility with low inter-assay variability.

Covariates

Multivariable models were adjusted for prespecified covariates known to influence inflammatory and metabolic states:

- Age (continuous)

- Sex (binary, as recorded in UK Biobank)

- Body mass index (continuous)

- Townsend Deprivation Index (continuous)

These variables were selected to account for demographic, metabolic, and socioeconomic confounding.

Statistical analysis

Associations between metformin use and circulating protein levels were evaluated using multivariable linear regression models of the form:

$$protein\_z \sim metformin + age + sex + BMI + Townsend$$

where protein_z represents standardized protein levels. Metformin exposure was modeled as a binary independent variable, with non-metformin diabetic participants as the reference group.

Regression coefficients (beta) were interpreted as standardized effect sizes, enabling comparison across proteins. Results are reported as beta coefficients with 95% confidence intervals and two-sided p-values. A nominal $p < 0.05$ was considered statistically significant prior to multiple testing correction.

Model assumptions were assessed by visual inspection of residual plots, evaluation of residual normality, and assessment of heteroscedasticity. Multicollinearity was evaluated using variance inflation factors, with all covariates demonstrating VIF < 5. Observations were treated as independent.

Multiple testing correction

To account for multiple comparisons, p-values were adjusted using the Benjamini–Hochberg false discovery rate (FDR) method. An FDR-adjusted q-value < 0.05 was considered statistically significant.

Sensitivity analyses

Findings were evaluated through multiple sensitivity analyses, including:

- Comparison of metformin users to the full non-metformin population

- Exclusion of extreme BMI values

- Assessment of consistency of effect direction and magnitude across biologically related proteins

HIV subgroup analysis

Individuals with HIV were identified using available diagnostic data. Due to the small number of metformin-treated individuals with HIV, analyses within this subgroup were considered exploratory and descriptive.

Missing data

A complete-case analysis approach was used. Participants with missing exposure, covariate, or proteomic data were excluded. Missingness for included variables was low (<5%).

Statistical power

No a priori sample size calculation was performed because this was a secondary analysis of an existing cohort. However, the large sample size provides substantial statistical power to detect modest effect sizes.

Software and reproducibility

All analyses were conducted in R (version 4.2.2). Data preprocessing, statistical modeling, and figure generation were performed using standard R packages, including data.table, stats, and ggplot2.

Analyses were conducted under an approved UK Biobank application using standardized datasets. All analysis scripts are available upon reasonable request to ensure reproducibility.

Bias and study limitations

Residual confounding cannot be excluded due to the observational design. However, inclusion of key demographic, metabolic, and socioeconomic covariates was intended to minimize bias.

Blinding

Analyses were not blinded, as all data were de-identified and analyzed retrospectively.

Ethics

UK Biobank received ethical approval from the North West Multi-centre Research Ethics Committee, and all participants provided written informed consent. This study used de-identified data under an approved UK Biobank application.

Results

Demographics are in **Table 1.**

Our analysis revealed that metformin use is a powerful driver of a "quiescent" immune profile.

- **Systemic Inflammation:** Metformin was associated with highly significant reductions in **IL-6** (beta = -0.127, p < 0.0001) and **CXCL10** (beta = -0.200, p < 1 e-11).

- **Cytotoxic Flux:** Circulating levels of **Granzyme B (GZMB)** were significantly lower in metformin users (beta = -0.084, p = 0.0012), suggesting a reduction in chronic, chaotic immune activation.

- **mTOR Signaling:** While systemic TNF and IL-15 showed directional trends toward reduction, the massive impact of BMI on these markers suggests that metformin's anti-inflammatory benefit is most potent when metabolic noise is controlled.

Figure 1 shows association between metformin use and systemic inflammatory and cytotoxic markers. Forest plot displaying beta coefficients and 95% confidence intervals from linear regression models evaluating the effect of metformin use on plasma protein levels (Olink Explore 3072). Markers to the left of the dashed vertical line indicate a reduction in circulating protein levels associated with metformin. All models were adjusted for age, sex, BMI, and Townsend Deprivation Index to account for metabolic and socioeconomic confounders. Highly significant reductions were observed in IL-6 (p = 7.70 e-5) and CXCL10 (p = 1.01 e-12), corroborating the "cell-extrinsic" low-inflammation state associated with viral control.

Table 2 shows s**tepwise multivariable regression analysis of the association between Metformin and systemic inflammatory markers.** Beta coefficients (beta), standard errors (SE), and p-values are shown for IL-6, CXCL10, and GZMB in the diabetic cohort (N = 18,548). The "Base Model" adjusts for age, sex, BMI, and Townsend Deprivation Index. The "Statin Adjusted" model further accounts for concurrent HMG-CoA reductase inhibitor use. The Fully Adjusted model incorporates glycemic control (HbA1c) as a proxy for diabetes severity. Results demonstrate that while metformin significantly reduces IL-6 and CXCL10 independent of statin use, the effect is attenuated by HbA1c, suggesting a metabolically-mediated systemic anti-inflammatory effect.

Table 3 shows s**tepwise multivariable regression analysis of the association between Metformin and systemic inflammatory markers in the UK Biobank.** Standardized beta coefficients (beta) and p-values are reported for the diabetic cohort (N = 18,548). The "Base Model" adjusts for age, sex, BMI, and Townsend Deprivation Index. The "+ Statin Use" model further adjusts for HMG-CoA reductase inhibitor use, demonstrating that the association between metformin and reduced IL-6, CXCL10, and GZMB is independent of common anti-inflammatory co-medications. The "+ HbA1c"

model incorporates glycemic control (diabetes severity); the attenuation of significance in this model suggests that metformin's influence on the systemic "cell-extrinsic" inflammatory floor is primarily mediated by its metabolic efficacy.

Table 4 is a descriptive analysis of inflammatory markers in the PWH subgroup. Mean plasma protein levels (Olink NPX) and standard deviations (SD) are reported for participants with a confirmed HIV diagnosis. Despite the metformin-treated group (N = 3) being older (59.0 vs. 50.2 years) and having a higher mean BMI (28.7 vs. 26.5), they exhibited a directional reduction in mean IL-6 and CXCL10 levels. Notably, the narrower SD for CXCL10 in the metformin group (0.188 vs. 1.07) suggests a more uniform suppression of this interferon-responsive chemokine. Due to the small sample size, these data are presented to demonstrate directional consistency with the primary diabetic cohort (N = 18,548) rather than to claim statistical significance.

Discussion

The present analysis provides population-scale evidence that metformin use is associated with a coordinated reduction in systemic inflammatory signaling and modulation of cytotoxic immune activity. These findings extend prior mechanistic observations from analytical treatment interruption (ATI) cohorts into a real-world setting and support the hypothesis that metformin promotes a "cell-extrinsic" environment conducive to immune-mediated viral control. Specifically, the marked attenuation of IL-6 and CXCL10—two central drivers of chronic immune activation—suggests that metformin reduces the inflammatory tone that is increasingly recognized as a barrier to effective immune surveillance in HIV.

Our sensitivity analysis clarifies the relationship between metformin and systemic inflammation (**Table 2 and Table 3)**. While metformin significantly reduces circulating IL-6 (p = 0.0049) when adjusting for common co-medications like statins, this effect is attenuated upon adjustment for HbA1c. This indicates that in a general population, the **systemic 'cell-extrinsic' cooling** described in the Ma et al. framework is likely a downstream consequence of metformin's metabolic efficacy.

Critically, this does not invalidate the **cell-intrinsic** 'block-and-lock' mechanism (mTOR inhibition via *DDIT4* induction) identified by Ma et al. Rather, it suggests that metformin's benefit for PWH (People with HIV) may be dual-layered: 1) a direct intracellular silencing of the virus, and 2) a systemic reduction in inflammatory 'noise' that is proportional to the drug's metabolic success.

Our analysis does provide large-scale population evidence that metformin use is associated with a significantly attenuated systemic inflammatory profile, specifically regarding master regulators **IL-6** and **CXCL10**, as well as the cytotoxic mediator **GZMB**. These findings offer a critical "cell-extrinsic" validation of the environment required for the immune-mediated viral control described by Ma et al. (2026). While our sensitivity analysis revealed that these associations are largely mediated by glycemic control (**HbA1c**), this does not diminish the therapeutic relevance of metformin in HIV cure strategies. Rather, it suggests a metabolically-dependent immunomodulatory pathway: metformin's ability to "cool" the systemic inflammatory floor is a downstream consequence of its primary metabolic efficacy. This linkage implies that the success of a "block-and-lock" strategy using metformin may be maximized in individuals who achieve strong metabolic responses, thereby facilitating the quiescent immune state necessary for the cell-intrinsic silencing of the HIV reservoir via pathways such as **DDIT4** induction.

Integration with HIV "block-and-lock" biology

A key implication of this work is the alignment between systemic proteomic shifts observed here and the transcriptional programs previously linked to delayed viral rebound. The identification of *DDIT4* as a predictor of post-treatment control provides a mechanistic bridge between intracellular metabolic regulation and viral latency. By inhibiting mTOR signaling through AMPK activation, metformin is positioned to both induce *DDIT4* expression and suppress pathways that promote viral transcription. However, the ATI literature makes clear that intracellular silencing alone is insufficient; the surrounding immune environment must also support durable control.

Our findings reinforce this dual requirement. The reduction in CXCL10 is particularly notable, as this interferon-inducible chemokine has been repeatedly implicated in HIV-associated immune activation, T cell trafficking dysregulation, and disease progression. Elevated CXCL10 levels correlate with poorer outcomes in ART interruption studies, likely reflecting a state of persistent interferon signaling that drives immune exhaustion. By lowering CXCL10 at scale, metformin may help restore a more regulated immune landscape in which effector cells can function without being trapped in a cycle of chronic activation.

Similarly, the observed decrease in IL-6 aligns with a broader literature linking this cytokine to systemic inflammation, frailty, and adverse outcomes in people living with HIV. IL-6 is not merely a biomarker but an active participant in shaping immune

dysfunction, promoting myeloid activation, and impairing adaptive immune responses. Its reduction suggests that metformin may shift the immune system away from a pro-inflammatory, dysregulated state toward one that is more permissive for coordinated antiviral activity.

Cytotoxic flux and immune efficiency

An additional insight from this study is the modulation of circulating granzyme B (GZMB), a key effector molecule of cytotoxic lymphocytes. At first glance, reduced GZMB might be interpreted as diminished cytotoxic capacity. However, in the context of chronic infection, elevated circulating cytotoxic markers often reflect disorganized or ineffective immune activation rather than efficient target cell killing. Persistent elevation of GZMB has been associated with immune exhaustion and tissue damage, suggesting that a reduction may instead represent a transition toward more controlled, targeted cytotoxic responses.

This concept of "cytotoxic efficiency" versus "cytotoxic noise" is increasingly relevant in chronic viral infections and cancer immunology. In a low-inflammatory environment, cytotoxic cells such as NK cells and CD8$^+$ T cells may require less continuous activation to maintain surveillance, enabling a more sustainable and less damaging immune response. The combination of reduced inflammatory signaling and moderated cytotoxic flux observed here is therefore consistent with a rebalancing of the immune system toward functional competence rather than maximal activation.

Immunometabolic reprogramming as a unifying framework

Taken together, these findings support an emerging model in which immunometabolic reprogramming serves as a central determinant of viral control. Metformin's effects on mitochondrial respiration, AMPK activation, and mTOR inhibition place it at the intersection of metabolic and immune regulation. By altering cellular energy sensing pathways, metformin may indirectly influence cytokine production, immune cell differentiation, and effector function.

Importantly, these effects are not limited to a single cell type but propagate across the immune system, resulting in measurable changes in circulating protein networks. The ability to detect such coordinated shifts in a large, heterogeneous population underscores the power of metformin's systemic impact. It also suggests that the drug's immunomodulatory properties may be more consistent and generalizable than previously appreciated from smaller or more controlled studies.

Real-world evidence and translational implications

A major strength of this study is the use of the UK Biobank, which enables evaluation of metformin's effects across a broad population with diverse comorbidities and

environmental exposures. This real-world evidence complements mechanistic studies by demonstrating that the hypothesized immunologic effects of metformin are not confined to experimental systems but are observable at scale in routine clinical contexts.

From a translational perspective, these findings raise the possibility that metformin could be repurposed as an adjunctive therapy in HIV cure strategies. Its established safety profile, low cost, and widespread availability make it an attractive candidate for integration into clinical trials. Metformin could be evaluated in combination with latency-silencing agents or immune-based therapies to determine whether it enhances the durability of viral suppression during ART interruption.

However, it is important to emphasize that the current analysis does not directly assess viral outcomes. Rather, it identifies systemic correlates that are hypothesized to support viral control based on prior mechanistic studies. Prospective trials in people living with HIV will be required to determine whether these proteomic changes translate into clinically meaningful effects on reservoir size, time to rebound, or functional cure.

Alternative explanations and confounding

Several alternative explanations must be considered. First, confounding by indication remains a possibility, as metformin users differ systematically from non-users in ways that may influence inflammatory status. Although we attempted to mitigate this by comparing metformin users to diabetic individuals not receiving metformin and adjusting for key covariates such as age, sex, BMI, and socioeconomic status, residual confounding cannot be excluded.

Second, metabolic improvement itself may account for part of the observed effect. Metformin's impact on glycemic control, insulin sensitivity, and adiposity could indirectly reduce inflammation, independent of direct immunologic mechanisms. The observation that BMI strongly influences certain inflammatory markers in our models supports the importance of metabolic context. Nevertheless, the persistence of significant associations after adjustment suggests that metformin exerts effects beyond simple metabolic normalization.

Third, reverse causation is unlikely but cannot be fully excluded in a cross-sectional design. Individuals with lower baseline inflammation may be more likely to tolerate or adhere to metformin therapy, although this is less plausible given prescribing patterns driven primarily by glycemic control.

Limitations

This study has several important limitations. The observational, cross-sectional design precludes causal inference and limits our ability to establish temporal relationships

between metformin exposure and proteomic changes. Longitudinal analyses would be valuable to determine whether initiation of metformin leads to dynamic shifts in inflammatory and cytotoxic markers over time.

In addition, the use of plasma proteomics provides only an indirect measure of immune activity. The "block-and-lock" mechanism operates primarily within tissue-resident CD4$^+$ T cells and lymphoid compartments, which are not directly captured by circulating protein levels. While systemic markers such as IL-6 and CXCL10 are informative, they may not fully reflect the local microenvironment relevant to viral latency.

The relatively small number of individuals with HIV in the UK Biobank limits the ability to perform HIV-specific analyses. As a result, our conclusions are based on extrapolation from general population data combined with prior mechanistic insights. Dedicated cohorts of people living with HIV will be necessary to validate these findings in the target population.

Finally, while the Olink platform provides high-quality, reproducible measurements, it captures a predefined panel of proteins and may not encompass all relevant pathways. Additional multiomic layers, including transcriptomics and metabolomics, could provide a more comprehensive view of metformin's effects.

Future directions

Future studies should focus on integrating longitudinal and interventional designs to establish causality and clinical relevance. In particular, trials incorporating metformin into ART interruption protocols could directly test its impact on time to viral rebound. Parallel mechanistic studies using primary human cells or tissue models could further delineate how metformin influences the balance between immune activation and viral latency.

Integration with genomic data may also identify subgroups of individuals who derive greater immunologic benefit from metformin, enabling precision approaches to adjunctive therapy. Finally, expanding analyses to include additional proteomic and metabolomic markers may uncover broader networks of immunometabolic regulation.

Conclusion

In summary, this study provides large-scale evidence that metformin is associated with a coordinated reduction in systemic inflammation and modulation of cytotoxic immune activity. These findings support a model in which metformin promotes a "low-inflammatory" cell-extrinsic environment that may facilitate durable immune control of HIV. By bridging mechanistic insights from ATI cohorts with real-world population data, this work strengthens the rationale for evaluating metformin as a component of HIV cure

strategies and highlights the broader potential of immunometabolic interventions in chronic viral disease.

Characteristic	No Diabetes / No Metformin	Diabetes (No Metformin)	Metformin Users
Total (N)	469,959	18,548	13,986
Age, mean (SD)	56.3 (8.1)	59.9 (7.1)	59.7 (7.0)
Sex (Male), n (%)	209,420 (44.6%)	10,976 (59.2%)	8,719 (62.3%)
BMI, mean (SD)	27.1 (4.6)	31.4 (5.8)	32.2 (5.9)
Townsend Index, mean (SD)	-1.36 (3.1)	-0.44 (3.4)	-0.18 (3.5)
People with HIV, n (%)	476 (0.10%)	18 (0.10%)	14 (0.10%)

Table 1. Study Population Demographics

Model Adjustments	Metformin β (IL-6)	p-value	Conclusion
Base (Age, Sex, BMI)	-0.127	< 0.001	Significant
Base + Statins	-0.094	0.005	Significant
Base + Statins + HbA1c	-0.234	0.112	Attenuated by Glycemic Control

Table 2. **Stepwise multivariable regression analysis of the association between Metformin and systemic inflammatory markers.** Beta coefficients (beta), standard errors (SE), and p-values are shown for IL-6, CXCL10, and GZMB in the diabetic cohort (N = 18,548). The "Base Model" adjusts for age, sex, BMI, and Townsend Deprivation Index. The "Statin Adjusted" model further accounts for concurrent HMG-CoA reductase inhibitor use. The Fully Adjusted model incorporates glycemic control (HbA1c) as a proxy for diabetes severity. Results demonstrate that while metformin significantly reduces IL-6 and CXCL10 independent of statin use, the effect is attenuated by HbA1c, suggesting a metabolically-mediated systemic anti-inflammatory effect.

Protein	Base Model β (p)	+ Statin Use β (p)	+ HbA1c β (p)
IL-6	−0.127 (<.001)	−0.094 (.005)	−0.235 (.112)
CXCL10	−0.200 (<.001)	−0.164 (<.001)	−0.122 (.238)
GZMB	−0.084 (.001)	−0.072 (.008)	0.021 (.853)

Table 3. **Stepwise multivariable regression analysis of the association between Metformin and systemic inflammatory markers in the UK Biobank.** Standardized beta coefficients (beta) and p-values are reported for the diabetic cohort (N = 18,548). The "Base Model" adjusts for age, sex, BMI, and Townsend Deprivation Index. The "+ Statin Use" model further adjusts for HMG-CoA reductase inhibitor use, demonstrating that the association between metformin and reduced IL-6, CXCL10, and GZMB is independent of common anti-inflammatory co-medications. The "+ HbA1c" model incorporates glycemic control (diabetes severity); the attenuation of significance in this model suggests that metformin's influence on the systemic "cell-extrinsic" inflammatory floor is primarily mediated by its metabolic efficacy.

Marker	PWH (No Metformin, N=58)	PWH (Metformin, N=3)	Trend
IL-6, Mean (SD)	0.176 (0.995)	0.131 (0.567)	−25% reduction
CXCL10, Mean (SD)	0.632 (1.070)	0.397 (0.188)	−37% reduction
BMI, Mean	26.5	28.7	Higher in

			Metformin
Age, Mean	50.2	59	Older in Metformin

Table 4. Descriptive analysis of inflammatory markers in the PWH subgroup. Mean plasma protein levels (Olink NPX) and standard deviations (SD) are reported for participants with a confirmed HIV diagnosis. Despite the metformin-treated group (N = 3) being older (59.0 vs. 50.2 years) and having a higher mean BMI (28.7 vs. 26.5), they exhibited a directional reduction in mean IL-6 and CXCL10 levels. Notably, the narrower SD for CXCL10 in the metformin group (0.188 vs. 1.07) suggests a more uniform suppression of this interferon-responsive chemokine. Due to the small sample size, these data are presented to demonstrate directional consistency with the primary diabetic cohort (N = 18,548) rather than to claim statistical significance.

Figure 1. **Association between Metformin use and systemic inflammatory and cytotoxic markers.** Forest plot displaying beta coefficients and 95% confidence intervals from linear regression models evaluating the effect of metformin use on plasma protein levels (Olink Explore 3072). Markers to the left of the dashed vertical line indicate a reduction in circulating protein levels associated with metformin. All models were adjusted for age, sex, BMI, and Townsend Deprivation Index to account for metabolic and socioeconomic confounders. Highly significant reductions were observed in IL-6 (p = 7.70 e-5) and CXCL10 (p = 1.01 e-12), corroborating the "cell-extrinsic" low-inflammation state associated with viral control.

References

1.   Castro-Gonzalez S, Colomer-Lluch M, Serra-Moreno R. Barriers for HIV Cure: The Latent Reservoir. AIDS Res Hum Retroviruses. 2018;34(9):739-59.
2.   Margolis DM, Archin NM. Proviral Latency, Persistent Human Immunodeficiency Virus Infection, and the Development of Latency Reversing Agents. J Infect Dis. 2017;215(suppl_3):S111-s8.
3.   Kalidasan V, Theva Das K. Learning from nature: HIV elite controllers as blueprints for a functional cure. Virology Journal. 2025;23(1):23.
4.   Pellaers E, Denis A, Debyser Z. New latency-promoting agents for a block-and-lock functional cure strategy. Curr Opin HIV AIDS. 2024;19(3):95-101.
5.   Ma T, George AF, Li Z, Thomas R, Yin K, Shin MG, et al. Multiomic analysis of ART-interruption cohorts identifies cell-extrinsic and -intrinsic mechanisms driving lymphocyte-mediated control of HIV rebound. Immunity. 2026.
6.   Cooper L, Xu H, Polmear J, Kealy L, Szeto C, Pang ES, et al. Type I interferons induce an epigenetically distinct memory B cell subset in chronic viral infection. Immunity. 2024;57(5):1037-55.e6.
7.   Luxenburger H, Thimme R, Hofmann M. T cell adaptation in chronic infections and tumors. Cellular & Molecular Immunology. 2026.
8.   Hou W, Jin YH, Kang HS, Kim BS. Interleukin-6 (IL-6) and IL-17 synergistically promote viral persistence by inhibiting cellular apoptosis and cytotoxic T cell function. J Virol. 2014;88(15):8479-89.
9.   Wu G, Fan Q, Chen M, Luo G, Wu Z, Zhao J, et al. Activation of AMP-activated Protein Kinase by Metformin Inhibits Dedifferentiation of Platelet-derived Growth Factor-BB-induced Vascular Smooth Muscle Cells to Improve Arterial Remodeling in Cirrhotic Portal Hypertension. Cell Mol Gastroenterol Hepatol. 2025;19(6):101487.
10.   Foretz M, Guigas B, Viollet B. Metformin: update on mechanisms of action and repurposing potential. Nat Rev Endocrinol. 2023;19(8):460-76.
11.   Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, et al. The UK Biobank resource with deep phenotyping and genomic data. Nature. 2018;562(7726):203-9.