medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

High occurrence of transportation and logistics occupations among vascular dementia patients: an observational study

A.C. van Loenhoud,[1]* C. de Boer,[1] K. Wols, Y.A. Pijnenburg,[1] A.W. Lemstra,[1] F.H. Bouwman,[1] N.D. Prins,[1] P. Scheltens,[1] R. Ossenkoppele,[1,3] W.M. van der Flier[1,2]

[1]Alzheimer Center Amsterdam, Department of Neurology, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, 1081 HV, Amsterdam, The Netherlands.

[2]Department of Epidemiology and Biostatistics, Vrije Universiteit Amsterdam, Amsterdam UMC, 1081 HV, Amsterdam, The Netherlands.

[3]Clinical Memory Research Unit, Lund University, 221 00, Lund, Sweden.

*corresponding author; address: room PK -1 Z 039, Vrije Universiteit Amsterdam, Amsterdam UMC, De Boelelaan 1118, 1081 HV, Amsterdam, The Netherlands; email: a.vanloenhoud@vumc.nl; telephone: +31 20 4448527; fax: +31 20 4448529.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

Abstract

Background: Growing evidence suggests a role of occupation in the emergence and manifestation of dementia. Occupations are often defined by complexity level, although working environments and activities differ in several other important ways. We aimed to capture the multi-faceted nature of occupation through its measurement as a categorical (instead of quantitative) variable and explored its relationship with different types of dementia.

Methods: We collected occupational history data of 2,121 dementia patients with various suspected etiologies from the Amsterdam Dementia Cohort (age: 67±8, 57% male, MMSE: 21±5). Our final sample included individuals with Alzheimer's disease (AD) dementia (n=1,467), frontotemporal dementia (n=281), vascular dementia (n=98), Lewy Body diseae (n=174) and progressive supranuclear palsy/corticobasal degeneration (n=101). Within the AD group, we used neuropsychological data to further characterize patients by clinical phenotypes. All participants were categorized into one of 11 occupational classes, across which we evaluated the distribution of dementia (sub)types with Chi^2 analyses. We gained further insight into occupation-dementia relationships through post-hoc logistic regressions that included various demographic and health characteristics as explanatory variables.

Results: There were significant differences in the distribution of dementia types across occupation groups (Chi^2=85.87, p<.001). Vascular dementia was relatively common in the Transportation/Logistics sector, and higher vascular risk factors partly explained this relationship. Alzheimer's disease occurred less in Transportation/Logistics and more in Health Care/Welfare occupations, which related to a higher/lower percentage of males. We found no relationships between occupational classes and clinical phenotypes of AD (Chi^2=53.65, n.s.).

Conclusions: Relationships between occupation and dementia exist beyond complexity level, which offers new opportunities for disease prevention and improvement of occupational health policy.

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

Keywords: occupation, vascular dementia, Alzheimer's disease, vascular risk factors, sex

Background

Dementia currently affects 46.8 million people worldwide [1], and this number is expected to increase as the aging population grows. Causes of dementia include Alzheimer's disease (AD), frontotemporal dementia (FTD), vascular dementia (VaD) and movement disorders such as Lewy Body disease (DLB), progressive supranuclear palsy (PSP) and corticobasal degeneration (CBD). Apart from age and genetic influences, lifestyle factors play a role in the emergence and manifestation of dementia. For example, a comprehensive body of research points to the influence of education and occupation on dementia incidence [2-5]. Occupation could be an especially strong contributing factor, as most individuals spend a major part of their life at work.

Studies on the relationship between occupation and dementia have generally focused on the role of occupational complexity (e.g. rated as "white versus blue collar" [6]). Results indicate that older individuals who engaged in greater levels of occupational complexity have a better current cognitive status [6-13], a lower risk to develop dementia [5,14-17] or a different rate of clinical progression after the onset of dementia [4,11,18]. However, these studies mostly investigated one particular form of dementia, or "dementia" in general without specifying its type. Moreover, the common quantification of occupation by complexity level represents an oversimplification of the multi-faceted quality of this lifestyle factor. Working environments and activities differ in several important ways beyond complexity level, such as the level of physical activity, stress, social demands and exposure to hazardous substances [19,20]. Although a broader evaluation of occupation could thus provide important new insights in the context of dementia, this approach is relatively rare.

To overcome this scarcity in the literature, we took a different approach by i) measuring occupation in a categorical rather than a quantitative manner, without making prior

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

assumptions about complexity or other potential protective or harmful factors, and ii) studying the relationships between occupation and various forms of dementia (i.e. different suspected etiologies and clinical phenotypes). Our sample included >2000 memory clinic patients with dementia due to AD, FTD, VaD or movement disorders. Occupation was categorized into several nominal groups (e.g. Pedagogical, Agricultural, Transportation/Logistics), across which we compared the distribution of these dementia types [21]. Within the AD subsample, we had sufficient data to additionally investigate relationships between occupation and specific cognitive profiles. As this study was exploratory in nature, we did not formulate a priori hypotheses. For relationships that emerged, we gained further insight through post-hoc analyses with additional demographic and health characteristics (i.e. age, sex, education and vascular risk factors [VRFs]).

Methods

Participants

Participants were selected from the Amsterdam Dementia Cohort (ADC [22,23]). The sample consists of individuals who visited the Alzheimer Center of the Amsterdam University Medical Center between 2000 and 2017. They underwent a standardized screening process that included the (patient and informant based) collection of demographic information, medical history, neurological examination, neuropsychological assessment, standard laboratory tests, brain magnetic resonance imaging (MRI) and lumbar puncture in a subsample. Diagnosis was established in a multidisciplinary meeting, based on common clinical and research criteria [22]. Dementia diagnoses in the present study included AD, FTD, VaD, DLB and PSP/CBD.

Assignment of occupational codes

Information on occupational history was collected by a physician during a semi-structured interview with the participants and their informants. We subsequently coded this information according to the Dutch ROA-CBS 2014 occupational categorization system (BRC 2014), a

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

derivative of the International Standard Classification of Occupations 2008 (ISCO 2008) [21]. The BRC 2014 consists of the following occupational classes: 1) Pedagogical, 2) Creative/Linguistic, 3) Commercial, 4) Business/Administrative, 5) Management, 6) Governmental/Law/Safety, 7) Technical, 8) Information and Communication Technology (ICT), 9) Agricultural, 10) Health Care/Welfare, 11) Service, 12) Transportation/Logistics (see Additional file 1). The BRC 2014 additionally includes a 13th "Other" category (mostly applicable when occupational descriptions are absent or unclear), and makes further divisions into segments and subgroups. For the present paper, we were only interested in the occupational classes (i.e. classification at the segment level would result in 40 groups of insufficient size). We excluded participants in the "Other" category and those classified as "no occupation". When multiple occupations were listed for a given participant, we used the first job title for consistency reasons.

The assignment of BRC 2014 codes was performed by two raters (A.C.v.L and K.W.). To promote interrater consistency, a subset of the sample was independently coded and subsequently compared. Discordant cases were discussed with R.O. to reach consensus. The remaining sample was divided among A.C.v.L. and K.W., and only ambiguous cases were discussed and coded together. In rare occasions where this did not resolve classification uncertainty, medical records were inspected for more details on occupational history. If this did not provide useful information, the participant was coded as 13 (unclear), hence excluded for further analysis.

Sample selection

We initially selected all ADC individuals with a dementia diagnosis at the baseline visit and information on occupational history. Exclusion criteria were: 1) age below 40 years, 2) presence of autosomal dominant mutations for a neurodegenerative disease. The original selection consisted of 2,310 participants, of which we excluded 142 individuals with occupational descriptions that were either non-informative, indicative of unemployment or not

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

detailed enough (even after inspection of medical records). Finally, we decided to omit the category ICT due to the small sample size (n=12). ICT is a relatively "young" sector that did not have a prominent position in the labor market during the period in which the majority of our sample was employed. Similarly, 35 patients with relatively rare forms of dementia (e.g. alcohol-related dementias, normal pressure hydrocephalus, Creutzfeldt–Jakob disease) were excluded (Figure 1). The final sample included 2,121 individuals, with a diagnosis of AD (n=1,467), FTD (n=281), VaD (n=98), DLB (n=174) or PSP/CBD (n=101).

Vascular risk factors

Complete data on VRFs were available for the majority of our sample (n=1,562/2,121), and included body mass index (BMI), smoking, alcohol use (units per day), and presence or absence of a myocardial infarction, hypertension, hypercholesterolemia and diabetes mellitus in the medical history. We dichotomized BMI (≤25=0, >25=1), smoking (never smoked=0, ever smoked=1) and alcohol (0-1 units per day=0, >1 units per day=1) in order to derive a VRF score based on the sum of these seven risk factors [24].

Neuropsychological assessment

Nearly all individuals underwent an elaborate neuropsychological assessment within three months of their diagnosis. For the AD dementia subsample, we therefore had a large dataset available (n=1,071/1,467) that consisted of participants who were cognitively tested in each of four cognitive domains: memory, attention/executive function, language and visuospatial function [25]. The memory domain included the total immediate and delayed recall as well as the recognition condition of the Rey Auditory Verbal Learning Test, and total recall in condition A of the Visual Association Test. In the attention/executive function domain, we used the Trail Making Test (part A and B), Stroop Test (condition I-III), 90-second Letter Digit Substitution Test [26], Digits Forward and Backwards, Letter Fluency and total score on the Frontal Assessment Battery. The language domain consisted of the Category Fluency, a short version of the Boston Naming Test and the total number of correctly named items of

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

the Visual Association Test. Finally, the visuospatial function domain was assessed with the Dot Counting, Number Location and Fragmented Letters subtests of the Visual Object and Space Perception battery.

We carried out multiple imputation in SPSS 22.0 for Windows (SPSS, Chicago, IL, USA) to account for missing data (ranging from 1-45% per test). This procedure was performed using the fully conditional specification method, an iterative Markov Chain Monte Carlo approach suitable for arbitrary patterns of missing data. We imputed 25 datasets and included demographic, clinical, neuropsychological variables, APOE status and cerebrospinal fluid biomarkers as predictors in the model (i.e. occupational code was not used). In the pooled dataset, we created standardized residuals (i.e. W-scores) for each neuropsychological test score based on regressions with age, sex and educational level (measured on a seven-item scale, [27]). We calculated average W-scores across tests measuring memory, attention/executive function, language and visuospatial function, to obtain cognitive domain scores.

These cognitive domain scores were subsequently used to divide AD participants into clinical phenotypes, according to a previously validated method [28]. Specifically, we averaged across cognitive domains to create a “global cognition” composite score for each participant. Next, each cognitive domain score was dichotomized based on whether they were below a certain threshold relative to the global cognition score. A score of 1 thus means that a given cognitive domain “stands out” by being more affected than (the combination of) other domains. Based on these dichotomized scores, we assigned each participant to one of the following cognitive profiles: 1) memory-dominant, 2) attention/executive-dominant, 3) language-dominant, 4) visuospatial-dominant, 5) multi-domain (i.e. either more than one or no cognitive domain=1). As this profile categorization relies on which threshold is used to dichotomize the cognitive domain scores, we systematically tested all values between .250 and .500, and selected .225 as the “optimal” value that yielded 1) the lowest number of

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

participants in the multi-domain profile, 2) the lowest sum of squared group sizes (Supplementary Figure 1). Most individuals had a memory-dominant cognitive profile (n=292), followed by a language-dominant (n=140), attention/executive-dominant (n=130) and visuospatial-dominant subgroup (n=126). The remaining participants fell into the multi-domain category (n=383).

Statistical analyses

First, we carried out a Chi^2 analysis in the total sample to investigate if the distribution of dementia types differed across occupational classes. In a second analysis, we reran the same model after exclusion of 93 persons with comorbid dementias (e.g. persons with AD as the primary diagnosis, but who also met NINDS-AIREN criteria for VaD) to create a sample with less overlap between diagnostic groups.

When Chi^2 results were significant, we post-hoc inspected the adjusted residuals of each combination of occupation and dementia type. Adjusted residuals reflect the difference between the expected and observed count (e.g. the number of AD participants in the Pedagogical occupational class). An absolute value of ~2 (i.e. 1.96 or -1.96) is considered significant at the $p<.05$ level. Since our contingency table was relatively large, we used a more stringent adjusted residual value threshold of ~3 (i.e. 3.29 or -3.29 corresponds to $p<.001$; [29]). In addition, to assess whether the main relationships observed in significant Chi^2 analyses could be explained by other demographic and health variables, we additionally performed logistic regressions. Specifically, we predicted dementia type from occupational class and evaluated age, sex, education and VRF score as potential explanatory variables. We used a forward selection procedure with a change-in-estimate (CIE) criterion of 10% [30,31] and ran these models in all participants with complete data (n=1,562/2,121).

Finally, we carried out Chi^2 analyses to investigate whether occupation related to cognitive profiles within the AD subsample, following the same approach as described above. In a

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

second analysis, we reran the same model using a different threshold for the dichotomization of cognitive domain scores (and subsequent categorization of participants into cognitive profiles; Supplementary Figure 1), and compared results to evaluate robustness of our findings.

Results

Participants

The mean age across diagnostic groups was 67±8 years (Table 1). Individuals with FTD were younger compared to other groups (63±7), and DLB participants were significantly older than the AD group (69±7). Overall, our sample included a somewhat larger proportion of males (57%), presumably reflecting the historically higher prevalence of males in the labor market (by comparison, the excluded "no occupation" sample was 82% female). The proportion of males was even higher among DLB participants (82%) in comparison with most other groups, while this was significantly lower for AD (52%) in contrast to all groups except PSP/CBD. Educational level was similar across dementia types (except for a DLB>VaD difference). Global cognitive impairment was most severe among AD participants (mean MMSE=20±5). Finally, VaD participants had a higher VRF score than most other diagnostic groups ($p<.05$; except FTD). Table 2 provides an overview of the number of participants in each occupational class for the total sample and according to dementia type. The largest occupation groups were Technical (n=429, 20%) and Business/Administrative (n=388, 18%), while the Agricultural class had the lowest number of participants (n=31, 1%).

Relationship between occupation and dementia types

There were significant differences in the distribution of dementia types across occupation groups ($Chi^2=85.87$, $p<.001$, Table 2). The adjusted residuals revealed three effects significant at $p<.001$. First, individuals from the Transportation/Logistics sector were more often diagnosed with VaD (adjusted residual: 4.0). Second, this occupational class had fewer AD participants (adjusted residual: -3.7). Third, in the Health Care/Welfare group, AD was

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

relatively common (adjusted residual: 3.6). These results are displayed in Figure 2, along with 11 other effects (e.g. a high proportion of PSP/CBD cases in the Agricultural sector) that were significant at p<.05 (adjusted residuals >2 or <-2). When we repeated the Chi^2 analysis with exclusion of 93 individuals with comorbid dementias (n=2,028), results remained essentially unchanged (Chi^2=78.73, p<.001; Supplementary Table 1).

Contribution of age, sex, education and vascular risk factors

In line with the previous results, uncorrected logistic regression models with either AD or VaD as a dependent variable (AD/VaD=1 in separate models, other diagnosis=0) confirmed Transportation/Logistics to be significantly related to both VaD (β=1.23, odds ratio [OR] = 3.41, p=.004) and AD (β=-.85, OR=.43, p=.001). Likewise, Health Care/Welfare was a significant determinant of AD (β=.55, OR=1.74, p=.003; Supplementary Table 2). Forward selection of explanatory variables in corrected models revealed that VRF score, but no other factors (i.e. age, sex, education), was positively associated with a VaD diagnosis (β=.43, OR=1.53, p<.001) and reduced the Transportation/Logistics effect by 15% (from β=1.23 to 1.04, OR=3.41 to 2.84). For AD, only the variable sex (range β=.68-.69, OR=1.96-1.98, p<.001) significantly reduced the effects of Transportation/Logistics (from β=-.85 to -.62, OR=.43 to .54; 27%) and Health Care/Welfare (from β=.55 to .31, OR=1.74 to 1.37; 43%). Figure 3 gives an overview of the number of VRFs and percentage of males for all dementia types and occupational classes.

Relationship between occupation and cognitive profiles of AD

We found only trend-significant differences between occupations in the distribution of cognitive profiles (Chi^2=53.65, p=.073; Table 3). Supplementary Figure 2 provides an overview of these findings. In a secondary analysis, we repeated the same analysis using cognitive profiles that were calculated based on a different threshold of -.442 (Supplementary Figure 1). Similar to the original findings, these results were not significant (Chi^2=38.24, p=.550).

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

253

254 Discussion

255 In this exploratory study, we investigated the association between occupation and dementia
256 (sub)types in a large sample of memory-clinic patients. The main finding was that VaD was
257 relatively common in participants with Transportation/Logistics occupations. Post-hoc
258 analyses suggested that this relationship was partly explained by a higher number of VRFs
259 among these individuals. In addition, the number of AD participants was relatively low in the
260 Transportation/Logistics sector, and high among Health Care/Welfare occupations. We found
261 that these effects were mostly attributable to differences between groups in sex distribution.
262 There was no relationship between occupation types and AD-related cognitive profiles.

263

264 Our finding that VRFs are more prominent among VaD participants is in line with previous
265 studies [32,33], but of particular interest was our observation of a higher number of VRFs in
266 Transportation/Logistics occupations. Although we cannot make any inferences about
267 dementia risk (due to the absence of a healthy control group), it raises the possibility that
268 certain job-related aspects or occupational habits contribute to the development of VRFs and
269 the emergence of VaD. Our data at least warrant an increased awareness and monitoring of
270 VRFs by occupational physicians in the Transport/Logistics sector. Furthermore, the lower
271 presence of AD cases in Transpiration/Logistics groups (male-predominant) versus higher
272 AD occurrence Healthcare/Welfare (female-predominant), is in line with evidence that
273 women have a higher risk for AD than men [34,35]. Importantly, while VRFs and sex partially
274 explained our main occupation-dementia relationships, these factors did not account for all
275 variance. This suggests there might be other important ways in which occupations are
276 related to the emergence and manifestation of dementia.

277

278 In addition, it is worth mentioning that a relatively large proportion of the Agricultural sector
279 consisted of PSP/CBD cases (i.e. 13% versus 5% in the total sample). This effect was
280 significant at $p<.05$ (not at $p<.001$), but as the Agricultural group was rather small compared

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

to other occupational classes (n=31), this may be due to a lack of statistical power. Although largely speculative, we propose that this effect could be related to exposure to pesticides [36]. There has been increasing evidence that pesticides are a risk factor for Parkinson's disease [37], and one study has also demonstrated a specific link between pesticide exposure and PSP [38].

In contrast to our present study, most previous studies operationalized occupation by complexity level. Apart from coarse ratings such as "white versus blue collar", more fine-grained classifications have been made in which the complexity of multiple occupational aspects was considered. An example is the use of questionnaires that differentiated between working with "people, data or things" (e.g. [12]). However, although these approaches have provided important insights, they did not consider other potential job-related risks or protective factors. The few studies that did take a more qualitative approach towards the measurement of occupation have used "job exposure matrices" or "occupational description databases" to show, among other results, that dementia risk and late-life cognition were influenced by the degree of work control (e.g. influence over work planning, variation in task content) [39], level of human interaction and physical activity [40] and mental demands (e.g. information processing, pattern detection, creativity) [41].

The novelty of the present work lies our "assumption-free" approach that did not involve any additional characterization of occupations besides their categorization across 11 occupational classes. This enabled examining relationships between occupation and dementia (sub)types in an exploratory manner. Importantly, there was no straightforward relationship between occupational classes and complexity level. Most classes comprised occupations that varied considerably with respect to complexity (e.g. the "Technical" group both contained engineers and assembly workers, see Additional file 1). Another strength of our study is the fact that our sample consisted of dementias with different suspected etiologies. Most earlier studies have either focused on the occurrence of dementia as a

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

general syndrome or one particular form of dementia. As our dataset included over two thousand carefully diagnosed participants, we were able to create several independent diagnostic groups and AD-related cognitive profiles that were sufficiently sized for statistical analysis. Finally, we consider use of the BRC 2014 classification system to be an important advantage, because it improves the replicability of our findings across other populations on an international scale.

Our study has several limitations. First, although we collected occupational data with a refined method that involved a semi-structured interview with the patient and caregiver, we did not have detailed enough information to capture a person's entire occupational timeline. For individuals who had multiple occupations listed, we often could not retrieve their sequence and respective durations. It is therefore possible that the first-mentioned job not always reflect the person's foremost/primary occupation, but rather the most recent occupation. This is non-optimal because of potential reverse causation: we cannot rule out that some patients switched jobs as a consequence of their dementia diagnosis (e.g. choosing less demanding work in a different sector). However, we have no reason to assume that reverse causation is more prominent for some dementia (sub)types than others, and thus the degree of bias seems limited. Second, we were unable to define occupation groups beyond the highest hierarchical level of the BRC 2014, due to a lack of detail in occupational descriptions and the fact that the resulting number of categories would be too large for our sample size (i.e. ~40 occupation groups). More detailed distinctions between occupations may have revealed additional relevant occupation-dementia relationships that were currently missed. Third, the absence of a healthy control group restricted our ability to make inferences about dementia risk or distinguish between protective and harmful effects of occupation (Supplementary Figure 3), which limits the interpretability of our results.

Conclusions

To conclude, we demonstrated relationships between occupation and dementia types. Our

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

findings suggest that these relationships emerged beyond occupational complexity level. Differences between occupational classes in the number of VRFs and sex distributions partially explained our results, but other - currently unknown - factors likely play a role as well. It is important to note that associations between occupation and dementia could exist for multiple reasons. Whilst it seems plausible that some occupational characteristics causally relate to the occurrence of specific dementia types, another possibility is that certain (genetic or early environmental) factors both influence career choice and dementia development later in life. This is consistent with theoretical frameworks that propose a neurodevelopmental component in the etiology of dementia [42-44]. Ultimately, as most people spend the majority of their life working, occupation could be an important lifestyle factor to consider in relation to preventive strategies for dementia. A better understanding of occupation-dementia relationships may improve occupational health policy through reduction of job hazards and more targeted health monitoring by occupational physicians.

List of abbreviations

AD – Alzheimer's disease, VaD – vascular dementia, FTD – frontotemporal dementia, DLB – Lewy Body disease, PSP – progressive supranuclear palsy, CBD – corticobasal degeneration, ADC – Amsterdam Dementia Cohort, MRI – magnetic resonance imaging, BRC 2014 – Dutch ROA-CBS 2014 occupational categorization system, ISCO 2008 – the International Standard Classification of Occupations 2008, ICT – Information and Communication Technology, BMI – body mass index, VRF – vascular risk factor, CIE – change-in-estimate, OR – odds ratio.

Declarations

Ethics approval and consent to participate

Participants gave written informed consent to use their medical data for scientific purposes. This procedure was approved by the local Medical Ethics Committee.

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

Consent for publication

Not applicable

Availability of data and material

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

Research of Alzheimer Center Amsterdam is part of the Neurodegeneration program of Amsterdam Neuroscience. The Alzheimer Center Amsterdam is supported by Alzheimer Nederland, Het Genootschap and Stichting VUMC funds. W.M. van der Flier holds the Pasman chair. This research was funded by the Internationale Stichting Alzheimer Onderzoek (ISAO) (to R.O.).

Authors' contributions

A.C.v.L. was involved in the methodological design of the research project and assignment of occupational codes; she furthermore conducted the statistical analyses, interpreted the data and wrote the manuscript. C.d.B. was responsible for the selection of the participant data from the ADC. K.W. took part in the methodological design and the assignment of occupational codes. Y.A.P., A.W.L., F.H.B., N.D.P., and P.S. were involved in the standardized screening process of participants and collection of occupational history data. R.O. was involved in the methodological design of the research project, interpretation of the data and writing the manuscript. W.M.v.d.F. is the coordinator of the ADC, and took part in the interpretation of the data and editing of the manuscript. All authors read and approved the final manuscript.

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

Acknowledgements

Not applicable

References

1. Prince M, Wilmo A, Guerchet M, Ali G, Wu Y, Prina M. World Alzheimer Report 2015. The Global Impact of Dementia: An Analysis of Prevalence, Incidence, Cost and Trends. Alzheimer's Disease International. 2015.
2. Bonaiuto S, Rocca WA, Lippi A, Giannandrea E, Mele M, Cavarzeran F, Amaducci L. Education and Occupation as Risk Factors for Dementia: A Population-Based Case-Control Study. Neuroepidemiology 1995; 14:101-109.
3. Stern Y, Albert S, Tang MX, Tsai WY. Rate of memory decline in AD is related to education and occupation. Neurology. 1999;53(9):1942-7.
4. Hughes TF, Ganguli M. Modifiable Midlife Risk Factors for Late-Life Cognitive Impairment and Dementia. Curr Psychiatry Rev. 2009; 5(2): 73–92.
5. Dekhtyar S, Wang HX, Fratiglioni L, Herlitz A. Childhood school performance, education and occupational complexity: a life-course study of dementia in the Kungsholmen Project. Int J Epidemiol. 2016; 45(4):1207-1215.
6. Ihle A, Ghisletta P, Ballhausen N, Fagot D, Vallet F, M Baeriswyl, et al. The role of cognitive reserve accumulated in midlife for the relation between chronic diseases and cognitive decline in old age: A longitudinal follow-up across six years. Neuropsychologia 2018, 121: 37-46.
7. Foubert-Samier A, Catheline G, Amieva H, Dilharreguy B, Helmer C, Allard M, Dartigues

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

JF. Education, occupation, leisure activities, and brain reserve: a population-based study. Neurobiol Aging. 2012; 33(2):423.e15-25.

8. Correa Ribeiro PC, Lopes CS, Lourenço RA. Complexity of lifetime occupation and cognitive performance in old age. Occupational Medicine 2013; 63(8): 556–562.
9. Vemuri P, Lesnick TG, Przybelski SA, Knopman DS, Preboske GM, Kantarci K, et al. Vascular and amyloid pathologies are independent predictors of cognitive decline in normal elderly. Brain 2015; 138(3): 761–771.
10. Placek K, Massimo L, Olm C, Ternes K, Firn K, Van Deerlin V, et al. Cognitive reserve in frontotemporal degeneration: Neuroanatomic and neuropsychological evidence. Neurology 2016: 87(17):1813-1819.
11. Pool LR, Weuve J, Wilson RS, Bültmann U, Evans DA, Mendes de Leon CF. Occupational cognitive requirements and late-life cognitive aging. Neurology 2016; 86(15):1386-1392.
12. Lane AP, Windsor TD, Andel R, Luszcz MA. Is Occupational Complexity Associated with Cognitive Performance or Decline? Results from the Australian Longitudinal Study of Ageing. Gerontology 2017; 63(6):550-559.
13. Kaup AR, Xia F, Launer LJ, Sidney S, Nasrallah I, Erus G, et al. Occupational cognitive complexity in earlier adulthood is associated with brain structure and cognitive health in midlife: The CARDIA study. Neuropsychology 2018; 32(8):895-905.
14. Dekhtyar S, Wang HX, Scott K, Goodman A, Koupil I, Herlitz A. A Life-Course Study of Cognitive Reserve in Dementia--From Childhood to Old Age. Am J Geriatr Psychiatry 2015;23(9):885-96.
15. Vuoksimaa E, Rinne JO, Lindgren N, Heikkilä K, Koskenvuo M, Kaprio J. Middle age self-report risk score predicts cognitive functioning and dementia in 20-40 years. Alzheimers Dement (Amst). 2016;4:118-125.
16. Rusmaully J, Dugravot A, Moatti JP, Marmot MG, Elbaz A, Kivimaki M, et al. Contribution of cognitive performance and cognitive decline to associations between socioeconomic factors and dementia: A cohort study. PLoS Med 2017;14(6):e1002334.

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

17. Wang HX, MacDonald SW, Dekhtyar S, Fratiglioni L. Association of lifelong exposure to cognitive reserve-enhancing factors with dementia risk: A community-based cohort study. PLoS Med 2017;14(3):e1002251.

18. Massimo L, Xie SX, Rennert L, Fick DM, Halpin A, Placet K, et al. McMillan, Occupational attainment influences longitudinal decline in behavioral variant frontotemporal degeneration. Brain Imaging Behav. 2018. doi: 10.1007/s11682-018-9852-x

19. Santibáñez M, Bolumar F, García AM. Occupational risk factors in Alzheimer's disease: a review assessing the quality of published epidemiological studies. Occup Environ Med. 2007;64(11):723-32.

20. Killin LO, Starr JM, Shiue IJ, Russ TC. Environmental risk factors for dementia: a systematic review. BMC Geriatr. 2016;16(1):175.

21. Fouarge D, Dijksman S. Beroepenindeling ROA-CBS 2014 (BRC 2014). (ROA Technical Reports; No. 005). Maastricht: Research Centre for Education and the Labour Market; 2015.

22. Van der Flier WM, Pijnenburg YA, Prins N, Lemstra AW, Bouwman FH, Teunissen CE, et al. Optimizing patient care and research: the Amsterdam Dementia Cohort. J Alzheimers Dis. 2014;41(1):313-327.

23. Van der Flier WM, Scheltens P. Amsterdam Dementia Cohort: Performing Research to Optimize Care. J Alzheimers Dis. 2018;62(3):1091-1111.

24. Groot C, Sudre CH, Barkhof F, Teunissen CE, Van Berckel BNM, Seo SW, et al. Clinical phenotype, atrophy, and small vessel disease in APOEε2 carriers with Alzheimer disease. Neurology 2018,91(20):e1851-e1859 .

25. Groot C, Van Loenhoud AC, Barkhof F, Van Berckel BNM, Koene T, Teunissen CE, et al. Differential effects of cognitive reserve and brain reserve on cognition in Alzheimer's disease. Neurology 2018; 90(2): e149-e156.

26. Van der Elst W, Van Boxtel MPJ, Van Breukelen GJP, Jolles J. The Letter Digit Substitution Test: Normative Data for 1,858 Healthy Participants Aged 24–81 from the

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

Maastricht Aging Study (MAAS): Influence of Age, Education, and Sex, Journal of Clinical and Experimental Neuropsychology 2006; 28(6):998-1009.

27. Verhage F. Intelligence and age (in Dutch). Assen: Van Gorcum; 1964.
28. Crane PK, Trittschuh E, Mukherjee S, Saykin AJ, Sanders RE, Larson EB, et al. Incidence of cognitively defined late-onset Alzheimer's dementia subgroups from a prospective cohort study. Alzheimers Dement. 2017;13(12):1307-1316.
29. Sharpe D. Your chi-square test is statistically significant: Now what? Practical Assessment, Research & Evaluation 2015.
30. Mickey RM, Greenland S. The impact of confounder selection criteria on effect estimation. Am J Epidemiol. 1989;129:125–37.
31. Maldonado G, Greenland S. Simulation study of confounder-selection strategies. Am J Epidemiol. 1993;138(11):923–36.
32. Fillit H, Nash DT, Rundek T, Zuckerman A. Cardiovascular risk factors and dementia. Am J Geriatr Pharmacother. 2008;6(2):100-118.
33. Tariq S, Barber PA. Dementia risk and prevention by targeting modifiable vascular risk factors. J. Neurochem. 2018; 144:565-581.
34. Viña J, Lloret A. Why women have more Alzheimer's disease than men: gender and mitochondrial toxicity of amyloid-beta peptide. J Alzheimers Dis. 2010;20 Suppl 2:S527-33.
35. Podcasy JL, Epperson CN. Considering sex and gender in Alzheimer disease and other dementias. Dialogues Clin Neurosci. 2016;18(4):437-446.
36. Gangemi S, Miozzi E, Teodoro M, Briguglio G, De Luca A, Alibrando C, et al. Occupational exposure to pesticides as a possible risk factor for the development of chronic diseases in humans (Review). Mol Med Rep. 2016;14(5):4475-4488.
37. Brouwer M, Huss A, Van der Mark M, Nijssen PCG, Mulleners WM, Sas AMG, et al. Environmental exposure to pesticides and the risk of Parkinson’s disease in the Netherlands. Environ Int. 2017; 107: 100-110.
38. Litvan I, Lees PS, Cunningham CR, Rai SN, Cambon AC, Standaert DG, et al.

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

Environmental and occupational risk factors for progressive supranuclear palsy: Case-control study. Movement disorders: official journal of the Movement Disorder Society 2016, 31(5), 644-52.

39. Hasselgren C, Dellve L, Ekbrand H, Zettergren A, Zetterberg H, Blennow K, et al. Socioeconomic status, gender and dementia: The influence of work environment exposures and their interactions with APOE □4. SSM Popul Health. 2018;5:171-179.
40. Potter GG, Helms MJ, Plassman BL. Associations of job demands and intelligence with cognitive performance among men in late life. Neurology.2007;70(19 Pt 2):1803-8.
41. Then FS, Luck T, Heser K, Ernst A, Posselt T, Wiese B, et al. Which types of mental work demands may be associated with reduced risk of dementia? Alzheimers Dement. 2017;13(4):431-440.
42. Miller ZA, Rosenberg L, Santos-Santos MA, Stephens M, Allen IE, Hubbard HI, et al. Prevalence of Mathematical and Visuospatial Learning Disabilities in Patients With Posterior Cortical Atrophy. JAMA Neurol. 2018;75(6):728-737.
43. Miller ZA, Mandelli ML, Rankin KP, Henry ML, Babiak MC, Frazier DT, et al. Handedness and language learning disability differentially distribute in progressive aphasia variants. Brain. 2013;136(11):3461-3473.
44. Geschwind DH, Robidoux J, Alarcón M, Miller BL, Wilhelmsen KC, Cummings JL, et al. Dementia and neurodevelopmental predisposition: cognitive dysfunction in presymptomatic subjects precedes dementia by decades in frontotemporal dementia. Ann Neurol. 2001;50(6):741-746.

Figure 1. Flow diagram of the study sample.

ADC=Amsterdam dementia cohort, BRC 2014= Dutch ROA-CBS 2014 occupational categorization system, a derivative of the Standard Classification of Occupations 2008 (ISCO 2008), ICT= Information and Communication Technology.

Figure 2. Proportions of dementia types for each occupational class.

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

AD=Alzheimer's disease dementia, FTD=frontotemporal dementia, VaD=vascular dementia, DLB=Lewy Body disease, PSP=progressive supranuclear palsy, CBD=corticobasal degeneration. *Chi^2 adjusted residual ≤ -2 or ≥2 (corresponding to p<.05), ***Chi^2 adjusted residual ≤ -3 or ≥3 (corresponding to p<.001).

Figure 3. Mean number of vascular risk factors and male percentage for occupational classes and dementia types.

Error bars represent standard errors. AD=Alzheimer's disease dementia, FTD=frontotemporal dementia, VaD=vascular dementia, DLB=Lewy Body disease, PSP=progressive supranuclear palsy, CBD=corticobasal degeneration. VRF=vascular risk factors, including included body mass index (BMI), smoking, alcohol use (units per day), and presence or absence of a myocardial infarction, hypertension, hypercholesterolemia and diabetes mellitus in the medical history.

Supplementary Figure 1. Thresholds used for the dichotomization of cognitive domains and division of participants across cognitive profiles.

Within the AD subsample (n=1,071), we dichotomized each cognitive domain score (i.e. memory, attention/executive functions, language and visuospatial functions), based on whether or not a domain's W-score was considerably lower compared to an individual's global cognition score. We used an optimal threshold between .250 and .500 for this dichotomization, and defined "optimality" in two different ways: 1) the lowest number of participants in the multi-domain profile and the lowest sum of squared group sizes (threshold=.255), 2) the highest number of atypical variant AD cases (n=85) categorized into the language (i.e. for logopenic aphasia) or visuospatial cognitive profile (i.e. for PCA; n=.442). Note that neither thresholds (i.e. .255 or .442) resulted in all atypical variants being categorized into these two cognitive profiles, presumably because in advanced disease stages, several cognitive domains become affected. For many participants with an initial logopenic aphasia or PCA diagnosis, visuospatial/language may no longer have been

medRxiv preprint doi: https://doi.org/10.1101/19005512; this version posted August 31, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
All rights reserved. No reuse allowed without permission.

predominant at study inclusion, causing them to be assigned to the multi-domain instead.

Supplementary Figure 2. Proportions of AD-related cognitive profiles for each occupational class.

There were no significant differences between groups (Chi^2 n.s.).

Supplementary Figure 3. Schematic representation of protective and harmful effects and resulting cells in the contingency table.

Each color (red, blue, green) represents a different occupational class. In scenario A, the red occupational class is relatively protected against dementia type 1, while the distribution of the remaining dementia types across occupations is equal. When the healthy control group is included in the analysis, this protective effect will be reflected in a higher proportion of individuals in red in the healthy group (4/6=67% versus 50% for blue/green) and a lower percentage of red persons in the type 1 dementia group (0% versus 1/6=17% for blue/green). All other proportions will be identical across occupational classes. However, when the healthy control group is not taken into account, the protective effect of the red occupation for dementia type 1 will create also an apparent “harmful” effect of the red occupation for dementia types 2 and 3 (i.e. 1/2=50%, compared to 1/3=33% for blue/green). In scenario B, the green occupational class shows a harmful effect on the development of dementia type 3, while no other differences between groups exist. Again, when the healthy control group is included, this only results in a lower percentage of healthy green individuals (1/6=17% versus 50% for red/blue) and a greater proportion of persons with green occupations in dementia type 3 (i.e. 50% versus 1/6=17% for red/blue). In the absence of a healthy control group, however, the green occupational class’ harmful effect for dementia type 3 leads to an apparent “protective” effect for dementia types 1 and 2 (1/5=20%, compared to 1/3=33% for red/blue).


2310 participants selected from the ADC cohort with dementia and occupational descriptions
140 participants excluded with occupational descriptions that were:
- non-informative
- indicative of unemployment
2170 participants eligible for assignment of BRC 2014 occupational codes
2 participants excluded with occupational descriptions that were not detailed enough to assign occupational codes
2168 participants eligible for statistical analysis
47 participants excluded because the size of their category was not sufficienctly large:
- 12 cases with ICT occupations
- 35 cases with rare forms of dementia
2121 participants included in the final sample


Transportation/Logistics
Service
Health Care/Welfare
Agricultural
Technical
Government/Law/Safety
Management
Business/Administrative
Commercial
Artistic/Linguistic
Pedagogical
0%
20%
40%
60%
80%
100%
Dementia type
AD
FTD
VaD
DLB
PSP/CBD


VRF score (mean)
3
2
1
0
AD
FTD
VaD
DLB
PSP/CBD
Pedagogical
Creative/Linguistic
Commercial
Business/Administrative
Management
Government/Law/Safety
Technical
Agricultural
Health Care/Welfare
Service
Logistics/Transportation
Male (%)
100
80
60
40
20
0


Cognitive profile
Memory
Attention/executive
Language
Visuospatial
Multi-domain
Multi-domain (n)
510
490
470
450
430
410
390
370
350
0.25
0.28
0.30
0.33
0.35
0.38
0.40
0.43
0.45
0.48
0.50
Threshold
Threshold: .255
Multi-domain: 383
Atypical variants correct: 34
292
130
140
126
383
Threshold: .442
Multi-domain: 465
Atypical variants correct: 39
256
106
125
119
465


Transportation/Logistics
Service
Health Care/Welfare
Agricultural
Technical
Government/Law/Safety
Management
Business/Administrative
Commercial
Artistic/Linguistic
Pedagogical
0%
20%
40%
60%
80%
100%
Cognitive profile
Memory
Attention/executive
Language
Visuospatial
Multi-domain


A)
Type 1
Type 2
Type 3
Healthy
Dementia
B)
Type 1
Type 2
Type 3
Healthy
Dementia