Prevalence of diabetic foot ulcer and its association with duration of illness and residence in Ethiopia: a systematic review and meta-analysis.

Authors

Henok Mulugeta, BSc, MSc^{1*} *Corresponding author

Lecturer of Nursing, Department of Nursing, College of Health Science, Debre Markos University

Address: P.O. Box 269, Debre Markos, Ethiopia

Email: mulugetahenok68@gmail.com

Fasil Wagnew, BSc, MSc¹

Lecturer of Nursing, Department of Nursing, College of Health Science, Debre Markos University

Address: P.O. Box 269, Debre Markos, Ethiopia

Email: fasilw.n@gmail.com

Haymanot Zeleke, BSc, MSc¹

Lecturer of Nursing, Department of Nursing, College of Health Science, Debre Markos University

Address: P.O. Box 269, Debre Markos, Ethiopia

Email: haymanotzeleke89@gmail.com

Bekele Tesfaye, BSc, MSc¹

Lecturer of Nursing, Department of Nursing, College of Health Science, Debre Markos University

Address: P.O. Box 269, Debre Markos, Ethiopia

Email: bekeletesfaye76@gmail.com

Haile Amha, BSc, MSc¹

Lecturer of Nursing, Department of Nursing, College of Health Science, Debre Markos University

Address: P.O. Box 269, Debre Markos, Ethiopia

Email: haileleul19@gmail.com

Cheru Tesema Leshargie, BSc, MPH²

Lecturer of Environmental Health, Department of Environmental Health, College of Health Science, Debre Markos University

Address: P.O. Box 269, Debre Markos, Ethiopia

Email: chertesema@gmail.com

Henok Biresaw, BSc, MSc³

Lecturer of Nursing, Department of Nursing, School of health science, College of Medicine and Health Science, Bahir Dar University

Address: P.O. Box 79, Bahir Dar, Ethiopia

Email: henok22@ymail.com

Getenet Dessie, BSc, MSc³

Lecturer of Nursing, Department of Nursing, School of health science, College of Medicine and Health Science, Bahir Dar University

Address: P.O. Box 79, Bahir Dar, Ethiopia

Email: ayalew.d16@gmail.com

Yihalem Abebe Belay, BSc, MPH⁴

Lecturer of Public Health, Department of Public Health, Debre Markos University

Address: P.O. Box 269, Debre Markos, Ethiopia

Email: yih2000ho@gmail.com

Tesfa Dejenie Habtewold, BSc, MSc⁵

Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands.

Email: tesfadej2003@gmail.com

Prevalence of diabetic foot ulcer and its association with duration of illness and residence in Ethiopia: a systematic review and meta-analysis.

Henok Mulugeta^{*1}, Fasil Wagnew¹, Haymanot Zeleke¹, Bekele Tesfaye¹, Haile Amha¹, Cheru Tesema Leshargie², Henok Biresaw³, Getenet Dessie³, Yihalem Abebe Belay⁴, Tesfa Dejenie Habtewold⁵

Abstract

Background: Diabetic foot ulcer (DFU), devastating complications of diabetes mellitus, is a major public health problem, and one of the leading reasons for hospital admission, amputations, and even death among diabetic patients in Ethiopia. Despite its catastrophic health consequences, the national burden of diabetic foot ulcer remains unknown in Ethiopia. Hence, the objective of this systematic review and meta-analysis was to estimate the national prevalence of diabetic foot ulcer and investigate the association with duration of illness and patient residence among diabetic patients.

Methods: We searched PubMed, Google Scholar, Cochrane Library, CINAHL, EMBASE, and PsycINFO databases for studies of diabetic foot ulcers prevalence that published from conception up to June 30, 2019. Quality of each article was assessed using a modified version of the Newcastle-Ottawa Scale for cross-sectional studies. All statistical analyses were done using STATA version 14 software for Windows, and meta-analysis was carried out using a random-effects method. The pooled national prevalence of diabetic foot ulcers was presented using a forest plot.

Results: A total of 10 studies with 3,029 diabetic patients were included. The pooled national prevalence of diabetic foot ulcers among Ethiopian diabetic patients was 11.27% (95% CI 7.22, 15.31%, $I^2=94.6$). Duration of illness (OR: 3.91, 95%CI 2.03, 7.52, $I^2=63.4\%$) and patients' residence (OR: 3.40, 95%CI 2.09, 5.54, $I^2=0.0\%$) were significantly associated with a diabetic foot ulcer.

Conclusion: In Ethiopia, at least one out of ten diabetic patients had diabetic foot ulcers. Healthcare policymakers (FMoH) need to improve the standard of diabetic care and should design effective preventive strategies to improve health care delivery for people with diabetes and reduce the risk of foot ulceration.

Keywords: *Diabetes mellitus, foot ulcers, prevalence, Ethiopia.*

Background

Diabetes is a chronic disorder characterized by hyperglycemia that occurs either due to insulin deficiency or resistance[1, 2] that has been increasing rapidly worldwide. According to the American Diabetes Association 2014 report, the global prevalence of diabetes among adults is 8.5%[2, 3]. In Africa, an estimated 12.1 million people were living with diabetes in 2010, and this figure is more likely to be 23.9 million by 2030[4].

Diabetes has several acute and chronic complications including diabetic foot ulceration. A diabetic foot, which is the formation of foot ulceration that manifested by neuropathy, ischemia, and infection, is one of the major public health problems affecting diabetic patients throughout the world[5, 6]. About 10% to 15% of diabetes patients are at risk of developing a foot ulcer during their lifetime[7]. The annual population-based incidence of diabetic foot ulceration ranges from 1.0% to 4.1%, and its prevalence among diabetic patients ranges from 4% to 10%[5, 8].

Diabetes foot problems are responsible for high morbidity and mortality of diabetes in Africa. In sub-Saharan Africa, the burden of a diabetic foot ulcer is increasing due to late diagnosis, poor awareness among patients, and poor access to diabetic care[9, 10]. The presence of foot ulcers in diabetes patient increases the risk of lower extremity amputations, worsens the physical, psychological, and social quality of life[11-13]. Furthermore, patients living with diabetic foot ulcers(DFUs) has a great economic burden and poor health-related quality of life[14].

Diabetic foot ulceration has been associated with various risk factors, such as longer duration of diabetes mellitus (DM), residence, lack of awareness, poor foot care practice, poor glycemic control, poor hygienic condition, foot trauma, peripheral neuropathy and presence of vascular

diseases[15-17] [8] [18] [19]. Interestingly, most of the risk factors can be prevented easily with health education, careful foot care, and glycemic control[20].

Despite the huge burden of diabetic foot ulceration and it is one of the challenging health concerns in Ethiopia, the overall prevalence of diabetic foot ulcer among diabetic patients at the national level remains unknown. Therefore, the objective of this systematic review and meta-analysis was to estimate the national prevalence of foot ulcers among diabetic patients and investigate the association with duration of illness and residence in Ethiopia. The findings of this study will be important to provide valuable information regarding the burden of the diabetic foot and to inform healthcare policymakers for areas of improvement in diabetic care in Ethiopia.

Methods

Search strategy and selection criteria

This systematic review and meta-analysis were conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guideline[21]. We searched articles reporting the prevalence of diabetic foot ulcers among diabetic patients from PubMed, Cochrane Library, Google Scholar, CINAHL, and EMBASE databases using the following search terms: “prevalence”, “diabetic”, “diabetic foot”, “ulcers”, “diabetic foot ulcers”, “patients”. Boolean operators like “AND” and “OR” were used to combine search terms. EndNote (version X8) software for Windows was used to search, download, organize, and cite the related articles. The reference lists of eligible articles were also searched to retrieve additional relevant articles. All published and unpublished studies that report the prevalence of diabetic foot ulcers among diabetic patients in Ethiopia were included. Studies, which investigated the association between duration of illness and patient residence with foot ulceration, were also included. Only these two factors such as diabetic duration and patient residence have been similarly assessed across the

included studies. Therefore, we estimate the pooled effects of these factors to determine their association with diabetic foot ulcers. The search for published studies was restricted to articles published in English language. Furthermore, due to limited literature, we reviewed all publications irrespective of publication year. Nevertheless, related articles which failed to report the prevalence of diabetic foot ulcers were excluded. The articles were searched from March 2019 to June 2019.

Data extraction

All the downloaded studies were independently screened for inclusion by two authors (HM and GD). From each included study, information on the name of the first author, year of publication, study area (region), health facility, study design, sample size, response rate, prevalence of diabetic foot ulcers and determinant factors (Diabetic duration and patient residence) were extracted using a pre-piloted template prepared in a Microsoft Excel spreadsheet. H.M. and H.A. conducted the primary data extraction and then B.T., F.W., and H.Z. examined the extracted data independently. Any disagreement and inconsistencies were resolved by discussion.

Quality assessment

We used the modified version of the Newcastle-Ottawa Scale for cross-sectional studies adapted from Modesti et al to assess the methodological and other quality of each relevant article. The key criteria in the Newcastle–Ottawa scale are representativeness of the sample, response rate, measurement tool used, comparability of the subject, appropriateness of the statistical test used to analyze the data[22]. Two authors (B.T, and H.B) independently assessed the quality of each article. Any disagreement was resolved through discussion and consensus.

Statistical Analysis

A random-effects model using the DerSimonian and Laird method[23] was employed to estimate the pooled national prevalence and OR of risk factors with a 95% confidence interval (CI). The result was presented using a forest plot. Heterogeneity across the studies was assessed by Cochrane's Q test and I^2 statistic. I^2 statistic ranges from 0 to 100 percent. We used the accepted I^2 statistic > 50% to define significant heterogeneity across the included studies[24, 25]. Moreover, we explored the source of between-study heterogeneity using meta-regression analysis. A funnel plot was used for visual assessment of publication bias. Asymmetry of the funnel plot indicates the presence of publication bias[26, 27]. Egger's tests were also used to assess the significance of publication bias with a p-value less than 0.05[28]. Finally, a sensitivity analysis was performed to assess whether the pooled prevalence estimates were influenced by individual studies. All data manipulation and statistical analyses were performed using Stata version 14.0 software for Windows.

Results

Study selection

The online search process initially yielded 1079 articles. Of which 21 articles duplicate records were identified and removed. After reviewing the title and abstract, we excluded 1016 irrelevant articles. From the remaining 42 articles, 32 articles were excluded since they failed to meet eligibility criteria and having poor quality based on the Newcastle-Ottawa Scale. Finally, a total of 10 relevant articles with 3,029 diabetic patients were included in the meta-analysis (Figure 1).

Study characteristics

The studies were conducted from 2000 to 2017 in different regions of the country. Of the 10 studies, three were conducted in Addis Ababa[29-31], three were in Amhara[32-34], two were in

164 Oromia[[35](#), [36](#)], and the other two[[37](#), [38](#)] were from other regions. All the included studies were
 165 cross-sectional by study design and their sample size ranged from 198 to 724 patients (Table1).
 166 Figure 1: PRISMA Flowchart diagram of the study selection

167 Table 1: Characteristics of studies included in the meta-analysis of foot ulcers among diabetic patients in Ethiopia

No	Author/s	Publication Year	Study area, Region	Health Facility Name	Study Design	Sample Size	Prevalence % (95%CI)	Risk factors	Funding
1	Yimam A.[31]	2017	Addis Ababa	Black Lion Hospital	Cross-sectional	198	26.0 (19.9,32.1)	-Residence -Duration of diabetes	Addis Ababa University
2	Ejigu A.[30]	2000	Addis Ababa	Menelik Hospital	Cross-sectional	283	1.70 (0.20,3.20)	-----	-----
3	Adem A. et al[29]	2011	Addis Ababa	Black Lion and St. Paul's Hospitals	Cross-sectional	724	9.7 (7.54,11.8)	-----	Addis Ababa University
4	Lebeta KR. et al[33]	2017	Bahir Dar, Amhara	Felege Hiwot Referral Hospital	Cross-sectional	344	21.2 (16.9,25.5)	-Duration of diabetes	Bahir Dar University
5	Abejew AA. et al[32]	2015	Dessie, Amhara	Dessie Referral Hospital	Cross-sectional	216	4.4 (1.7,7.1)	-----	Wollo University
6	Mariam TG. et al[34]	2017	Gondar, Amhara	University of Gondar Hospital	Cross-sectional	279	13.6 (9.6,17.6)	-Residence	University of Gondar
7	Worku D. et al[36]	2010	Jimma, Oromia	Jimma University Specialized Hospital	Cross-sectional	305	4.6 (2.3,6.9)	-----	Jimma University
8	Tilahun A. et al[35]	2017	Jimma, Oromia	Jimma University Specialized Hospital	Cross-sectional	236	8.5 (4.9,12.1)	-----	Jimma University
9	Gebrekiros K. et al[38]	2015	Mekelle, Tigray	Ayder Specialized Hospital	Cross-sectional	228	12.0 (7.8,16.2)	-----	Mekelle University
10	Deribe B. et al[37]	2014	Arba Minch, SNNP	Arba Minch General Hospital	Cross-sectional	216	14.8 (10.1,19.5)	-Residence -Duration of diabetes	Jimma University

Prevalence of diabetic foot ulcers

The pooled national prevalence of diabetic foot ulcers among diabetic patients was 11.27% (95% CI (7.22, 15.31%), $I^2=94.6$). The overall pooled effect size of foot ulceration in diabetes mellitus patients presented using forest plot (Figure2).

Figure 2: Forest plot showing the pooled prevalence of diabetic foot ulcers

Assessment of Heterogeneity

In this meta-analysis, the presence of heterogeneity was checked by I^2 statistics. The result of this meta-analysis using the random-effects model revealed a high heterogeneity across the included studies ($I^2 = 94.6\%$, $P=0.001$). The sources of variation were assessed using a meta-regression model using publication year and sample size as covariates. The result of the meta-regression analysis showed that both covariates were not statistically significant for the presence of heterogeneity (Table 2).

Table 2: Meta-regression analysis of factors with heterogeneity

Heterogeneity source	Coefficients	Std. Err.	P-value
Sample size	0.009	0.020	0.66
Publication Year	0.397	0.229	0.13

Assessment of publication bias

Publication bias was examined by using a funnel plot and Egger's test. Visual inspection of the funnel plot suggests asymmetry, as three studies lay on the left side and seven studies on the right side of the line (Figure 3). Besides, asymmetry of the funnel plot was statistically significant as evidenced by Egger's test ($P=0.001$), which indicated the presence of publication

bias. Furthermore, trim and fill analysis, which is a nonparametric method for estimating the number of missing studies[39], was performed to reduce and adjust publication bias(Figure 4).

Figure 3: Funnel plot to test the publication bias of the 10 studies

Figure 4: Result of trim and fill analysis after adjusting for publication bias

Association diabetic foot ulcers with diabetic duration and patient residence

Diabetic patients with duration of diabetes more than 10 years were 3.91 times more likely to develop diabetic foot ulcers as compared with patients with duration of diabetes less than 10 years (OR, 3.91, 95% CI 2.03, 7.52, $I^2=63.4\%$) (Figure 5). Likewise, those diabetic patients who lived in the rural area were 3.40 times more likely to develop diabetic foot ulcers than those who lived in the urban area [OR = 3.40; 95% CI: 2.09, 5.54, $I^2=0.0\%$] (Figure 6).

Figure 5: Forest plot showing the association of patient residence and diabetic foot ulcers

Figure 6: Forest plot showing the association of diabetic duration and diabetic foot ulcers

Discussion

The global prevalence of diabetic complications is rising rapidly[40]. Diabetic foot ulcer (DFU) is one of the major causes of morbidity, and it is responsible for 50% of diabetes-related hospital admissions in hospitals[41]. Moreover, a systematic literature review revealed that diabetes and related complications including foot ulceration are a common problem in Africa[42]. Early diagnosis and management of diabetic foot are critical to prevent its serious consequences such as infection, gangrene, and death[41].

In this study, the pooled national prevalence of diabetic foot among diabetic patients in Ethiopia was 11.27%. This finding was comparable with reports from other similar studies in Asia[43, 44] and Africa [45-47]. This might be due to similarities in the socio-demographic characteristics of

the study participants. On the other hand, our finding was higher than the global (i.e. 6.3%) and national or continental (7.2% in Africa) prevalence of diabetic foot ulcers[19]. Likewise, our finding is higher than other similar studies[48, 49]. One possible explanation for the high prevalence of diabetic foot in Ethiopian diabetic patients might be due to variations in risk factors such as poor diabetic-related knowledge and poor foot self-care practice. In spite of that, the current study finding was lower than the study finding in Nigeria and Netherlands where the prevalence of diabetic foot ulcers was 41.5% and 20.4% respectively[50, 51]. It could be due to a difference in sample size and study setting.

Even though several factors have been reported as a risk factor of DFU in different literature[15, 44, 52-54], we have identified only two similar factors (duration of diabetes and patient residence) across the included studies for analysis. In agreement with studies conducted in Egypt, Iran and India[43, 54, 55], our study revealed that the duration of diabetes and patients' residence had a significant association with a diabetic foot ulcer. Diabetic patients with long duration are exposed to multiple complications like neuropathy, which may increase the risk of foot ulceration due to trauma and high focal foot pressures[56]. Regarding residence, the possible reason might be since individuals in rural areas in Ethiopia are farmers and are often barefoot, which can result in damage to the foot and it leads to ulcers. Besides, most of the people in rural areas of the country are poor with low educational level so that they have no adequate knowledge and practice of foot care. However, the controversial result was reported from Sudan where diabetic patients from the urban area have been shown significantly associated with the development of DFU[57].

The findings of this meta-analysis have implications for clinical practice. Determining the prevalence of DFU provides up-to-date evidence to develop prevention, public health planning

and management strategy for diabetic foot ulcers. It also used to improve patients' quality of life and reduce the economic burden. Moreover, identifying determinant factors help clinicians to consider during their routine clinical practice. However, some limitations should be considered in future researches. It was difficult to analyze some important major factors since they were not examined consistently across the studies. Moreover, we identified significant heterogeneity across the included studies and the presence of publication bias so that the interpretation of the results has to be taken cautiously.

Conclusion

Available data suggest that the prevalence of active DFU in people with diabetes in Ethiopia was high, but it is all derived from selected populations. Duration diabetes and Patients' residence were significant factors for foot ulceration. Diabetes care in Ethiopia needs to develop a high standard of treatment guideline and preventive strategies to reduce the burden of DFU and prevent its risk factors. Health education should be given to patients with a longer duration of diabetes and those living in rural areas to reduce the progression of the disease and improve health outcomes. Moreover, there is a clear need for a large-scale prospective study to determine the extent of this important health care problem.

Abbreviations

CI: Confidence Interval, **FMoH:** Federal Ministry of Health, **OR:** Odds Ratio, **PRISMA:** Preferred Reporting Items for Systematic Reviews and Meta-Analyses, **SNNP:** Southern Nations, Nationalities, and Peoples, **DFUs:** Diabetic Foot Ulcers.

Declarations

Consent for publication

253 Not applicable.

254 **Availability of data and materials**

255 The data analyzed during the current systematic review and meta-analysis is available from the
256 corresponding author on reasonable request.

257 **Competing interests**

258 The authors declare that they have no competing interests.

259 **Funding**

260 Not applicable.

261 **Ethics approval and consent to participate**

262 Not applicable.

263 **Authors' contributions**

264 HM developed the protocol and involved in the design, selection of study, data extraction, and
265 statistical analysis and developing the initial drafts of the manuscript. GD, FW, HZ, BT, and HB
266 involved in data extraction, quality assessment, statistical analysis and revising subsequent
267 drafts. HM, FW and TDH prepared the final draft of the manuscript. All authors read and
268 approved the final draft of the manuscript.

269 **Acknowledgment**

270 we would like to thank all authors of studies included in this systematic review and meta-
271 analysis.

272 **References**

- 273 1. Lakshmi N, Patel N, Parmar P, Garasiya K, Chaudhary M: **Study the foot care practice among**
274 **diabetic patients in Ahmedabad city, Gujarat.** *International Journal of Medical Science and*
275 *Public Health* 2018, **7**(5):333-338.
- 276 2. Teshome HM, Ayalew GD, Shiferaw FW, Leshargie CT, Boneya DJ: **The Prevalence of Depression**
277 **among Diabetic Patients in Ethiopia: A Systematic Review and Meta-Analysis, 2018.**
278 *Depression Research and Treatment* 2018, **2018**.

3. Wild S, Roglic G, Green A, Sicree R, King H: **Global prevalence of diabetes: estimates for the year 2000 and projections for 2030.** *Diabetes care* 2004, **27**(5):1047-1053.
4. Hall V, Thomsen RW, Henriksen O, Lohse N: **Diabetes in Sub Saharan Africa 1999-2011: epidemiology and public health implications. A systematic review.** *BMC public health* 2011, **11**(1):564.
5. Alexiadou K, Doupis J: **Management of diabetic foot ulcers.** *Diabetes Therapy* 2012, **3**(1):4.
6. Pendsey SP: **Understanding diabetic foot.** *International journal of diabetes in developing countries* 2010, **30**(2):75.
7. Boulton AJ, Vileikyte L, Ragnarson-Tennvall G, Apelqvist J: **The global burden of diabetic foot disease.** *The Lancet* 2005, **366**(9498):1719-1724.
8. Singh N, Armstrong DG, Lipsky BA: **Preventing foot ulcers in patients with diabetes.** *Jama* 2005, **293**(2):217-228.
9. Agwu E, Dafiewhare EO, Ekanem PE: **Possible diabetic-foot complications in Sub-Saharan Africa.** In: *Global Perspective on Diabetic Foot Ulcerations.* edn.: InTech; 2011.
10. Levitt N: **Diabetes in Africa: epidemiology, management and healthcare challenges.** *Heart* 2008.
11. Jeffcoate WJ, Harding KG: **Diabetic foot ulcers.** *The lancet* 2003, **361**(9368):1545-1551.
12. Nabuurs-Franssen M, Huijberts M, Kruseman AN, Willems J, Schaper N: **Health-related quality of life of diabetic foot ulcer patients and their caregivers.** *Diabetologia* 2005, **48**(9):1906-1910.
13. Ribu L, Hanestad BR, Moum T, Birkeland K, Rustoen T: **A comparison of the health-related quality of life in patients with diabetic foot ulcers, with a diabetes group and a nondiabetes group from the general population.** *Quality of Life Research* 2007, **16**(2):179-189.
14. Kibachio J, Omolo J, Muriuki Z, Juma R, Karugu L, Ng'ang'a Z: **Risk factors for diabetic foot ulcers in type 2 diabetes: A case control study, Nyeri, Kenya.** *African Journal of Diabetes Medicine* 2013, **21**(1).
15. Desalu O, Salawu F, Jimoh A, Adekoya A, Busari O, Olokoba A: **Diabetic foot care: self reported knowledge and practice among patients attending three tertiary hospital in Nigeria.** *Ghana medical journal* 2011, **45**(2).
16. Lavery LA, Armstrong DG, Wunderlich RP, Mohler MJ, Wendel CS, Lipsky BA: **Risk factors for foot infections in individuals with diabetes.** *Diabetes care* 2006, **29**(6):1288-1293.
17. Lipsky BA, Berendt AR, Deery HG, Embil JM, Joseph WS, Karchmer AW, LeFrock JL, Lew DP, Mader JT, Norden C: **Diagnosis and treatment of diabetic foot infections.** *Clinical Infectious Diseases* 2004:885-910.
18. Wikblad K, Smide B, Bergström A, Kessi J, Mugusi F: **Outcome of clinical foot examination in relation to self-perceived health and glycaemic control in a group of urban Tanzanian diabetic patients.** *Diabetes research and clinical practice* 1997, **37**(3):185-192.
19. Zhang P, Lu J, Jing Y, Tang S, Zhu D, Bi Y: **Global epidemiology of diabetic foot ulceration: a systematic review and meta-analysis.** *Annals of medicine* 2017, **49**(2):106-116.
20. Wu SC, Driver VR, Wrobel JS, Armstrong DG: **Foot ulcers in the diabetic patient, prevention and treatment.** *Vascular health and risk management* 2007, **3**(1):65.
21. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D: **The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration.** *PLoS medicine* 2009, **6**(7):e1000100.
22. Modesti PA, Reboldi G, Cappuccio FP, Agyemang C, Remuzzi G, Rapi S, Perruolo E, Parati G: **Panethnic differences in blood pressure in Europe: a systematic review and meta-analysis.** *PloS one* 2016, **11**(1):e0147601.

23. McFarland LV: **Meta-analysis of probiotics for the prevention of antibiotic associated diarrhea and the treatment of Clostridium difficile disease.** *The American journal of gastroenterology* 2006, **101**(4):812.
24. Higgins JP, Thompson SG, Deeks JJ, Altman DG: **Measuring inconsistency in meta-analyses.** *BMJ: British Medical Journal* 2003, **327**(7414):557.
25. Islam MM, Iqbal U, Walther B, Atique S, Dubey NK, Nguyen P-A, Poly TN, Masud JHB, Li Y-CJ, Shabbir S-A: **Benzodiazepine use and risk of dementia in the elderly population: a systematic review and meta-analysis.** *Neuroepidemiology* 2016, **47**(3-4):181-191.
26. Ried K: **Interpreting and understanding meta-analysis graphs: a practical guide.** 2006.
27. Tang J-L, Liu JL: **Misleading funnel plot for detection of bias in meta-analysis.** *Journal of clinical epidemiology* 2000, **53**(5):477-484.
28. Egger M, Smith GD, Phillips AN: **Meta-analysis: principles and procedures.** *Bmj* 1997, **315**(7121):1533-1537.
29. Adem A, Demis T, Feleke Y: **Trend of diabetic admissions in Tikur Anbessa and St. Paul's University Teaching Hospitals from January 2005-December 2009, Addis Ababa, Ethiopia.** *Ethiopian medical journal* 2011, **49**(3):231-238.
30. Ejigu A: **Brief communication: Patterns of chronic complications of diabetic patients in Menelik II Hospital, Ethiopia.** *Ethiopian Journal of health development* 2000, **14**(1):113-116.
31. Yimam A: **Prevalence od Diabetic foot ulcer and associated factors among diabetic patient attending Tikur Anbesa Specialized Hospital Diabetic clinic, Addis Ababa, Ethiopia, 2017.** 2017.
32. Abejew AA, Belay AZ, Kerie MW: **Diabetic complications among adult diabetic patients of a tertiary hospital in Northeast Ethiopia.** *Advances in Public Health* 2015, **2015**.
33. Kidist Reba Lebeta, Zeleke Argaw, Birhane BW: **Prevalence of Diabetic Complications and Its Associated Factors Among Diabetes Mellitus Patients Attending Diabetes Mellitus Clinics; Institution Based Cross Sectional Study.** *American Journal of Health Research* 2017, **5**(2):38-43.
34. Mariam TG, Alemayehu A, Tesfaye E, Mequannt W, Temesgen K, Yetwale F, Limenih MA: **Prevalence of Diabetic Foot Ulcer and Associated Factors among Adult Diabetic Patients Who Attend the Diabetic Follow-Up Clinic at the University of Gondar Referral Hospital, North West Ethiopia, 2016: Institutional-Based Cross-Sectional Study.** *Journal of diabetes research* 2017, **2017**.
35. Tilahun A, Waktola C, Tewodros G, Sadik G, Amare D: **Major Microvascular Complications and Associated Risk Factors among Diabetic Outpatients in Southwest Ethiopia.** *Endocrinol Metab Syndr* 2017, **6**(272):2161-1017.10002.
36. Worku D, Hamza L, Woldemichael K: **Patterns of diabetic complications at jimma university specialized hospital, southwest ethiopia.** *Ethiopian journal of health sciences* 2010, **20**(1).
37. Deribe B, Woldemichael K, Nemera G: **Prevalence and factors influencing diabetic foot ulcer among diabetic patients attending Arbaminch Hospital, South Ethiopia.** *J Diabetes Metab* 2014, **5**(1):1-7.
38. Gebrekirstos K, Gebrekiros S, Fantahun A: **Prevalence and Factors Associated With Diabetic Foot Ulcer among Adult Patients in Ayder Referral Hospital Diabetic Clinic Mekelle, North Ethiopia, 2013.** *J Diabetes Metab* 2015, **6**(579):2.
39. Duval S, Tweedie R: **Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis.** *Biometrics* 2000, **56**(2):455-463.
40. Susan van D, Beulens JW, Yvonne T. van der S, Grobbee DE, Nealb B: **The global burden of diabetes and its complications: an emerging pandemic.** *European Journal of Cardiovascular Prevention & Rehabilitation* 2010, **17**(1_suppl):s3-s8.

41. Kasiya MM, Mang'anda GD, Heyes S, Kachapila R, Kaduya L, Chilamba J, Goodson P, Chalulu K, Allain TJ: **The challenge of diabetic foot care: Review of the literature and experience at Queen Elizabeth Central Hospital in Blantyre, Malawi.** *Malawi Medical Journal* 2017, **29**(2):218-223.
42. Bos M, Agyemang C: **Prevalence and complications of diabetes mellitus in Northern Africa, a systematic review.** *BMC public health* 2013, **13**(1):387.
43. Shahi SK, Kumar A, Kumar S, Singh SK, Gupta SK, Singh T: **Prevalence of diabetic foot ulcer and associated risk factors in diabetic patients from North India.** *The Journal of Diabetic Foot Complications* 2012, **4**(3):83-91.
44. Yusuf S, Okuwa M, Irwan M, Rassa S, Laitung B, Thalib A, Kasim S, Sanada H, Nakatani T, Sugama J: **Prevalence and risk factor of diabetic foot ulcers in a regional hospital, eastern Indonesia.** *Open Journal of Nursing* 2016, **6**(01):1.
45. Ndip EA, Tchakonte B, Mbanya J-C: **A study of the prevalence and risk factors of foot problems in a population of diabetic patients in cameroon.** *The international journal of lower extremity wounds* 2006, **5**(2):83-88.
46. Neuhaus H, Warter-Neuhaus C, Lyaruu I, Msuya L: **Diabetes care in Kilimanjaro region: clinical presentation and problems of patients of the diabetes clinic at the regional referral hospital—an inventory before structured intervention.** *Diabetic medicine* 2002, **19**(6):509-513.
47. Ogbera A, Fasanmade O, Ohwovoriole A, Adediran O: **An assessment of the disease burden of foot ulcers in patients with diabetes mellitus attending a teaching hospital in Lagos, Nigeria.** *The international journal of lower extremity wounds* 2006, **5**(4):244-249.
48. Nyamu P, Otieno C, Amayo E, McLigeyo S: **Risk factors and prevalence of diabetic foot ulcers at Kenyatta National Hospital, Nairobi.** *East African Medical Journal* 2003, **80**(1):36-43.
49. Walters D, Catling W, Mullee M, Hill R: **The distribution and severity of diabetic foot disease: a community study with comparison to a non-diabetic group.** *Diabetic Medicine* 1992, **9**(4):354-358.
50. Bouter K, Storm A, Uitslager R, Erkelens D, Diepersloot R: **The diabetic foot in Dutch hospitals: epidemiological features and clinical outcome.** *The European journal of medicine* 1993, **2**(4):215-218.
51. Ogbera A, Adedokun A, Fasanmade O, Ohwovoriole A, Ajani M: **The foot at risk in Nigerians with diabetes mellitus—the Nigerian scenario.** *International journal of Endocrinology and Metabolism* 2005, **2005**(4, Autumn):165-173.
52. Adler AI, Boyko EJ, Ahroni JH, Stensel V, Forsberg RC, Smith DG: **Risk factors for diabetic peripheral sensory neuropathy: results of the Seattle Prospective Diabetic Foot Study.** *Diabetes care* 1997, **20**(7):1162-1167.
53. Ahmad W, Khan IA, Ghaffar S, Al-Swailmi FK, Khan I: **Risk factors for diabetic foot ulcer.** *Journal of Ayub Medical College Abbottabad* 2013, **25**(1-2):16-18.
54. Yazdanpanah L, Shahbazian H, Nazari I, Arti HR, Ahmadi F, Mohammadianinejad SE, Cheraghian B, Latifi SM: **Prevalence and related risk factors of diabetic foot ulcer in Ahvaz, south west of Iran.** *Diabetes & Metabolic Syndrome: Clinical Research & Reviews* 2018.
55. El Din SA, Mekawy MM, Besely WN, Azer SZ: **Prevalence of Risk Factors for Egyptian Diabetic Foot Ulceration.** *Journal of Nursing and Health Science* 2016; **5** (2): 45, 57.
56. Frykberg RG, Zgonis T, Armstrong DG, Driver VR, Giurini JM, Kravitz SR, Landsman AS, Lavery LA, Moore JC, Schuberth JM: **Diabetic foot disorders: a clinical practice guideline (2006 revision).** *The journal of foot and ankle surgery* 2006, **45**(5):S1-S66.
57. Awadalla H, Noor SK, Elmadhoun WM, Almobarak AO, Elmak NE, Abdelaziz SI, Sulaiman AA, Ahmed MH: **Diabetes complications in Sudanese individuals with type 2 diabetes: Overlooked problems in sub-Saharan Africa?** *Diabetes & Metabolic Syndrome: Clinical Research & Reviews* 2017, **11**:S1047-S1051.

420

Figure 1: PRISMA Flowchart diagram of the study selection

Figure 2: Forest plot showing the pooled prevalence of diabetic foot ulcers

Figure 3: Funnel plot to test the publication bias of the 10 studies

Figure 4: Result of trim and fill analysis for adjusting publication bias of the 10 studies

Figure 5: Forest plot showing the association of residence and diabetic foot ulcer

Figure 6: Forest plot showing the association of diabetic foot ulcer with duration of the diseases