1 Original article

2 **Harmonizing Palliative Care: National Survey to Evaluate the Knowledge** 3 **and Attitude of Emergency Physicians towards Palliative Care**

4 Ameena Mohammed Al-Ansari¹, MBBCh, MRCP, Palliative Care Medicine Fellowship, Saleem
5 Nawaf Suroor, ²MBBCh, MRCP, Palliative Care Medicine Fellowship, Sobhi Mostafa AboSerea,³
6 MBBCh, MSc, Wafaa Mostafa Abd-El-Gawad,^{4*} MBBCh, MSc, MD

7 ¹ Palliative Care Center, Al-Sabah Health Area, Al-Shuwaikh, Kuwait. dr_alansari@hotmail.com

8 ² Palliative Care Center, Al-Sabah Health Area, Al-Shuwaikh, Kuwait. bonawaf77@gmail.com

9 ³ Palliative Care Center, Al-Sabah Health Area, Al-Shuwaikh, Kuwait. sobhymaster@yahoo.com

10 ^{4*} Geriatrics and Gerontology department, Faculty of Medicine, Ain Shams University, Al-
11 Abbasseya, Cairo, Egypt

12 E-mail: wafaager@hotmail.com, wafaager@med.asu.edu.eg

13 Telephone: +201010031624, +96599920836

14 Running title: Knowledge and Attitude of Emergency Physicians towards Palliative Care

15 **Funding:** This research received a grant from Kuwait Foundation for the Advancement of Sciences
16 (KFAS) under grant agreement no. P116-13NO-01.

17 **Authors' contributions:** Ameena Al-Ansari, Saleem Nawaf Suroor, Sobhi Mostafa AboSerea,
18 **Wafaa Mostafa AbdElGawad:** study concept, study design, acquisition of subjects and data,
19 interpretation of data, preparation of the manuscript. All authors had approved the final article.

20 **Competing interests:** The author(s) declared no potential conflicts of interest with respect to the
21 research, authorship and/or publication of this article.

22

23 **Background and Aim:** Although the challenges of integrating palliative care practices
24 across care settings are real and well recognized until now little is known about palliative
25 care practice of emergency physicians (EPs) and their accessibility to palliative care services
26 in Kuwait. So the aim of this study was to explore the attitude, and knowledge encountered
27 by EPs in providing palliative care in all general hospitals in Kuwait.

28 **Method:** A cross-sectional survey was performed in the emergency rooms of all general
29 hospitals in Kuwait using Palliative Care Attitude and Knowledge (PCAK) questionnaire.

30 **Results:** Of the total number of physicians working in emergency rooms (n=156), 104
31 (66.67%) had completed the survey. 76.9% (n=80) of the EPs had either uncertain attitude
32 toward palliative care. Most of the EPs (n=73, 70.28%) didn't discuss the need of the patients
33 to palliative care either with the patients or their families. Only 16 (15.4%) of the EPs
34 responded correctly to the most of the questions while nearly half of the EPs (n=51, 49%) had
35 poor knowledge especially in the most effective management of refractory dysnea (n=18,
36 17.3%). Experience $\geq$ 11yrs and better knowledge scores were independent predictors of
37 positive attitude after adjustment of age, sex, qualifications, specialty, position, and
38 nationality [OR: 5.747 (CI: 1.031-25.00), 1.458(CI: 1.148-1.851); p-value: 0.021, 0.002
39 respectively]..

40 **Conclusions:** Despite recognizing palliative care as an important competence, the majority of
41 the emergency physicians in Kuwait had uncertain attitude and poor knowledge towards
42 palliative care. Lack of knowledge, direct accessibility to palliative care services and lack of
43 support from palliative medicine specialists were the main reasons for uncertain and negative
44 attitude. Efforts should be done to enhance physician training and provide palliative care
45 resources in order to improve the quality of care given to patients visiting emergency
46 departments.

48 **Keywords:** Palliative care, Palliative Care Attitude and Knowledge (PCKA), Emergency
49 physicians, opioids, Kuwait.

What this paper adds

- Studies proved that the emergency room may be a suitable place for early referral of patients who may benefit from palliative care especially old age to prevent upcoming undesired admissions and hospital deaths.
- The integration of palliative care concepts and consultation teams into emergency medicine may help to avoid unnecessary and burdensome treatments, tests, and procedures that are not aligned with patients' goals of care.
- Although the challenges of integrating palliative care practices across care settings are real and well recognized until now little is known about palliative care practice of emergency physicians and their accessibility to palliative care services in Kuwait.
- Recently, a newly developed tool called Palliative Care Attitude and Knowledge (PCKA) questionnaire was created to assess the attitude and knowledge of non-palliative physicians toward palliative care. So the aim of this study was to explore the attitude, and knowledge encountered by emergency physicians in providing palliative care using PCKA⁸ in emergency departments in all general
- Studies showed that early palliative care consultation was shown to improve quality of life for cancer patients and may even lengthen their survival.

50

What this study adds

- Despite recognizing palliative care as an important competence, the majority of the emergency physicians in Kuwait had uncertain attitude and poor knowledge towards palliative care. Lack of knowledge, direct accessibility to palliative care services and lack of support from palliative medicine specialists were the main reasons for uncertain and negative attitude.
- Efforts should be done to enhance physician training and provide palliative care resources in order to improve the quality of care given to patients visiting emergency departments.

51

52

53

54

55

56

57

58 **Introduction:**

59 Emergency medicine is a medical specialty that is concerned with stabilization of
60 patients with acute illness or injury for definitive care.¹ in this specialty, emergency
61 physicians are trained to provide acute treatments for emergency medical conditions aiming
62 at preserving patients' life regardless of their wellbeing.^{1,2}

63 While patients and their families under palliative care can experience stressful and
64 overwhelming moments during the disease trajectory such as sudden respiratory distress,
65 severe pain, vomiting, confusion and many other symptom,³ the first point of access is often
66 the emergency room. Unfortunately, many of them die in the hospitals despite their wish to
67 die at home.⁴ Hospitals; where palliative medicine specialists are available; can provide the
68 opportunity for the patients and their families to establish their preferences for care and to
69 coordinate the aggressiveness of treatment, symptom management, and place of death.^{3,4}

70 Moreover, early palliative care consultation was shown to improve quality of life for
71 cancer patients and may even lengthen their survival.⁵ Most recently, studies proved that the
72 emergency room may be a suitable place for early referral of patients who may benefit from
73 palliative care ⁶ especially old age to prevent upcoming undesired admissions and hospital
74 deaths.⁷

75 The integration of palliative care concepts and consultation teams into emergency
76 medicine may help to avoid unnecessary and burdensome treatments, tests, and procedures
77 that are not aligned with patients' goals of care. ² Although the challenges of integrating
78 palliative care practices across care settings are real and well recognized until now little is known
79 about palliative care practice of emergency physicians and their accessibility to palliative care
80 services in Kuwait. Recently, a newly developed tool called Palliative Care Attitude and Knowledge
81 (PCAK) questionnaire was created to assess the attitude and knowledge of non-palliative physicians

toward palliative care.⁸ So the aim of this study was to explore the attitude, and knowledge encountered by emergency physicians in providing palliative care using PCAK⁸ in emergency departments in all general hospitals in Kuwait.

Method:

Study design and setting:

A cross-sectional survey was performed in the emergency rooms of all general hospitals under the Ministry of Health of Kuwait. This includes six general hospitals; Al-Sabah, Al-Jahra, Al-Amiri, Al-Adan, Al-Farawanyia and Mubarak Al-Kabeer hospitals.

Measurements and intervention:

Palliative Care Attitude and Knowledge Questionnaire (PCAK):⁸

It is a newly developed questionnaire composed of three sections. Section one includes demographic data such as age, sex, level of education, work experience, workplace, medical subspecialty and palliative care experience. Section two is assessing the attitude and formed of 11 items. The tool had a 5 point Likert scale ranging from strongly disagree (1) to strongly agree (5). Negative or unfavorable attitude was considered if the participant scored ≤ 25 , uncertain attitude if scored > 25 but < 41 , positive or favorable attitude if scored ≥ 41 . The third section inquiries about knowledge. It includes 2 parts; the first part was about the self-reported knowledge (3 questions) and the second part is 12 clinical questions. Regarding self-knowledge, 5 points likert scale was used ranging from excellent response (5) to none (1). Regarding basic knowledge scoring; Poor knowledge was calculated if participant scored less than 50% of the total score (12 points) (≤ 5 points), fair knowledge if $\geq 50\%$ to $< 75\%$ (6-9 points), good knowledge if scored $> 75\%$ (≥ 10 points).⁸

Selection of Participants:

The questionnaires were distributed to all emergency physicians in all general hospital in Kuwait (Figure 1) and recollected back by hand to hand and revised within 48 hours. The flowchart of the sampling procedure was presented in Figure 1.

Ethical statement:

The approval of the Institutional Review Board (IRB) of the Ministry of Health was taken prior to the study (No.210/2016, March 2016). Informed consent was obtained from all participants. Participants' anonymity and confidentiality were kept throughout all phases of the study. Approvals were also obtained from participating emergency departments of all general hospitals.

Statistical analysis:

All data manipulation and analysis were performed using the SPSS (Statistical Package for Social Science) SPSS version 20. P-values less than 0.05 were regarded as a sign of statistical significance. Independent sample t-test was used for comparison between quantitative data. Chi-square test or Fisher's Exact when appropriate were used to compare between qualitative variables. Generalized linear method was used to find out the independent predictors of positive attitude toward palliative care with adjustment of any confounding factors. We used the consolidated criteria for reporting observational studies (RECORD) checklist as far as this was applicable to our study.

Results:

Characteristics of study subjects:

Of the total number of physicians working in emergency rooms in Kuwait (n=156 physicians), 19 physicians were on annual leave and 11 physicians refused to participate. The remaining 126 emergency physicians who responded to the questionnaire, 104 (82.54%) had completed the survey in all general hospitals. (Figure1). The number of participants was 15 (14.2%) from AlSabah hospital, 20 (19.2%) from Al-Amiri hospital, 18 (20%) from Mubarak Al-Kabeer hospital, 20 (17.33%) from Al-Farawanyia hospital, 14 (13.5%) from Al-Adan hospital and 17 (20.44%) from Al-Jahrah hospital. (Figure 1).

86.6% (n=90) were males and 13.5% (n=14) were females and mean age of the respondents was 38.83 ± 9.5 years old. The median years of experience were 11 (IQR: 7-13) years. Due to the diversity of population demographics, only 20.19% (n=21) were Kuwaiti physicians. Most of respondents had postgraduate emergency medicine studies (n=68; 65.4%), followed by internal medicine studies (12; 11.5%). (Table 1).

Main results:

Only 18.3% (n=19) of the emergency physicians had a favorable attitude towards palliative care. Most of them had uncertain attitude (76.9%, n=80) and 4.8% (n=5) had an unfavorable attitude. Most of them agreed about the need of all adults and children who are terminally ill for palliative care services, not just those with cancer (Q7: n =65; 62.5%) and approved also that palliative care benefits include enhanced quality of life for the patient and family (Q9:n=81; 77.9%), skilled care for terminally ill patients (Q10: n=82; 78.8%), and expert pain and symptom management (Q11: n=85; 81.7%). Unfortunately, they emphasized on the lack of timely communication between palliative care providers and themselves (Q2: n=71; 68.3%). Many of them dissatisfied with palliative care services in the past (Q1: n=61; 58.7%), reported unfamiliarity with the currently available palliative care services (Q3: n=65; 62.5%), and its types (Q5:n=54; 51.9%). Moreover, the uncertainty about its length of

coverage was 53.8% (Q4: n=56). Generally, the emergency physicians with experience ≥ 11 years had statistically significant better attitude scores than the emergency physicians with experience < 11 (p value=0.019, Figure 2) but in most of the individualized items of the attitude, no statistically significant between them.

Most of the emergency physicians (n=73, 70.28%) didn't discuss the need of the patients to palliative care either with the patients or their families. 44 (42.3%) of them reported no or little knowledge in pain assessment and management, while 58 (55.8%) and 41 (39.4%) reported good knowledge in other symptoms assessment and management, and the counseling respectively. Neither years of experience nor nationality had any statically significant effect on self-assessment of their knowledge and experience of the emergency physicians towards palliative care. (Table 1).

The overall percentage of the emergency physicians that responded correctly to the most of the questions about the basic knowledge of palliative care was only 16 (15.4%) while 37 (35.6%) had a fair knowledge and unfortunately many of them (n=51; 49%) responded to $> 50\%$ of the questions incorrectly. The median number of total corrected answers was 6 (IQR: 4-8) out of 12. Surprisingly, none of the emergency physicians answered all the questions correctly. An overview of the questions and their answers was presented in Table 3. Q11 for the symptoms of cord compression was the one with the highest percentage of correct answers (Q11: n=94; 90.4%) while Q8 about the most effective treatment for the refractory dyspnea question had the lowest rate of correct answers (Q8:n=18;17.3%). They had mostly poor to fair knowledge regarding opioids handling; such as types of opioids (Q3:n=57;54.8%), World Health Organization ladder for pain management (Q7: n=39;37.5%), opioids toxicity (Q8:n=31; 29.8%), use of oral opioids (Q15:n=19; 18.3%); and

delirium (Q4:n=62; 59.65%) and management of catastrophic bleeding in palliative setting (Q10:n=20; 19.2%).

Most of the emergency physicians had good knowledge scores regarding the differences between traditional and palliative care (Q1: n=79; 76%) and the multidisciplinary team role in palliative care (Q2:n=83; 79.8%). No statistically significant differences between years of experience of the emergency physicians and their knowledge scores (Figure 3) in most of the questions except palliative care definitions and signs of opioid toxicity in which the emergency physicians with less experience answered significantly better (86% vs 66.7%, 40% vs 20.4%; p-value: 0.024, 0.029 respectively) while in diagnosis of superior venous obstruction, the emergency physicians with more years of experience answered better (12% vs 35.2%; p-value: 0.011). No statistically significant differences between the emergency physicians from different hospitals in the attitude or knowledge scores or demographic characteristics. (Table 4).

By using generalized linear method to find out the independent predictors of positive attitude toward palliative care, we found that years of experience ≥ 11 yrs and better knowledge scores were independent predictors of positive attitude after adjustment of age, sex, qualifications, specialty, position, and nationality [OR: 5.747 (CI: 1.031-25.00), 1.458(CI: 1.148-1.851); p-value: 0.021, 0.002 respectively]. (Table 5)

Discussion

To the best of our knowledge, this is the first survey of its kind conducted in Kuwait to evaluate palliative care attitude and knowledge targeting the emergency physicians anywhere in the entire region. We used PCAK which is a newly developed validated and reliable tool⁸ to assess the attitude and knowledge among physicians. There are more than 150 physicians working in the emergency rooms in 6 hospitals distributed all over Kuwait.

Most of them had uncertain attitude and poor knowledge scores. Better knowledge scores and experience ≥ 11 yrs were positive predictors of favorable attitude. Although more than 10 different nationalities are existing in Kuwait like Egyptians, Syrians, Pakistanis, Indians, and others from different cultures, emergency rooms demographic, medical education, specialties, years of experience but no statically significant differences between them in their knowledge or attitude toward palliative care. This differed from many studies which reported that clinician attitudes or knowledge were affected by nationality, physicians' position, staff type, years of experience, emergency rooms demographic and hospital type.⁹

In our study, 76.9% of the emergency physicians had either uncertain attitude toward palliative care while only 18.3% had a favorable or positive attitude. This was mainly due to either dissatisfaction about PC services, types, accessibility, length of coverage and lastly lack of timely communication between them and palliative care providers as reported in our study. These results is similar to the original results of PCAK questionnaire.⁸ Many factors affecting the attitude of the emergency physicians depending on their diverse experiences, which ranged from positive, encouraging to a negative and distressing expiernce.^{9,10} This includes sometimes a sense of incompetence to alter the course of the disease^{3, 11} or feeling anxious and uncomfortable about discussing death and dying with the terminally ill patient.^{12,13} This negative attitude can hinder the quality of care provision especially at the end of life.

Most of the emergency physicians (n=73, 70.28%) didn't discuss the need to palliative care either with the patients or their families. The willingness to engage in end-of-life discussions becomes key to improve palliative care in the emergency department.^{14, 15} However, palliative care referrals remain largely underutilized in all areas of healthcare and are characteristically initiated late in the disease trajectory.^{16,17} Furthermore, most of the

emergency physicians feel that it is not their obligation to discuss the need for palliative care and they feel that the family physicians or the oncologists are the one who is responsible about this discussion.^{3,18}

Nearly half of the emergency physicians (49%) had poor knowledge. Although many of them subjectively reported good knowledge in counseling and other symptoms (rather than pain) assessment and management and they strongly agreed that palliative care is an important competence for emergency physicians^{3,18} but it wasn't necessarily reflected in their effectiveness of care especially in view of their poor knowledge at most. These results is similar to the original results of PCAK questionnaire.⁸

They had mostly poor knowledge in opioids handling, delirium and management of catastrophic bleeding in the palliative setting. This is similar to many studies that reported a lack of knowledge in palliative care, especially pain management proficiency.¹⁸⁻²⁰

The setting of emergency rooms; as crowded spaces, noisy environment, compromised privacy, frequent interruptions, time constraints, illness complexity, and medico-legal threats;^{21,22} make the initiation of palliative care discussions less than ideal in these settings.^{3,22} Unfortunately, being not confident with the diagnosis and concerns of diminishing patients' or families' hope, or believing that patients are not prepared to hear forthcoming information make physicians often hesitate to discuss patients prognosis either with the patients or their families.²² Subsequently, aggressive interventions often initiated may ultimately be misaligned with overall goals of care although retrospectively viewed as futile, harmful, and painful.²³ In reality, the culture of emergency setting that provides stabilization of acute medical emergencies was sometimes at odds with the culture of palliative care, which balances quality of life with the burdens of invasive treatments.^{2,18}

In our study, good knowledge was positively associated with the positive attitude of the emergency physicians toward palliative care. This was agreed with Thulesius et al. who revealed that good knowledge of staff significantly improved attitudes toward the end of life care and the authors suggested initiatives to educate physicians working in ED's will help them to change their negative attitudes and contribute better quality of patient care.²⁴

Many studies reported a significant correlation between the level of knowledge and attitudes towards palliative care. This is highlighting that as participants' level of knowledge increased, attitudes become more positive in hospitals such as in Lebanon,²⁵ Ethiopia,²⁶ Saudi Arabia,²⁷ and India.²⁸ As a part of human nature, the degree and complexity of their knowledge affect their attitudes and in turn their behavior.²⁹

The effective measurement of knowledge and attitude of the physicians is an important component of the evaluation of education and practice especially if used prior to any educational activity or program. This would result in more willingness to contribute in discussion and generate greater receptivity to the educational materials.^{30,31} Again it can be applied post activity to document the change in their knowledge and attitude and at the same time to evaluate the efficiency of the training and educational program.

Strength and limitation of the study:

It is the first study to assess the knowledge and attitude of the emergency physicians toward palliative care in the entire region. All the emergency physicians in Kuwait were involved in the research with no selection bias. The scarcity of similar studies carried out in Kuwait, Eastern Mediterranean region and in other parts of the world made the comparison and discussion difficult. Palliative care is recently introduced in Kuwait and it is still in the phase of development with the shortage of the number of the available palliative medicine specialists, this may explain the uncertain or the negative attitude towards palliative care.

Despite recognizing palliative care as an important competence, the majority of the emergency physicians in Kuwait had uncertain attitude and poor knowledge towards palliative care. Lack of knowledge, direct accessibility to palliative care services and lack of support from palliative medicine specialists were the main reasons for uncertain and negative attitude.

Conclusion:

Despite recognizing palliative care as an important competence, the majority of the emergency physicians in Kuwait had uncertain attitude and poor knowledge towards palliative care. Lack of knowledge, direct accessibility to palliative care services and lack of support from palliative medicine specialists were the main reasons for uncertain and negative attitude. Efforts should be done to enhance physician training and provide palliative care resources in order to improve the quality of care given to patients visiting emergency departments. Attention should be given to palliative care by the national health policy. There is an urgent need to be incorporated into the national curriculum of medical students and the emergency physicians' education. Further efforts are needed to push the palliative care across different healthcare sectors.

Ethics approval and consent to participate: The research project has been approved by the Institutional Review Board (IRB) of the Ministry of Health, Kuwait (No.210/2016, March 2016) within which the work was undertaken and that it conforms to the provisions of the Declaration of Helsinki. All subjects gave informed consent and their anonymity was preserved.

Consent for publication: not applicable.

Availability of data and materials: The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests: The author(s) declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

Funding: This research received a grant from Kuwait Foundation for the Advancement of Sciences (KFAS) under grant agreement no. P116-13NO-01.

Authors' contributions: Aameena Al-Ansari, Saleem Nawaf Suroor, Sobhi Mostafa AboSerea, Wafaa Mostafa AbdelGawad: study concept, study design, acquisition of subjects and data, interpretation of data, preparation of the manuscript. All authors had approved the final article.

Acknowledgements

The authors would like to thank all the participants for their valuable contributions to this study.

References:

- 1- Schneider SM1, Hamilton GC, Moyer P, Stapczynski JS. Definition of emergency medicine. Acad Emerg Med 1998;5(4):348-51.
- 2- Grudzen CR1, Richardson LD, Hopper SS, Ortiz JM, Whang C, Morrison RS. Does palliative care have a future in the emergency department? Discussions with attending emergency physicians. J Pain Symptom Manag. 2012;43(1):1-9.
- 3- Rivera MR, Torres FS. Physician Attitudes on the Provision of Palliative Care in Puerto Rican Emergency Departments. J Palliat Care Med 2015; 5: 201.
- 4- Pritchard RS, Fisher ES, Teno JM, Sharp SM, Reding DJ, et al. Influence of patient preferences and local health system characteristics on the place of death. SUPPORT Investigators. Study to understand prognoses and preferences for risks and outcomes of treatment. Journal of the American Geriatric Society 1998; 46: 1242-1250.
- 5- Temel JS, Greer JA, Muzikansky A, et al. Early palliative care for patients with metastatic non-small-cell lung cancer. N Engl J Med 2010; 363:733–742. [PubMed: 20818875]
- 6- Lipinski M, Eagles D, Fischer LM, Mielniczuk L, Stiell IG. Heart failure and palliative care in the emergency department. Emerg Med J 2018; 35(12):726-729. doi: 10.1136/emermed-2017-207186.

- 7- Beynon T, Gomes B, Murtagh FEM, Glucksman Ed, Parfitt A, Burman R, et al. How common are palliative care needs among older people who die in the emergency department? *Emerg Med J* 2011;28(6):491-5. doi: 10.1136/emj.2009.090019.
- 8- Al-Ansari AM, Suroor SN, AboSerea SM, Abd-El-Gawad WM. Development of palliative care attitude and knowledge (PCAK) questionnaire for physicians in Kuwait. *BMC Palliat Care*. 2019;18(1):49. doi:10.1186/s12904-019-0430-9. PubMed PMID: 31170968; PubMed Central PMCID: PMC6555752.
- 9- Cheung KY, Chan KC. Experiences of healthcare professionals in providing palliative end-of-life care to patients in emergency departments: a systematic review protocol. *JBIC Database System Rev Implement Rep* 2016;14(10):9-14.
- 10- Jackson V, Mack J, Matsuyama R, Lakoma M, Sullivan A, Arnold R, et al. A qualitative study of oncologists' approaches to end-of-life care. *J Palliat Med* 2008; 11(6):893–906.
- 11- Peters L, Cant R, Payne S, O'Connor M, McDermott F, Hood K, et al. How death anxiety impacts nurses' caring for patients at the end of life: a review of literature. *Open Nurs J* 2013;7:14–21.
- 12- Tait V, Higgs M, Magann L, Dixon J, Davis J, Fernandez R. Attitudes of nonpalliative care nurses towards palliative care. *Intl J Palliat Care* 2015;2015:1–6.
- 13- Hui D, Bansal S, Park M, Reddy A, Cortes J, Fossella F, et al. Differences in attitudes and beliefs toward end-of-life care between hematologic and solid tumor oncology specialists. *Ann Oncol* 2015;26(7): 1440–6.
- 14- Shearer FM, Rogers IR, Monterosso L, Ross-Adjie G, Rogers JR. Understanding emergency department staff needs and perceptions in the provision of palliative care. *Emerg Med Australas* 2014;26(3):249–55.
- 15- Heaston S, Beckstrand RL, Bond AE, Palmer SP. Emergency nurses' perceptions of obstacles and supportive behaviors in end-of-life care. *J Emerg Nurs* 2006;32(6):477–85.
- 16- Kistler EA, Sean Morrison R, Richardson LD, Ortiz JM, Grudzen CR. Emergency department-triggered palliative care in advanced cancer: proof of concept. *Acad Emerg Med* 2015;22(2):237–9.
- 17- Zimmermann C, Swami N, Krzyzanowska M, Hannon B, Leighl N, Oza A, et al. Early palliative care for patients with advanced cancer: a cluster-randomised controlled trial. *Lancet* 2014;383(9930):1721–30.
- 18- Stone SC, [Mohanty S](#), [Grudzen CR](#), [Shoenberger J](#), Asch S, [Kubrick K](#), Lorenz KA **Emergency medicine physicians' perspectives of providing palliative care in an emergency department.** *J Palliat Med*. 2011; 14(12):1333-8. doi: 10.1089/jpm.2011.0106.
- 19- Goldonowicz JM, Runyon MS, Bullard MJ. Palliative care in the emergency department: an educational investigation and intervention. *BMC Palliative Care* 2018; 17:43. <https://doi.org/10.1186/s12904-018-0293-5>
- 20- Chan GK. End-of-life models and emergency department care. *Acad Emerg Med* 2004;11(1):79–86.
- 21- Jesus J, Geiderman J, Venkat A, et al. Physician orders for life-sustaining treatment and emergency medicine: ethical considerations, legal issues, and emerging trends. *Ann Emerg Med*. 2014;64(2):140–4.
- 22- Limehouse WE, Feeser VR, Bookman KJ, Derse A. A model for emergency department end-of-life communications after acute devastating events—part II: moving from resuscitative to end-of-life or palliative treatment. *Acad Emerg Med* 2012;19(11):1300–8.

- 23- 7. Forero R, McDonnell G, Gallego B, et al. A Literature Review on Care at the End-of-Life in the Emergency Department. *Emergency Medicine International*. 2012;2012:486516.
- 24- Thulesius H, Petersson C, Petersson K, Håkansson A. Learnercentred education in end-of-life care improved well being in home care staff: a prospective controlled study. *Palliat Med* 2002;16: 347-354.
- 25- Huijjer AH, Dimassi H. Palliative care in Lebanon: knowledge, attitudes and practices of physicians and nurses. *J Med Liban* 2007;55(3):121-8.
- 26- Kassa H, Murugan R, Zewdu F, Hailu M, Woldeyohannes D. Assessment of knowledge, attitude and practice and associated factors towards palliative care among nurses working in selected hospitals, Addis Ababa, Ethiopia. *BMC Palliat Care* 2014;4:13(1):6.
- 27- Abudari G, Zahreddine H, Hazeim H, Assi MA, Emara S. Knowledge of and attitudes towards palliative care among multinational nurses in Saudi Arabia. *Int J Palliat Nurs* 2014;20(9):435-41.
- 28- Das A, Haseena T. Knowledge and Attitude of Staff Nurses Regarding Palliative Care. *International Journal of Science and Research* 2015; 4 (11):1790-5.
- 29- Fabrigar LR, Petty RE, Smith SM, Crites SL Jr. Understanding knowledge effects on attitude-behavior consistency: the role of relevance, complexity, and amount of knowledge. *J Pers Soc Psychol* 2006;90(4):556-77.
- 30- Kawaguchi S, Mirza R, Nissim R, & Ridley J. Internal Medicine Residents' Beliefs, Attitudes, and Experiences Relating to Palliative Care: A Qualitative Study. *American Journal of Hospice and Palliative Medicine* 2017; 34(4), 366–372.
- 31- von Gunten C F, Twaddle M, Preodor M, Neely K J, Martinez J, & Lyons J. Evidence of improved knowledge and skills after an elective rotation in a hospice and palliative care program for internal medicine residents. *Am J Hosp Palliat Care* 2005;22(3):195-203.

Table 1: General description of the emergency physicians in Kuwait

		Total (N=104)	experience <11 yrs(n=50)	experience ≥11 yrs(n=54)	p- value*
Age		38.83(±9.5)	32.12(3.88)	45.03(8.87)	<0.001
Sex	Males	90(86.6%)	39(78%)	51(94.4%)	0.014
	females	14(13.5%)	11(22%)	3(5.6%)	
Nationality	Kuwaiti	21(20.2%)	18(38%)	3(5.6%)	<0.001
	Non-Kuwaiti	83(79.8%)	32(64%)	51(94.4%)	
Qualification	MBBS	30(28.8%)	21(42%)	9(16.7%)	0.202**
	Master	59 (56.7%)	24(48%)	35(64.8%)	
	MD, MRCP/MRCS	15(14.5%)	5(10%)	10(18.6%)	
Specialty	ER	68(65.4%)	32(64%)	36(66.7%)	0.007**
	Internal Medicine	12(11.5%)	2(4%)	10(18.5%)	
	Surgery	10(9.6%)	4(8%)	6(11.1%)	
	Family Medicine	3(2.9%)	3(6%)	0	
	Others	11(10.6%)	9(18%)	2(3.7%)	
Position	Assistant registrar	6(5.8%)	6(12%)	0	0.020**
	Registrar	86(82.7%)	40(80%)	46(85.2%)	
	Senior registrar	8(7.7%)	4(8%)	4(7.4%)	
	Specialist/ Consultant	4(3.8%)	0	4(7.4%)	
Discussion about palliative care	No patients	73(70.2%)	34(68%)	39(72.2%)	0.726**
	1 to 5 patients	23(22.1%)	11(22%)	12(22.2%)	
	6 to 10 patients	6(5.8%)	4(8%)	2(3.7%)	
	11 to 15 patients	1(1%)	1(2%)	0	
	> 15 patients, families	1(1%)	0	1(1.9%)	
Self-assessment of his knowledge in:					
1-Pain	Excellent	3(2.9%)	1(2%)	2(3.7%)	0.311**
	very good	15(14.4%)	8(16%)	7(13%)	
	good	42(40.4%)	21(42%)	21(38.9%)	
	weak	29(27.9%)	10(20%)	19(35.2%)	
	none	15(14.4%)	10(20%)	5(9.3%)	
2-Other Symptoms	Excellent	4(3.8%)	2(4%)	2(3.7%)	0.683**
	very good	21(20.2%)	10(20%)	11(20.4%)	
	good	58(55.8%)	28(56%)	30(55.6%)	
	weak	14(13.5%)	5(10%)	9(16.7%)	
	none	7(6.7%)	5(10%)	2(3.7%)	
3-Counseling	Excellent	6(5.8%)	2(4%)	4(7.4%)	0.402**
	very good	29(27.9%)	15(30%)	14(25.6%)	
	good	41(39.4%)	20(40%)	21(38.9%)	
	weak	18(17.3%)	6(12%)	12(22.2%)	
	none	10(9.6%)	7(14%)	3(5.6%)	

*p- value < 0.05 is significant.

** Fisher Exact

Table 2: Distributions of the emergency physicians' attitude towards palliative care in Kuwait

		Total (n=104)	experience <11yrs(n=50)	experience ≥11yrs(n=54)	p- value*
Q1-I have been dissatisfied with PC services in the past	Positive attitude	16(15.4%)	5(10%)	11(20.4%)	0.232
	Negative attitude	61(58.7%)	29(58%)	32(59.3%)	
	Not sure	27(26%)	16(32%)	11(20.4%)	
Q2-I feel there is a lack of timely communication between PC providers and myself.	Positive attitude	10(9.6%)	5(10%)	5(9.35)	0.992
	Negative attitude	71(68.3%)	34(68%)	37(68.5%)	
	Not sure	22(21.2%)	11(22%)	12(22.2%)	
Q3- I am not familiar with PC services in this community.	Positive attitude	17(16.3%)	6(12%)	11(20.4%)	0.358
	Negative attitude	65(62.5%)	31(62%)	34(63%)	
	Not sure	22(21.2%)	13(26%)	9(16.7%)	
Q4-I am uncertain of the length of coverage under the PC benefit.	Positive attitude	13(12.5%)	4(8%)	9(16.7%)	0.006
	Negative attitude	56(53.8%)	35(70%)	21(38.9%)	
	Not sure	35(33.7%)	11(22%)	24(44.4%)	
Q5- I am uncertain of the type of services covered under the PC benefit.	Positive attitude	13(12.5%)	7(14%)	6(11.1%)	0.925
	Negative attitude	54(51.9%)	26(52%)	28(51.9%)	
	Not sure	37(35.6%)	17(34%)	20(37%)	
Q6- Patients or families are unwilling or unready to elect PC services.	Positive attitude	29(27.9%)	15(30%)	14(25.9%)	0.910
	Negative attitude	33(31.7%)	15(30%)	18(33.3%)	
	Not sure	42(40.4%)	20(40%)	22(40.7%)	
Q7- All adults and children who are terminally ill are candidates for PC services, not just those with cancer.	Positive attitude	65(62.5%)	29(58%)	36(66.7%)	0.561
	Negative attitude	11(10.6%)	5(10%)	6(11.1%)	
	Not sure	28(26.9%)	16(32%)	12(22.2%)	
Q8-Patients receiving radiation for palliation of symptoms are not candidates for PC.	Positive attitude	49(47.1%)	10(20%)	13(24.1%)	0.691
	Negative attitude	23(22.1%)	26(52%)	23(42.6%)	
	Not sure	32(30.8%)	12(28%)	18(33.3%)	
Q9- PC benefits include enhanced quality of life for the patient and family.	Positive attitude	81(77.9%)	39(78%)	42(77.8%)	0.763**
	Negative attitude	6(5.8%)	2(4%)	4(7.4%)	
	Not sure	17(16.3%)	9(18%)	8(14.8%)	
Q10-PC benefits include skilled care for terminally ill patients.	Positive attitude	82(78.8%)	42(84%)	40(74.1%)	0.091**
	Negative attitude	5(4.8%)	0(0%)	5(9.35)	
	Not sure	17(16.3%)	8(16%)	9(16.7%)	
Q11-PC benefits include expert pain and symptom management.	Positive attitude	85(81.7%)	40(80%)	45(83.3%)	0.608**
	Negative attitude	4(3.8%)	3(6%)	1(1.9%)	
	Not sure	15(14.4%)	7(14%)	8(14.8%)	

PC: palliative care.* p- value < 0.05 is significant.

** Fisher Exact

Table 3: Distributions of the emergency physicians' knowledge towards palliative care in Kuwait

		Total(N=104)	experience <11yrs(n=50)	experience ≥11yrs(n=54)	p- value*
Knowledge	Total score	5.68(2.5)	5.92(2.20)	5.46(2.68)	0.347
Objective assessment					
4- PC Definition	True	79(76%)	43(86%)	36(66.7%)	0.024
	False	25(24%)	7(14%)	18(33.3%)	
5- Members of PC team	True	83(79.8%)	43(86%)	40(74.1%)	0.149
	False	21(20.2%)	7(14%)	14(25.9%)	
6- Weak Opioids	True	57(54.8%)	28(56%)	29(46.3%)	0.846
	False	47(45.2%)	22(44%)	25(46.3%)	
7- Delirium	True	62(59.6%)	33(66%)	29(53.7%)	0.234
	False	42(40.4%)	17(34%)	25(46.3%)	
8- Dyspnea	True	18(17.3%)	9(18%)	9(16.7%)	0.857
	False	86(82.7%)	41(82%)	45(83.3%)	
9- Hypercalcaemia	True	65(62.5%)	29(58%)	36(66.7%)	0.420
	False	39(37.5%)	21(42%)	18(33.3%)	
10- WHO pain management	True	39(37.5%)	21(42%)	18(33.3%)	0.420
ladder	False	65(62.5%)	29(58%)	36(66.7%)	
11- Opioid toxicity	True	31(29.8%)	20(40%)	11(20.4%)	0.029
	False	73(70.2%)	30(60%)	43(79.6%)	
12- SVC obstruction	True	25(24%)	6(12%)	19(35.2%)	0.011
	False	79(75.9%)	44(88%)	35(64.8%)	
13- Catastrophic bleeding	True	20(19.2%)	10(20%)	10(18.5%)	0.848
	False	84(80.8%)	40(80%)	44(81.5%)	
14- Spinal cord	True	94(90.4%)	45(90%)	49(90.7%)	0.999**
compression	False	10(9.6%)	5(10%)	5(9.3%)	
15-Oral opioids	True	19(18.3%)	9(18%)	10(18.5%)	0.945
	False	85(81.7%)	41(82%)	44(81.5%)	

PC: palliative care, WHO: World Health Organization, SVC: superior venacaval obstruction. * p-value < 0.05 is significant

** : Fisher Exact

Table (5): Generalized linear model of predictors of positive attitude:

	B	SE	OR	95% CI		p- value
				Lower	Upper	
Sex (male)	0.641	0.805	1.899	0.392	9.194	0.426
Age	0.025	0.038	1.026	0.953	1.104	0.501
Nationality (Kuwaiti)	0.393	0.809	1.482	0.304	7.233	0.627
Qualification (master, MBBS)	0.621	1.01	0.538	0.075	3.877	0.538
Position (assistant registrar or registrar)	0.419	1.19	0.658	0.064	6.752	0.724
Subspecialty	0.399	0.567	0.538	0.075	3.877	0.538
Years of experience	1.75	0.756	5.747	1.031	25.00	0.021
Basic knowledge score	0.377	0.122	1.458	1.148	1.851	0.002

* *p* -value < 0.05 is significant

Table 4: Comparison between the emergency physicians' characteristics working in different hospitals in Kuwait

	Mubarak AlKabeer (N=18)	AlAdan (N=14)	AlSabah (N=15)	Al-Jahra (N=17)	Al-Farawanyia (N=20)	Al-An (N=20)
Age	40.11(±11.6)	39.64(±10.4)	43.11(±8.8)	34.25(±5.2)	39.4(±9.8)	36.95(±10.0)
Sex						
• Males	14(77.8%)	13(92.9%)	14(93.3%)	17(100.0%)	16(80.0%)	16(80.0%)
• Females	4(22.2%)	1(7.1%)	1(6.7%)	0	4(20.0%)	4(20.0%)
Nationality						
• Kuwaiti	5(27.8%)	2(14.3%)	2(13.3%)	1(5.9%)	5(25.0%)	6(30.0%)
• Others	13(72.2%)	12(85.7%)	13(86.7%)	16(94.1%)	15(75.0%)	14(70.0%)
Qualification						
• MBBS	4(22.2%)	2(14.3%)	1(6.7%)	3(17.6%)	7(35.0%)	11(55.0%)
• Master	12(66.7%)	8(57.1%)	8(53.3%)	14(82.4%)	12(60.0%)	7(35.0%)
• MD	2(11.1%)	4(28.6%)	6(40.0%)	0	1(5.0%)	2(10.0%)
Position						
• Assistant registrar	2(11.1%)	0	0	1(5.0%)	1(5.0%)	2(10.0%)
• Registrar	14(77.8%)	10(71.4%)	11(73.3%)	16(94.1%)	18(90.0%)	17(85.0%)
• Senior registrar	1(5.6%)	4(28.6%)	1(6.7%)	0	1(5.0%)	1(5.0%)
• Specialist/ Consultant	1(5.6%)	0	3(20%)	0	0	0
Specialty						
• Emergency Medicine	8(44.4%)	12(85.7%)	8(53.3%)	10(58.8%)	16(80.0%)	14(70.0%)
• Internal Medicine	5(27.8%)	1(7.1%)	2(13.3%)	2(11.8%)	1(5.0%)	1(5.0%)
• Surgery	0	0	3(20.0%)	1(5.9%)	3(15.0%)	3(15.0%)
• Others	5(27.8%)	1(7.1%)	2(13.3%)	4(23.5%)	0	2(10.0%)
Experience yrs)	14.11(±9.6)	14.79(±10.5)	17.47(±8.5)	9.38(±5.1)	15.05(±10.3)	10.9(±10.0)
Attitude						
• Positive attitude	7(38.9%)	7(50.0%)	5(33.3%)	4(23.5%)	5(25.0%)	6(30.0%)
• Negative attitude	2(11.1%)	2(14.3%)	4(26.7%)	5(29.4%)	3(15.0%)	5(25.0%)
• Not sure	9(50.0%)	5(35.7%)	6(40.0%)	8(47.1%)	12(60.0%)	9(45.0%)
Knowledge						
Total score	4.78(2.25)	6.0(2.51)	6.07(2.81)	6.12(2.48)	5.65(2.43)	5.6(2.43)
• Good knowledge	1(5.6%)	3(21.4%)	4(26.7%)	3(17.6%)	3(15.0%)	2(10.0%)
• Fair knowledge	6(33.3%)	4(28.6%)	6(40.0%)	7(41.2%)	7(35.0%)	7(35.0%)
• Poor knowledge	11(61.1%)	7(50.0%)	5(33.3%)	7(41.2%)	10(50.0%)	11(55.0%)

* P value < 0.05 is significant **; Fisher Exact

Figure (2): Knowledge of Emergency Physicians toward Palliative Care

Figure 1: The Flowchart for the sampling procedure

EPs: emergency physicians

Figure (2): Attitude of Emergency Physicians toward Palliative Care