Clinical and Inflammatory Factors Influencing Constipation and Quality of Life in Cerebral Palsy

Ana Cristina Fernandes Maria Ferreira¹, Ryan J. Eveloff², Marcelo Freire^{*2}, Maria Teresa Botti Rodrigues Santos^{*1,3}

¹ Cruzeiro do Sul University, Rua Galvão Bueno, 868 - Liberdade, São Paulo, SP, 01506-000, Brazil

² J. Craig Venter Institute, 4120 Capricorn Lane, La Jolla, California, USA

³ Associação de Assistência à Criança Deficiente, AACD, São Paulo, BRAZIL

***Corresponding Author:**

Email: maria.botti@cruzeirosul.edu.br (UNICSUL)

Email: mfreire@jcv.org (JCVI)

Acknowledgments

This study was supported by the São Paulo Research Foundation (Fundação de Amparo à Pesquisa do Estado de São Paulo, FAPESP #2017/15160-4).

1 Abstract

2 **Introduction:** Intestinal constipation is a clinical consequence, secondary to neuromotor
3 disturbances, which acts on subjects with cerebral palsy (CP). The aim of this study was to
4 investigate the factors influencing constipation and the quality of life (QOL) of CP subjects.

5 **Materials and methods:** We recruited a total of 63 subjects aging from 5-17 years with spastic
6 CP who received physical rehabilitation. The subjects were divided into two groups including
7 with and without constipation. Subjects were assigned into one of the 4 groups (G1-4) based
8 on the prevalence of 1) CP and 2) Constipation. Subjects were assigned as CP with
9 constipation (G1), CP without constipation (G2), and controls without CP with constipation
10 (G3) and without CP and without constipation (G4). Subjects' demographics, use of
11 antiepileptic drugs (AEDs), motor function, caregiver priorities and child health index of life
12 with disabilities (CPCHILD) were included. In addition to medical metadata, a subset of
13 subjects was evaluated for oral and systemic inflammation through gingival bleeding and
14 serum cytokine levels (TNF- α , IL-1 β , IL-6, IL-8, IL-10) respectively. Statistical significance was
15 evaluated by ANOVA One-Way (parametric data) and Kruskal Wallis (non-parametric data).

16 **Results:** A significant relationship was found between the type of medication and constipation.
17 Subjects taking GABA and GABA+ (GABA in association with other medication) were more
18 likely to be constipated than the other groups ($P < 0.01$). Additionally, quality of life was directly
19 correlated with constipation; subjects in G1 presented the lower mean score of CPOCHILD (49.0
20 ± 13.1) compared to G2 (71.5 ± 16.7), when compared to G3 (88.9 ± 7.5), and G4 (95.5 ± 5.0)
21 ($P < 0.01$). Inflammation was more severe for patients in G1 ($P < 0.001$). There were no
22 differences among groups regarding gender ($P = 0.332$) and age ($P = 0.292$).

23 **Conclusions:** Our results suggest that constipation was mostly affected by the use of certain
24 antiepileptic drugs, namely GABA. This showed direct detrimental effect on CP quality of life,
25 which was influenced by inflammatory cytokines and the dosage and type of AEDs.

26 **Keywords:** Cerebral palsy, Constipation; Antiepileptic drugs, Quality of Life, Caregiver
27 Priorities and Child Health Index of Life with Disabilities (CPOCHILD), Cytokines

Introduction

Cerebral palsy (CP) is a life-limiting and costly disability [1]. CP is characterized by permanent neuromotor disorder affecting movement and by non-progressive degeneration of the brain [2]. As the major etiological factor for severe disability, the estimated prevalence of CP ranges from 2.3 to 2.9 per 1000 live births observed in the United States (2011–2012 National Survey of Children’s Health and the 2011–2013 National Health Interview Survey) [3]. Children with CP experience limitations in everyday activities, all of which are factors influencing their quality of life (QOL). It can be accompanied by difficulties related to perception, sensation, behavior, cognition, communication, and epilepsy. And, their QOL can be impacted by the development of musculoskeletal problems such as oral-gut motor impairment through muscle spasticity [2].

Due to an abnormal increase in muscle tone and injury to neural pathways, spasticity acts as a negative factor in the lives of 85-90% of individuals with CP [2,4,5]. The severity of movement disability is classified by Gross Motor Function Classification System (GMFCS) [6]. GMFCS is stratified into five levels of body mobility: Level I (walks without limitations), Level II (walks with limitations), Level III (walks using a hand-held mobility device), Level IV (self-mobility with limitations) and Level V (transported in a wheelchair) [6]. Our previous studies demonstrated that out of 254 subjects, 50 (19.7%) subjects had GMFCS I, II or III while 204 (80.3%) presented GMFCS IV or V [7–9]. These proportion results reflect on the population of this current cohort. It is worthy mentioning, that epilepsy prevalence increases in spastic CP subjects as the levels of GMFCS rise (IV and V) [10]. In fact, epilepsy affects 77% of the subjects with CP, and standard clinical treatment for epileptic CP patients is based on therapy with antiepileptic drugs (AEDs) [11]. As the principal inhibitory neurotransmitter of the central nervous system and also regulatory signal for muscle tone, Gamma Aminobutyric Acid (GABA) is used for neuronal excitability control (e.g.: benzodiazepines, phenobarbital, topiramate, vigabatrin, gabapentin, enacarbil and ezogabine). Side effects of AEDs can range from

subjective reports of mild drowsiness to life-threatening neurologic and gastrointestinal consequences [12].

The prevalence of severe constipation and lack of gastrointestinal control varies from 25% to 74% [3]. Constipation may be caused by multiple components including: 1) reduced intake of fiber and liquids (responsible for the digestive system functioning) [13], 2) central nervous system damage [14], 3) mobility reduction [15], 4) and/or the use of AEDs [13,16]. In subjects using AEDs the incidence of constipation is elevated [7]. Other side effects co-occurring with the gastrointestinal complications include oral dysbiosis, gingival bleeding (GB), and increase of systemic inflammation [17]. The exact clinical phenomic factors that are influenced by the use of AEDs and the impact on QOL remains elusive.

Here, we aimed to investigate the clinical factors that mostly influenced QOL in cerebral palsy subjects. Results demonstrated positive relationships of gastrointestinal constipation and the use of medication impacted the quality of life of CP patients. Of the AEDs tested, GABA was the strongest factor that influenced constipation, with significant results to gut and oral compartments. We further investigated the mechanism involved in the oral-gut phenotypes and assayed the inflammation as a catalyst for the clinical observations. For a subset of 37 participants, all of whom were CP subjects, we analyzed the salivary cytokines IL-6, IL-8, IL-10, IL1 β , and TNF- α . Overall, the results demonstrated that, *via* elevated inflammation, the most constipated subjects had increased frequency of non-communicable diseases (oral and gut), leading to a decrease in QOL.

Materials and Methods

Study Design

This study was reviewed and approved by the Research Ethics Committee of the Cruzeiro do Sul University-Brazil Platform, São Paulo, Brazil (IRB #2,452,626). Written informed consent

was obtained from the guardian of each child or adolescent after they were informed about the study. A cross-sectional study was performed with subjects with spastic CP diagnosis, who received a physical rehabilitation treatment at a referral center in São Paulo, Brazil, at the time of data collection.

Participants

One hundred subjects with a medical diagnosis of CP were invited to participate in this study at AACD (Disabled Child Care Association) in São Paulo, Brazil. Inclusion criteria were a medical diagnosis of spastic CP ranging from 5–17 years, both male and female, and presence or absence of constipation. Participants who presented progressive or neurodegenerative lesions or uncooperative behavior were excluded during clinical oral examinations. Data was collected from year of 2018 until 2019. Both demographic and clinical data were collected for each subject.

Gastrointestinal (GI) Constipation

This study adopted the clinical constipation definition proposed by the Rome III criteria for constipation [18]. The individuals must present two or more of the following symptoms, in the previous three months, to be classified as presenting constipation including: (a) makes force to evacuate during at least 25% of defecations; (b) hard or irregular stools in at least 25% of defecations; (c) sensation of incomplete evacuation for at least 25% of defecations; (d) sensation of obstruction for at least 25% of defecations; (e) use of manual maneuvers in at least 25% of defecations; (f) less than three defecations per week.

The final sample of the study was composed of 30 subjects with CP with constipation (G1), 33 subjects with CP without constipation (G2), 7 subjects without CP with constipation (G3) and 23 subjects without CP and without constipation (G4). Data regarding gender, age, race (white, black and others), caregiver occupation and education, family income, medical diagnosis of

CP according to the type of movement disorder (spastic), clinical pattern (tetraplegia, diplegia and hemiplegia), Gross Motor Function Classification System [6] (levels I to V) and use of AEDs were collected from their medical records. Caregivers answered the Caregiver Priorities and Child Health Index of Life with Disabilities [19] that consists of 36 items rated in six sections. CPCHILD is represented by the following domains: (1) Personal Care (eight items); (2) Positioning, Transfer, and Mobility (eight items); (3) Communication and Social Interaction (seven items); (4) Comfort, Emotions, and Behavior (nine items); (5) Health (three items); and (6) Overall Quality Of Life (one item).

Saliva Collection

Unstimulated saliva samples were collected in dental assessment sessions. Subjects were asked to refrain from eating, drinking liquids or brushing their teeth for at least 1 hour prior to saliva collection. The collection was performed with the subjects sitting comfortably in a bright and ventilated room. Whole saliva was collected by passive flow for 5 minutes. After collection, the Salivette® was centrifuged at 5,000 rpm for five minutes at 4°C (Hettich Centrifuge, model Universal 320R, Tuttlingen, Germany) and frozen in a freezer at -80°C.

Biomarker Sub Analysis

The analysis of cytokines in saliva was performed using a CBA Cytokine Inflammatory Kit (Becton Dickinson, CA, USA) for the detection of TNF- α , IL-1 β , IL-6, IL-8, IL-10. All analyses were performed in duplicate. Briefly, 25 μ L of fluorescent particles conjugated to antibodies specific for each cytokine were added to 25 μ L of the saliva and incubated for one hour at room temperature away from light. Subsequently, 25 μ L of the secondary antibody conjugated to a fluorochrome was added to the mixture and incubated for two hours at room temperature. The results were compared to a standard curve with serially diluted cytokines. The particles were washed to remove unbound antibodies, resuspended in wash buffer and analyzed using a BD

Accuri (BD Biosciences). Data acquisition was performed using BD-Accuri C6 Software, and concentrations were determined using FCAP software v.3.0 (BD Biosciences).

Gingival Index (GI)

The evaluation of oral dysbiosis was assayed through the GI [20] by using a millimeter plastic periodontal probe (HuFriedy's Colorvue PerioScreen Kit probe, Chicago, IL, USA), which was gently passed in the gingival margin of all teeth, in reference to the distobuccal papilla, the buccal margin, the mesiobuccal papilla and the lingual / palliative margin. Partially erupted teeth and residual roots were excluded without replacement. The index was calculated by the percentage of the sum of the individual values of each tooth divided by the number of faces examined. Classified as positive for gingivitis were the individuals that presented gingival marginal bleeding more than 20% of the total sites evaluated [20].

Statistical Analyses

Analyses of descriptive statistics were performed to characterize the sample, calculate measures of central tendency and variability for the quantitative variables. The normality assumption of the quantitative variables was evaluated using the Shapiro-Wilks test. When normal distribution was observed, parametric tests were performed. Otherwise, non-parametric tests were selected to determine the significance of intergroups differences. The Chi-square test was used to analyze the variables of gender, presence of gingival bleeding, clinical type of CP, GMFCS, and AEDs used.

ANOVA One-Way (parametric data) was used to determine significant intergroups differences in relation to age and the periodontal condition studied. Kruskal Wallis(nonparametric data) was used to determine significant differences in relation intergroups cytokines levels. IBM SPSS Statistics (SPSS for Windows, Version 20.0, Armonk, NY: IBM Corp.) was used for all

analyses, with a significance level at $P < 0.05$. Data was visualized using Python 3.7 Anaconda with PyPlot and Seaborn and RStudio with ggplot2.

Results

1-Participants Demographics

The goal of our study was to identify the main factors affecting QOL in CP subjects. We focused on first collecting subject information including demographics, clinical factors, and AEDs as potential influencers on a patient's wellbeing and QOL (Table 1). In our population, 47 of our 93 subjects were female and 46 were male, accurately representing the relatively equal ratio of women to men. The sample power was calculated using means and standard deviations of overall domains among subjects with CP with constipation (51.0 ± 13.1), subjects with CP without constipation (28.5 ± 16.7), subjects without CP with constipation (11.1 ± 7.5) and subjects without CP and without constipation (4.5 ± 5.0) (OpenEpi online; www.openepi.com). The results showed that the G*Power at the 95% confidence interval was 96.88%.

There were no significant differences among groups regarding gender ($P = 0.332$), age ($P = 0.291$), race ($P = 0.08$), and education ($P = 0.5628$), yet significant differences were found on caregivers occupation ($P = 0.0017$) and family income ($P = 0.018$). G1 was composed of 30 non-ambulatory subjects (GMFCS levels IV and V) and G2 of 14 ambulatory subjects (GMFCS levels I, II, and III) ($P = 0.0002$). Worse QOL measures were observed for constipated subjects presenting the clinical pattern tetraplegia when compared to non-constipated CP ($P < 0.001$) (Table 1). Interestingly, gender was significantly associated with constipation ($P < 0.05$) with females appearing more likely to be constipated in our population. Distribution of age in our population shows a peak at ~6 years of age, with a slight steady dropoff in population representation until approximately age 14, with a steep dropoff in population representation from ages 14-19 (Figure 1A).

2- CP Diagnosis

When diagnosing CP, there are three widely accepted presentations: 1) hemiplegic, or Increased muscle tone in a hemibody; 2) diplegic, Increased muscle tone in lower limbs (may affect upper limbs, but to a lesser extent); and 3) tetraplegic, a system-wide decay of neuromotor function, represented by increased muscle tone in all four limbs and trunk involvement. In our population, the distribution of subjects is made up of these three groups as well as healthy individuals (control). Hemiplegia is the least prominent in our population, with 9.68% of subjects presenting hemiplegia, 26.88% diplegia, and 31.18% tetraplegia (32.26% of our population was healthy at examination) (Figure 1B). CP diagnosis was found to be significant in our population ($P < 0.001$) in determining QOL (Table 1).

Another mechanism to classify subjects with spastic CP is according to the motor function ability. The Gross Motor Function Classification System (GMFCS) classify as ambulatory that ones who present levels I-III, while levels IV and V are reserved for non-ambulatory subjects. In our population, levels I (37.63%), IV (29.03%) and V (23.66%) make up the vast majority of our population (Figure 1C). Level I is mostly populated by our control group, while groups IV and V are reserved for subjects with severe CP, namely diplegic and tetraplegic CP (Figure 1D). More specifically, group IV is made up of diplegics subjects who can typically operate a wheelchair manually, while tetraplegic subjects require caregiver-assisted or motorized forms of transportation.

We found that a subject's GMFCS level was statistically significant in determining their QOL ($P < 0.001$). Functional motor independence of subjects with CP is directly related to the extent of cortical injury. Diplegic and quadriplegic subjects (GMFCS IV and V) have significant motor impairment that limits daily living activities. Therefore, the diagnosis of CP and greater motor damage are two strong factors that influence the quality of life of the subjects.

3- Constipation Impact on Quality of Life

According to CPCHILD, the groups differed significantly ($P < 0.05$). G1 presented higher values (lower QOL) for all sections of the questionnaire while compared to G2, G3 and G4. The CPCHILD questionnaire measures QOL presented 6 subscores: 1) Positioning, Transferring, and Mobility; 2) Caregiver-Reported QOL; 3) Hospitalization and Caregiver-Reported Health; 4) Comfort and Emotions; 5) Personal Care; and 6) Communication and Social Interaction. These subscores are scaled from 0 to 100 and were averaged to determine overall QOL. The combination of constipation and cerebral palsy, showed to lower the quality of life of each individual. The inability to effectively and routinely have bowel movements and the inhibition or even incapability to move freely leads to fewer opportunities and consequently lower overall QOL.

In our initial analysis, we uncovered the aspects of subject's life that determine their overall QOL. In figure 2, each subscore is plotted in relation to the remaining 5 subscores. For example, figures 2A-F show Hospitalization and Caregiver-Reported Health (HCRH) compared to all other subscores. A striking trend is observed wherein a particular subscore (in this case) was not particularly influential in determining overall QOL, as a rating of 80 in HCRH can still yield QOL scores as low as 20. Conversely, Communication and Social Interaction (CSI) scores much more closely reflect QOL scores as a whole. Gender was not significantly correlated with QOL in our population. Further analysis into what physical and psychological factors that most directly pertain to a subject's QOL is required to better understand and treat CP subjects.

3- Medication Affecting Constipation

The key factor affecting QOL and constipation in our population was revealed to be medication types. As the common thread linking all CP subjects, pharmacological treatment is an essential point of emphasis of our study. Subjects that use of GABA (the principal inhibitory

neurotransmitter in the central nervous system and regulate muscle tone) and GABA+ presented the worst quality of life. The use of sodium inhibitors (Na inhibitor) were positively correlated with a higher quality of life, potentially due to the removal of sodium from the digestive tract, yielding softer stool (Figure 1 A-C). Specifically, the choice of AED used for treatment of CP and epilepsy greatly influenced the physical and mental health of our subjects. GB is seen in figures 3D-F to be higher in subjects taking GABA and GABA+, or GABA in association with other AEDs. Conversely, subjects taking sodium inhibitors were clustered at lower mean GB.

Sodium inhibitors were associated with a higher QOL and GABA (the chief inhibitory neurotransmitter in our body) was associated with a lower QOL. Similarly, GABA+ was associated with a significantly lower QOL. In Figure 3G, there is a clear trend wherein GABA subjects are clustered at a lower QOL, while sodium inhibitor subjects, particularly in our non-constipated subcohort (Figure 3H), exhibit a clustering at a much higher QOL. Calcium inhibitors were only taken by 2 subjects in our population, so further studies must be performed with larger populations to fully understand their effect on the QOL and bowel health of CP and epilepsy subjects. With this information, it is clear that the adverse effects of GABA on a subjects' body and mental wellbeing must be observed further; similarly, sodium inhibitors should be studied to determine their efficacy as an AED as a replacement for or synergistically with GABA.

4- Oral and Systemic Inflammation are Elevated in Constipated Patients

From a clinical perspective, CP is considered a neurological disease. However, its symptoms can manifest themselves throughout the body. In many subjects, the lack of an ability to care for oneself leads to the development of oral and gum disease. These diseases manifest themselves through bleeding of the gingiva. If a large number of teeth have bleeding gingiva,

there is a high chance of oral disease manifesting itself as gingival inflammation and periodontitis.

In figure 4A, Density for males (green) appears more concentrated at lower levels of Gingival bleeding (GB). Females (blue) are more stratified, but still show a higher density where GB is < 20%. The average GB observed for males was $8.37 \pm 12.18\%$, while the same for females was $16.70 \pm 22.26\%$. Overall mean GB was $12.08 \pm 18.27\%$. In figure 4B, a slight visual trend is observed with the upper limit of mean GB decreasing with age. Higher densities of high gingival index occur in younger subjects, while higher densities of low gingival bleeding occur in older subjects. Healthy individuals (those who do not suffer from CP) have markedly lower mean GB (23%) than any other group, with 2.79% mean hemiplegic GB, 12.19% mean diplegic GB, and 27.11% mean tetraplegic GB (Figure 4C).

Indirect symptoms of CP are not limited to the oral cavity; gastrointestinal problems can often arise as a reaction to medications or from an inability to properly digest and pass food. Of our 63 subjects who present CP, 30 were constipated at examination, representing a drastically higher prevalence of constipation (47.62%) compared to the estimated prevalence of 16% in the global population [21].

For a subset of our population, we collected salivary cytokine measures to get an accurate idea of systemic inflammation and its relationship with constipation and medication. For all cytokines for which data was collected (IL-6, IL-8, IL-10, IL-1 β , and TNF- α), mean levels approximately doubled in constipated subjects. In figures 5A-E, there is a clear visual trend of higher resting systemic inflammation in individuals with severe CP. Of these cytokines, only IL1 β was significantly correlated with constipation ($P < 0.05$, Figure 5D).

Figure 6 reveals the positive and negative correlations in our study cohort among clinical factors and demographics. Naturally, each subscore is correlated with the overall QOL of an individual as they are part of its calculation. Key trends to notice are: 1) The presence of constipation significantly lowers QOL; 2) GMFCS and CP Type are strongly correlated, both to one another and to QOL; 3) GB exhibits a strong negative correlation with QOL that is statistically significant; and 4) Overall QOL is more strongly correlated to the self-reported mental health of -subject than to caregiver-reported metrics.

Discussion

The results provided novel information related to the severity of CP and the consequences of treatment on constipation, oral health, and quality of life. To the best of our knowledge, this is the first study to evaluate the association between intestinal constipation and quality of life of CP using the Caregiver Priorities and Child Health Index of Life with Disabilities. According to the motor type disorder, subjects with CP are classified as at least one of spastic, dyskinetic (dystonic, choreic and athetotic), or ataxic. The subjects presented spastic CP, the most prevalent type of motor disorder, with tetraplegia the clinical pattern most prevalent (32.26%) (Figure 1B). Consistent with our results, tetraplegia was also observed in a 3,294 children study developed in 18 states in the USA (32.9%) [22]. Early brain damage triggers complex adaptive neuroplasticity processes involving multiple functional systems. Different factors interact with brain plasticity potentials, influencing the natural history of cerebral palsy in the early years of childhood [23]. The later the interdisciplinary intervention, the smaller the gain during brain plasticity, reducing the potential for improvement in the individual's condition.

Here, we observed a shortage of national instruments, which appears to reflect the difficulty of the scientific community to develop tools for assessing quality of life that apply to the socio-cultural diversity of the country. For all questionnaires revised, only Caregiver Priorities and

Child Health Index of Life with Disabilities (CPCHILD) [19] and Cerebral Palsy Quality of Life Questionnaire for Children Child report (CPQoL-Child) [24] are specific to Cerebral Palsy subjects. CPQOL-Child is the individual's own report, since it was mostly used in children with GMFCS level I or II and were excluded the individual with any intellectual deficits and lack of an efficient communication system. In this study, the largest number of individuals belonged to GMFCS level IV and V (Figure 1C) and due to the difficulty of communication, their caregivers answered the questionnaire as their proxy; in this case the CPCHILD is the best choice of questionnaire. The other questionnaires were for children and adapted for a wide range of disabilities, while CPCHILD is more specific to our study population. The Cronbach's alpha value estimated for the Caregiver Priorities and Child Health Index of Life with Disabilities was 0.879. It was observed that G1 presented a higher mean score of CPCHILD (51.0 $\pm$ 13.1) compared to G2 (28.5 $\pm$ 16.7), G3 (11.1 $\pm$ 7.5), and G4 (4.5 $\pm$ 5.0) ($P < 0.05$) (Table 1). Seventy percent of caregivers of CP subjects in this study had less than 8 years of study (Table1). Intellectual capital, represented by education, increases the likelihood of access to information, health-related issues [25] facilitating understanding and recognizing the importance of self-care. A low level of caregiver education influences the child's overall health, represented by self-care of general and oral health [26].

The possible explanation for the lower education level of caregivers for individuals with CP is the 'caregiver martyr syndrome' wherein a caregiver neglects their own health, studies, and finances to provide more frequent and consistent [27] care [27]. With a lower educational level and fewer paid employment opportunities, there is a direct logical connection linking significantly lower income to caregivers of individuals with CP (Table 1). Caregivers' potentially lower standard of education prevents them from entering the skilled labor market. Thus, they dedicate themselves exclusively to household chores and taking care of the individual with CP [28]. Emerging concepts of quality of life related to CP are emerging, and in the last five years, a total of 469 articles were found in the literature.

347

348 We noticed that the severity of pain and discomfort has been associated with increased motor
349 impairment and other comorbidities, particularly constipation and spasticity. In individuals with
350 minor GMFCS, pain was observed during voluntary movements, whereas in individuals with
351 larger GMFCS, pain was directly related to movement performed by the caregiver or therapist
352 [29]. Higher intensity of pain and discomfort had a negative effect on physical and
353 psychological quality of life (Figure 2A–F). Communication and social interactions in
354 individuals with cerebral palsy in GMFCS level V with intellectual disability develop less
355 favorably and show large variation [30] (Figure 2F). Fairhust et al. concluded in 2018 that
356 increasing awareness of CP pain and comorbidities, particularly overall health and
357 constipation, can help prevent and treat the disorder more effectively, increasing patients'
358 overall quality of life.

359

360 Medication such as AEDs are frequently used for the control of epileptic seizures [31]. These
361 drugs can be administered in monotherapy or polytherapy, depending on the response to
362 seizure control [32]. Results demonstrated that individuals with a lower quality of life were
363 positively correlated to those who use polytherapeutic GABA treatments (GABA+), while those
364 who use monotherapy, especially sodium channel inhibitors, have a better quality of life (Figure
365 3C) when compared with GABA and GABA+ (Figure 3B). AEDs have an effect on voltage-
366 dependent modulation of sodium, calcium, and potassium channels, and the Aminobutyric
367 Gamma Acid (GABA) is considered the primary inhibitory neurotransmitter not only in the
368 central nervous system, but also on enteric nerves [32]. However, AEDs aims to achieve
369 seizure control are usually limited by toxicity and adverse effects that impair an individual's
370 quality of life [33] (Figure 3B-D). The subjects taking GABA or GABA association presented a
371 negative correlation with quality of life and also present intestinal constipation not only for the
372 inhibitory GABA action, [34] but may also be related to dysbiosis.

A higher prevalence of gingivitis was observed in constipated subjects from this study, that were in use of GABA/or GABA+ (Figure 3E-G, *Supplementary Figure 1*). G1 presented significantly higher percentages of subjects with gingivitis compared to all other groups ($P < 0.001$). Additionally, CP subjects using GABA or GABA+ who were constipated had higher levels of gingival bleeding than those who did not ($P = 0.006$, Table 1). Current literature indicates that intestinal disorders play a prominent role in inflammatory responses to neurological conditions [35]. This line of evidence is fundamental to identifying the effects of dysbiosis on mucosal inflammation throughout the digestive tract. Significantly higher levels of IL1 β , IL6, IL8, and IL10 were found in constipated subjects with gingival bleeding from this study (Figure 5), indicating an ongoing inflammatory process and the progression of periodontal disease [36–40]. Since the use of these medications cause CP subjects to present a reduced salivary flow rate, an increased value of salivary osmolality, dry mouth and gingivitis, which is represented by higher levels of inflammatory cytokines in quadriplegics [17]. Sodium inhibitors, however, have a less severe correlation with gingival bleeding ($R_2=.684$), but have markedly lower effects on a patient's QOL and they are associated with lower levels of constipation, it can be assumed that sodium inhibitors carry less of a systemic reaction in our study population.

Constipation is a common comorbidity described in CP subjects [13], and current literature agrees that constipation is the causal factor of gut dysbiosis [41,42]. Constipation prevalence observed in this study was higher (47.62%), with higher prevalence in females CP subjects (Figure 4). Females without CP present constipation and almost 2 times more likely to suffer from constipation than males [43]. Constipation and intestinal dysbiosis leads to increased mucosal permeability (leaky-gut) [42] of the Gut-Brain Axis [44] increasing serum endotoxin concentration. These endotoxins activate the immune system and promote IL1 β production as observed in individuals with Autism Syndrome Disorder [45]. Moreover, there has been no published research to date that evaluated the effect of dysbiosis on gingivitis development in individuals with cerebral palsy.

In addition to GI and oral inflammation, we have observed gender to be an important variable in regard to inflammation. Females presented higher pro-inflammatory cytokine expression when compared to males (Figure 4). This gender bias has not been addressed in the literature and will require further studies. The possible explanation for these findings may be related to the phenomenon that CP occurs more frequently in male [46] categorized as GMFCS V [47]. Due to this limitation, these subjects require more advanced care, notably oral care and hygiene performed by the caregiver. Perhaps the females were able to perform oral hygiene on their own due to fewer neuromotor limitations, which in turn may have resulted in greater gingival bleeding. Additionally, hormonal changes in puberty should be examined as a factor for this observed gender dichotomy [48].

Inflammatory biomarkers can be evaluated both in serum [49] or in saliva [8]. Blood collection in subjects with CP is a hard task, because these subjects present sympathetic nervous system predominance [50], resulting in vasoconstriction, and consequently making peripheral venous access more difficult. On the other hand, saliva is readily available and its components are easy to be collected, and that has been used to evaluate salivary parameters in subjects with CP [17]. Ideal for future longitudinal monitoring.

The comparison between the subjects constipated and non-constipated from this study demonstrated that constipation acts as a damaging factor causing approximately doubled mean cytokine production (Figures 5A-E). In the presence of dysbiosis, virulence factors are released from pathogenic microorganisms present in the oral cavity, activating host's immune-inflammatory responses [51–53]. IL1- β is an inflammatory cytokine associated with innate immune response, inflammation, pathogenesis and progression of periodontal disease [54,55]. Higher levels of IL1 β ($P < 0.05$) were found in this study's subjects, who were constipated and presented higher values of gingival bleeding (Figure 5B, *Supplementary*

Figure 2B). Previous studies showed higher levels of IL-1 β IL-6, IL-8 in spastic CP subjects presenting GMCFS level V [17].

Significantly higher levels of IL-6, IL-8 and IL-10 were found in constipated subjects with gingival bleeding (*Figure 5C-E; Supplementary Figure 2C-E*). These cytokines represent the action of pro-inflammatory and anti-inflammatory cytokines [55], critical biomarkers of periodontal inflammation corresponding to the clinical severity of the disease [37]. IL-6 and IL-8 were also described as potential predictors for oral diseases, reinforcing the importance of evaluating the salivary levels of these biomarkers [56]. The presence of the chemokine IL-8 induces the secretion of lymphocytes, monocytes, epithelial cells, fibroblasts, tumor cells, bone resorption and IL-1 β , indicating an ongoing inflammatory process and the progression of periodontal disease [38,39]. IL-10 acts as an anti-inflammatory cytokine, inhibiting pro-inflammatory cytokines IL-1 and IL-6 associated with improvements in periodontal clinical parameters [57]. However, the degree of mucosal inflammation of constipated CP subjects represented by gingival bleeding was so high that IL-10 action did not reduce inflammatory process of dysbiosis [40]. Only one study reported the effect of mechanical treatment on gingivitis control measures by cytokines levels, and the results showed that, although occurs reduction of inflammatory process, the levels of IL-1 β , IL-6 and IL-8 remain high compared to subjects without CP [8]. In conclusion, constipation has a direct detrimental effect on quality of life, which is influenced by antiepileptic drugs intake, type of antiepileptic drugs, and inflammatory signals.

Acknowledgments

This study was supported by the São Paulo Research Foundation (Fundação de Amparo à Pesquisa do Estado de São Paulo, FAPESP #2017/15160-4).

References

1. Grether JK, Nelson KB. Maternal infection and cerebral palsy in infants of normal birth weight. *JAMA*. 1997;278: 207–211.
2. Rosenbaum P, Paneth N, Leviton A, Goldstein M, Bax M, Damiano D, et al. A report: the definition and classification of cerebral palsy April 2006. *Dev Med Child Neurol Suppl*. 2007;109: 8–14.
3. Maenner MJ, Blumberg SJ, Kogan MD, Christensen D, Yeargin-Allsopp M, Schieve LA. Prevalence of cerebral palsy and intellectual disability among children identified in two U.S. National Surveys, 2011–2013. *Ann Epidemiol*. 2016;26: 222–226.
4. Wimalasundera N, Stevenson VL. Cerebral palsy. *Practical Neurology*. 2016. pp. 184–194. doi:10.1136/practneurol-2015-001184
5. Sewell MD, Eastwood DM, Wimalasundera N. Managing common symptoms of cerebral palsy in children. *BMJ*. 2014;349: g5474.
6. Palisano R, Rosenbaum P, Walter S, Russell D, Wood E, Galuppi B. Development and reliability of a system to classify gross motor function in children with cerebral palsy. *Dev Med Child Neurol*. 1997;39: 214–223.
7. Ferreira ACFM, Ana Cristina Fernandes, Mayer MPA, Kawamoto D, Maria Teresa Botti. Constipation, antiepileptic drugs, and gingivitis in children and adolescents with cerebral palsy. *International Journal of Paediatric Dentistry*. 2019. pp. 635–641. doi:10.1111/ipd.12488
8. Yoshida RA, Gorjão R, Mayer MPA, Corazza PFL, Guare RO, Ana Cristina Fernandes, et al. Inflammatory markers in the saliva of cerebral palsy individuals with gingivitis after periodontal treatment. *Brazilian Oral Research*. 2019. doi:10.1590/1807-3107bor-2019.vol33.0033
9. Gutierrez G-M, Siqueira V-L, Loyola-Rodriguez J-P, Diniz M-B, Guaré R-O, Ferreira A-C-F-M, et al. Effects of treatments for drooling on caries risk in children and adolescents with cerebral palsy. *Med Oral Patol Oral Cir Bucal*. 2019;24: 204–210.
10. Delacy MJ, Reid SM. Profile of associated impairments at age 5 years in Australia by cerebral palsy subtype and Gross Motor Function Classification System level for birth years 1996 to *Medicine & Child ...*. 2016. Available: <https://onlinelibrary.wiley.com/doi/abs/10.1111/dmcn.13012>
11. Bearden DR, Monokwane B, Khurana E, Baier J, Baranov E, Westmoreland K, et al. Pediatric Cerebral Palsy in Botswana: Etiology, Outcomes, and Comorbidities. *Pediatr Neurol*. 2016;59: 23–29.
12. Selai C, Bannister D, Trimble M. Antiepileptic drugs and the regulation of mood and quality of life (QOL): the evidence from epilepsy. *Epilepsia*. 2005;46 Suppl 4: 50–57.
13. Veugelers R, Benninga MA, Calis EAC, Willemsen SP, Evenhuis H, Tibboel D, et al. Prevalence and clinical presentation of constipation in children with severe generalized cerebral palsy. *Dev Med Child Neurol*. 2010;52: e216–21.

- 496 14. Johanson JF, Sonnenberg A, Koch TR, McCarty DJ. Association of constipation
497 with neurologic diseases. *Dig Dis Sci*. 1992;37: 179–186.
- 498 15. Park ES, Park CI, Cho S-R, Na S-I, Cho YS. Colonic transit time and constipation
499 in children with spastic cerebral palsy. *Arch Phys Med Rehabil*. 2004;85: 453–456.
- 500 16. Jahromi SR, Togha M, Fesharaki SH, Najafi M, Moghadam NB, Kheradmand JA,
501 et al. Gastrointestinal adverse effects of antiepileptic drugs in intractable epileptic
502 patients. *Seizure*. 2011. pp. 343–346. doi:10.1016/j.seizure.2010.12.011
- 503 17. Santos MTBR, Maria Teresa Botti, Diniz MB, Guaré RO, Ferreira MCD, Gutierrez
504 GM, et al. Inflammatory markers in saliva as indicators of gingival inflammation in
505 cerebral palsy children with and without cervical motor control. *International*
506 *Journal of Paediatric Dentistry*. 2017. pp. 364–371. doi:10.1111/ipd.12270
- 507 18. Longstreth GF, Grant Thompson W, Chey WD, Houghton LA, Mearin F, Spiller
508 RC. Functional Bowel Disorders. *Gastroenterology*. 2006. pp. 1480–1491.
509 doi:10.1053/j.gastro.2005.11.061
- 510 19. Pellegrino LA, Ortolan EVP, Magalhães CS, Viana AA, Narayanan UG. Brazilian
511 Portuguese translation and cross-cultural adaptation of the “Caregiver Priorities
512 and Child Health Index of Life with Disabilities” (CPCHILD©) questionnaire. *BMC*
513 *Pediatrics*. 2014. doi:10.1186/1471-2431-14-30
- 514 20. Chaves ES, Wood RC, Jones AA, Newbold DA, Manwell MA, Kornman KS.
515 Relationship of “bleeding on probing” and “gingival index bleeding” as clinical
516 parameters of gingival inflammation. *Journal of Clinical Periodontology*. 1993. pp.
517 139–143. doi:10.1111/j.1600-051x.1993.tb00328.x
- 518 21. Mugie SM, Benninga MA, Di Lorenzo C. Epidemiology of constipation in children
519 and adults: a systematic review. *Best Pract Res Clin Gastroenterol*. 2011;25: 3–
520 18.
- 521 22. Pulgar S, Bains S, Gooch J, Chambers H, Noritz GH, Wright E, et al. Prevalence,
522 Patterns, and Cost of Care for Children with Cerebral Palsy Enrolled in Medicaid
523 Managed Care. *J Manag Care Spec Pharm*. 2019;25: 817–822.
- 524 23. Fiori S, Staudt M, Boyd RN, Guzzetta A. Neural Plasticity after Congenital Brain
525 Lesions. *Neural Plast*. 2019;2019: 9154282.
- 526 24. Bracciali LMP, Almeida VS, Sankako AN, Silva MZ, Bracciali AC, Carvalho SMR,
527 et al. Translation and validation of the Brazilian version of the Cerebral Palsy
528 Quality of Life Questionnaire for Children – child report. *Jornal de Pediatria*
529 (Versão em Português). 2016. pp. 143–148. doi:10.1016/j.jpedp.2015.12.005
- 530 25. Ickes SB, Hurst TE, Flax VL. Maternal Literacy, Facility Birth, and Education Are
531 Positively Associated with Better Infant and Young Child Feeding Practices and
532 Nutritional Status among Ugandan Children. *The Journal of Nutrition*. 2015. pp.
533 2578–2586. doi:10.3945/jn.115.214346
- 534 26. Liu H-Y, Chen J-R, Hsiao S-Y, Huang S-T. Caregivers’ oral health knowledge,
535 attitude and behavior toward their children with disabilities. *J Dent Sci*. 2017;12:
536 388–395.
- 537 27. Sullivan AB, Miller D. Who is Taking Care of the Caregiver? *J Patient Exp*. 2015;2:
538 7–12.

- 539 28. Barros ALO, de Gutierrez GM, Barros AO, Santos MTBR. Quality of life and
540 burden of caregivers of children and adolescents with disabilities. Spec Care
541 Dentist. 2019;39: 380–388.
- 542 29. Fairhurst C, Shortland A, Chandler S, Will E, Scrutton D, Simonoff E, et al. Factors
543 associated with pain in adolescents with bilateral cerebral palsy. Developmental
544 Medicine & Child Neurology. 2019. pp. 929–936. doi:10.1111/dmcn.14113
- 545 30. Tan SS, van Gorp M, Voorman JM, Geytenbeek JJ, Reinders-Messelink HA,
546 Ketelaar M, et al. Development curves of communication and social interaction in
547 individuals with cerebral palsy. Dev Med Child Neurol. 2019.
548 doi:10.1111/dmcn.14351
- 549 31. Devinsky O, Vezzani A, O'Brien TJ, Jette N, Scheffer IE, de Curtis M, et al.
550 Epilepsy. Nature Reviews Disease Primers. 2018. doi:10.1038/nrdp.2018.25
- 551 32. Kwan P, Arzimanoglou A, Berg AT, Brodie MJ, Allen Hauser W, Mathern G, et al.
552 Definition of drug resistant epilepsy: Consensus proposal by the ad hoc Task
553 Force of the ILAE Commission on Therapeutic Strategies. Epilepsia. 2009. pp.
554 1069–1077. doi:10.1111/j.1528-1167.2009.02397.x
- 555 33. Blond BN, Detyniecki K, Hirsch LJ. Assessment of Treatment Side Effects and
556 Quality of Life in People with Epilepsy. Neurologic Clinics. 2016. pp. 395–410.
557 doi:10.1016/j.ncl.2015.11.002
- 558 34. Hyland NP, Cryan JF. A Gut Feeling about GABA: Focus on GABA(B) Receptors.
559 Front Pharmacol. 2010;1: 124.
- 560 35. Ming X, Chen N, Ray C, Brewer G, Kornitzer J, Steer RA. A Gut Feeling: A
561 Hypothesis of the Role of the Microbiome in Attention-Deficit/Hyperactivity
562 Disorders. Child Neurol Open. 2018;5: 2329048X18786799.
- 563 36. Finoti LS, Nepomuceno R, Pigossi SC, Corbi SC, Secolin R, Scarel-Caminaga
564 RM. Association between interleukin-8 levels and chronic periodontal disease: A
565 PRISMA-compliant systematic review and meta-analysis. Medicine . 2017;96:
566 e6932.
- 567 37. Batool H, Nadeem A, Kashif M, Shahzad F, Tahir R, Afzal N. Salivary Levels of IL-
568 6 and IL-17 Could Be an Indicator of Disease Severity in Patients with Calculus
569 Associated Chronic Periodontitis. Biomed Res Int. 2018;2018.
570 doi:10.1155/2018/8531961
- 571 38. Ertugrul AS, Sahin H, Dikilitas A, Alpaslan N, Bozoglan A. Comparison of CCL28,
572 interleukin-8, interleukin-1 β and tumor necrosis factor-alpha in subjects with
573 gingivitis, chronic periodontitis and generalized aggressive periodontitis. Journal of
574 Periodontal Research. 2013. pp. 44–51. doi:10.1111/j.1600-0765.2012.01500.x
- 575 39. Gamonal J, Acevedo A, Bascones A, Jorge O, Silva A. Characterization of cellular
576 infiltrate, detection of chemokine receptor CCR5 and interleukin-8 and RANTES
577 chemokines in adult periodontitis. Journal of Periodontal Research. 2001. pp. 194–
578 203. doi:10.1034/j.1600-0765.2001.360309.x
- 579 40. Costa FO, Cortelli SC, Silva TA, Costa AA, Lima RPE, Cortelli JR, et al. Cytokine
580 levels in crevicular fluid associated with compliance during periodontal
581 maintenance therapy. Clin Oral Investig. 2019;23: 3517–3526.

- 582 41. Zhu L, Liu W, Alkhouri R, Baker RD, Bard JE, Quigley EM, et al. Structural
583 changes in the gut microbiome of constipated patients. *Physiol Genomics*.
584 2014;46: 679–686.
- 585 42. Khalif IL, Quigley EMM, Konovitch EA, Maximova ID. Alterations in the colonic
586 flora and intestinal permeability and evidence of immune activation in chronic
587 constipation. *Dig Liver Dis*. 2005;37: 838–849.
- 588 43. Meinds RJ, van Meegdenburg MM, Trzpis M, Broens PMA. On the prevalence of
589 constipation and fecal incontinence, and their co-occurrence, in the Netherlands.
590 *Int J Colorectal Dis*. 2017;32: 475–483.
- 591 44. Li Q, Han Y, Dy ABC, Hagerman RJ. The Gut Microbiota and Autism Spectrum
592 Disorders. *Front Cell Neurosci*. 2017;11: 120.
- 593 45. Jyonouchi H, Sun S, Le H. Proinflammatory and regulatory cytokine production
594 associated with innate and adaptive immune responses in children with autism
595 spectrum disorders and developmental regression. *Journal of Neuroimmunology*.
596 2001. pp. 170–179. doi:10.1016/s0165-5728(01)00421-0
- 597 46. Romeo DM, Sini F, Brogna C, Albamonte E, Ricci D, Mercuri E. Sex differences in
598 cerebral palsy on neuromotor outcome: a critical review. *Dev Med Child Neurol*.
599 2016;58: 809–813.
- 600 47. Chen M, Li T, Lin S, Bi D, Zhu D, Shang Q, et al. Association of Interleukin 6 gene
601 polymorphisms with genetic susceptibilities to spastic tetraplegia in males: A case-
602 control study. *Cytokine*. 2013. pp. 826–830. doi:10.1016/j.cyto.2013.01.011
- 603 48. Nakagawa S, Fujii H, Machida Y, Okuda K. A longitudinal study from prepuberty to
604 puberty of gingivitis. Correlation between the occurrence of *Prevotella intermedia*
605 and sex hormones. *J Clin Periodontol*. 1994;21: 658–665.
- 606 49. Hu L, Mauro TM, Dang E, Man G, Zhang J, Lee D, et al. Epidermal Dysfunction
607 Leads to an Age-Associated Increase in Levels of Serum Inflammatory Cytokines.
608 *J Invest Dermatol*. 2017;137: 1277–1285.
- 609 50. Ferreira MC, Pastore C, Imada R, Guaré R, Leite M, Poyares D, et al. Autonomic
610 nervous system in individuals with cerebral palsy: a controlled study. *J Oral Pathol*
611 *Med*. 2011;40: 576–581.
- 612 51. Meyle J, Chapple I. Molecular aspects of the pathogenesis of periodontitis.
613 *Periodontol 2000*. 2015;69: 7–17.
- 614 52. Huang N, Gibson FC 3rd. Immuno-pathogenesis of Periodontal Disease: Current
615 and Emerging Paradigms. *Curr Oral Health Rep*. 2014;1: 124–132.
- 616 53. Cekici A, Kantarci A, Hasturk H, Van Dyke TE. Inflammatory and immune
617 pathways in the pathogenesis of periodontal disease. *Periodontology 2000*. 2014.
618 pp. 57–80. doi:10.1111/prd.12002
- 619 54. Rudick CP, Lang MS, Miyamoto T. Understanding the pathophysiology behind
620 chairside diagnostics and genetic testing for IL-1 and IL-6. *Oral Dis*. 2019.
621 doi:10.1111/odi.13030
- 622 55. Degasperis GR, Etchegaray A, Marcelino L, Sicard A, Villalpando K, Pinheiro SL.
623 Periodontal disease: General aspects from biofilm to the immune response driven

by periodontal pathogens. *Advances in Microbiology*. 2018;8: 1.

56. Rezaei F, Mozaffari HR, Tavasoli J, Zavattaro E, Imani MM, Sadeghi M. Evaluation of Serum and Salivary Interleukin-6 and Interleukin-8 Levels in Oral Squamous Cell Carcinoma Patients: Systematic Review and Meta-Analysis. *J Interferon Cytokine Res*. 2019. doi:10.1089/jir.2019.0070
57. Zhang Q, Chen B, Yan F, Guo J, Zhu X, Ma S, et al. Interleukin-10 inhibits bone resorption: a potential therapeutic strategy in periodontitis and other bone loss diseases. *Biomed Res Int*. 2014;2014: 284836.

Figure Legends:

Table 1. Population-Wide Demographics Table. Numbers and relevant statistics for study groups. Groups are defined as G1 (CP with constipation), G2 (CP without constipation), G3 (control with constipation), and G4 (control without constipation).

Figure 1. Distribution patterns of the CP study population. **A)** Subjects from the cohort aged from 3-19, with a mean age of 9.57 ± 4.20 years (n=93). **B)** The study population was composed of healthy individuals (controls, 32.26%), tetraplegics (32.26%), diplegics (26.88%), and hemiplegics (9.68%). **C)** Distribution is measured along the GMFCS scale from I-V. Stages I-III are considered ambulatory, while stages IV and V require significant assistance (typically a wheelchair or mobility scooter). Stages II and III are less represented (3.23% and 6.45% respectively), while stages I (37.63%), IV (29.03%) and V (23.66%) make up the vast majority of our population. **D)** Clustering shows that hemiplegics and healthy individuals are more likely to be ambulatory, most without assistance. However, diplegic and tetraplegic individuals are by definition not ambulatory, due to the increased muscle tone. Therefore, diplegics are frequently classified as GMFCS stage IV and tetraplegics as stage V.

Figure 2. Effects of CPCHILD Subscores on Overall QOL. Distributions of each individual CPCHILD subscore A) Positioning, Transferring, and Mobility; B) Hospitalization and Caregiver-Reported Health; C) Caregiver-Reported Quality of Life; D) Comfort and Emotions; E) Personal Care; and F) Communication and Social Interaction) is measured on the x-axes, while the y-axis marks the average of the remaining subscores. A lighter green represents a lower density within the distribution, while a darker blue represents a higher density.

Figure 3. Correlation of Clinical Factors in CP patients. A) Each KDE plot shows clustering by the type of medication taken by subjects. Green figures show constipated subjects, blue figures show non-constipated subjects, and Green-to-Blue figures show the full population. Constipation is measured and compared to the medication types. In **B-D)** QOLS is scored from 0-100 based on the average of CPCHILD subscores. **E-G)** gingival bleeding is measured by the percentage of teeth after periodontal probing.

Figure 4. Factors impacting gingival bleeding. A) Violin plots depicting gingival bleeding patterns stratified by gender. Box plot within the violin represent the inner quartiles, and the whiskers show the 5% confidence interval (CI). KDE clustering demonstrating gingival bleeding in relation to age distribution. Data represented as means $\pm$ sd, and significant level at p-value < 0.05 (n=93). **C)** Gingival bleeding according to CP diagnosis. Boxes represent inner quartiles, center line represents the mean, while whiskers represent 5% CI (***, $P < 0.001$).

Figure 5. Constipation actions on Inflammatory Cytokine Levels. Distributions of each measured cytokine [A) TNF- α , B) IL-1 β , C) IL-6, D) IL-8, and E) IL-10] in constipated subjects (blue) and non-constipated subjects (orange) are mapped using a violin plot. White dots mark the means, bars show the inner quartiles, and whiskers mark the 5% confidence interval.

Figure 6. Correlation of Clinical Factors in CP patients. Correlogram of Pearson's correlation of clinical factors influencing CP patients. Each axis is labeled by relevant factors to form a lower-triangle correlogram. Each dot represents a significant correlation coefficient of two factors assigned ($P < 0.05$), where size and saturation represent the magnitude of the correlation, and the color itself represents whether the correlation is positive (blue) or negative (red), following the scale bar. Dots are overlaid by an X represent associations and are not significant regardless of the strength. (GB–Gingival Bleeding, CR-QOL–Caregiver-Reported Quality of Life, CE–Comfort and Emotions, CSI–Communication and Social Interaction, HCR-H–Hospitalization and Caregiver-Reported Health, GMFCS–Movement Ability, PC–Personal Care, PTM–Positioning, Transferring, and Mobility).

Supplementary Figure 1: Quality of Life association with specific cytokines. Linear Regressions of collected cytokine levels (TNF- α , IL-1 β , IL-6, IL-8, and IL-10 were levels were collected from subjects) and QOL. R- and p-values are displayed, and distribution for each axis is displayed in the opposite margins.

Supplementary Figure 2: Gingival bleeding association with specific cytokines. Linear Regressions of collected cytokine levels (TNF- α , IL-1 β , IL-6, IL-8, and IL-10 were levels were collected from subjects) and gingival bleeding. R- and p-values are displayed, and distribution for each axis is displayed in the opposite margins.

702

703

Ferreira et al.

Figures

Figure 1.

A)

B)

C)

D)

Figure 2.

Figure 3.

Figure 4.

A)

B)

C)

Figure 5.

A)

B)

C)

D)

E)

Figure 6.

Supplemental Figure 1.

Supplemental Figure 2.

TABLE 1.

	G1 - Constipated CP	G2-No Constipated CP	G3- Constipated Control ^a	G4- No Constipated Control ^a	p - value *
DEMOGRAPHICS					
Total Subjects	30 (32.25%)	33 (35.48%)	7 (7.52%)	23 (24.73%)	
Gender					
Female	19 (66.3%)	14 (42.4%)	4 (57.1%)	10 (43.5%)	0.332
Male	11 (36.7%)	19 (57.6%)	3 (42.9%)	13 (56.5%)	
Age (mean±SD)	8.5 ± 4.1	10.3 ± 3.7	8.8 ± 4.5	10.2 ± 4.3	0.291
Race					
White	19 (63.4%)	15(45.5%)	6(85.7%)	18(78.3%)	
Others	9(30%)	16(48.5%)	0(0%)	5(21.7%)	0.08
Black	2(6.4%)	2(6%)	1(14.3%)	0(0%)	
Occupation (Caregiver)					
Employed	11 (36,66%)	13 (39,39%)	5 (71,42%)	19 (82,60)	0.0017*
Education					
≤ 8 years	21 (70%)	23 (69.69%)	4 (57.15%)	14 (60.86%)	0.5628
> 8 years	9 (30%)	10 (30.31%)	3 (42.85%)	9 (39.14%)	
Family (Income)					
≥3 MW	4 (13.34%)	9 (27.28%)	3 (42.86%)	11 (47.82%)	0.018*
<3 MW	26 (86.66%)	24 (72.72%)	4 (57.14%)	12 (52.17%)	
ORAL HEALTH- PERIODONTAL PARAMETERS					
Gingival Bleeding	32.4 ± 19.8	4.1 ± 3.5	0.6 ±1.4	0.1 ± 0.6	<0.001*
Gingivitis					
Presence	23 (76.7%)	2 (6.1%)	0 (0%)	0 (0%)	<0.001*
Absence	7 (23.3%)	31 (93.9%)	7 (100%)	23 (100%)	
SYSTEMIC HEALTH- CEREBRAL PALSY					
Clinical Pattern					
Tetraplegia	20 (66.66%)	9 (27.27%)	0 (0%)	0 (0%)	0.0009*
Diplegia	10 (33.33%)	15 (45.45%)	0 (0%)	0 (0%)	
Hemiplegia	0 (0%)	9 (27.27%)	0 (0%)	0 (0%)	
GMFCS					
I - II - III	0 (0%)	14 (42.4%)	7 (100%)	23 (100%)	0.0002*
IV - V	30 (100%)	19 (57.6%)	0 (0%)	0 (0%)	
Medication					
GABA	5 (16.7%)	1 (3%)	0 (0%)	0 (0%)	
GABA +	9 (30%)	2 (6.1%)	0 (0%)	0 (0%)	
Sodium	5 (16.7%)	4 (12.1%)	0 (0%)	0 (0%)	0.0063*
Calcium	1 (3.3%)	1 (3%)	0 (0%)	0 (0%)	
No medication	10 (33.3%)	25 (75.8)	7 (100%)	23 (100%)	
Quality of Life					
Personal Care	79.8 ± 21.6A	46.4 ± 37.5B	20.0 ±23.8C	3.8 ± 8.4C	<0.05*
Positioning, Transfer ar	70.7 ± 26.1A	42.1 ± 34.4B	0.0 ±0.0C	0.1 ± 0.5C	<0.05*
Communication and So	44.4 ± 28.3A	17.9 ± 19.3B	0.0 ±0.0C	0.0 ±0.0C	<0.05*
Comfort, Emotions and	40.6 ± 22.7A	18.8 ± 16.00B	8.7 ±12.8C	3.9 ± 4.9C	<0.05*
Health	26.4 ± 18.2A	16.2 ± 12.9B	9.3±10.2C	3.5 ± 5.7C	<0.05*
Overall Quality of Life	44.3 ± 24.3A	29.7± 15.2B	28.6 ±19.5C	18.3 ± 20.8C	<0.05*
Domains Means	51.0 ± 13.1A	28.5 ±16.7B	11.1 ±7.5C	4.5 ±5.0C	<0.05*

^a Values represent mean ± SD or percentage (%); * p-values for group comparisons were significant at 0.05; Chi-square, ANOVA, One-Way, Kruskal-Wallis tests. Different letters denote statistically significant differences (p<0.05). Uppercase letters compare values horizontally (intragroup evaluation).

Abbreviations : SD-standard deviation; BOP-Bleeding on probing; G1-CP with constipation; G2-CP without constipation; G3- without CP with constipation; G4- without CP without constipation; MW- minimum wage, unit for measuring income in Brazil, broadly corresponding to \$251 U.S. during the study period