Tracking the spread of novel coronavirus (2019-nCoV) based on big data

Zhao, Xumao^[1]; Liu, Xiang^[1]; Li, Xinhai^{[2,3],*}

^[1] State Key Laboratory of Grassland Agro-Ecosystem/Institute of Innovation Ecology, Lanzhou University, Lanzhou, 730020, China

^[2] Key Laboratory of Animal Ecology and Conservation Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing, 100101, China

^[3] University of Chinese Academy of Sciences, Yuquan Road, Beijing 100049, China

*Correspondence to: lixh@ioz.ac.cn ; zhaoxm@lzu.edu.cn

Abstract: The novel coronavirus (2019-nCoV) appeared in Wuhan in late 2019 have infected 34,598 people, and killed 723 among them until 8th February 2020. The new virus has spread to at least 316 cities (until 1st February 2020) in China. We used the traffic flow data from Baidu Map, and number of air passengers who left Wuhan from 1st January to 26th January, to quantify the potential infectious people. We developed multiple linear models with local population and air passengers as predicted variables to explain the variance of confirmed cases in every city across China. We found the contribution of air passengers from Wuhan was decreasing gradually, but the effect of local population was increasing, indicating the trend of local transmission. However, the increase of local transmission is slow during the early stage of novel coronavirus, due to the super strict control measures carried out by government agents and communities.

Key words: Baidu Map, big data, quarantine, pandemic, traffic flow

Introduction

Since the first case of pneumonia named 2019 Novel Coronavirus (2019-nCoV) being identified in in Wuhan, China, a total of 34,598 confirmed infections were reported in the country until 8th February 2020, which caused 723 deaths (1). Guo et al (2020) have found that 2019-nCoVs was similar with bat coronaviruses (2), which occasionally transmitted to human due to wildlife consumption. Understanding the pattern of transmission characteristics of 2019-nCoV is important for effective

preventing and controlling the ongoing pandemic disease. The value of R_0 (basic reproduction number) was estimated as 2.2, inferring a median size outbreak (3). However, based on the epidemic transmission model, the number of actual infections would be much larger than the number of confirmed cases (4).

At present, tracking the passengers from Wuhan in January 2020 is still the top task for preventing the further spread of novel coronavirus (2019-nCoV). To accurately estimate the risk of the novel coronavirus, we compiled the detailed daily traffic data outbound Wuhan from a big-data source, Baidu Map, before the lockdown of Hubei Province, in order to provide information for risk assessment of 2019-nCoV at the province level and the city level (Supp. Fig. 1).

Methods

The traffic flow data outbound Wuhan from 1st January to 26th January 2020 was downed from Baidu Map Huiyan platform (5). The number of air passengers from Wuhan from 30th December 2019 to 20 January 2020 was released by Aviationtalk (6). The time series data of confirmed 2019-nCoV cases from 10th January to 30 January 2020 was obtained from People's daily-Dingxiangyuan (1), which was released by China National Health Commission.

We did Spearman correlation analysis for the daily traffic from Wuhan (from 1st January to 26th January) and the total traffic in this period with the number of confirmed cases (from 25th January to 30th January). To explain the variance of confirmed cases in all infected provinces, we developed multiple linear models including population, GDP, population density, and mean temperature as independent variables. All analysis was performed using R (version 3.6.2).

Results

From 20th December 2019 to 20 January 2020, 854,424 air passengers left Wuhan Tianhe Airport to 49 cities in China (Fig .1). From 1st to 26th January, about three million domestic passengers travelled from Wuhan to other cities. Among the passengers, a few thousands had been confirmed to infected by the novel coronavirus (Fig. 2). The distribution of air passengers from Wuhan to other cities in China had high correlation coefficients (0.71) with the number of confirmed infection cases in those cities on 22nd January. The correlation coefficient drops to 0.56 on 24th January. Then the number of confirmed infection cases was positive correlation with local

62 population size.

63 We used a multiple regression model to explain the variance of the number of
64 cases in the infected cities. After model selection, only two variables remained, the
65 number of passengers and local population (Fig. 3). Overall, the population of the
66 provinces explains near half of the variance in the number of confirmed cases across
67 34 provinces and province-level municipalities, whereas the number of passengers
68 from Wuhan explained around 10% (Fig. 3).

69 Correlation coefficients of number of confirmed that the number of cases in the
70 cities ($n=97$) from 25th January to 30th January match the number of passengers from
71 Wuhan during the period from 1st January to 26th January. The highest correlation
72 appears on 5th January, inferring a long incubation period up to two weeks (Supp.
73 Table 1).

74 Discussion

75 In the beginning of the spread of the pneumonia, there is a high correlation (0.71)
76 between the number of confirmed infection cases and air passengers from Wuhan,
77 proofed Wuhan the source of the pneumonia (7). As time going, local population
78 played a more dominant role, because the local spread of 2019-nCoV is likely to
79 happen. The basic reproductive number of the infection (R_0) to be estimated as 3.8
80 means 72-75% of transmissions must be prevented to stop the outbreak (4).
81 Fortunately, the transmission of the virus was really controlled due to strict prevention
82 measures carried out by Chinese government. Restricted population movements ban
83 was enforced upon 16 cities in Hubei Province since 23rd January 2020 (8), resulting
84 in significant decrease in passengers from Wuhan and adjacent cities, which
85 effectively reduces the spread of the pneumonia. However, 3-5 million people had left
86 Wuhan for numerous cities in China before the province lockdown (Supp. Fig.2), and
87 among them a number of infected people have no clinical symptom yet infectious to
88 others. We believe this is the highest challenge against the current national level
89 antiviral campaign.

90 Currently the first-level response to major public health emergencies has been
91 initiated in 30 provinces, municipalities and autonomous regions in China on 25th
92 January 2020 (9), so that strict control procedures are carried out to prevent the spread
93 of the virus. According to an infectious disease model, it was estimated that the actual

94 number of infected people in Wuhan on 4th February 2020 may reach 250 thousand
 95 (prediction interval, 164,602 to 351,396) without any control (3). Whereas only 5,142
 96 confirmed infections were reported in China on 2nd February 2020. Starting from 3rd
 97 February, all suspected case in Wuhan will be taken in to medical care, and the virus
 98 spread in Wuhan can be controlled gradually.

99 The correlation coefficients between the number of confirmed cases from 26th
 100 January to 30th January and number of passengers from Wuhan were higher than that
 101 on 25th January (Supp. Table 1). We think the reason is that the confirmed cases on
 102 25th is too low due to lack of virus detection kit. After 25th January the supply of virus
 103 detection kit was enough and the number of confirmed cases reflect the real situation,
 104 which have very high correlation with the number of passengers from Wuhan to these
 105 cities during 1st January to 26th January. We notice the highest correlation appear on
 106 5th January, two to three weeks ahead of the confirmed cases in those cities
 107 (confirmation also needs several days to complete at that time), which infers the long
 108 latent period of 2019-nCoV.

109 Combining the two variables (local population and passengers from Wuhan) to
 110 interpret the virus outbreak risk, we recommend that the preventing and controlling
 111 measures can be divided into two different stages. In the early stage, checking the
 112 passengers from Wuhan is more important. Some cities (e.g., Wenzhou in Zhejiang
 113 Province) with a large number of returnees from Wuhan have many cases even when
 114 they are far from Wuhan in space. The database of travel routes of confirmed patients
 115 has been developed and published for free use (10), in order to tracking and warning
 116 close contactors. In the late stage, local population become the most important factor
 117 in predicting the number of confirmed infection cases. This suggests us that the
 118 densely populated metropolitan areas, such as Shanghai, Beijing, Guangzhou, and
 119 Shenzhen should pay special attentions to preventing the second-generation infections.
 120 Moreover, the densely populated rural areas around Wuhan may face double threats of
 121 the spread of infectious people from Wuhan and a large number of local susceptible
 122 people. These areas, including areas of Hubei Province except for Wuhan, and
 123 surrounding area of Henan (Nanyang, Xinyang) and Chongqing, need to prepare for a
 124 surge in infection. The rural medical facilities in these areas are scarcer than in cities,
 125 clustering cases are more likely to happen in these places.

Fig. 1 The distribution of 854,424 air passengers from Wuhan Tianhe Airport to 49 cities in China during the period from 30th December 2019 to 20th January 2020. The sizes of the points indicate the number of passengers, which are also represented in color series: blue, green, yellow, brown, and red.

130 Fig. 2 The spatial distribution of 24,281 confirmed cases (incomplete data) of 2019-nCoV on 5th
 131 February in 299 cities in China. The sizes of red points represent the number of confirmed cases.
 132 The sizes of grey points show the population of the cities.

133 Fig. 3 The proportion of explained variance in linear regression: Number of confirmed cases in 34
 134 provinces ~ Air passengers from Wuhan * Population of the province for the date from 24th
 135 January to 1st February.

136 Supplementary documents

137 Supp. Fig. 1 The daily passenger (include all vehicles) numbers from Wuhan to top 50 cities in
 138 China from 1st January to 26th January are showed by an animated GIF file
 139 2019-nCoV_spread_1-26 Jan.gif (Supp. Fig. 1).

143 Supp. Fig. 3 Estimated daily number of passengers from Wuhan to 12 major cities in China (upper
144 panel) and 12 major cities in Hubei Province (lower panel) base on Baidu Bigdata server. The red
145 lines show the daily accumulated confirmed case in each city from 25th January to 30th January.

146 References

- 147 (1) Dingxiangyuan: https://ncov.dxy.cn/ncovh5/view/pneumonia_peopleapp?scene=126&clicktime=1579832412&from=timeline&isappinstalled=0
- 148
- 149 (2) Guo Q, Li M, Wang C, Wang P, Fang Z, Tian J, Wu S, Xiao Y, Zhu H. Host and infectivity
- 150 prediction of Wuhan 2019 novel coronavirus using deep learning algorithm. 2020, bioRxiv.
- 151 <http://dx.doi.org/10.1101/2020.01.21.914044>
- 152 (3) Riou J, Althaus CL. Pattern of early human-to-human transmission of Wuhan 2019-nCoV.
- 153 2020, bioRxiv. <http://dx.doi.org/10.1101/2020.01.23.917351>
- 154 (4) Read JM, Bridgen JRE, Cummings DAT, Ho A, Jewell C. Novel coronavirus 2019-nCoV:
- 155 early estimation of epidemiological parameters and epidemic predictions. 2020, medRxiv.
- 156 (5) Baidu Map Huiyan platform: <https://qianxi.baidu.com/?city=420100> (in Chinese)
- 157 (6) Aviationtalk : <https://mp.weixin.qq.com/s/7ynWYxB-s7nfz7rmjBLSpQ> (in Chinese)
- 158 (7) Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R, Niu P, Zhan
- 159 F. A novel coronavirus from patients with pneumonia in China, 2019. 2020, New England
- 160 Journal of Medicine. DOI: 10.1056/NEJMoa2001017.
- 161 (8) China news : <http://www.chinanews.com/gn/2020/01-25/9069668.shtml> (in Chinese)
- 162 (9) Xinhua net: http://www.xinhuanet.com/2020-01/26/c_1125503530.htm (in Chinese)
- 163 (10) Close contactor detection tool:
- 164 <http://2019ncov.nosugartech.com/?from=timeline&isappinstalled=0> (in Chinese)

Jan. 1