Clinical characteristics of 50466 patients with 2019-nCoV infection

Pengfei Sun¹, Shuyan Qie², Zongjan Liu², Jizhen Ren³, Jianing Xi^{□4}

¹Department of Plastic Surgery, Zibo Central Hospital, 255000, Zibo, China

²Department of Rehabilitation, Beijing Rehabilitation Hospital of Capital Medical University, 100044, Beijing, China

³Department of Plastic Surgery, Affiliated Hospital of Qingdao University, 266000, Qingdao, China

⁴Department of Respiratory Rehabilitation, Beijing Rehabilitation Hospital of Capital Medical University, 100044, Beijing, China

□ Corresponding author

Abstract

Objective: We aim to summarize reliable evidences of evidence-based medicine for the treatment and prevention of the 2019 novel coronavirus (2019-nCoV) by analyzing all the published studies on the clinical characteristics of patients with 2019-nCoV.

Methods: PubMed, Cochrane Library, Embase, and other databases were searched. Several studies on the clinical characteristics of 2019-nCoV infection were collected for Meta-analysis.

Results: Ten studies were included in Meta-analysis, including a total number of 50466 patients with 2019-nCoV infection. Meta-analysis shows that, among these patients, the incidence of fever was 89.1%, the incidence of cough was 72.2%, and the incidence of muscle soreness or fatigue was 42.5%. The incidence of acute respiratory distress syndrome (ARDS) was 14.8%, the incidence of abnormal chest computer tomography (CT) was 96.6%, the percentage of severe cases in all infected cases was 18.1%, and the case fatality rate of patients with 2019-nCoV infection was 4.3%.

Conclusion: Fever and cough are the most common symptoms in patients with 2019-nCoV infection, and most of these patients have abnormal chest CT examination. Several people have muscle soreness or fatigue as well as ARDS. Diarrhea, hemoptysis, headache, sore throat, shock, and other symptoms only occur in a small number of patients. The case fatality rate of patients with 2019-nCoV infection is lower than that of Severe Acute Respiratory Syndrome (SARS) and Middle East Respiratory Syndrome (MERS).

Keywords 2019-nCoV, Coronavirus, Meta-analysis, Clinical Symptoms

1 Introduction

Since December 2019, the epidemic of 2019 novel coronavirus (2019-nCoV) infectious pneumonia in Wuhan, China. The Chinese government and researchers have taken rapid measures to control the epidemic ^[1]. On January 30, 2020, WHO declared that the epidemic of 2019-nCoV was a public health emergency of international concern (PHEIC). At present, the number of patients with 2019-nCoV infection is still rising, and its harm to human beings has exceeded the outbreak of Severe Acute Respiratory Syndrome (SRAS) in China, 2002 ^[2]. The clinical characteristics and the case fatality rate after 2019-nCoV infection have always been concerned by people, and it is also the focus of medical workers' research at present. However, due to the different design of different clinical studies and insufficient sample size, the conclusions of the published studies were different. In order to acquire

more accurate conclusions on the clinical characteristics and mortality of patients with 2019-nCoV infection, we searched the relevant literatures and carried out single arm Meta-analysis^[3]. Our findings provide important guidance for current clinical work on the prevention and treatment of 2019-nCoV infection.

2 Material and Methods

2.1 Search strategy

Three popular medical databases including PubMed, Cochrane Library, and Embase databases were searched for related literatures, using the following keywords: "2019-nCoV", "Coronavirus", "COVID-19", "SARS-CoV-2" and "Wuhan Coronavirus". Articles reviewed were dated up to February 2020. In this Meta-analysis, there was no language restriction. Only available data from published articles were collected. Data from unpublished papers were not included.

2.2 The inclusive and exclusive criteria

2.2.1 Inclusive criteria: studies that include randomized controlled trials, non-randomized controlled trials, case-control studies, cohort studies, cross-sectional studies, and also case reports on the clinical characteristics of patients with 2019-nCoV infection. Clinical characteristics of patients with 2019-nCoV infection include fever, cough, muscle soreness or fatigue, acute respiratory distress syndrome (ARDS), abnormal chest computer tomography (CT) detection, patients in critical condition, as well as death due to 2019-nCoV infection.

2.2.2 Exclusive criteria: articles were published repeatedly; studies did not include the research indicators needed for Meta-analysis; research data were missing.

2.3 Data extraction and paper quality evaluation

First of all, we screened the literature according to the literature abstract, excluding the articles that obviously do not meet the inclusive criteria, and then read the full article for re-screening. If any disagreement on the choice of the literature exists, a third evaluator will join to make the decision. All included literature were evaluated using the Newcastle-Ottawa Scale (NOS) ^[4]. The highest quality of the literature was 9 stars and the lowest 0 stars.

2.4 Statistical analysis

The statistical software *Stata* version 12.0 was used to carry out the single arm Meta-analysis. Original data included in the literature were first transformed by double arcsine method to make them conform to normal distribution and then analyzed in *Stata*. The initial conclusion obtained by Meta-analysis was then restored using formula $(P=(\sin(tp/2))^2)$ to reach final conclusion. In order to objectively evaluate the publication bias of the included literature, the Egger test with $P < 0.05$ as the existence of publication bias was performed, the values larger than which were considered as no publication bias.

3 Results

3.1 Literature inclusion

A total of 284 articles were retrieved, among which 39 papers were removed due to repeated retrieval, 212 papers were removed after reading abstracts, 23 were eliminated after reading the full text. At the end, a total of 10 articles ^[5-14] were included in this Meta-analysis, including data from 50466 patients. The specific operation flow is shown in Figure 1. The characteristics of the literature are shown in Table 1.

3.2 Meta-analysis results

Through Meta-analysis, we found that among all the clinical characteristics of patients with 2019-nCoV infection, the incidence of fever was 89.1% (Figure 2), that of cough was 72.2% (Figure 3), and that of muscle soreness or fatigue was 42.5% (Figure 4). The incidence of ARDS was 14.8% (Figure 5), correspondingly that of abnormal chest CT detection was 96.6% (Figure 6), severe cases in all infected cases occupied a percentage of 18.1% (Figure 7). The case fatality rate of patients with 2019-nCoV infection was 4.3% (Figure 8). Detailed results of Meta-analysis are shown in Table 2.

3.3 Publication bias detection

The results of Egger test in Table 3. There was publication bias in the Meta-analysis of the ARDS group($P=0.008$)..

4 Discussion

2019-nCoV is one type of coronaviruses which belongs to β -coronavirus cluster. It causes the third kind of zoonotic coronavirus disease after Severe Acute Respiratory Syndrome (SARS) and Middle East Respiratory Syndrome (MERS). Wei Ji *et al.* ^[15] found that 2019-nCoV seemed to be a recombinant virus between bat coronavirus and coronavirus of another unknown origin. Kangpeng Xiao *et al.* ^[16] confirmed that malayan pangolins were the most likely intermediate host of 2019-nCoV. The research of Domenico Benvenuto *et al.* ^[17] found that 2019-nCoV was only closely related to the coronavirus isolated from chrysanthemum-headed bat in 2015. Their research suggests that bats may be the reservoir host for 2019-nCoV. According to the research by Peng Zhou *et al.* ^[18] and Fan Wu *et al.* ^[19], it was found that the sequence homology between 2019-nCoV and SARS-CoV was 79.5%. The homology between 2019-nCoV and bat coronavirus at the genetic level was 96%. Therefore, very likely it can be confirmed that 2019-nCoV comes from bats.

According to the results of Meta-analysis, we find that the most common symptoms of patients with 2019-nCoV infection were fever and cough, and most of the patients had abnormal chest CT. Some patients with 2019-nCoV infection had muscle soreness or fatigue, ARDS, and other symptoms. Diarrhea, hemoptysis, headache, sore throat,

shock, and other symptoms only occurred among a small number of patients. The proportion of severe cases in all infected cases was 18.1%, and the case fatality rate was 4.3%. At present, the Chinese government has announced that the case fatality rate of patients with 2019-nCoV infection in China is 3.36%, which is lower than that of another two widely contagious zoonotic coronavirus diseases, SARS and MERS. Infection mortality rates of SARS and MERS are 9.6%^[20] and 35%^[21], respectively.

Gaohong Sheng *et al.*^[22] have shown that viral infection could increase the risk of pulmonary fibrosis. Xie *et al.*^[23] found that 45% of patients showed signs of pulmonary fibrosis within one month after being infected with SARS-CoV. Hui *et al.*^[24] revealed that 36% and 30% of patients infected with SARS-CoV developed pulmonary fibrosis 3 and 6 months after infection. These studies consistently suggest that pulmonary fibrosis will become one of the serious complications in patients with 2019-nCoV infection. How to prevent and reduce the occurrence of pulmonary fibrosis in patients with 2019-nCoV infection are urgent problems for medical workers in the treatment of 2019-nCoV.

Yu Zhao *et al.* ^[25] found that angiotensin converting enzyme 2 (ACE2) was the receptor of 2019-nCoV. In normal lung tissue, ACE2 is mainly expressed by type I and type II alveolar epithelial cells. It was reported that 83% of II-type alveolar cells expressed ACE2. Therefore, 2019-nCoV infection causes damages to most II type alveolar cells. In addition, the use of mechanical ventilation in the treatment of patients with 2019-nCoV infection can also aggravate the injury of alveolar cells. After alveolar cell injury, transforming growth factor- β (TGF- β) released in tissue promotes the lung repair. Virus infection often leads to excessive activation of TGF- β pathway, which leads to the occurrence of pulmonary fibrosis.

Based on the mechanism of pulmonary fibrosis after virus infection, we can formulate corresponding prevention and treatment proposals. The first is the treatment of 2019-nCoV infection. At present, a variety of drugs have been found to inhibit 2019-nCoV. Michelle L. Holshue *et al.* ^[26] achieved remarkable results in the treatment of patients with 2019-nCoV infection with remdesivir. This drug is currently in clinical trials. Second, it is necessary

to inhibit the inflammatory reaction and reduce the exudation. On this direction, antibiotics not only have traditional antibacterial effects, but also have immunomodulatory and anti-inflammatory properties ^[27]. Macrolide antibiotics such as clarithromycin, azithromycin, and erythromycin prevent the production of pro-inflammatory cytokines and immune mediators ^[28]. The third is to inhibit the fibrogenic effect of TGF- β . Luo *et al.* ^[29] found that arsenic trioxide could inhibit the signal transduction of TGF- β and play a role in anti-fibrosis. The fourth is the rational use of ventilator to avoid unnecessary lung damages. In addition, pirfenidone and nintedanib as drugs for the treatment of idiopathic pulmonary fibrosis can also be used to prevent and treat pulmonary fibrosis in patients with 2019-nCoV infection ^[30].

In all, a total of 10 articles were covered in the Meta-analysis, including 50466 patients with 2019-nCoV infection. By far, it is the first Meta-analysis with the largest sample size. The quality of the literature included in this study is high, the analysis is rigorous, and the conclusions drawn by the study are highly credible. However, this Meta-analysis also has some limitations: 1) All studies included in this Meta-analysis are retrospective studies

with large heterogeneity; 2) Most patients in our Meta-analysis are Chinese, and we aimed to use the conclusions of this study to predict patients in general, including other countries and races; 3) There was publication bias in the Meta-analysis of the ARDS group. The analytical conclusion of the ARDS group may be influenced by publication bias. Therefore, based on the above limitations, the conclusions of this Meta-analysis still need to be verified by more relevant studies with more careful design, more rigorous execution, and larger sample size.

Contributors: Jianing Xi and Pengfei Sun had the idea for and designed the study and had full access to all data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Pengfei Sun and Shuyan Qie contributed to writing of the report. Jianing Xi contributed to critical revision of the report. Shuyan Qie, Zongjan Liu and Jizhen Ren contributed to the statistical analysis. All authors contributed to data acquisition, data analysis, or data interpretation, and reviewed and approved the final version.

Acknowledgements: The authors declare no conflict of interest in this study. We acknowledge TopEdit LLC for the linguistic editing and proofreading during the preparation of this manuscript.

References

- [1] Lancet T. Emerging understandings of 2019-nCoV. Lancet 2020. 395(10221): 311.
- [2] Lu H, Stratton CW, Tang YW. Outbreak of Pneumonia of Unknown Etiology in Wuhan China: the Mystery and the Miracle. J Med Virol 2020. doi: 10.1002/jmv.25678.
- [3] Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. J Clin Epidemiol 2009. 62(10):1006-1012.
- [4] Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. Eur J Epidemiol 2010. 25(9):603-605.
- [5] Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet 2020. pii: S0140-6736(20)30183-5.
- [6] Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. JAMA 2020. doi: 10.1001/jama.2020.1585.

- [7] Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. Lancet 2020. pii: S0140-6736(20)30211-7.
- [8] Weijie Guan, Zhengyi Ni, Yu Hu, et al. Clinical characteristics of 2019 novel coronavirus infection in China, medRxiv 2020. doi: <https://doi.org/10.1101/2020.02.06.20020974>.
- [9] Chen L, Liu HG, Liu W, et al, 2020. Analysis of clinical features of 29 patients with 2019 novel coronavirus pneumonia. Zhonghua Jie He He Hu Xi Za Zhi.43(0):E005.
- [10] Kaiyuan Sun, Jenny Chen, Cecile Viboud. Early epidemiological analysis of the 2019-nCoV outbreak based on a crowdsourced data. medRxiv 2020. doi: <https://doi.org/10.1101/2020.01.31.20019935>.
- [11] Yang Yang, Qingbin Lu, Mingjin Liu, et al. Epidemiological and clinical features of the 2019 novel coronavirus outbreak in China. medRxiv 2020. doi: <https://doi.org/10.1101/2020.02.10.20021675>.
- [12] Jie Li, Shilin Li, Yurui Cai, et al. Epidemiological and Clinical Characteristics of 17 Hospitalized Patients with 2019 Novel Coronavirus Infections Outside Wuhan, China. medRxiv 2020. doi: <https://doi.org/10.1101/2020.02.11.20022053>.

- [13] The Noval Coronavirus Pneumonia Emergency Response Epidemiology Team, Chinese Center for Disease Control and Prevention. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. Zhong Hua Liu Xing Bing Xue Za Zhi 2020. 41(2):145-151.
- [14] Xu XW, Wu XX, Jiang XG, et al. Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-Cov-2) outside of Wuhan, China: retrospective case series. BMJ 2020;368:m606.
- [15] Ji W, Wang W, Zhao X, et al. Homologous recombination within the spike glycoprotein of the newly identified coronavirus may boost cross-species transmission from snake to human. J Med Virol 2020. doi: 10.1002/jmv.25682.
- [16] Kangpeng Xiao, Junqiong Zhai, Yaoyu Feng, et al. Isolation and Characterization of 2019-nCoV-like Coronavirus from Malayan Pangolins. medRxiv 2020. doi: <https://doi.org/10.1101/2020.02.17.951335>.
- [17] Benvenuto D, Giovannetti M, Ciccozzi A, et al. The 2019-new coronavirus epidemic: evidence for virus evolution. J Med Virol 2020.doi: 10.1002/jmv.25688.
- [18] Peng Zhou, XingLou Yang, XianGuang Wang, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature 2020. doi:<https://doi.org/10.1038/s41586-020-2012-7>.

- [19] Fan Wu, Su Zhao, Bin Yu, et al. A new coronavirus associated with human respiratory disease in China. Nature 2020.doi:<https://doi.org/10.1038/s41586-020-2008-3>.
- [20] Hui DSC, Zumla A. Severe Acute Respiratory Syndrome: Historical, Epidemiologic, and Clinical Features. Infect Dis Clin North Am 2019. 33(4):869-889.
- [21] Azhar EI, Hui DSC, Memish ZA, et al. The Middle East Respiratory Syndrome (MERS). Infect Dis Clin North Am 2019. 33(4):891-905.
- [22] Sheng G, Chen P, Wei Y, et al. Viral Infection Increases the Risk of Idiopathic Pulmonary Fibrosis: A Meta-Analysis. Chest 2019. doi: 10.1016/j.chest.2019.10.032.
- [23] Xie L, Liu Y, Xiao Y, et al. Follow-up study on pulmonary function and lung radiographic changes in rehabilitating severe acute respiratory syndrome patients after discharge. Chest 2005. 127(6):2119-24.
- [24] Hui DS, Joynt GM, Wong KT, et al. Impact of severe acute respiratory syndrome (SARS) on pulmonary function, functional capacity and quality of life in a cohort of survivors. Thorax 2005. 60(5):401-9.
- [25] Yu Zhao, Zixian Zhao, Yujia Wang, et al. Single-cell RNA expression profiling of ACE2, the putative receptor of

Wuhan 2019-nCov. bioRxiv 2020. doi: <https://doi.org/10.1101/2020.01.26.919985>.

- [26] Holshue ML, DeBolt C, Lindquist S, et al. First Case of 2019 Novel Coronavirus in the United States. N Engl J Med 2020. doi: 10.1056/NEJMoa2001191.
- [27] Buret AG. Immuno-modulation and anti-inflammatory benefits of antibiotics: the example of tilmicosin. Can J Vet Res 2010. 74(1):1-10.
- [28] Wuyts WA, Willems S, Vos R, et al. Azithromycin reduces pulmonary fibrosis in a bleomycin mouse model. Exp Lung Res 2010. 36(10):602-14.
- [29] Luo F, Zhuang Y, Sides MD, et al. Arsenic trioxide inhibits transforming growth factor- β 1-induced fibroblast to myofibroblast differentiation in vitro and bleomycin induced lung fibrosis in vivo. Respir Res 2014. 15:51.
- [30] Wang Chunbin ,Zhang Shuang ,Zhang Qian, et al. Research Progress of Two New Antifibrotic Drugs for Idiopathic Pulmonary Fibrosis. Chin Pharm J 2019. 09:682-686.

Figure legends

Figure 1. PRISMA flow diagram

Figure 2. Meta-analysis of the incidence of fever

Figure 3. Meta-analysis of the incidence of cough

Figure 4. Meta-analysis of the incidence of muscle soreness or fatigue

Figure 5. Meta-analysis of the incidence of ARDS

Figure 6. Meta-analysis of the incidence of abnormal chest CT

Figure 7. Meta-analysis of the proportion of severe cases in all infected cases

Figure 8. Meta-analysis of the mortality rate of patients with 2019-nCoV infection

Table 1: The characteristics of the literature

[illegible]

Weijie Guan ^[8]	2020	China	28	1099	Male:640	47	Retrospective Study	8	Fever ARDS Muscle Soreness or Fatigue Cough Abnormal Chest CT Patient in Critical Condition Death of Patient
					Female:459				
Lei Chen ^[9]	2020	China	15	29	Male:21	56	Retrospective Study	6	Fever Muscle Soreness or Fatigue Cough Patient in Critical Condition Death of Patient
					Female:8				
Kaiyuan Sun ^[10]	2020	America	42	288	Male:182	44	Retrospective Study	5	Death of Patient
					Female:106				
Yang Yang ^[11]	2020	China	51	4021	Male:2211	49	Retrospective Study	5	Patient in Critical Condition Death of Patient
					Female:1810				
Jie Li ^[12]	2020	China	21	17	Male:9	45	Retrospective Study	6	Fever Muscle Soreness or Fatigue Cough Abnormal Chest CT
					Female:8				
China CDC ^[13]	2020	China	43	44672	Male:22981	--	Retrospective Study	6	Patient in Critical Condition Death of Patient
					Female:21691				

Xiaowei Xu ^[14]	2020	China	16	62	Male:36	41	Retrospective Study	6	Fever Muscle Soreness or Fatigue Cough Abnormal Chest CT Patient in Critical Condition Death of Patient
					Female:26				

China CDC: Chinese Center for Disease Control and Prevention.

Table 2:Meta-analysis results

Clinical Symptom	Results of Meta-analysis	Adjusted Results
Fever	2.47(2.26,2.67)	0.891(0.818,0.945)
ARDS	0.79(0.43,1.15)	0.148(0.046,0.296)
Muscle Soreness or Fatigue	1.42(0.96,1.88)	0.425(0.213,0.652)

Cough	2.03(1.89,2.17)	0.722(0.657,0.782)
Abnormal Chest CT	2.77(2.57,2.97)	0.966(0.921,0.993)
Patient in Critical Condition	0.88(0.73,1.03)	0.181(0.127,0.243)
Death of Patient	0.42(0.33,0.50)	0.043(0.027,0.061)

Table 3: Results of Egger test

Clinical Symptom	Fever	ARDS	Muscle Soreness or Fatigue	Cough	Abnormal Chest CT	Patient in Critical Condition	Death of Patient
P 值	0.866	0.008	0.090	0.278	0.908	0.826	0.258