Polygenic modulation of lipoprotein(a)-associated cardiovascular risk.

Brief title: Lipoprotein(a)-associated cardiovascular risk.

Mark Trinder^{a,b}, MSc, Liam R. Brunham^{a,b,c,d}, MD, PhD.

^a Centre for Heart Lung Innovation, University of British Columbia, Vancouver, BC, Canada.

^b Experimental Medicine Program, University of British Columbia, Vancouver, British Columbia, Canada

^c Department of Medicine, University of British Columbia, Vancouver, British Columbia, Canada

^d Department of Medical Genetics, University of British Columbia, Vancouver, British Columbia, Canada

Funding: This project was supported by the Providence Health Care Research Institute. M.T. is supported by a Vanier Canada Graduate Scholarship. L.R.B. is supported by a Michael Smith Foundation for Health Research Scholar Award and is a Canada Research Chair in Precision Cardiovascular Disease Prevention.

Disclosures: None.

*** Address for correspondence:**

Dr. Liam R. Brunham
Centre for Heart Lung Innovation
Room 166-1081 Burrard Street
Vancouver, BC V6Z 1Y6
Phone: (604) 682-2344 x63929
Fax: (604) 806-9274
Email: liam.brunham@ubc.ca
Twitter handle: @LiamBrunham

ABSTRACT

Aims: Elevated levels of lipoprotein(a) are one of the strongest inherited risk factors for coronary artery disease (CAD). However, there is variability in cardiovascular risk among individuals with elevated lipoprotein(a). The sources of this variability are incompletely understood. We assessed the effects of a genomic risk score (GRS) for CAD on risk of myocardial infarction among individuals with elevated lipoprotein(a).

Methods: We calculated CAD GRSs for 408,896 individuals of British white ancestry from the UK Biobank using 6.27 million common genetic variants. Lipoprotein(a) levels were measured in 310,020 individuals. The prevalence and risk of myocardial infarction versus CAD GRS percentiles were compared for individuals with and without elevated lipoprotein(a) defined as ≥ 120 or 168 nmol/L (≈ 50 or 70 mg/dL, respectively).

Results: Individuals with elevated lipoprotein(a) displayed significantly greater CAD GRSs than individuals without elevated lipoprotein(a), which was largely dependent on the influence of genetic variants within or near the *LPA* gene. Continuous levels of CAD GRS percentile were significantly associated with risk of myocardial infarction for individuals with elevated lipoprotein(a). Notably, the risk of myocardial infarction for males with elevated lipoprotein(a) levels, but a CAD GRS percentile in the lower quintile ($< 20^{\text{th}}$ percentile), was less than the overall risk of myocardial infarction for males with non-elevated lipoprotein(a) levels (hazard ratio [95% CI]: 0.79 [0.64 - 0.97], $p=0.02$). Similar results were observed for females.

Conclusion: These data suggest that CAD genomic scores influence cardiovascular risk among individuals with elevated lipoprotein(a) and may aid in identifying candidates for preventive therapies.

53 **Keywords:** Lp(a), genomic risk score, myocardial infarction, genetic risk, atherosclerosis,
54 polygenic

INTRODUCTION

Lipoprotein(a) is a plasma lipoprotein composed of a low-density lipoprotein particle that is covalently linked to apolipoprotein(a) by a disulfide bond. Lipoprotein(a) levels are among the strongest inherited risk factors for cardiovascular disease¹⁻⁹, and Mendelian randomization studies suggest that elevated levels of lipoprotein(a) are causal for aortic stenosis¹⁰, heart failure¹¹, ischemic stroke¹², and coronary artery disease (CAD)^{1,13}. Plasma levels of lipoprotein(a) are largely determined by genetic factors which include single-nucleotide variants and copy number variants in the kringle IV type 2 (KIV-2) domain of the *LPA* gene^{5,6,14,15}.

Elevated lipoprotein(a) is a common condition that is estimated to affect 1 in 5 individuals of European ancestry (plasma lipoprotein(a) levels greater than 50 mg/dL)¹⁶. Although elevated lipoprotein(a) is associated with increased cardiovascular risk, the optimal treatment of elevated lipoprotein(a) is uncertain. While there are currently no approved therapies for lowering lipoprotein(a), antisense oligonucleotides that target apolipoprotein(a) and reduce lipoprotein(a) levels by up to 80% have been developed and are currently being studied in clinical trials¹⁷⁻¹⁹. At the individual level, there is substantial variability in cardiovascular risk among individuals with elevated lipoprotein(a)^{1,20,21}, and it remains unclear whether isolated elevated lipoprotein(a) in the absence of other cardiovascular risk factors merits treatment to reduce cardiovascular risk.

Recent work suggests that genomic risk scores (GRSs), which summarize the association of millions of common genetic variants with risk of CAD into a single value, may be a useful tool for improving risk prediction of CAD^{7,22-25}. Here we tested the hypothesis that, among individuals with elevated lipoprotein(a), the risk of CAD would be modulated by a GRS for CAD, and that this may be useful for risk stratification.

METHODS

Study population.

We studied participants from the UK Biobank, a large population-based prospective cohort study from the United Kingdom that aims to improve the prevention, diagnosis and treatment of disease by following the health and well-being of approximately 500 000 individuals. Individuals were enrolled between 2006-2010 and were between 40-69 years-of-age. Individuals underwent deep phenotyping at study enrollment and DNA was collected for genotyping, as previously described²⁶. The UK Biobank resource was approved by the UK Biobank Research Ethics Committee, and all participants provided written informed consent to participate in the study. This study was approved by the UK Biobank (application ID: 42857) and by the Clinical Research Ethics Board of the University of British Columbia (H18-02181).

Biochemical measurements.

Biochemical measurements were assessed at the time of study enrollment (Supplemental Methods; Supplemental Table 1). Lipoprotein(a) was measured using an immuno-turbidimetric method on the Beckman Coulter AU5800 platform (Randox Bioscience, UK), which is essentially isoform insensitive²⁷. Where indicated, lipoprotein(a) concentration was converted from nmol/L to mg/dL by dividing values by 2.4^{3,28}. Elevated lipoprotein(a) was defined as lipoprotein(a) levels ≥ 120 nmol/L (≈ 50 mg/dL) or ≥ 168 nmol/L (≈ 70 mg/dL)^{27,29,30}. For individuals taking cholesterol-lowering medication, total cholesterol and low-density lipoprotein cholesterol levels were adjusted by multiplying on-treatment lipid levels by 1.43, corresponding to an estimated 30% reduction in low-density lipoprotein cholesterol^{31,32}.

Calculation of coronary artery disease genomic risk scores.

The weightings used to calculate CAD GRSs were obtained from the Cardiovascular Disease Initiative Knowledge Portal and can be accessed at www.broadcvdi.org^{7,22}. CAD GRSs were calculated for individuals of British white ancestry using imputed genetic data that were not flagged as outliers for excess missingness or heterozygosity²⁶. CAD GRSs were calculated using 6.27 million of the 6.63 million potential single-nucleotide variants described by Khera et al. (2018). Variants that displayed deviation from Hardy-Weinberg equilibrium ($p < 1 \times 10^{-6}$), a genotyping rate $< 95\%$, or minor allele frequency $< 1\%$ were removed. The PLINK score function was used to multiply the number of alleles associated with adjusted β -coefficient for increased risk of CAD at each single-nucleotide variant by its respective linkage disequilibrium-adjusted weight and sums these products across all available variants to generate a GRS for each individual^{23,33}. CAD GRSs were also calculated after excluding genetic variants located within 7.5 million base pairs upstream and downstream of the *LPA* gene. CAD GRSs were converted into percentiles relative to the distribution of CAD GRSs in the UK Biobank study population.

The top 100 genetic variants used in the calculation of CAD GRSs were manually curated for association with known cardiovascular risk factors using the NHGRI-EBI Genome-Wide Association Study Catalog³⁴.

Definition of cardiovascular events.

Phenotypes, including the primary outcome of myocardial infarction, were defined using reports from medical history interviews occurring at enrollment, International Classification of Diseases (ICD)-9th and -10th Revision codes, and death registry records (Supplemental Methods, Supplemental Table 2). Events occurring before and after enrolment were included unless otherwise stated. Events occurring prior to enrolment were identified by either self-reported medical history and/or previous hospital admission within an electronic health record. Incident

events were defined by hospital admission with an electronic health record entry or death records. Events were censored on the date of loss-to-follow-up or if individuals remained event-free up to March 31, 2017.

Statistical analyses.

All statistical analyses were performed using R version 3.6.0 software (R Core Team, 2019). Individuals with missing values were excluded from analyses.

Chi-square tests were used for contingency analyses. For comparison of 2 groups, data were analyzed with an unpaired t-test or Mann–Whitney U test as appropriate. For comparison of more than 2 groups, data were analyzed with one-way analysis of variance test (with Tukey's multiple comparison post hoc tests) or Kruskal-Wallis test, as appropriate (with Dunn's multiple comparison post hoc tests).

The distributions of CAD GRS percentiles were compared between individuals with traditional cardiovascular risk factors determined at study enrollment that included: hypertension, severe hypercholesterolemia (low-density lipoprotein cholesterol levels ≥ 4.9 mmol/L), diabetes mellitus, obesity (body mass index ≥ 30 kg/m²), current smoker status, insufficient physical activity (classification of low activity level according to the International Physical Activity Questionnaire), insufficient fruit and vegetable intake (< 3 servings per day; sum of cooked vegetable intake, fresh fruit intake, and salad/raw vegetable intake), and strata of lipoprotein(a) levels (< 72 , 72–120, 120–168, and ≥ 168 nmol/L).

The prevalence of cardiovascular events of interest were calculated for each decile of CAD GRS percentile. Linear regression models were used to assess the correlation between cardiovascular disease prevalence and CAD GRS percentile using elevated lipoprotein(a) status as a covariate or interactive term

Time-to-event analyses were analyzed with the “survival” version 2.43-3 package for R with Log-rank tests. Hazard ratios (HRs) and 95% confidence intervals (CIs) were calculated using Cox regression models stratified by sex with age of event as a time scale. All Cox regression models were adjusted for the first 4 principal components of ancestry and genotyping array.

Statistical significance was claimed when two-sided p-values were <0.05.

RESULTS

Characteristics of individuals from the UK Biobank.

This study included 408,896 individuals of British white ancestry from the UK Biobank study, of whom 310,020 had lipoprotein(a) levels measured at enrollment (Table 1). The average age of enrollment was 56.9 years-of-age (standard deviation: 8.0 years) and 54.1% were female.

Genomic risk scores for coronary artery disease associate with the prevalence of traditional cardiovascular risk factors.

First, we assessed how CAD GRSs were associated with traditional cardiovascular risk factors. Individuals with hypertension, severe hypercholesterolemia, diabetes mellitus, obesity, and a current smoker status displayed significantly higher CAD GRS percentiles than those without the cardiovascular risk factor of interest (Mann-Whitney U test: $p=0.0006$ for smoking status, $p<0.0001$ for others; Figure 1a-e; Supplemental Table 3). There was also a significant stepwise increase in the median CAD GRS percentile as the number of traditional CAD risk factors per an individual increased (Kruskal-Wallis test: $p<0.0001$; Figure 1f). Alternatively, CAD GRSs did not display an association with insufficient physical activity (Mann-Whitney U

test: $p=0.40$) or an insufficient daily intake of fruits and vegetables (Mann–Whitney U test: $p=0.40$) (Supplemental Figure 1).

Individuals with elevated lipoprotein(a) have higher genomic risk scores for coronary artery disease.

CAD GRS percentiles displayed a significant and strong association with lipoprotein(a) levels (Kruskal–Wallis test: $p<0.0001$; Figure 2A). Notably, individuals with lipoprotein(a) levels greater or equal to 120 nmol/L (≈ 50 mg/dL) displayed a median CAD GRS of 65.2 percentile. Individuals with elevated lipoprotein(a) tended to have CAD GRSs that were comparable to individuals with multiple traditional cardiovascular risk factors at the time of study enrollment (*i.e.* the median CAD GRS percentile for individuals with hypertension, severe hypercholesterolemia, diabetes mellitus, obesity, and current smoker status was 65.5 percentile). The higher CAD GRS percentile observed among individuals with elevated lipoprotein(a) was dependent on genetic loci within and around the *LPA* gene. When CAD GRSs were re-calculated after the exclusion of DNA variants located 7.5 million base pairs up- or downstream of the *LPA* gene, the difference in CAD GRSs across strata of lipoprotein(a) levels was attenuated (Kruskal–Wallis test: $p=0.02$; Figure 2B; Supplemental Table 4).

The strong association between CAD GRSs and elevated lipoprotein(a) reflects both the high heritability and atherogenicity of lipoprotein(a) particles. Specifically, the locus in the CAD GRS with the greatest standardized effect size (rs186696265; GRCh37 position 6:161111700:C:T), and hence the greatest influence on CAD GRSs by weight, is known to associate with lipoprotein(a) levels (Figure 2C). Moreover, 6 of the top 100 loci with the largest standardized effect sizes in the CAD GRSs have known associations with lipoprotein(a) levels and include: rs186696265, rs10455872 (6:161010118:A:G), rs55730499 (6:161005610:C:T),

rs118039278 (6:160985526:G:A), rs56393506 (6:161089307:C:T), and rs4252185 (6:161123451:T:C) (Figure 2C).

Association of lipoprotein(a) levels with risk of myocardial infarction.

Consistent with previous studies^{1,27}, lipoprotein(a) levels were right-skewed among individuals of British white ancestry from the UK Biobank (Figure 3A). The median and interquartile range of lipoprotein(a) levels was 20.1 and 50.9 nmol/L, respectively. The risk of myocardial infarction was significantly associated with lipoprotein(a) levels for both male (adjusted HR [95% CI] 1.06 [1.06-1.07] per 20 nmol/L increase, $p < 0.0001$; Figure 3B) and female UK Biobank participants (adjusted HR [95% CI] 1.03 [1.02-1.05] per 20 nmol/L increase, $p < 0.0001$; Figure 3C).

Risk of myocardial infarction is modulated by coronary artery disease genomic risk score among individuals with elevated lipoprotein(a).

Next, we assessed if CAD GRSs may help explain some of the heterogeneity in risk of CAD that is observed in individuals with elevated lipoprotein(a), defined as ≥ 120 nmol/L (≈ 50 mg/dL). The prevalence of myocardial infarction was significantly associated with deciles of CAD GRS percentile in both individuals with and without elevated lipoprotein(a) (Figure 4A). Individuals with elevated lipoprotein(a) displayed a significantly greater prevalence of myocardial infarction relative to individuals with non-elevated lipoprotein(a) levels across all deciles of CAD GRS (β [SE] = 0.786 [0.194], $R^2 = 0.913$, $p = 0.0009$; Figure 4A; Supplemental Table 5). This effect was even more striking when deciles of CAD GRS percentile were calculated after excluding loci associated within and in proximity to the *LPA* gene (β [SE] = 1.494 [0.216], $R^2 = 0.920$, $p < 0.0001$; Figure 4B). Individuals with elevated lipoprotein(a) but a CAD GRS in the first decile had a lower prevalence of myocardial than individuals with non-

elevated lipoprotein(a) and a CAD GRS in the tenth decile (2.98% vs 6.61%, Figure 4B). These data indicate that the CAD GRS modulates the risk of myocardial infarction associated with elevated lipoprotein(a). There was no significant interaction between lipoprotein(a) status and CAD GRS that included or excluded loci in and nearby *LPA*.

We next examined the cumulative risk of myocardial infarction across time in males and females with elevated lipoprotein(a) as a function of CAD GRS. We observed a significant effect of the CAD GRS on risk of myocardial infarction in these individuals with elevated lipoprotein(a). The adjusted HR for males was 1.28 (95% CI: 1.23-1.33, $p < 0.0001$) and for females was 1.16 (95% CI: 1.08-1.25, $p < 0.0001$) per 20 unit increase in continuous CAD GRS percentile. Notably, the risk of myocardial infarction for individuals with elevated lipoprotein(a) levels, but a CAD GRS percentile in the lower quintile ($< 20^{\text{th}}$ percentile), was less than or equivalent to the overall risk of myocardial infarction for those with non-elevated lipoprotein(a) levels (adjusted HR [95% CI]: 0.79 [0.64-0.97] and 0.91 [0.66-1.26] with $p = 0.02$ and $p = 0.58$ for males and females, respectively; Figure 4C). Similar results were observed when the risk of myocardial infarction was compared between quintiles of CAD GRS percentiles that excluded loci surrounding the *LPA* gene (adjusted HR [95% CI] were 0.85 [0.73-0.98] and 0.83 [0.64 - 1.07] with $p = 0.03$ and $p = 0.15$ for males and females, respectively; Figure 3D).

Even among individuals with higher lipoprotein(a) levels of ≥ 168 nmol/L (≈ 70 mg/dL and comprising the top 3.5 percent of the population distribution), we observed a significant effect of the CAD GRS percentiles on risk of myocardial infarction (Supplemental Figure 2). Specifically, among these individuals, the adjusted HR for myocardial infarction was 1.31 for males (95% CI: 1.22-1.41, $p < 0.0001$) and 1.19 for females (95% CI: 1.04-1.34, $p = 0.009$) per 20 percentile increase in continuous CAD GRS. These findings highlight that genomic background

can still modulate the risk of myocardial infarction for individuals with elevated lipoprotein(a) and suggest that the CAD GRS may be useful to stratify cardiovascular risk in individuals identified to have elevated lipoprotein(a).

Effect of coronary artery disease genomic risk score on other cardiovascular outcomes associated with lipoprotein(a).

Lipoprotein(a) is also likely a causal risk factor for peripheral vascular disease and aortic stenosis^{10,35}, and, to a lesser extent, ischemic stroke^{12,36}. Since there is considerable overlap of risk factors between these cardiovascular conditions and CAD³⁷, we sought to assess if CAD GRSs also modulate the risk of these conditions among individuals with elevated lipoprotein(a). We observed a trend towards increased prevalence of peripheral vascular disease (β [SE]: 0.120 [0.055], $p=0.002$, $R^2=0.712$), aortic stenosis (β [SE]: 0.241 [0.056], $p=0.0005$, $R^2=0.615$), and ischemic stroke (β [SE]: 0.182 [0.072], $p=0.02$, $R^2=0.361$) for individuals with elevated lipoprotein(a), relative to individuals with non-elevated lipoprotein(a) (Supplemental Figure 3; Supplemental Tables 6-8). The strength of the association between prevalence of ischemic stroke and deciles of CAD GRS percentile was weak due to fewer events relative to the other cardiovascular conditions.

DISUCSSION

Here we report that common genetic factors modify the risk of myocardial infarction and other adverse cardiovascular outcomes among individuals with elevated lipoprotein(a). These data highlight the atherogenicity and high heritability of lipoprotein(a) levels, as reflected by the substantial increases in CAD GRSs observed among individuals with elevated lipoprotein(a). The increased CAD GRSs observed in individuals with elevated lipoprotein(a) were almost

entirely explained by single-nucleotide variants near the *LPA* gene having large weighted effect sizes, and thus, a large influence on the CAD GRS.

This work adds to the growing literature highlighting the role that polygenic factors play in modifying the penetrance or expressivity of monogenic conditions, such as familial hypercholesterolemia^{22,38–40}. To our knowledge, this is the first study to describe the polygenic modulation of myocardial infarction and other adverse cardiovascular outcomes by CAD GRSs among individuals with elevated lipoprotein(a).

Elevated lipoprotein(a) is a common monogenic condition resulting from genetic variation in and around the *LPA* gene, which includes single-nucleotide variants and copy number variation in the KIV-2 DNA sequence. Variation in the number of KIV-2 repeats results in plasma lipoprotein(a) particles that display considerable interindividual variation in apolipoprotein(a) isoform size (ranging from 300-800 kDa). The number of KIV-2 repeats are also inversely associated with lipoprotein(a) levels. However, lipoprotein(a) levels (*i.e.* molar concentration), rather than apo(a) isoform size, appears to be most strongly associated with risk of CAD^{20,41,42}.

Approximately 1 in 5 individuals of European ancestry have lipoprotein(a) levels greater than 120 nmol/L ($\approx$ 50 mg/mL), which is a common cut-off for estimation of cardiovascular risk and consideration of therapy³⁰. Lipoprotein(a) levels greater than 120 nmol/L are also common among individuals of African (1 in 4), South and South East Asian ($\sim$ 1 in 10), Arab ($\sim$ 1 in 10), and Latin American ($\sim$ 1 in 7) ethnicity and also associated with risk of CAD in these populations^{20,43,44}. However, the optimal management of patients with elevated lipoprotein(a) is uncertain. Antisense inhibitors that potentially lower lipoprotein(a) are currently being studied in clinical trials in subjects with elevated lipoprotein(a) and established cardiovascular disease^{17–19}.

Our study suggests that the CAD GRS may be useful for identifying high risk patients that are likely to benefit most from lipoprotein(a)-lowering therapy, or other therapies that reduce cardiovascular risk among individuals with elevated lipoprotein(a). In support of this hypothesis, previous studies have shown that individuals with high genetic risk for CAD demonstrate greater absolute risk reduction of cardiovascular events from statins^{24,45} or PCSK9 inhibitors⁴⁶ relative to individuals at lower genetic risk. In addition, the data presented here suggest that a GRS may be useful to de-risk patients with isolated elevated lipoprotein(a), as we observed that a low GRS in such individuals was associated with a risk of myocardial infarction that was similar to, or lower than, that of individuals with non-elevated lipoprotein(a) and a median GRS. The extent to which CAD GRSs can improve risk prediction beyond established clinical risk scores requires further investigation⁴⁷. However, a major advantage of assessing germline genetic variation for cardiovascular risk assessment is that it only needs to be measured once and can be measured early in life, allowing for earlier intervention before other clinical risk factors may emerge^{7,25,46}.

This study has some notable strengths and limitations worthy of consideration. Firstly, we were able to use a very large population of more than 300,000 individuals with standardized lipoprotein(a) measurements using an isoform insensitive assay, and genome-wide genotype data. As such, this represents one of the largest population-based studies of lipoprotein(a) to date. Limitations are that the population studied was comprised of individuals of European ancestry and may not be representative of other global populations. While this aspect of the study reduces the risk of bias or confounding due to population stratification, it also limits the generalization of these results to other ancestral populations. Future studies using large populations of individuals from diverse ancestral backgrounds are therefore essential. Indeed, as the calculation and calibration of the CAD GRSs depend on linkage disequilibrium^{7,23}, which varies substantially

across ancestries, future studies will be needed to derive and validate CAD GRSs in other ancestral groups. The UK Biobank population also displays a healthy volunteer selection bias relative to the overall health and well-being of individuals living across the United Kingdom⁴⁸. Furthermore, the CAD GRS was constructed to best identify individuals at risk of CAD and is likely not as sensitive at identifying individuals at risk of the other cardiovascular conditions described, which include peripheral vascular disease, aortic stenosis, and ischemic stroke. GRSs calibrated specifically for these conditions will likely improve risk prediction.

In summary, genetic factors were shown to significantly modulate the risk of CAD in individuals with elevated lipoprotein(a). Assessment of background polygenic factors may help to personalize risk assessment in individuals with elevated lipoprotein(a) and identify candidates for preventative therapy.

319 **ACKNOWLEDGEMENTS**

320 We acknowledge the participants and individuals involved in the UK Biobank studies; the
321 developers of software tools used in this study; and Compute Canada for establishing
322 computational resources and providing support.

REFERENCES

1. Kamstrup PR, Tybjaerg-Hansen A, Steffensen R, Nordestgaard BG. Genetically elevated lipoprotein(a) and increased risk of myocardial infarction. *JAMA* 2009;301:2331–2339.
2. Holmes DT, Schick BA, Humphries KH, Frohlich J. Lipoprotein(a) is an independent risk factor for cardiovascular disease in heterozygous familial hypercholesterolemia. *Clin Chem* 2005;51:2067–2073.
3. Khera AV, Everett BM, Caulfield MP, Hantash FM, Wohlgemuth J, Ridker PM, Mora S. Lipoprotein(a) concentrations, rosuvastatin therapy, and residual vascular risk: an analysis from the JUPITER Trial (Justification for the Use of Statins in Prevention: An Intervention Trial Evaluating Rosuvastatin). *Circulation* 2014;129:635–642.
4. Emerging Risk Factors Collaboration, Erqou S, Kaptoge S, Perry PL, Di Angelantonio E, Thompson A, White IR, Marcovina SM, Collins R, Thompson SG, Danesh J. Lipoprotein(a) concentration and the risk of coronary heart disease, stroke, and nonvascular mortality. *JAMA* 2009;302:412–423.
5. Clarke R, Peden JF, Hopewell JC, Kyriakou T, Goel A, Heath SC, Parish S, Barlera S, Franzosi MG, Rust S, Bennett D, Silveira A, Malarstig A, Green FR, Lathrop M, Gigante B, Leander K, Faire U de, Seedorf U, Hamsten A, Collins R, Watkins H, Farrall M; PROCARIS Consortium. Genetic variants associated with Lp(a) lipoprotein level and coronary disease. *N Engl J Med* 2009;361:2518–2528.
6. Boerwinkle E, Leffert CC, Lin J, Lackner C, Chiesa G, Hobbs HH. Apolipoprotein(a) gene accounts for greater than 90% of the variation in plasma lipoprotein(a) concentrations. *J Clin Invest* 1992;90:52–60.
7. Khera AV, Chaffin M, Aragam KG, Haas ME, Roselli C, Choi SH, Natarajan P, Lander ES, Lubitz SA, Ellinor PT, Kathiresan S. Genome-wide polygenic scores for common diseases identify individuals with risk equivalent to monogenic mutations. *Nat Genetics* 2018;50:1219–1224.
8. Wei WQ, Li X, Feng Q, Kubo M, Kullo IJ, Peissig PL, Karlson EW, Jarvik GP, Lee MTM, Shang N, Larson EA, Edwards T, Shaffer CM, Mosley JD, Maeda S, Horikoshi M, Ritchie M, Williams MS, Larson EB, Crosslin DR, Bland ST, Pacheco JA, Rasmussen-Torvik LJ, Cronkite D, Hripesak G, Cox NJ, Wilke RA, Stein CM, Rotter JI, Momozawa Y, Roden DM, Krauss RM, Denny JC. LPA variants are associated with residual cardiovascular risk in patients receiving statins. *Circulation* 2018;138:1839–1849.
9. Emdin CA, Khera AV, Natarajan P, Klarin D, Won H-H, Peloso GM, Stitzel NO, Nomura A, Zekavat SM, Bick AG, Gupta N, Asselta R, Duga S, Merlini PA, Correa A, Kessler T, Wilson JG, Bown MJ, Hall AS, Braund PS, Samani NJ, Schunkert H, Marrugat J, Elosua R, McPherson R, Farrall M, Watkins H, Willer C, Abecasis GR, Felix JF, Vasani RS, Lander E, Rader DJ, Danesh J, Ardisson D, Gabriel S, Saleheen D, Kathiresan S; CHARGE-Heart Failure Consortium; CARDIOGRAM Exome Consortium. Phenotypic characterization of genetically lowered human lipoprotein(a) levels. *J Am Coll Cardiol* 2016;68:2761–2772.

10. Kamstrup PR, Tybjaerg-Hansen A, Nordestgaard BG. Elevated lipoprotein(a) and risk of aortic valve stenosis in the general population. *J Am Coll Cardiol*. 2014;63:470–477.
11. Kamstrup PR, Nordestgaard BG. Elevated lipoprotein(a) levels, LPA risk genotypes, and increased risk of heart failure in the general population. *JACC Heart Fail* 2016;4:78–87.
12. Langsted A, Nordestgaard BG, Kamstrup PR. Elevated lipoprotein(a) and risk of ischemic stroke. *J Am Coll Cardiol* 2019;74:54–66.
13. Saleheen D, Haycock PC, Zhao W, Rasheed A, Taleb A, Imran A, Abbas S, Majeed F, Akhtar S, Qamar N, Zaman KS, Yaqoob Z, Saghir T, Rizvi SNH, Memon A, Mallick NH, Ishaq M, Rasheed SZ, Memon F-R, Mahmood K, Ahmed N, Frossard P, Tsimikas S, Witztum JL, Marcovina S, Sandhu M, Rader DJ, Danesh J. Apolipoprotein(a) isoform size, lipoprotein(a) concentration, and coronary artery disease: a mendelian randomisation analysis. *Lancet Diabetes Endocrinol* 2017;5:524–533.
14. Kettunen J, Demirkan A, Würtz P, Draisma HHM, Haller T, Rawal R, Vaarhorst A, Kangas AJ, Lyytikäinen LP, Pirinen M, Pool R, Sarin AP, Soininen P, Tukiainen T, Wang Q, Tiainen M, Tynkkynen T, Amin N, Zeller T, Beekman M, Deelen J, van Dijk KW, Esko T, Hottenga JJ, van Leeuwen EM, Lehtimäki T, Mihailov E, Rose RJ, de Craen AJM, Gieger C, Kähönen M, Perola M, Blankenberg S, Savolainen MJ, Verhoeven A, Viikari J, Willemsen G, Boomsma DI, van Duijn CM, Eriksson J, Jula A, Järvelin MR, Kaprio J, Metspalu A, Raitakari O, Salomaa V, Slagboom PE, Waldenberger M, Ripatti S, Ala-Korpela M. Genome-wide study for circulating metabolites identifies 62 loci and reveals novel systemic effects of LPA. *Nat Commun* 2016;7:11122.
15. Coassin S, Erhart G, Weissensteiner H, Eca Guimarães de Araújo M, Lamina C, Schönherr S, Forer L, Haun M, Losso JL, Köttgen A, Schmidt K, Utermann G, Peters A, Gieger C, Strauch K, Finkenstedt A, Bale R, Zoller H, Paulweber B, Eckardt K-U, Hüttenhofer A, Huber LA, Kronenberg F. A novel but frequent variant in LPA KIV-2 is associated with a pronounced Lp(a) and cardiovascular risk reduction. *Eur Heart J* 2017;38:1823–1831.
16. Nordestgaard BG, Chapman MJ, Ray K, Borén J, Andreotti F, Watts GF, Ginsberg H, Amarencu P, Catapano A, Descamps OS, Fisher E, Kovanen PT, Kuivenhoven JA, Lesnik P, Masana L, Reiner Z, Taskinen M-R, Tokgözoğlu L, Tybjaerg-Hansen A. Lipoprotein(a) as a cardiovascular risk factor: current status. *Eur Heart J* 2010;31:2844–2853.
17. Tsimikas S, Karwatowska-Prokopczuk E, Gouni-Berthold I, Tardif JC, Baum SJ, Steinhagen-Thiessen E, Shapiro MD, Stroes ES, Moriarty PM, Nordestgaard BG, Xia S, Guerriero J, Viney NJ, O’Dea L, Witztum JL. Lipoprotein(a) reduction in persons with cardiovascular disease. *N Engl J Med* 2020;382:244–255.
18. Viney NJ, van Capelleveen JC, Geary RS, Xia S, Tami JA, Yu RZ, Marcovina SM, Hughes SG, Graham MJ, Croke RM, Croke ST, Witztum JL, Stroes ES, Tsimikas S. Antisense oligonucleotides targeting apolipoprotein(a) in people with raised lipoprotein(a): two randomised, double-blind, placebo-controlled, dose-ranging trials. *Lancet* 2016;388:2239–2253.

19. Tsimikas S, Viney NJ, Hughes SG, Singleton W, Graham MJ, Baker BF, Burkey JL, Yang Q, Marcovina SM, Geary RS, Crooke RM, Witztum JL. Antisense therapy targeting apolipoprotein(a): a randomised, double-blind, placebo-controlled phase 1 study. *Lancet* 2015;386:1472–1483.
20. Paré G, Çaku A, McQueen M, Anand SS, Enas E, Clarke R, Boffa M, Koschinsky M, Wang X, Yusuf S, INTERHEART Investigators. Lipoprotein(a) levels and the risk of myocardial infarction among 7 ethnic groups. *Circulation* 2019;139:1472–1482.
21. Langsted A, Kamstrup PR, Benn M, Tybjaerg-Hansen A, Nordestgaard BG. High lipoprotein(a) as a possible cause of clinical familial hypercholesterolaemia: a prospective cohort study. *Lancet Diabetes Endocrinol* 2016;4:577–587.
22. Khera AV, Chaffin M, Zekavat SM, Collins RL, Roselli C, Natarajan P, Lichtman JH., D’Onofrio G, Mattera J, Dreyer R, Spertus JA., Taylor KD, Psaty PM, Rich SS, Post W, Gupta N, Gabriel S, Lander E, Ida Chen YD, Talkowski ME, Rotter JJ, Krumholz HM, Kathiresan S. Whole-genome sequencing to characterize monogenic and polygenic contributions in patients hospitalized with early-onset myocardial infarction. *Circulation* 2019;139:1593–1602.
23. Vilhjálmsson BJ, Yang J, Finucane HK, Gusev A, Lindström S, Ripke S, Genovese G, Loh PR, Bhatia G, Do R, Hayeck T, Won HH; Schizophrenia Working Group of the Psychiatric Genomics Consortium, Discovery, Biology, and Risk of Inherited Variants in Breast Cancer (DRIVE) study, Kathiresan S, Pato M, Pato C, Tamimi R, Stahl E, Zaitlen N, Pasaniuc B, Belbin G, Kenny EE, Schierup MH, De Jager P, Patsopoulos NA, McCarroll S, Daly M, Purcell S, Chasman D, Neale B, Goddard M, Visscher PM, Kraft P, Patterson N, Price AL. Modeling linkage disequilibrium increases accuracy of polygenic risk scores. *Am J Hum Genet* 2015;97:576–592.
24. Natarajan P, Young R, Stitzel NO, Padmanabhan S, Baber U, Mehran R, Sartori S, Fuster V, Reilly DF, Butterworth A, Rader DJ, Ford I, Sattar N, Kathiresan S. Polygenic risk score identifies subgroup with higher burden of atherosclerosis and greater relative benefit from statin therapy in the primary prevention setting. *Circulation* 2017;135:2091–2101.
25. Pereira A, Mendonca MI, Borges S, Sousa AC, Freitas S, Henriques E, Rodrigues M, Freitas AI, Guerra G, Freitas C, Pereira D, Brehm A, Reis RPD. Additional value of a combined genetic risk score to standard cardiovascular stratification. *Genet Mol Biol* 2018;41:766–774.
26. Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, Motyer A, Vukcevic D, Delaneau O, O’Connell J, Cortes A, Welsh S, Young A, Effingham M, McVean G, Leslie S, Allen N, Donnelly P, Marchini J. The UK Biobank resource with deep phenotyping and genomic data. *Nature* 2018;562:203–209.
27. Verbeek R, Boekholdt SM, Stoeckenbroek RM, Hovingh GK, Witztum JL, Wareham NJ, Sandhu MS, Khaw KT, Tsimikas S. Population and assay thresholds for the predictive value of lipoprotein (a) for coronary artery disease: the EPIC-Norfolk Prospective Population Study. *J Lipid Res* 2016;57:697-705.

28. Brown WV, Ballantyne CM, Jones PH, Marcovina S. Management of Lp(a). *J Clin Lipidol* 2010;4:240–247.
29. Burgess S, Ference BA, Staley JR, Freitag DF, Mason AM, Nielsen SF, Willeit P, Young R, Surendran P, Karthikeyan S, Bolton TR, Peters JE, Kamstrup PR, Tybjaerg-Hansen A, Benn M, Langsted A, Schnohr P, Vedel-Krogh S, Kobylecki CJ, Ford I, Packard C, Trompet S, Jukema JW, Sattar N, Di Angelantonio E, Saleheen D, Howson JMM, Nordestgaard BG, Butterworth AS, Danesh J; European Prospective Investigation Into Cancer and Nutrition–Cardiovascular Disease (EPIC-CVD) Consortium. Association of LPA variants with risk of coronary disease and the implications for lipoprotein(a)-lowering therapies: a Mendelian randomization analysis. *JAMA Cardiol* 2018;3:619–627.
30. Madsen CM, Kamstrup PR, Langsted A, Varbo A, Nordestgaard BG. Lipoprotein(a)-lowering by 50 mg/dL (105 nmol/L) may be needed to reduce cardiovascular disease 20% in secondary prevention. *Arterioscler Thromb Vasc Biol* 2020;40:255–266.
31. Benn M, Watts GF, Tybjaerg-Hansen A, Nordestgaard BG. Mutations causative of familial hypercholesterolaemia: screening of 98 098 individuals from the Copenhagen General Population Study estimated a prevalence of 1 in 217. *Eur Heart J* 2016;37:1384–1394.
32. Jones PH, Davidson MH, Stein EA, Bays HE, McKenney JM, Miller E, Cain VA, Blasetto JW; STELLAR Study Group. Comparison of the efficacy and safety of rosuvastatin versus atorvastatin, simvastatin, and pravastatin across doses (STELLAR* Trial). *Am J Cardiol* 2003;92:152–160.
33. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de Bakker PIW, Daly MJ, Sham PC. PLINK: A tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;81:559–575.
34. Buniello A, MacArthur JAL, Cerezo M, Harris LW, Hayhurst J, Malangone C, McMahon A, Morales J, Mountjoy E, Sollis E, Suveges D, Vrousseau O, Whetzel PL, Amode R, Guillen JA, Riat HS, Trevanion SJ, Hall P, Junkins H, Flicek P, Burdett T, Hindorf LA, Cunningham F, Parkinson H. The NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res* 2019;47:D1005–D1012.
35. Thanassoulis G, Campbell CY, Owens DS, Smith JG, Smith AV, Peloso GM, Kerr KF, Pechlivanis S, Budoff MJ, Harris TB, Malhotra R, O'Brien KD, Kamstrup PR, Nordestgaard BG, Tybjaerg-Hansen A, Allison MA, Aspelund T, Criqui MH, Heckbert SR, Hwang SJ, Liu Y, Sjogren M, van der Pals J, Kalsch H, Muhleisen TW, Nothen MM, Cupples LA, Caslake M, Di Angelantonio E, Danesh J, Rotter JJ, Sigurdsson S, Wong Q, Erbel R, Kathiresan S, Melander O, Gudnason V, O'Donnell CJ, Post WS; CHARGE Extracoronary Calcium Working Group. Genetic associations with valvular calcification and aortic stenosis. *N Engl J Med*. 2013;368:503–512.
36. Helgadottir A, Gretarsdottir S, Thorleifsson G, Holm H, Patel RS, Gudnason T, Jones GT, van Rij AM, Eapen DJ, Baas AF, Tregouet DA, Morange PE, Emmerich J, Lindblad B, Gottsater A, Kiemeny LA, Lindholt JS, Sakalihasan N, Ferrell RE, Carey DJ, Elmore JR,

- Tsao PS, Grarup N, Jørgensen T, Witte DR, Hansen T, Pedersen O, Pola R, Gaetani E, Magnadottir HB, Wijmenga C, Tromp G, Ronkainen A, Ruigrok YM, Blankensteijn JD, Mueller T, Wells PS, Corral J, Soria JM, Souto JC, Peden JF, Jalilzadeh S, Mayosi BM, Keavney B, Strawbridge RJ, Sabater-Lleal M, Gertow K, Baldassarre D, Nyssönen K, Rauramaa R, Smit AJ, Mannarino E, Giral P, Tremoli E, de Faire U, Humphries SE, Hamsten A, Haraldsdottir V, Olafsson I, Magnusson MK, Samani NJ, Levey AI, Markus HS, Kostulas K, Dichgans M, Berger K, Kuhlenbäumer G, Ringelstein EB, Stoll M, Seedorf U, Rothwell PM, Powell JT, Kuivaniemi H, Onundarson PT, Valdimarsson E, Matthiasson SE, Gudbjartsson DF, Thorgeirsson G, Quyyumi AA, Watkins H, Farrall M, Thorsteinsdottir U, Stefansson K. Apolipoprotein(a) genetic sequence variants associated with systemic atherosclerosis and coronary atherosclerotic burden but not with venous thromboembolism. *J Am Coll Cardiol*. 2012;60:722–729.
37. Klarin D, Lynch J, Aragam K, Chaffin M, Assimes TL, Huang J, Lee KM, Shao Q, Huffman JE, Natarajan P, Arya S, Small A, Sun YV, Vujkovic M, Freiberg MS, Wang L, Chen J, Saleheen D, Lee JS, Miller DR, Reaven P, Alba PR, Patterson OV, DuVall SL, Boden WE, Beckman JA, Gaziano JM, Concato J, Rader DJ, Cho K, Chang KM, Wilson PWF, O'Donnell CJ, Kathiresan S; VA Million Veteran Program, Tsao PS, Damrauer SM. Genome-wide association study of peripheral artery disease in the Million Veteran Program. *Nature Med* 2019;25:1274–1279.
38. Trinder M, Li X, DeCastro ML, Cermakova L, Sadananda S, Jackson LM, Azizi H, Mancini GBJ, Francis GA, Frohlich J, Brunham LR. Risk of premature atherosclerotic disease in patients with monogenic versus polygenic familial hypercholesterolemia. *J Am Coll Cardiol*. 2019;74:512–522.
39. Fahed AC, Wang M, Homburger JR, Patel AP, Bick AG, Neben CL, Lai C, Brockman D, Philippakis A, Ellinor PT, Cassa CA, Lebo M, Ng K, Lander ES, Zhou AY, Kathiresan S, Khera AV. Polygenic background modifies penetrance of monogenic variants conferring risk for coronary artery disease, breast cancer, or colorectal cancer. *medRxiv* 2019;19013086.
40. Oetjens MT, Kelly MA, Sturm AC, Martin CL, Ledbetter DH. Quantifying the polygenic contribution to variable expressivity in eleven rare genetic disorders. *Nat Commun* 2019;10:4897.
41. Gudbjartsson DF, Thorgeirsson G, Sulem P, Helgadottir A, Gylfason A, Saemundsdottir J, Bjornsson E, Norddahl GL, Jonasdottir A, Jonasdottir A, Eggertsson HP, Gretarsdottir S, Thorleifsson G, Indridason OS, Palsson R, Jonasson F, Jonsdottir I, Eyjolfsson GI, Sigurdardottir O, Olafsson I, Danielsen R, Matthiasson SE, Kristmundsdottir S, Halldorsson BV, Hreidarsson AB, Valdimarsson EM, Gudnason T, Benediktsson R, Steinthorsdottir V, Thorsteinsdottir U, Holm H, Stefansson K. Lipoprotein(a) concentration and risks of cardiovascular disease and diabetes. *J Am Coll Cardiol* 2019;74:2982–2994.
42. Willeit P, Kiechl S, Kronenberg F, Witzum JL, Santer P, Mayr M, Xu Q, Mayr A, Willeit J, Tsimikas S. Discrimination and net reclassification of cardiovascular risk with

- lipoprotein(a): prospective 15-year outcomes in the Bruneck Study. *J Am Coll Cardiol* 2014;64:851–860.
43. Virani SS, Brautbar A, Davis BC, Nambi V, Hoogeveen RC, Sharrett AR, Coresh J, Mosley TH, Morrisett JD, Catellier DJ, Folsom AR, Boerwinkle E, Ballantyne CM. Associations between lipoprotein(a) levels and cardiovascular outcomes in black and white subjects: the Atherosclerosis Risk in Communities (ARIC) Study. *Circulation* 2012;125:241–249.
44. Gupta R, Vasisht S, Bahl VK, Wasir HS. Correlation of lipoprotein (a) to angiographically defined coronary artery disease in Indians. *Int J Cardiol* 1996;57:265–270.
45. Mega JL, Stitzel NO, Smith JG, Chasman DI, Caulfield MJ, Devlin JJ, Nordio F, Hyde CL, Cannon CP, Sacks FM, Poulter NR, Sever PS, Ridker PM, Braunwald E, Melander O, Kathiresan S, Sabatine MS. Genetic risk, coronary heart disease events, and the clinical benefit of statin therapy: an analysis of primary and secondary prevention trials. *Lancet* 2015;385:2264–2271.
46. Marston NA, Kamanu FK, Nordio F, Gurmu Y, Roselli C, Sever PS, Pedersen TR, Keech AC, Wang H, Lira Pineda A, Giugliano RP, Lubitz SA, Ellinor PT, Sabatine MS, Ruff CT. Predicting Benefit From Evolocumab Therapy in Patients With Atherosclerotic Disease Using a Genetic Risk Score: Results From the FOURIER Trial. *Circulation*. doi: 10.1161/CIRCULATIONAHA.119.043805.
47. Elliott J, Bodinier B, Bond TA, Chadeau-Hyam M, Evangelou E, Moons KGM, Dehghan A, Muller DC, Elliott P, Tzoulaki I. Predictive accuracy of a polygenic risk score–enhanced prediction model vs a clinical risk score for coronary artery disease. *JAMA* 2020;323:636–645.
48. Fry A, Littlejohns TJ, Sudlow C, Doherty N, Adamska L, Sprosen T, Collins R, Allen NE. Comparison of sociodemographic and health-related characteristics of UK Biobank participants with those of the general population. *Am J Epidemiol* 2017;186:1026–1034.

FIGURES

Figure 1. Genomic scores for coronary artery disease associate with traditional cardiovascular risk factors. The distribution of coronary artery disease genomic score percentiles are displayed for traditional cardiovascular risk factors assessed at study enrollment which include: (A) hypertension, (B) high low-density lipoprotein cholesterol (LDL-C $\geq$ 4.9 mmol/L), (C) diabetes mellitus, (D) obesity (body mass index $\geq$ 30 kg/m²), (E) current smoking status, and (F) a sum of the aforementioned risk factors. Boxplots display the median and interquartile range. The boxplot's whiskers show the range and the accompanying density distribution. *** p<0.001, **** p<0.0001.

Figure 2. Elevated lipoprotein(a) is a major determinant of coronary artery disease

genomic risk scores. The distribution of coronary artery disease genomic risk score percentiles are displayed across strata of lipoprotein(a) levels calculated (A) with and (B) without genetic variants in proximity to the *LPA* gene. Boxplots display the median and interquartile range. The boxplot's whiskers show the range and the accompanying density distribution. (C) The standardized effect sizes for the top 100 genetic variants used in calculating coronary artery disease genomic risk scores are shown coloured by reported associations with traditional or emerging cardiovascular risk factors in the NHGRI-EBI Genome-Wide Association Study Catalog. Lipoprotein(a) [Lp(a)].

Figure 3. Elevated lipoprotein(a) associates with increased risk of myocardial infarction.

(A) The density distribution of lipoprotein(a) levels for UK Biobank participants of British white ancestry is right-skewed. The risk of myocardial infarction versus continuous lipoprotein(a) levels are shown for individuals of (B) male and (C) female sex. Hazard ratios were calculated relative to the median lipoprotein(a) level in this population (20 nmol/L) and adjusted for the

576 first 4 components of genetic ancestry and genotyping array/batch. The darker and lighter blue
577 shading represent the standard error and 95% confidence interval, respectively.
578

Figure 4. Coronary artery disease genomic risk scores are a modifier of risk of myocardial infarction for individuals with elevated lipoprotein(a). The prevalence of myocardial infarction for each decile of coronary artery disease genomic risk score percentile is shown for scores calculated (A) with and (B) without genetic variants in proximity to the *LPA*. Solid lines depict linear regression analyses and gray shading indicate the associated 95% confidence interval stratified by non-elevated versus elevated lipoprotein(a) levels. Time-to-first-event analyses are displayed for risk of myocardial infarction events stratified by quintiles of coronary artery disease genomic risk score percentile among individuals with elevated lipoprotein(a) (≥ 120 nmol/L) calculated (C) with and (D) without genetic variants in proximity to the *LPA* gene. The solid black line depicts the risk of myocardial infarction among a reference group comprised of all individuals with non-elevated lipoprotein(a) (Lp(a)-, < 120 nmol/L). The horizontal, dotted black line depicts the myocardial infarction-free probability of the non-elevated lipoprotein(a) reference group at 70 years-of-age.

TABLES

Table 1. Enrollment characteristics and cardiovascular history of the study group. High-density lipoprotein cholesterol (HDL-C), interquartile range (IQR), low-density lipoprotein cholesterol (LDL-C) standard deviation (SD).

Characteristic	Measurement	Values
n	no.	408896
Age	mean (SD)	56.9 (8.0)
Female sex	no. (%)	221082 (54.1)
Hypertension	no. (%) / n	110888 (27.2) / 408256
Severe hypercholesterolemia	no. (%) / n	31978 (8.2) / 389158
Diabetes mellitus	no. (%) / n	19773 (4.8) / 408003
Obesity	no. (%) / n	98879 (24.3) / 407600
Current smoker	no. (%) / n	41320 (10.1) / 407461
Total cholesterol (mmol/L)	median (IQR) / n	5.8 (1.55) / 389875
LDL-C (mmol/L)	median (IQR) / n	3.61 (1.16) / 389158
Triglycerides (mmol/L)	median (IQR) / n	1.50 (1.11) / 389567
HDL-C (mmol/L)	median (IQR) / n	1.40 (0.50) / 356840
Lipoprotein(a) (nmol/L)	median (IQR) / n	20.1 (50.88) / 310020
Hemoglobin A1c (mmol/mol)	median (IQR) / n	35.2 (5.1) / 389771
C-reactive protein (mg/L)	median (IQR) / n	1.33 (2.1) / 389036