Clinical Data on Hospital Environmental Hygiene Monitoring and Medical Staff Protection during the Coronavirus Disease 2019 Outbreak

Yanfang Jiang^{1*}, Haifeng Wang^{1*}, Yukun Chen², Jiaxue He¹, Ligu Chen¹, Yong Liu¹, Xinyuan Hu¹, Ang Li¹, Siwen Liu¹, Peng Zhang¹, Hongyan Zou¹, Shucheng Hua^{1,3 #}

1. Key Laboratory of Organ Regeneration & Transplantation of the Ministry of Education, Genetic Diagnosis Center, The First Hospital of Jilin University, Changchun, China

2. Hospital-Acquired Infection Control Department, The First Hospital of Jilin University, Changchun, China

3. Department of Respiratory Medicine, The First Hospital of Jilin University, Jilin University, Changchun, Jilin, China

Corresponding Author: Shucheng Hua, Department of Respiratory Medicine, The First Hospital of Jilin University, Jilin University, Changchun, Jilin, China

E-mail address: huashucheng2020@163.com

* Yanfang Jiang and Haifeng Wang made equal contributions to this study.

24 Yanfang Jiang yanfangjiang@hotmail.com

25 Haifeng Wang wanghaifeng1994@163.com

26 Yukun Chen cyk0503@qq.com

27 Jiaxue He jiaxue-he@163.com

28 Ligu Chen 123043714@qq.com

29 Yong Liu 530440980@qq.com

30 Xinyuan Hu 1586878478@qq.com

31 Ang Li li987001500@qq.com

32 Siwen Liu 530229652@qq.com

33 Peng Zhang 123126672@qq.com

34 Hongyan Zou 2361044502@qq.com

35

36

37

38

39

40

41

42

43

44

45

Abstract

Background: The outbreak of coronavirus disease 2019 (COVID-19) has placed unprecedented challenges on hospital environmental hygiene and medical staff protection. It is crucial to assess hospital environmental hygiene to understand the most important environmental issues for controlling the spread of COVID-19 in hospitals.

Objective: To detect the presence of COVID-19 in the samples from the area at risk of contamination in the First Hospital of Jilin University.

Methods: Viruses in the air were collected by natural sedimentation and air particle sampler methods. Predetermined environmental surfaces were sampled using swabs at seven o'clock in the morning before disinfection. The real-time reverse-transcription PCR method was used to detect the existence of COVID-19 pathogens.

Results: Viruses could be detected on the surfaces of the nurse station in the isolation area with suspected patients and in the air of the isolation ward with an intensive care patient.

Conclusion: Comprehensive monitoring of hospital environmental hygiene during pandemic outbreaks is conducive to the refinement of hospital infection control. It is of great significance to ensure the safety of medical treatment and the quality of hospital infection control through the monitoring of environmental hygiene.

Key words COVID-19; hospital environmental hygiene; hospital-associated infection, medical staff protection

1.Introduction

The outbreak of COVID-19 in Wuhan in December 2019 has led to a serious public health event ^[1-3]. Meanwhile, the outbreak of this novel virus has placed unprecedented challenges on hospital environmental hygiene. The occurrence of medical staff-associated infections is closely related to long-lived pathogens in the hospital environment ^[4,5]. Thus, it is crucial to assess hospital environmental hygiene to understand the most important environmental issues for controlling the spread of COVID-19 in hospitals. The Chinese government quickly adopted quarantine measures for confirmed and suspected patients to restrain the spread of the pandemic^[6]. Comprehensive monitoring of hospital environmental hygiene during the outbreak of the pandemic is conducive to the refinement of hospital infection control ^[5,7]. It also increases understanding of the environmental challenges corresponding to the reemergence of COVID-19 or similar viruses. Therefore, it is of great significance to ensure the safety of medical treatment and the quality of hospital infection control through the monitoring of environmental hygiene. According to a report from the China Centers for Disease Control and Prevention, as of February 11, 2020 medical staff members were infected with the new coronavirus (including confirmed cases, suspected cases, clinically diagnosed cases, and asymptomatic infections, of which 1716 were confirmed cases), indicating that infection through no occupational exposure may occur ^[8], but more evidence is needed to support this hypothesis. Meanwhile, it is extremely important to ensure the safety of medical treatments and the quality of hospital infection control by monitoring hospital environmental hygiene.

However, sensitive and effective methods for monitoring hospital-acquired infection control are still limited. Here, by monitoring COVID-19 and detecting nucleic acid, we reveal clinical data indicating hospital environmental hygiene levels and provide a way to monitor pathogenic microorganism contamination and nosocomial infections.

2.Methods

2.1 Participant Characteristics

Fifteen suspected patients and one confirmed intensive care patient (ORF1ab and N were positive according to RT-PCR) were hospitalized in two isolation areas at the First Hospital of Jilin University, a unit for COVID-19 detection. The hospital carries out district management for these two types of patients, and each area includes a nursing station. We detected 158 samples of COVID-19 from the area at risk of contamination. This study was approved by the Ethics Committee of the First Hospital of Jilin University (Changchun, China).

Table 1: Collection of 158 samples from the area at risk of contamination

Risk level	Classification	Sample collection points	
High-risk area	CONTAMINATED AREA	Isolation ward area 1 (2F)	Isolation wards 1-15 door handle
			Isolation wards 1-15 general subject surface
			Isolation wards 1-15 indoor air
			Isolation wards 1-15 window frames
			Isolation wards 1-15 sides of a toilet handle
			Nurse station general subject surface
			Nurse station indoor air
			Corridor exit general subject surface
			General subject surface of the refuse storage area near the nurse station
		Isolation ward area 2 (B1)	Isolation ward 16 door handle
			Isolation ward 16 general subject surface
			Isolation ward 16 indoor air
			Isolation ward 16 window frames

			Isolation ward 16 sides of a toilet handle
			Nurse station indoor air
		Fever clinic 2 consulting rooms	Door handle (inside and outside)
			General subject surface
			Indoor air
			Computer keyboard
		Fever clinic observation room	Door handle (inside and outside)
			General subject surface
			Indoor air
			Computer keyboard
		Fever clinic laboratory	Door handle (inside and outside)
			General subject surface
			Indoor air
			Computer keyboard
		Fever clinic blood collection room	General subject surface
			Indoor air
Medium-risk area	SEMICONTAMINATED AREA	Isolation ward area buffer room	Door handle
			General subject surface
			Indoor air
			Doctor's office window frames
			Refuse storage area wall surface
			Disinfection bucket surface
			Dressing room ground
			Medical staff of infection department protective clothing
		Pre-check triages	General subject surface
			Infrared thermometer
Low-risk area	CLEAN AREA	Isolation ward area working area	Entrance corridor general subject surface
			Indoor air
			Entrance corridor window frames
			Physician's office door handle
			Physician's office general subject surface
			Director's office door handle
			Door handle

104

105 2.2 Sampling and Sample Processing

106 The environmental monitoring methods referenced the hospital sanitation standards

107 (GB15982-2012). All air was collected by two methods: natural sedimentation^[9] and

a microbial air sampler (MAS-100 ECO), for which the stream of air was set to exactly 100 liters/minute (Merck, Germany) ^[10]. Environmental surfaces were sampled using swabs.

2.3 Primer and Probe Sequences

Two sequence regions (ORF1ab and N) that are highly conserved among sarbecoviruses were selected for primer and probe design. The primer and probe sequences for the ORF1ab gene assay were 5'-TGGGGYTTTACRGGTAACCT-3' (forward; Y = C/T, R = A/G), 5'-AACRCGCTTAACAAAGCACTC-3' (reverse; R = A/G) and 5'-TAGTTGTGATGCWATCATGACTAG-3' (probe, in 5'-FAM/ZEN/3'-IBFQ format; W = A/T), and the primer and probe sequences for the N gene assay were 5'-TAATCAGACAAGGAACTGATTA-3' (forward), 5'-CGAAGGTGTGACTTCCATG-3' (reverse) and 5'-GCAAATTGTGCAATTTGCGG-3' (probe, in 5'-FAM/ZEN/3'-IBFQ format). The expected amplicon sizes of the ORF1ab and N gene assays were 132 bp and 110 bp, respectively^[11]. All primers and probes were purchased from a commercial source (Integrated DNA Technologies). The primer and probe sequences were subsequently confirmed to have perfect matches with other COVID-19 genome sequences available from the Global Initiative on Sharing All Influenza Data (GISAID; <https://www.gisaid.org/>; accession numbers: EPI_ISL_402119, EPI_ISL_402120, EPI_ISL_402121, EPI_ISL_402123 and EPI_ISL_402124; accessed 12 January 2020).

2.4 RNA Extraction

All the collected samples were inactivated by heating at 56 °C for 30 minutes. For RNA extraction from 600 µL of 0.9% NaCl solution in a 2.0 mL tube, the samples

were centrifuged at 12000 g for 10 min and incubated for 30 min, the supernatant was discarded, 50 µL of RNA release agent was added to the tube, and the sample was mixed and incubated for 10 min.

2.5 Real-time Reverse-Transcription PCR

A 50-µL reaction contained 20 µL of RNA and 30 µL of 1× reaction buffer consisting of 26 µL of COVID-19 PCR mix (containing primers (4.62%), probes (1.15%), dNTPs (3.85%), MgCl₂ (0.77%), RNasin (0.48%), and PCR buffer (89.13%) (Shengxiang, Hunan, China)) and 4 µL each of COVID-19 PCR enzyme mix (containing RT enzyme (62.5%) and Taq enzyme (37.5%) (Shengxiang, Hunan, China)). Thermal cycling was performed at 50 °C for 30 min for reverse transcription, followed by 95 °C for 1 min and then 45 cycles of 95 °C for 15 s and 60 °C for 30 s. Participating laboratories used a Hongshi SLAN 96 S instrument (Hongshi, Shanghai, China). A sample was considered positive when the qPCR Ct value was ≤40.

3. Results

A total of 158 samples were collected from the area at risk of contamination. The two positive areas were the surfaces of the nurse station in the isolation area with suspected patients and the air of the isolation ward with an intensive care patient (Tables 2 and 3). We found that the virus was present both on surfaces and in the air. The total positive rate was 1.26% (2/158). The positive rates of the air and surface samples were 3.57% (1/28) and 0.77% (1/130), respectively.

Table 2: Air monitoring results for the different risk areas

Type		Negative	Positive	Positive rate
High-risk area	Isolation ward	19	1	5%
	Fever clinic consulting rooms	2		
	Fever clinic observation room	1		
	Fever clinic laboratory	1		
	Fever clinic blood collection room	1		
Middle-risk area	Isolation ward nurse station	1	0	0
	Isolation ward buffer room	1	0	0
Low-risk area	Isolation ward clean area	1	0	0
Total		27	1	3.57%

Table 3: Surface monitoring results for the different risk areas

Type		Negative	Positive	Positive rate
High-risk area	Isolation ward door handle, general surface	51	0	0%
	Isolation ward toilet handle	6	0	0
	Fever clinic consulting rooms	3	0	0
	Fever clinic observation room	3	0	0
	Fever clinic laboratory	6	0	0
	Fever clinic blood collection room	16	0	0
	Protective clothing for medical staff in the infection department	1	0	0
Middle-risk area	Pre-check triages	11	0	0
	Isolation ward nurse station (keyboard)	5	1	16.67%
	Isolation ward buffer room	3	0	0
Low-risk area	Isolation ward clean area	12	0	0
	Clean area keyboards	9	0	
Total		129	1	0.77%

Based on the original 24 hours of ultraviolet air filtering and 1000-2000 mg/L chlorine-containing disinfectant for ambient air and floor disinfection, the frequencies and duration times of air disinfections were extended. Key surfaces such as computer keyboards that were easily overlooked were clearly noted and carefully disinfected. The samples from the positive area and indoor air were collected 24 hours later, and the test results were negative.

4. Discussion

Our positive results in nucleic acid tests indicated that viruses were present in the air of an isolation ward with intensive care patients. The hermetic spaces of isolation wards are weak in air flow, which may cause high contact with the virus. Additionally, the confirmed patient underwent tracheal intubation the day before the samples were collected, and this procedure produced large amounts of aerosols that spread in the wards and seriously polluted the air. Siegel et al. reported that medical ventilators might generate respiratory aerosols that have been associated with an increased risk of occupationally acquired infections among healthcare personnel ^[12]. Therefore, it is necessary to increase the intensity of disinfection for isolation wards with intensive care patients. For example, we should enhance ventilator exhaust management by adding ventilator exhaust port filters.

The suspected and confirmed patients stay a relatively long time in the high-risk area and may influence the environment. The intrusive and other operations that can easily produce aerosols may cause serious air pollution. The sites under observation without patients and operations have less impact on the environment. The samples from the same collection points were tested after a series of infection prevention and control measures were taken, such as continuous use of an air disinfection machine for the disinfection of ward air; extending the frequency of disinfection for ground and object surfaces; changing gloves and cleaning hands after operations and leaving the ward; and covering computer keyboards and changing the cover each day. Following the implementation of these measures, the results were negative, showing the

effectiveness of disinfection. In the high-risk areas, such as hospital pre-check triages and the fever clinic, due to the open environment and large amount of air flow, the results were negative.

The previous research reports that the routine cleaning and disinfection procedures are the most likely cause of the deficiency of disinfection in the hospital environment, leading to the spread of pathogens ^[13]. In our study, viruses were detected on the surfaces of the nurse station in the isolation area with suspected patients, suggesting that we need to strengthen the surface disinfection of nurse stations, especially focusing on computer keyboards, mice and types of equipment that are not easy to disinfect. A large number of studies have shown that the hands or gloves of medical staff members may be contaminated by contact with environmental surfaces contaminated by pathogens^[13-15]. In the process of diagnosis and treatment, pathogens may be transmitted to patients and cause hospital-associated infections. Therefore, it is also very important to further strengthen the hand hygiene of the medical staff.

We need to formulate refinement measures based on environmental hygiene monitoring data to improve the quality of hospital infection control. Due to fact that “uncultivable” microbes widely exist, traditional methods used to measure the effect of hospital infection control, such as sedimentation (exposing a microbial growth plate to the environment) have shown limited effects and poor sensitivity. For laboratories and testing organizations, high concentrations of nucleic acid may exist in aerosols, which may influence the test results and the operator’s safety and may even cause spread of the disease. Nucleic acid detection provides an effective method with

which to monitor the environment and to evaluate the effectiveness of disinfection, especially for highly contagious diseases or pathogenic microorganisms with a potential aerosol risk, such as COVID-19. These nucleic acid detection results are very important for standardized hospital infection control. This study also increases our understanding of the environmental challenges corresponding to the reemergence of COVID-19 or similar viruses.

The data obtained in this study suggest that COVID-19 exists in the air of isolation wards with intensive care patients and existed on the surfaces of nurse stations and that medical staff encounter this virus in the air and on surfaces during patient care activities. An effective disinfection procedure may reduce the biosafety risk.

For hospital infection control, we suggest:

1. Different people should be responsible for the isolation observation area and the isolation ward area; crossing the areas should be strictly prohibited.
2. Objects in each area should be used exclusively; equipment that must be used for special purposes should be sterilized before being used in other areas.
3. Hand hygiene rules should be observed strictly, thoroughly and at all times.
4. Additional gloves, disposable isolators, protective screens or hoods should be worn when an operation may cause the spattering of blood, bodily fluid or aerosol, and these items should be discarded immediately to avoid the pollution of other areas.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (Nos. 30972610 and 81273240), the Jilin Province Science and Technology Agency (Nos. 20160101037JC, 20170622009JC, 2017C021, and 2017J039), the Norman Bethune Program of Jilin University (2012206), and Open Project Funding From the State Key Laboratory of Kidney Diseases. We thank the patients and their families for participating in this study. We would also like to acknowledge the professional manuscript services of American Journal Experts.

Disclosure Statement

The authors have no conflicts of interest to declare.

References

1. Benvenuto, D., et al., The global spread of 2019-nCoV: a molecular evolutionary analysis. *Pathog Glob Health*. 2020, 12: 1-4.
2. Huang, C., et al., Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020, 395(10223): 497-506.
3. The, L., Emerging understandings of 2019-nCoV. *Lancet*. 2020, 395(10221): 311.
4. Anderson, D.J., et al., Enhanced terminal room disinfection and acquisition and infection caused by multidrug-resistant organisms and *Clostridium difficile* (the Benefits of Enhanced Terminal Room Disinfection study): a cluster-randomised, multicentre, crossover study. *Lancet*. 2017, 389(10071): 805-814.
5. Beggs, C., et al., Environmental contamination and hospital-acquired infection: factors that are easily overlooked. *Indoor air*. 2015, 25(5): 462-74.
6. Wang, D., et al., Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA*. 2020.
7. Phan, L.T., et al., Respiratory viruses in the patient environment. *Infect Control Hosp Epidemiol*. 2020, 41(3):259-266.
8. Chinese Center for Disease Control and Prevention, Beijing 102206, China. Epidemiology group of new coronavirus pneumonia emergency response mechanism, Chinese center for disease control and prevention, The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China, *Zhonghua Liu Xing Bing Xue Za Zhi*. 2020, 41(2):145-151.

- 271 9. Odebode A, et al., Airborne fungi spores distribution in various locations in Lagos,
272 Nigeria. Environ Monit Assess. 2020, 192(2):87.
- 273 10. Irie K, et al., Investigation of the detection ability of an intrinsic
274 fluorescence-based bioaerosol detection system for heat-stressed bacteria. PDA
275 J Pharm Sci Technol. 2014, 68(5):478-93
- 276 11. Chu DKW, et al., Molecular Diagnosis of a Novel Coronavirus (2019-nCoV)
277 Causing an Outbreak of Pneumonia. Clin Chem. 2020, pii: hvaa029. doi:
278 10.1093/clinchem/hvaa029.
- 279 12. Siegel, J.D., et al., 2007 Guideline for Isolation Precautions: Preventing
280 Transmission of Infectious Agents in Health Care Settings. Am J Infect Control.
281 2007, 35(10 Suppl 2): S65-164.
- 282 13. Pirincci, E. and B. Altun, An analysis of hospital cleaning staff's attitudes and
283 conduct regarding hand hygiene and cleaning. Int J Occup Saf Ergon. 2016, 22(2):
284 241-5.
- 285 14. Pittet, D., et al., Evidence-based model for hand transmission during patient care
286 and the role of improved practices. Lancet Infect Dis. 2006, 6(10): 641-52.
- 287 15. Christensen, T.E., J.S. Jorgensen, and H.J. Kolmos, [The importance of hygiene for
288 hospital infections]. Ugeskr Laeger. 2007, 169(49): 4249-51