1 **Restoration of leukomonocyte counts is associated with viral** 2 **clearance in COVID-19 hospitalized patients**

3

4 Xiaoping Chen¹, Jiabin Ling², Pingzheng Mo¹, Yongxi Zhang¹, Qunqun
5 Jiang¹, Zhiyong Ma¹, Qian Cao¹, Wenjia Hu¹, Shi Zou¹, Liangjun Chen³,
6 Lei Yao⁴, Mingqi Luo¹, Tielong Chen¹, Liping Deng¹, Ke Liang¹, Shihui
7 Song¹, Rongrong Yang¹, Ruiying Zheng¹, Shicheng Gao¹, Xien Gui¹,
8 Hengning Ke¹, Wei Hou⁴, Åke Lundkvist², Yong Xiong¹

9

10 ¹ Department of Infectious Diseases, Zhongnan Hospital of Wuhan
11 University, No.169, Donghu Road, Wuchang District, 430071 Wuhan,
12 Hubei Province, China

13 ² Department of Medical Biochemistry and Microbiology, Zoonosis
14 Science Center, University of Uppsala, Uppsala, Sweden

15 ³ Department of Laboratory Medicine, Zhongnan Hospital of Wuhan
16 University, Wuhan, People's Republic of China

17 ⁴ State Key Laboratory of Virology/ Institute of Medical Virology/ Hubei
18 Province Key Laboratory of Allergy and Immunology, School of Basic
19 Medical Sciences, Wuhan University, 185 Donghu Road, Wuhan 430071,
20 Hubei Province, China

21

22

23 Corresponding to:

24 Xiaoping Chen, MD & PhD

25 E-mail address; alackcn@126.com

26 Tel: 86-27-67812880

27 Address: Department of infectious diseases, Zhongnan Hospital of

28 Wuhan University, No.169, Donghu Road, Wuchang District, Wuhan

29 City, Hubei Province, P.R China.

30

31 Yong Xiong, MD & PhD

32 E-mail address; yongxiong64@163.com

33 Tel: 86-27-67812880

34 Address: Department of infectious diseases, Zhongnan Hospital of

35 Wuhan University, No.169, Donghu Road, Wuchang District, Wuhan

36 City, Hubei Province, P.R China.

37

38

39 Summary

40 *Background:* Viral clearance is one important indicator for the recovery
41 of SARS-CoV-2 infected patients. Previous studies have pointed out that
42 suboptimal T and B cell responses can delay viral clearance in MERS-
43 CoV and SARS-CoV infected patients. The role of leukomonocytes in
44 viral clearance of COVID-19 patients is not yet well defined.

45 *Methods:* From January 26 to February 28, 2020, an observational study
46 was launched at the Department of Infectious Diseases, Zhongnan
47 Hospital of Wuhan University, Wuhan, China. We enrolled 25
48 laboratory-confirmed COVID-19 patients, whose throat-swab specimens
49 were tested positive for SARS-CoV-2 infection by qRT-PCR. To
50 investigate the factors that contribute to the viral clearance, we
51 comprehensively analyzed clinical records, counts of lymphocyte subsets
52 including CD3+, CD4+, CD8+ T cells, B cells and NK cells in the
53 patients who successfully cleared SARS-CoV-2, and compared to those
54 that failed to, after a standardized treatment of 8-14 days.

55 *Findings:* In 25 enrolled COVID-19 patients, lymphopenia was a
56 common feature. After the treatment, 14 out of the 25 enrolled patients
57 were tested negative for SARS-CoV-2. The patients that cleared the
58 infection had restored the numbers of CD3+, CD4+, CD8+ T cells and B

59 cells as compared to the still viral RNA positive patients, while the
60 recovered patients had a higher count of leukomonocytes.

61 *Conclusions:* By comparison of leukomonocytes counts in COVID-19
62 patients at different stages of the disease, we found that CD3+, CD4+,
63 CD8+ T cells and B cells appear to play important roles in viral clearance.
64 The restoration of leukomonocytes counts from peripheral blood can be
65 used as prognosis for the recovery of an COVID-19 infection. We
66 propose that restoration of leukomonocytes counts can be added to the
67 COVID-19 diagnostic guidance as a criterion for releasing and
68 discharging patients.

69 **Keywords.** COVID-19; SARS-CoV-2, leukomonocytes; viral clearance.

70 **Funding**

71 This study was funded by the Zhongnan Hospital of Wuhan University
72 Science, Technology and Innovation Seed Fund, grant number
73 znp2018007 and the Swedish Research Council, grant number 2017-
74 05807. The funders had no role in study design, data collection or
75 analysis, decision to publish or preparation of the manuscript. The authors
76 declared no competing interests.

77 **Introduction**

78 In December 2019, a cluster of novel coronavirus-associated pneumonia
 79 cases was emerging in Wuhan, China [1]. The disease rapidly spread
 80 from Wuhan and caused a wide epidemic in China. The World Health
 81 Organization (WHO) has recently named the disease as Coronavirus
 82 Disease-2019 (COVID-19) [2]. The etiology of the disease is a novel β -
 83 coronavirus, taxonomically named as severe acute respiratory syndrome
 84 coronavirus 2 (SARS-CoV-2). Until February 29, 2020, SARS-CoV-2
 85 causes an epidemic worldwide. More than 80,000 people in over 60
 86 countries or territories have been contracted of COVID-19, and more
 87 than 3000 people have died [3].

88 The clinical manifestations of COVID-19 cases range from an
 89 asymptomatic course to severe acute respiratory symptoms and even acute
 90 respiratory failure [4-7]. Transmission occurs from human to human with
 91 nosocomial transmissions. The incubation time of COVID-19 has been
 92 reported from 0 to 24 days and also asymptomatic carriers can be
 93 infectious [8].

94 As a zoonotic virus, the natural host of SARS-CoV-2 is believed to be
 95 bats [9], however, the process of cross-species transmission(s) is still not
 96 known. The very first cases were associated with the Huanan Seafood
 97 Wholesale Market in Wuhan, China [10]. There is no effective antiviral

98 drug available yet. Although the pathogenesis of COVID-19 is still
 99 unclear, lymphopenia has been found in most COVID-19 patients [10-12].
 100 This feature has also been found in the other two fatal human coronavirus
 101 infections caused by severe acute respiratory syndrome coronavirus
 102 (SARS-CoV) and the Middle East respiratory syndrome coronavirus
 103 (MERS-CoV) [13, 14]. Previous studies showed that T cells, especially
 104 CD4⁺ and CD8⁺ T cells, play a significant antiviral role in balancing the
 105 combat against MERS-CoV or SARS-CoV infections [15]. These results
 106 suggested that leukomonocytes counts could be critical indicators
 107 associated with severity and the disease outcome. In this study, we
 108 retrospectively analyzed 25 COVID-19 cases to elucidate the dynamic of
 109 lymphocyte subset changes in the peripheral blood and their roles during
 110 the viral clearance.

111

112 **Methods**

113 *Study design*

114 From January 26 to February 28, 2020, 25 COVID-19 patients who had
 115 been repeatedly analyzed for lymphocyte subsets and SARS-CoV-2 RNA
 116 were enrolled in the study. Informed consents were obtained from all

117 patients upon admission to the Department of Infectious Diseases,
118 Zhongnan Hospital of Wuhan University, Wuhan, China.

119

120 *Data collection*

121 The medical records, including epidemiological and demographic
122 information, clinical manifestations, laboratory data, and outcome of
123 disease, were collected. To avoid subjective biases, two investigators
124 reviewed the data independently and made a final consensus.

125 Peripheral blood (100 µl) was collected from patients at admission to
126 hospital and after 7 – 10 days of standard therapy (Version 6, 2020 Feb
127 18,

128 [http://www.nhc.gov.cn/yzygj/s7653p/202002/8334a8326dd94d329df351](http://www.nhc.gov.cn/yzygj/s7653p/202002/8334a8326dd94d329df351d7da8aefc2.shtml)
129 [d7da8aefc2.shtml](http://www.nhc.gov.cn/yzygj/s7653p/202002/8334a8326dd94d329df351d7da8aefc2.shtml)). Lymphocyte subsets such as CD3+, CD4+, CD8+ T
130 cells, B cells and NK cells were stained by using the BD Multitest™ IMK
131 kit (BD Ltd., San Jose, CA, USA) according to the manufacturer's
132 instruction. The lymphocyte subset analyses were performed by flow
133 cytometry (BD FACSCanto™ II Flow Cytometer).

134 Serial throat-swab samples were taken from patients at the admission to
135 hospital during 8 – 14 days of standardized treatment (see Table 1)
136 according to the recommendation in the guideline. The swab samples

137 were placed in 150 μ L of virus preservation solution and viral RNA
138 subsequently extracted as described earlier [5, 16]. Two target genes of
139 the SARS-CoV-2 genome, including the open reading frame 1ab
140 (ORF1ab) and the nucleocapsid protein (N) gene, were simultaneously
141 amplified and tested in the real-time RT-PCR assay by using a SARS-
142 CoV-2 nucleic acid detection kit according to the manufacturer's protocol
143 (Shanghai Bio-Germ Medical Technology Co Ltd, Shanghai, China). The
144 reaction mixture consisted of 12 μ L reaction buffer, 4 μ L of enzyme
145 solution, 3 μ L of diethyl pyrocarbonate-treated water, 2 μ L of RNA
146 template, and 4 μ L of probe/primers solution, which contained two sets of
147 probe primers: one set targets towards to ORF1ab (forward prime 5'-
148 CCCTGTGGGTTTTACACTTAA-3'; reverse primer 5'-
149 ACGATTGTGCATCAGCTGA-3'; and the probe 5'-VIC-
150 CCGTCTGCGGTATGTGGAAAGGTTATGG-BHQ1-3'), and another
151 targets towards to N (forward primer 5'-
152 GGGGAAGTTCTCCTGCTAGAAT-3'; reverse primer 5'-
153 CAGACATTTTGCTCTCAAGCTG-3'; and the probe 5'-FAM-
154 TTGCTGCTGCTTGACAGATT-TAMRA-3').

155 The results of the real-time RT-PCR analyzes were interpreted as: a
156 cycle threshold value (Ct-value) less than 37 was defined as a positive
157 result; a Ct-value of 40 or more was defined as a negative result; a

158 medium load result, defined as a Ct-value of 37 to less than 40, required
159 another test for confirmation. The patients who repeatedly tested SARS-
160 CoV-2 RNA negative for at least two times with an interval of more than
161 1 day were regarded as viral negative. These diagnostic criteria were
162 based on the recommendation by the National Institute for Viral Disease
163 Control and Prevention, China
164 (<http://ivdc.chinacdc.cn/gjhz/jldt/202002/P020200209712430623296.pdf>).

165 *Statistical analysis*

166 We used the SPSS 17.0 software package for the statistical analyses. We
167 used X^2 tests or Fisher's exact tests for categorical variables. For
168 measurement of the data, a normal distribution was tested first and then
169 proceeded to t-test, expressed as the mean $\pm$ standard deviations;
170 otherwise the Mann-Whitney U test was used for non-normal distribution
171 data, expressed in terms of median (25%–75% interquartile range, IQR).
172 A p value of < 0.05 was considered statistically significant.

173

174 *The principle of medical ethics*

175 This study was approved by the ethics board in Zhongnan Hospital of
176 Wuhan University, Wuhan, China (No.2020011).

177

178 **Results**

179 *Baseline Characteristics of COVID-19 patients*

180 A total of 25 patients, 11 men and 14 women were enrolled in this study
 181 (Table 1). The ages of the patients spanned from 25 to 80 years (average
 182 51.4 ± 16.6 years). Fourteen people described a previous exposure history
 183 to the source of infection, either from the Huanan Seafood Wholesale
 184 Market or through direct contacts with COVID-19 patients. The most
 185 common symptoms at the onset of illness were: fatigue (80.0%), fever
 186 ($37.4\text{--}39.1^\circ\text{C}$, 75.0%), myalgia (68.0%), cough (64.0%), and less
 187 common: dyspnea (28.0%) and diarrhea (20.0%). Seven cases had
 188 underlying diseases such as hypertension, cardiovascular disease,
 189 diabetes, malignancy or chronic liver disease (Table 1).

190 **Laboratory analysis before and after treatment**

191 Laboratory analysis were performed before the treatment. The
 192 biochemical tests included alanine aminotransferase, aspartate
 193 aminotransferase, aspartate aminotransferase, creatinine, and D-dimer,
 194 which all were found normal (Table 2). However, blood counts of the
 195 patients showed leucopenia ($4.5 \pm 1.9 \times 10^9/\text{L}$) and lymphopenia (< 1.3
 196 $\times 10^9/\text{L}$). The counts of CD3+ T cells, CD4+ T cells, CD8+ T cells, B
 197 cells, and NK cells were found lower than normal values. Interleukin-6

198 had increased to a mean of 16.1 (3.7 – 31.4) pg/ml. Five patients had an
199 increase of procalcitonin (≥ 0.05 ng/mL) (Table 2).

200 The treatment was mainly of supportive care (Table 1). Twenty-four
201 patients were given antiviral therapy including arbidol (orally, 200 mg,
202 three times per day), and oxygen support. Antibiotic therapy, both orally
203 and intravenous, were given as described in Table 1. Fourteen patients
204 received corticosteroids to suppress an excessive inflammatory activation.
205 After standardized treatment of 8–14 days, according to the laboratory
206 analysis of viral RNA, fourteen patients showed viral clearance and 11
207 patients were still positive for SARS-CoV-2 viral RNA (Table 1).

208 When comparing the group that showed viral clearance and the group that
209 failed, we found no significant association to sex, epidemiological
210 exposure history, comorbidities, onsets of signs and symptoms, or
211 treatment, but with the age and days of hospitalization. The mean days of
212 hospitalization in the group that showed viral clearance were 12 days
213 (10.0 – 15.0), which was shorter as compared to the group that failed to
214 clear the virus, for which the average days were 21 (18.0 – 23.0) days.
215 Age can play an important role during the viral clearance: patients
216 without viral clearance were older (60.2 ± 16.9 , y) than patients with viral

clearance (44.5 ± 13.0 y), showing a statistically significant difference (Table 1).

Association between peripheral lymphocyte counts and disease outcomes

Since lymphopenia was found in all the patients in the current study and also earlier reported elsewhere [10], we further analyzed the association between peripheral lymphocyte counts and disease outcomes. In the 14 patients with viral clearance after the 8 – 14 days of treatment, the counts of CD3+ T cells, CD4+ T cells, CD8+ T cells, and B cells were restored close to normal levels and with a significant difference as compared to the counts at admission in 12 patients. In the remaining 2 patients with viral clearance the leukomonocytes counts were still at low levels (Figure 1, A-E). However, in the patients without viral clearance (No.=11), no differences were seen before and after treatment regarding the CD3+, CD4+, CD8+, B, and NK cell counts (Figure 1, F-J). The association between peripheral lymphocyte counts and viral detection suggested that higher counts of CD3+ T cells, CD4+ T cells, CD8+ T cells, and B cells had a significant impact on the viral clearance.

236 To further monitor the disease outcome of those 11 patients who were
237 still positive for SARS-CoV-2 virus RNA, we extended the same
238 treatment to 15 – 23 days (Figure 2). At 15 –23 days, 7 patients were
239 negative for viral RNA detection (data for leukomonocyte subsets was
240 available for 6 of these 7 patients), but 4 patients were still positive (all
241 these four had been tested for leukomonocyte subsets). The 4 viral RNA
242 positive patients had low counts of CD3+ T cells ($< 1200 \times 10^9/L$), CD4+
243 T cells ($< 700 \times 10^9/L$), CD8+ T cells ($< 500 \times 10^9/L$), B cells (< 400
244 $\times 10^9/L$), and NK cells ($< 400 \times 10^9/L$). Due to the small sample size, no
245 statistically significant differences for the cell counts were observed,
246 except for the NK cell counts, which showed a significant decrease in the
247 4 viral RNA positive patients (Figure 2, J). However, we found a similar
248 trend that the increased counts of CD3+ T cells and CD8+ T cells
249 corresponded to cases which were negative for viral RNA detection (5
250 out of 6). One case that still had low counts of CD3+ T cells, CD4+ T
251 cells, CD8+ T cells, B cells and NK cells was found negative for viral
252 RNA.

253

254 **Clinical outcome**

255 We observed the clinical course of 25 COVID-19 patients within 23 days
256 of treatment. During the first phase of treatment, at 8 – 14 days, fourteen

257 patients showed viral clearance and 12 of them were discharged from the
 258 hospital according to the guideline. The treatment of oxygen support
 259 therapy and clinical observation was continued for the remaining 2
 260 patients in this group. Eleven viral RNA positive patients received a
 261 second phase of treatment between 15– 23 days. Until February 28, 7 of
 262 these 11 patients turned out to be viral RNA negative. Four of these 7
 263 patients were then discharged from the hospital, while the other 3 (one
 264 had a liver disease and the other two still had significant clinical
 265 symptoms), together with the 4 patients who were still positive for viral
 266 RNA, were kept under treatment (Table 1). In total, 9 out of the 25
 267 COVID-19 patients were still hospitalized after 23 days of clinical
 268 treatment.

269

270 **DISCUSSION**

271 In this report, we retrospectively investigated the changes of CD3+,
 272 CD4+, CD8+, B, and NK cell counts in peripheral blood of 25 COVID-
 273 19 patients during the viral infection and the mechanisms for viral
 274 clearance. We found that restoration of CD3+, CD4+, CD8+ T cells, B
 275 cells and NK cells was associated the viral clearance in COVID-19
 276 patients.

277 According to the Guideline of the treatment of COVID-19 (Version 6,
278 2020 Feb 18,
279 [http://www.nhc.gov.cn/yzygj/s7653p/202002/8334a8326dd94d329df351](http://www.nhc.gov.cn/yzygj/s7653p/202002/8334a8326dd94d329df351d7da8aefc2.shtml)
280 [d7da8aefc2.shtml](http://www.nhc.gov.cn/yzygj/s7653p/202002/8334a8326dd94d329df351d7da8aefc2.shtml)), the diagnosis of COVID-19 infection is based on an
281 epidemiological history of contact with COVID-19 patients or travel to
282 endemic areas, combined with laboratory testing including viral RNA
283 detection and radiological findings. COVID-19 patients have similarities
284 of the clinical features with other coronavirus infections. Most patients
285 had fever, cough, myalgia and fatigue, and less often showed symptoms
286 of dyspnoea, haemoptysis and diarrhea [10]. Half of the cases had
287 comorbidities such as diabetic, hypertension and cardiovascular diseases
288 [10]. In our study, we also found some cases that had underlying diseases
289 but without any association with the disease outcome. Age is a risk factor
290 for a more severe disease outcome, because of the generally inferior
291 function of the immune system among older people. Based on the
292 guideline, the patients who have normal body temperature for more than
293 three days, mitigation of respiratory symptoms, improvement of
294 radiological evidences for lesions, and had been tested negative for
295 specific SARS-CoV-2 RT-PCR at least twice, can be discharged from the
296 hospital. The average hospitalized time was 17 days (IQR 11.5 – 21.5),
297 and even as long as 23 days for the patients that were released from our
298 study [5], suggesting the time for SARS-CoV-2 clearance is also around

299 17 days or longer. This was also been observed in an earlier study where
300 the shedding of SARS-CoV-2 in saliva continued up to 11 days [17].

301 Until now, there is no effective antivirals available against SARS-CoV-2.
302 To eliminate the virus an effective immunological response with a
303 minimum of immunopathological effects is required [18]. The antigen
304 presentation requires inhibitory alveolar macrophages and dendritic cells
305 (DCs). Successful stimulation can produce viral specific CD4 T cells,
306 which are involved in the development of a specific humoral response,
307 including neutralizing antibodies that can block the viral entry to the cells
308 [19]. CD8+ cytotoxic T cells are required for recognition and killing of
309 the infected cells, further playing a crucial role in SARS-CoV clearance
310 [20, 21]. However, in the biopsy samples from the patients who died of
311 COVID-19, histological examination showed pulmonary oedema with
312 hyaline membrane formation, and interstitial mononuclear inflammatory
313 infiltrates in the lung tissue, suggesting that acute respiratory failure was
314 the cause of death, underlying mechanism that an over-activated
315 immunity injured the lung tissue [12]. Among SARS patients, high
316 expression of dysregulated chemokines and cytokines were found in
317 blood and lungs [22], due to the reason of aberrant immune response
318 caused by the viruses [13, 14]. The delayed or suboptimal immune
319 responses would make chances for SARS-CoV and MERS-CoV to

320 escape from the immune responses, and survive and replicate in the host
 321 cells, leading a further delay of the viral clearance [15, 23]. In our study,
 322 COVID-19 patients at older ages needed a longer time for the recovery,
 323 likely due to the fact that aging is associated with a set of functional and
 324 structural alterations in the immune system [24]. Immune-modulators, as
 325 corticosteroids, have been empirically used in both SARS and COVID-19
 326 patients [10, 25]. Since corticosteroids might have a decline effect of
 327 circulating specific B and T cell subsets [26], the usage on COVID-19
 328 patients should be carefully evaluated before considered [12, 27].

329 A rapid and generalized lymphopenia has been observed as a prominent
 330 part in SARS-CoV and MERS infections [28]. Lymphopenia has also
 331 been observed in other viral infections such as measles virus and avian
 332 influenza virus, swine foot-and-mouth disease virus, respiratory syncytial
 333 virus, and HIV [29-33]. The possible reasons would be lymphocyte
 334 sequestration in the lung tissues, or immune-mediate lymphocyte
 335 destruction. In MERS infections, the function of bone marrow or thymus
 336 suppressed, and furthermore, T cells underwent apoptosis [34-36]. In
 337 COVID-19 patients, peripheral CD4 and CD8 T cell counts were found
 338 decreased but, they were also hyperactivated, which could result in severe
 339 immune injuries [12].

340 In the present study, we followed the current guidelines and used a
 341 specific SARS-CoV-2 realtime RT-PCR for viral detection. Surprisingly,
 342 one lymphopenia patient (37y, male, chronic liver disease) was found
 343 negative by the realtime RT-PCR and his symptoms had been improved
 344 but he is still in hospital. Until now, specific viral RNA detection based
 345 on realtime RT-PCR is commonly used in the clinical diagnosis. The
 346 usage of realtime RT-PCR is versatile in the detection of SARS-CoV-2 in
 347 the respiratory tract specimen or throat swab, but we must keep in mind
 348 that those methods always are followed by the probability of false
 349 positive/negative. Besides, around 14% of the discharged COVID-19
 350 patients were tested as positive again on follow up samples
 351 ([https://www.caixinglobal.com/2020-02-26/14-of-recovered-covid-19-](https://www.caixinglobal.com/2020-02-26/14-of-recovered-covid-19-patients-in-guangdong-tested-positive-again-101520415.html)
 352 [patients-in-guangdong-tested-positive-again-101520415.html](https://www.caixinglobal.com/2020-02-26/14-of-recovered-covid-19-patients-in-guangdong-tested-positive-again-101520415.html)), showing
 353 the recurrence of SARS-CoV-2. This imply that viral detection cannot be
 354 used as a single criterion for COVID-19 diagnosis. As a highly
 355 contiguous disease with a fatality of approximately 2.3% and up to 49.0%
 356 in critical cases [7], every hospitalized COVID-19 patient must be
 357 evaluated carefully before discharged. The peripheral leukomonocytes
 358 counts could provide more information in the evaluation of COVID-19
 359 patients.

360 In conclusion, we retrospectively analyzed 25 COVID-19 cases and
 361 found that restoration of peripheral leukomonocytes could help the
 362 clearance of SARS-CoV-2. We propose here that leukomonocytes counts
 363 may serve as a valid prognosis of immune reconstitution for recovery of
 364 COVID-19 infection and that can update the current COVID-19 diagnosis
 365 guideline.

366

367 Acknowledgement

368 An especially strong appreciation goes to all the clinical physicians,
369 health caregivers, clinical laboratory personals, epidemiologists, and
370 researchers who have been fighting against the SARS-CoV-2 epidemic.
371 We also acknowledge the supports from both nationwide and
372 international resources.

373

374 Reference

- 375 1. Li Q, Guan X, Wu P, et al. Early Transmission Dynamics in Wuhan, China, of
376 Novel Coronavirus-Infected Pneumonia. *N Engl J Med* **2020**.
- 377 2. Naming the coronavirus disease (COVID-2019) and the virus that causes it.
378 Available at: [https://www.who.int/emergencies/diseases/novel-coronavirus-](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-causes-it)
379 [2019/technical-guidance/naming-the-coronavirus-disease-\(covid-2019\)-and-](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-causes-it)
380 [the-virus-that-causes-it](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-causes-it).
381 Available at: <https://promedmail.org/>.
- 382 3. Pan Y, Guan H, Zhou S, et al. Initial CT findings and temporal changes in
383 patients with the novel coronavirus pneumonia (2019-nCoV): a study of 63
384 patients in Wuhan, China. *Eur Radiol* **2020**.
- 385 4. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized
386 Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China.
387 *JAMA* **2020**.
- 388 5. Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with
389 Pneumonia in China, 2019. *N Engl J Med* **2020**; 382(8): 727-33.
- 390 6. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the
391 Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a
392 Report of 72314 Cases From the Chinese Center for Disease Control and
393 Prevention. *JAMA* **2020**.
- 394 7. Bai Y, Yao L, Wei T, et al. Presumed Asymptomatic Carrier Transmission of
395 COVID-19. *JAMA* **2020**.
- 396 8. Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a
397 new coronavirus of probable bat origin. *Nature* **2020**.
- 398 9. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019
399 novel coronavirus in Wuhan, China. *Lancet* **2020**; 395(10223): 497-506.
- 400 10. Chan JF, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated
401 with the 2019 novel coronavirus indicating person-to-person transmission: a
402 study of a family cluster. *Lancet* **2020**; 395(10223): 514-23.
- 403 11. Xu Z, Shi L, Wang Y, et al. Pathological findings of COVID-19 associated with
404 acute respiratory distress syndrome. *Lancet Respir Med* **2020**.

13. Wong RS, Wu A, To KF, et al. Haematological manifestations in patients with severe acute respiratory syndrome: retrospective analysis. *BMJ* **2003**; 326(7403): 1358-62.
14. Lau SK, Lau CC, Chan KH, et al. Delayed induction of proinflammatory cytokines and suppression of innate antiviral response by the novel Middle East respiratory syndrome coronavirus: implications for pathogenesis and treatment. *J Gen Virol* **2013**; 94(Pt 12): 2679-90.
15. Liu J, Zheng X, Tong Q, et al. Overlapping and discrete aspects of the pathology and pathogenesis of the emerging human pathogenic coronaviruses SARS-CoV, MERS-CoV, and 2019-nCoV. *J Med Virol* **2020**.
16. Chen H, Guo J, Wang C, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *The Lancet* **2020**.
17. To KK, Tsang OT, Chik-Yan Yip C, et al. Consistent detection of 2019 novel coronavirus in saliva. *Clin Infect Dis* **2020**.
18. Newton AH, Cardani A, Braciale TJ. The host immune response in respiratory virus infection: balancing virus clearance and immunopathology. *Semin Immunopathol* **2016**; 38(4): 471-82.
19. Chen J, Lau YF, Lamirande EW, et al. Cellular immune responses to severe acute respiratory syndrome coronavirus (SARS-CoV) infection in senescent BALB/c mice: CD4+ T cells are important in control of SARS-CoV infection. *J Virol* **2010**; 84(3): 1289-301.
20. Janice Oh HL, Ken-En Gan S, Bertoletti A, Tan YJ. Understanding the T cell immune response in SARS coronavirus infection. *Emerg Microbes Infect* **2012**; 1(9): e23.
21. Zhao J, Zhao J, Perlman S. T cell responses are required for protection from clinical disease and for virus clearance in severe acute respiratory syndrome coronavirus-infected mice. *J Virol* **2010**; 84(18): 9318-25.
22. Jiang Y, Xu J, Zhou C, et al. Characterization of cytokine/chemokine profiles of severe acute respiratory syndrome. *Am J Respir Crit Care Med* **2005**; 171(8): 850-7.
23. Min CK, Cheon S, Ha NY, et al. Comparative and kinetic analysis of viral shedding and immunological responses in MERS patients representing a broad spectrum of disease severity. *Sci Rep* **2016**; 6: 25359.
24. Sadighi Akha AA. Aging and the immune system: An overview. *J Immunol Methods* **2018**; 463: 21-6.
25. Stockman LJ, Bellamy R, Garner P. SARS: systematic review of treatment effects. *PLoS Med* **2006**; 3(9): e343.
26. Olnes MJ, Kotliarov Y, Biancotto A, et al. Effects of Systemically Administered Hydrocortisone on the Human Immune. *Sci Rep* **2016**; 6: 23002.
27. Wu F, Zhao S, Yu B, et al. A new coronavirus associated with human respiratory disease in China. *Nature* **2020**.
28. Li T, Qiu Z, Zhang L, et al. Significant changes of peripheral T lymphocyte subsets in patients with severe acute respiratory syndrome. *J Infect Dis* **2004**; 189(4): 648-51.
29. Okada H, Kobune F, Sato TA, et al. Extensive lymphopenia due to apoptosis of uninfected lymphocytes in acute measles patients. *Arch Virol* **2000**; 145(5): 905-20.
30. Zitzow LA, Rowe T, Morken T, Shieh WJ, Zaki S, Katz JM. Pathogenesis of avian influenza A (H5N1) viruses in ferrets. *J Virol* **2002**; 76(9): 4420-9.

- 455 31. Bautista EM, Ferman GS, Golde WT. Induction of lymphopenia and inhibition
456 of T cell function during acute infection of swine with foot and mouth disease
457 virus (FMDV). *Vet Immunol Immunopathol* **2003**; 92(1-2): 61-73.
- 458 32. O'Donnell DR, Carrington D. Peripheral blood lymphopenia and neutrophilia in
459 children with severe respiratory syncytial virus disease. *Pediatr Pulmonol*
460 **2002**; 34(2): 128-30.
- 461 33. Li CX, Li YY, He LP, et al. The predictive role of CD4(+) cell count and
462 CD4/CD8 ratio in immune reconstitution outcome among HIV/AIDS patients
463 receiving antiretroviral therapy: an eight-year observation in China. *BMC*
464 *Immunol* **2019**; 20(1): 31.
- 465 34. He Z, Zhao C, Dong Q, et al. Effects of severe acute respiratory syndrome
466 (SARS) coronavirus infection on peripheral blood lymphocytes and their
467 subsets. *Int J Infect Dis* **2005**; 9(6): 323-30.
- 468 35. Yang Y, Xiong Z, Zhang S, et al. Bcl-xL inhibits T-cell apoptosis induced by
469 expression of SARS coronavirus E protein in the absence of growth factors.
470 *Biochem J* **2005**; 392(Pt 1): 135-43.
- 471 36. Mubarak A, Alturaiki W, Hemida MG. Middle East Respiratory Syndrome
472 Coronavirus (MERS-CoV): Infection, Immunological Response, and Vaccine
473 Development. *J Immunol Res* **2019**; 2019: 6491738.

474

475

Table 1. Demographics, baseline characteristics, treatment, and clinical outcomes of 25 patients infected with SARS-CoV-2

	Total (n=25)	Viral clearance (n=14) ^a	without viral clearance (n=11) ^a	P value
Sex				
female	14(56.0%)	9(64.3%)	5(45.5%)	0.435
male	11(44.0%)	5(35.7%)	6(54.5%)	
Huanan Seafood Wholesale	14 (56.0%)	9 (64.3%)	5 (45.5%)	0.697
Market or patient infected with SARS-CoV-2 exposure				
Age, mean $\pm$SD, y	51.4 $\pm$ 11.6	44.5 $\pm$ 13.0	60.2 $\pm$ 16.9	0.015
Comorbidities				
Hypertension	3(12.0%)	0(0.0%)	3(25.0%)	0.072
Cardiovascular disease	2(8.0%)	1(7.0%)	1(9.1%)	1.000
Diabetes	2(8.0%)	0(0.0%)	2(18.2%)	0.183
Malignancy	1(4.0%)	0(0.0%)	1(9.1%)	0.440
COPD	0(0.0%)	0(0.0%)	0(0.0%)	-
Chronic liver disease	3(12.0%)	0(0.0%)	3(25.0%)	0.072
Signs and symptoms				
Fever	18 (75.0%)	8 (57.1%)	10 (90.9%)	0.090
Fatigue	20 (80.0%)	11 (78.6%)	9 (81.8%)	1.000
Myalgia	17 (68.0%)	8 (57.1%)	9 (81.8%)	0.234
Cough	16 (64.0%)	9 (64.3%)	5 (45.5%)	0.697
Dyspnea	7 (28.0%)	4 (28.6%)	3 (27.3%)	1.000
Diarrhea	5 (20.0%)	2 (14.3%)	3 (27.3%)	0.623

Headache	2 (8.0%)	0 (0.0%)	2 (13.3%)	0.183
Heart rate, mean $\pm$ SD, bpm	88.2 $\pm$ 15.1	86.8 $\pm$ 11.8	89.9 $\pm$ 18.9	0.617
Respiratory rate, median (IQR)	20.0 (19.0, 22.0)	20.0 (19.0, 22.0)	20.0 (19.0, 22.0)	0.845
Mean systolic blood pressure, mean $\pm$ SD, mm Hg	130.0 $\pm$ 24.4	126.8 $\pm$ 22.6	134.2 $\pm$ 27.1	0.464
Days from illness onset to hospital, median (IQR), d	6 (3,11)	8 (3,11)	6 (3,11)	1.000
Hospital stays, median (IQR), d	17.0 (11.5, 21.5)	12.0 (10.0,15.0)	21.0 (18.0,23.0)	0.001
Treatment				
Oxygen support	15 (60%)	6 (42.9%)	9 (81.8%)	0.099
Antiviral therapy	24(96.0%)	13(92.9%)	11(100.0%)	1.000
Antibiotic therapy	25(100.0%)	14(100.0%)	11(100.0%)	-
Use of corticosteroid	14 (56.0%)	6(42.9%)	8(72.7%)	0.227
Clinical outcome^b				
Remained in hospital	9(36.0%)	2(14.3%)	7(63.6%)	0.017
Discharged	16(64.0%)	12(85.7%)	4(36.4%)	

478

479 ^a After 7-14 days, viral clearance or not was determined by viral RNA detections.

480 ^b After 18-25 days, patients were discharged according to the guideline: 1) a series negative result of
481 viral RNA detection; 2) no clinical symptoms; 3) improved chest CT imaging.

482

483

484

485

486

Table 2. Laboratory results of 25 patients with the outcome of viral clearance.

	Normal Range	Total	Viral clearance(n=14)	without viral clearance (n=11)	P value
White blood cell Count ($\times 10^9$ /L)	3.5-9.5	4.5 $\pm$ 1.9	4.5 $\pm$ 1.7	4.5 $\pm$ 2.2	0.919
Lymphocyte count ($\times 10^9$ /L)	1.1-3.2	0.9 (0.6, 1.3)	0.8 (0.6, 1.4)	0.9 (0.6, 1.2)	0.565
Monocyte count ($\times 10^9$ /L)	0.1-0.6	0.37 $\pm$ 0.15	0.33 $\pm$ 0.15	0.42 $\pm$ 0.14	0.117
Neutrophil count ($\times 10^9$ /L)	1.8-6.3	2.8 (1.5, 4.5)	3.1 (1.6, 4.8)	2.6 (1.3, 4.3)	0.805
Platelet count ($\times 10^9$ /L)	125-350	179.5 $\pm$ 68.8	191.5 $\pm$ 70.4	164.3 $\pm$ 66.7	0.336
Alanine aminotransferase (U/L)	9-50	24 (20.5, 38.5)	21 (19, 50.5)	25 (21.0, 35.0)	0.583
Aspartate aminotransferase (U/L)	15-40	26 (21.0, 33.0)	23.0(17.0, 34.8)	28.0(26.0,33.0)	0.066
Total bilirubin (mmol/L)	5-21	9.6 (8.0, 12.7)	9.6 (7.6, 12.2)	10.7 (8.2, 16.7)	0.511
Lactate dehydrogenase (U/L)	125-243	197(164.5, 286.5)	197.0(155.0, 340.0)	201.0(174.0, 279.0)	0.956
Prothrombin time (s)	9.4-12.5	12.3 (11.7, 13.0)	12.6 (11.7,13.5)	12.3 (12.1,12.4)	0.641
Activated partial thromboplastin time (s)	25.1-36.5	29.9 $\pm$ 3.0	29.8 $\pm$ 2.6	30.0 $\pm$ 3.6	0.860
D-dimer, (mg/L)	0-500	187.0 (125.0,309.5)	181.0 (104.5,271.0)	233.0 (145.0,630.0)	0.179
Creatinine (μ mol/L)	64-104	65.5 $\pm$ 14.6	64.3 $\pm$ 14.3	67.0 $\pm$ 15.6	0.659
Procalcitonin, ng/mL $\geq$ 0.05, No. (%)	<0.05	5 (20.0%)	3 (21.4%)	2 (18.2%)	1.000
Interleukin-6 (pg/ml)	0-7	16.1 (3.7, 31.4)	13.8 (5.3, 30.2)	17.6 (2.8, 49.9)	0.805
CD3+ T cell count ($\times 10^9$ /L)	805-4459	561.0 (338.0, 770.0)	639.0 (400.3, 1038.0)	485.0 (384.0, 732.0)	0.311
CD4+ T cell count ($\times 10^9$ /L)	345-2350	297.0 (171.5, 447.5)	327.0 (182.8, 560.5)	220.0 (148.0, 420.0)	0.262
CD8+ T cell count ($\times 10^9$ /L)	345-2350	272.9$\pm$129.7	291.4 $\pm$ 149.3	249.5 $\pm$ 101.6	0.435
B cell count ($\times 10^9$ /L)	240-1317	111.5$\pm$75.8	133.2 $\pm$ 84.8	83.8 $\pm$ 54.2	0.107
NK cell count ($\times 10^9$ /L)	210-1514	170.0 (67.5, 245.0)	193.5 (75.0, 286.0)	170.0 (43.0, 215.0)	0.511

491

492 **Figure legends**

493 **Figure 1.** Comparison of peripheral changes in CD3+, CD4+, CD8+ T,
 494 B, and NK cell counts in patients who were negative for the SARS-CoV-
 495 2 (A-E) and positive (F-J) after 8-14 days of treatment. The medium
 496 number of peripheral lymphocyte counts are represented in the solid line.
 497 The threshold of normal peripheral lymphocyte counts is represented in
 498 the dash line. NS means no significant difference tested in statistical
 499 analysis: Mann-Whitney U test was used for A, B, E, F since these data
 500 are non-normal distributed and t-test was used for the other normal
 501 distributed data).

502

503 **Figure 2.** Comparison of peripheral changes in CD3+, CD4+, CD8+ T,
 504 B, and NK cell counts in six patients who were negative for the SARS-
 505 CoV-2 (A-E) and four patients who were positive (F-J) after 15-23 days
 506 of treatment. The medium number of peripheral lymphocyte counts are
 507 represented in the solid line. The threshold of normal peripheral
 508 lymphocyte counts is represented in the dash line. NS means no
 509 significant difference tested in statistical analysis and all data used Mann-
 510 Whitney U test.