

Extended SIR prediction of the epidemics trend of COVID-19 in Italy and compared with Hunan, China

1 Jia Wangping ^{1,2*}, Han Ke ^{1*}, Song Yang ¹, Cao Wenzhe ¹, Wang Shengshu ¹, Yang Shanshan ¹,
2 Wang Jianwei ¹, Kou Fuyin ¹, Tai Penggang ¹, Li Jing ¹, Liu Miao ¹, He Yao ¹

3 ¹Institute of Geriatrics, Beijing Key Laboratory of Aging and Geriatrics, National clinical research
4 center for geriatrics diseases, Second Medical Center of Chinese PLA General Hospital, 28 Fuxing
5 Road, Beijing, 100853, China;

6 ²Department of Military Medical Technology Support, School of Non-commissioned Officer, Army
7 Medical University, Shijiazhuang, Hebei, China.

8 * Correspondence:

9 Prof. He Yao, MD, PhD

10 E-mail: yhe301@x263.net

11 Liu Miao, MD, PhD

12 Email: liumiaolmbxb@163.com

13 **Keywords: COVID-19, coronavirus, Italy, prediction, epidemics trend.**

14 Abstract

15 **Background:** Coronavirus Disease 2019 (COVID-19) is currently a global public health threat.
16 Outside of China, Italy is one of the most suffering countries with the COVID-19 epidemic. It is
17 important to predict the epidemics trend of COVID-19 epidemic in Italy to help develop public
18 health strategies.

19 **Methods:** We used time-series data of COVID-19 from Jan 22,2020 to Mar 16,2020. An infectious
20 disease dynamic extended susceptible-infected-removed (eSIR) model, which covers the effects of
21 different intervention measures in dissimilar periods, was applied to estimate the epidemic trend in
22 Italy. The basic reproductive number was estimated using Markov Chain Monte Carlo methods and
23 presented using the resulting posterior mean and 95% credible interval (CI). Hunan, with similar total
24 number of populations in Italy, was used as a comparative item.

25 **Results:** In the eSIR model, we estimated that the basic reproductive number for COVID-19 was
26 respectively 4.10 (95% CI: 2.15–6.77) in Italy and 3.15(95% CI: 1.71–5.21) in Hunan. There would
27 be totally 30 086 infected cases (95%CI:7920-81 869) under the current country blockade and the
28 endpoint would be Apr 25 (95%CI: Mar 30 to Aug 07) in Italy. If the country blockade is imposed 5
29 day later, the total number of infected cases would expand the infection scale 1.50 times.

30 **Conclusion:** Italy's current strict measures can efficaciously prevent the further spread of COVID-19
31 and should be maintained. Necessary strict public health measures be implemented as soon as

32 possible in other European countries with a high number of COVID-19 cases. The most effective
33 strategy needs to be confirmed in further studies.

34 **1 Introduction**

35 The Corona Virus Disease 2019 (COVID-19) started in Wuhan, China in December and quickly
36 spread to China and many countries and regions in the world(1-3). The COVID-19 outbreak made
37 assessment as a pandemic by the World Health Organization (WHO) on March 11. It is currently a
38 global public health threat and more than 100 countries including Italy, Iran, the United States, South
39 Korea, and Japan are suffering from COVID-19. Outside of China, Italy is one of the most suffering
40 countries with the COVID-19 epidemic. As of Mar 16, the cumulative number of confirmed cases in
41 Italy reached 27980, ranking second in the world, the total confirmed deaths, 2158, and the fatality
42 rate, as high as 7.71%, which has become one of the highest among the major epidemic countries.
43 However, few studies have assessed the epidemic status in Italy(4, 5).

44 Global public health measures are required to cope with the rapid spread of the epidemic. China has
45 taken precise and differentiated strategies, including self-quarantine of residents in Wuhan and other
46 areas, community-based prevention and control. These measures have played an important role in
47 preventing and controlling the epidemic. Previous studies have shown that due to the isolation of
48 Wuhan, the overall epidemiological progress in mainland China has been delayed by three to five
49 days and the number of internationally transmitted cases has been reduced by nearly 80%(6). Italy
50 detected the first two cases of imported COVID-19 on Jan 31. After that, Italy was the first country to
51 declare a state of emergency. Since then, various measures have been implemented to control the
52 spread of COVID-19. It is vital to evaluate the role of Italian quarantine measures for decision-
53 making.

54 Mathematical modeling is helpful to predict the possibility and severity of disease outbreak and
55 provide key information for determining the type and intensity of disease intervention. The SIR
56 model and its modifications such as SEIR model have been widely applied to the current outbreak of
57 COVID-19. Tang et al. estimated the infectivity of COVID-19 based on a classical susceptible-
58 exposed-infected-removed (SEIR) epidemiological model(7). Wu et al proposed an extended SEIR
59 model to forecast the spread of 2019-nCoV both within and outside of mainland China(3). However,
60 these studies assumed that the exposed population were not infectious, which may be not suitable in
61 COVID-19. Yang Z et al. predicted that China's epidemic will peak in late February and end in late
62 April by a combination of SEIR model and a machine-learning artificial intelligence (AI)
63 approach(8). However, this study and the above studies did not consider the phase-adjusted
64 preventive measures and time-varying parameters, which may affect the accuracy of predictions.

65 We adopted extended susceptible-infected-removed (eSIR) model(9), which covers the effects of
66 different epidemic prevention measures in different periods and helps to achieve the following
67 specific objectives:

68 AIM 1: Compare the epidemic development of COVID-19 in Italy with provinces with similar total
69 number of populations in China.

70 AIM 2: Predict the epidemiological trend of COVID-19 in Italy via a modified and calibrated model.

71 AIM 3: Assess the effectiveness of Italian existing and assumed precautionary quarantine measures.

72 **2 Methods**

73 2.1 Data sources

74 In this study, we used the publicly available dataset of COVID-19 provided by the Johns Hopkins
75 University(10). This dataset includes many countries' daily count of confirmed cases, recovered
76 cases and deaths. As time-series data, it is available from 22 January 2020. Besides, we also gathered
77 and cross-checked data in DXY.cn(11), a website providing real-time data of the COVID-19.

78 These data are collected through public health authorities' announcements and are directly reported
79 public and unidentified patient data, so ethical approval is not required.

80 2.2 Prediction models

81 The reproduction number, R_0 , reflects the transmissibility of a virus spreading under no control,
82 representing the average number of new infections generated by each infected person(12). The
83 COVID-19 is likely to decline and eventually disappear if $R_0 \leq 1$. To estimate trends and calculate the
84 R_0 , we used an extended SIR model (eSIR model) with a time-varying transmission rate(9). In brief,
85 this model is one extension to the basic epidemiological model—susceptible-infected-removed
86 (SIR) model. S, I and R represent the number of susceptible, infected, and recovered individuals
87 accordingly, and $N = S + I + R$ is the total population. The transmission rate is constant in the SIR
88 model. It should be noted that in actual situations, the speed of transmission can be changed through
89 many interventions, such as personal protective measures, community-level isolation and city
90 blockade. As is shown below, the eSIR model adds transmission modifier $\pi(t)$ to the SIR model, so it
91 allows a time-varying probability of the transmission rate.

92 An extended SIR model with time-varying transmission rates.



97

$$98 \quad \frac{dS}{dt} = -\pi(t) \frac{\beta SI}{N}$$

99

$$\frac{dS}{dt} = \pi(t) \frac{\beta SI}{N} - \gamma I$$

100

$$\frac{dR}{dt} = \gamma I$$

101

$$R_0 = \frac{\beta}{r}$$

102 The transmission rate modifier $\pi(t)$ can be specified according to actual interventions in different
103 times and regions. According to Chinese government isolation measures and previous study, we set
104 $\pi(t)=0.9$ if $t \in$ (Jan 23, Feb 04], city blockade; $\pi(t)=0.5$ if $t \in$ (Feb 4, Feb 8], enhanced quarantine;

105 $\pi(t)=0.1$ if $t > \text{Feb } 8$, more enhanced quarantine in Hunan. In the opinion of Italy government
106 isolation measures, we set $\pi(t)=0.9$ if $t < \text{Mar } 4$, no concrete quarantine protocols; $\pi(t)=0.5$ if $t \in$ (
107 $\text{Mar } 04, \text{Mar } 09]$, some cities blockade and enhanced quarantine; $\pi(t)=0.1$ if $t > \text{Mar } 09$, country
108 blockade and more enhanced quarantine in Italy. We also assumed that the Italy government took
109 country blockade and intensified quarantine on $\text{Mar } 05$ or $\text{Mar } 15$.

110 We did prediction with an R software package—eSIR which can output the Markov Chain Monte
111 Carlo (MCMC) estimation, inference, and prediction under the extended SIR models with time-
112 varying transmission modifier $\pi(t)$. The model can also yield the turning points of the
113 epidemiological trend of COVID-19. The first turning point was defined as the mean predicted time
114 when the daily proportion of infected cases becomes smaller than the previous ones. The second
115 turning point was defined as the mean the predicted time when the daily proportion of removed cases
116 (i.e. both recovered and dead) becomes larger than that of infected cases. Besides, an end point was
117 defined as the time when the median proportion of current infected cases turn to zero.

118 We did all analyses in R (version 3.6.2).

119 **3 Results**

120 **3.1 Epidemic development of COVID-19 in Italy compared with Hunan**

121 Figure 1 demonstrates daily new COVID-19 cases and epidemic distribution of COVID-19 in Hunan,
122 China and Italy. The number of new cases and confirmed cases show an exponential trend since Feb
123 21 in Italy from while the number of new cases turns to zero from Feb 29 in Hunan.

124 **3.2 Prediction of the epidemics trend of COVID-19 in Italy compared with Hunan**

125 Table 1 summarizes the posterior mean and credible intervals of R_0 , incidence, and endpoint in
126 Hunan and Italy according to the eSIR model. R_0 for COVID-19 was respectively 3.15(95% CI:
127 1.71–5.21) in Hunan and 4.10 (95% CI: 2.15–6.77) in Italy. There would be totally 3 988 infected
128 cases (95% CI:786-9 065) in Hunan. There would be totally 30 086 infected cases (95% CI:7 920-81
129 869) under the current country blockade in Italy. Figure 2 indicates an epidemiological trend of
130 COVID-19 under existing preventions in Hunan, China and Italy. The first and second turning point
131 in Hunan appeared on Feb 04 and Feb 09. The first and second turning point in Italy is Mar 10 and
132 Mar 11. The predictions suggest that the endpoints of the COVID-19 epidemics in Hunan and Italy
133 will come on Feb 23 (95% CI: Feb 19 to Mar 23) and Apr 25 (95% CI: Mar 30 to Aug 07), separately.

134 **3.3 Prediction of the epidemics trend of COVID-19 in Italy assumed precautionary 135 quarantine measures.**

136 Table 2 summarizes the posterior mean and credible intervals of R_0 , incidence, and endpoint
137 according to different beginning of country blockade in Italy. The mean number of COVID-19
138 infected cases in Italy would be 10636(95% CI: 2357-23326) and the endpoint would arrive on Mar
139 16 (95% CI: Mar 09 to Apr 22) if the beginning time of country blockade were Mar 05. Furthermore,
140 that would be 44993(95% CI: 16676-96759) and Apr 27 (95% CI: Apr 01 to Jul 27) if the beginning
141 of country blockade were Mar 15.

142

143 **4 Discussion**

144 This impact of the COVID-19 response (overall quarantine regulations , social distancing, and
145 isolation of infections) in China is encouraging for the many other countries(13).We compared the
146 situation in Hunan, China, which has the similar population as Italy to predict. The spread of COVID-
147 19 in Hunan Province appeared relatively early and has now entered a phase of no inflections, which
148 helps to observe the entire course of the epidemic. Besides, due to the similarity of population size and
149 geographical location adjacent to Hubei, Hunan's public health measures can provide useful guidance
150 for Italy in preventing the further spread of COVID-19.

151 In our study, the eSIR model with R software package was used to evaluate the impact of intervention
152 measures on Italian COVID-19 epidemic. In previous studies, estimation of the epidemic of an
153 infectious disease is often performed using constant parameter(14-17). The advantage of eSIR model is
154 that it combines time-varying isolation measures and expands SIR model to adapt to the time-varying
155 transmission rate in the population. Lili Wang et al. found that COVID-19 outside Hubei in China has
156 been so far much less severe.(9) But they did not perform each province's analyses. The first and
157 second points in our study are respectively Feb 04 and Feb 09, which are the same as these outside
158 Hubei in China. Furthermore, the actual number of infected cases (1018) is included in the predicted
159 number of infected cases (786-9065) and the endpoint(Feb 29) is included in the predicted
160 endpoint(Feb 19 to Mar 23) in our study, which also reflects stability and accuracy of the eSIR model.
161 Combining the above data and methods, these findings show that eSIR model is more suitable for
162 predicting the epidemic trend of COVID-19.

163 Our results showed that R_0 was estimated to be respectively 3.12 (95% CI, 1.83-3.01) and 3.74 (95%
164 CI, 2.09-6.02) in SIR model and eSIR model in Italy; 2.97 (95% CI, 1.80-5.01) and 3.15 (95% CI, 1.71-
165 5.21) in SIR model and eSIR model in Hunan. Li Qun et al estimated R_0 to be 2.2 (95% CI, 2.09-6.02)
166 among the first 425 patients in Wuhan, China(18). Other studies estimated R_0 to be 1.4-2.5 (19),
167 2.68(95% CI 2.47~2.68) (3), 3.6-3.8(20), 6.47(95% CI 5.71-7.23) (7) . Ying Liu et al found that the
168 estimated mean R_0 for COVID-19 is around 3.28, with a median of 2.79 and IQR of 1.16 by reviewing
169 R_0 of COVID-19 in 12 studies. (21) We estimated R_0 to be 3.15(95% CI 1.71-5.21) in Hunan, which
170 is in agreement with these findings. But R_0 in Italy was estimated to be 4.10 (95% CI: 2.15–6.77),
171 which is larger than that in Hunan. This needs to be confirmed by further studies.

172 This study showed that COVID-19 spread rapidly throughout Italy after Feb 21. Possible reasons for
173 such rapid growth of infections include:(1) more timely caution and preventative measures were not
174 taken, (2) The number of infections during Jan 31-Feb 20 could be under-reported due to
175 underdiagnosis, given subclinical or asymptomatic cases. The incubation period for COVID-19 is
176 thought to be within 14 days following exposure, with most cases occurring approximately four to five
177 days after exposure(18, 22, 23). So it seems impossible to maintain totally two or three cases during Jan
178 31-Feb 20 in Italy. In addition, the rapid increase in the number of infections after Feb 21 might reflect
179 a belated realization of the spread of COVID-19.

180 Previous studies have shown that more rigorous government control policies were associated with a
181 slower increase in the infected population(6, 16, 24-27) . Our report also showed that rigorous
182 measures can effectively prevent the further spread of COVID-19 in Hunan. Based on our model,
183 Italy should still maintain all levels of quarantines like China by Apr 25(Mar 30-Aug 07).
184 Furthermore, Tianyi Qiu et al found that delaying the lockdown by 1-6 days in Wuhan would expand
185 the infection scale 1.23-4.94 times and the epidemic would be finally out of control if lockdown
186 imposed 7 days later(17). Our study also shows that taking government control earlier can decrease
187 the number of infected cases. In addition, from China's experience, various control measures,
188 including the early detection and isolation of individuals with symptoms, traffic restrictions, medical

189 tracking, and entry or exit screening, can well prevent the further spread of COVID-19. These
190 measures are in line with the latest recommendations by the World Health Organization and a
191 previous study in Spain(28). But the most effective strategy still needs to be confirmed by further
192 studies. Consequently, it is better and necessary to apply strict public health measures in other
193 European countries with a high number of COVID-19 cases.

194 Our study has some limitations. Firstly, it is based on the assumption that rigorous measures like
195 China have been taken in Italy, although this study uses the new model to obtain dynamic results,
196 which is instructive for the prevention and control of the epidemic in Italy. Secondly, the suspected
197 cases and the daily number of hospitalized cases are not available, so they are not considered in the
198 eSIR model. Thirdly, some unforeseeable factors may affect these estimated data in our study such as
199 super-spreaders exist.

200 In conclusion, the current study is the first to provide a prediction for epidemic trend after strict
201 prevention and control measures were implemented in Italy. Our study suggests that rigorous
202 measures like China should still be maintained in Italy by Apr 25(Mar 30-Aug 07) to prevent further
203 spread of COVID-19.

204

205 **5 Conflict of Interest**

206 We declare no competing interests.

207 **6 Author Contributions**

208 JW, HK, LM and HY contributed to the study design. JW, HK, and LM contributed to the writing of
209 the manuscript. JW, HK, SY, CW contributed to the data analysis. SY, CW and WJ contributed to
210 the data compilation. WJ, YS, WS and HY contributed to critical review. TP, KF and LJ contributed
211 to the literature search. YS, WJ, and KF contributed to the design of tables and figures.

212 **7 Funding**

213 The study was funded by Army Logistics Emergency Scientific Research Project; Emergency
214 scientific research of the army and the emergency scientific research of Chinese PLA General
215 Hospital (20EP008).

216 **8 Acknowledgments**

217 We greatly appreciated the technical assistance provided by Zhenxing Cheng; Institute of Blue and
218 Green Development, Shandong University, Weihai, 264209, PR China.

219 **9 Reference**

- 220 1. Benvenuto D, Giovanetti M, Salemi M, Prosperi M, De Flora C, Junior Alcantara LC, et al. The
221 global spread of 2019-nCoV: a molecular evolutionary analysis. *Pathog Glob Health* (2020):1-4. doi:
222 10.1080/20477724.2020.1725339. PubMed PMID: 32048560.
- 223 2. Liao X, Wang B, Kang Y. Novel coronavirus infection during the 2019-2020 epidemic:
224 preparing intensive care units-the experience in Sichuan Province, China. *Intensive Care Med* (2020).
225 doi: 10.1007/s00134-020-05954-2. PubMed PMID: 32025779.

- 226 3. Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and
227 international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study.
228 *Lancet* (2020). doi: 10.1016/S0140-6736(20)30260-9. PubMed PMID: 32014114.
- 229 4. Giovanetti M, Benvenuto D, Angeletti S, Ciccozzi M. The first two cases of 2019-nCoV in Italy:
230 Where they come from? *J Med Virol* (2020):10.1002/jmv.25699. doi: 10.1002/jmv.25699. PubMed
231 PMID: 32022275.
- 232 5. Porcheddu R, Serra C, Kelvin D, Kelvin N, Rubino S. Similarity in Case Fatality Rates (CFR)
233 of COVID-19/SARS-COV-2 in Italy and China. *J Infect Dev Ctries* (2020) 14(2):125-8. doi:
234 10.3855/jidc.12600. PubMed PMID: 32146445.
- 235 6. Chinazzi M, Davis JT, Ajelli M, Gioannini C, Litvinova M, Merler S, et al. The effect of travel
236 restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. *Science (New York, NY)*
237 (2020).
- 238 7. Tang B, Wang X, Li Q, Bragazzi NL, Tang S, Xiao Y, et al. Estimation of the Transmission Risk
239 of the 2019-nCoV and Its Implication for Public Health Interventions. *J Clin Med* (2020) 9(2). doi:
240 10.3390/jcm9020462. PubMed PMID: 32046137.
- 241 8. Yang Z, Zeng Z, Wang K, Wong S-S, Liang W, Zanin M, et al. Modified SEIR and AI prediction
242 of the epidemics trend of COVID-19 in China under public health interventions. *2020* (2020).
- 243 9. Song PX, Wang L, Zhou Y, He J, Zhu B, Wang F, et al. An epidemiological forecast model and
244 software assessing interventions on COVID-19 epidemic in China. *medRxiv*
245 (2020):2020.02.29.20029421. doi: 10.1101/2020.02.29.20029421.
- 246 10. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time.
247 *The Lancet Infectious diseases* (2020). Epub 2020/02/23. doi: 10.1016/s1473-3099(20)30120-1.
248 PubMed PMID: 32087114.
- 249 11. Sun K, Chen J, Viboud C. Early epidemiological analysis of the coronavirus disease 2019
250 outbreak based on crowdsourced data: a population-level observational study. *The Lancet Digital*
251 *Health* (2020). doi: 10.1016/s2589-7500(20)30026-1.
- 252 12. Imai, N. et al. Report 3: Transmissibility of 2019-nCoV. [https://www.imperial.ac.uk/mrc-global-](https://www.imperial.ac.uk/mrc-global-infectious-disease-analysis/news-wuhan-coronavirus/)
253 [infectious-disease-analysis/news-](https://www.imperial.ac.uk/mrc-global-infectious-disease-analysis/news-wuhan-coronavirus/)
254 [wuhan-coronavirus/](https://www.imperial.ac.uk/mrc-global-infectious-disease-analysis/news-wuhan-coronavirus/) (2020).
- 255 13. Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will country-based
256 mitigation measures influence the course of the COVID-19 epidemic? *Lancet* (2020):S0140-
257 6736(20)30567-5. doi: 10.1016/S0140-6736(20)30567-5. PubMed PMID: 32164834.
- 258 14. Zhao S, Lin Q, Ran J, Musa SS, Yang G, Wang W, et al. Preliminary estimation of the basic
259 reproduction number of novel coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven
260 analysis in the early phase of the outbreak. *International Journal of Infectious Diseases* (2020) 92:214-
261 7. doi: <https://doi.org/10.1016/j.ijid.2020.01.050>.
- 262 15. Roosa K, Lee Y, Luo R, Kirpich A, Rothenberg R, Hyman JM, et al. Real-time forecasts of the
263 COVID-19 epidemic in China from February 5th to February 24th, 2020. *Infectious Disease Modelling*
264 (2020) 5:256-63. Epub 2020/02/29. doi: 10.1016/j.idm.2020.02.002. PubMed PMID: 32110742;
265 PubMed Central PMCID: PMC7033348.
- 266 16. Fang Y, Nie Y, Penny M. Transmission dynamics of the COVID-19 outbreak and effectiveness
267 of government interventions: A data-driven analysis. *J Med Virol* (2020). Epub 2020/03/07. doi:
268 10.1002/jmv.25750. PubMed PMID: 32141624.
- 269 17. Wan H, Cui J-a, Yang G-J. Risk estimation and prediction by modeling the transmission of the
270 novel coronavirus (COVID-19) in mainland China excluding Hubei province. *medRxiv*
271 (2020):2020.03.01.20029629. doi: 10.1101/2020.03.01.20029629.
- 272 18. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in Wuhan,
273 China, of Novel Coronavirus-Infected Pneumonia. *The New England journal of medicine* (2020). doi:
274 10.1056/NEJMoa2001316. PubMed PMID: MEDLINE:31995857.
- 275 19. Mahase E. China coronavirus: what do we know so far? *BMJ (Clinical research ed)* (2020)

- 275 368:m308. Epub 2020/01/26. doi: 10.1136/bmj.m308. PubMed PMID: 31980434.
- 276 20. Read, J. M., Bridgen, J. R. E., Cummings, D. A. T., Ho, A. & Jewell, C. P. Novel coronavirus
277 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions. medrxiv.
278 <https://www.medrxiv.org/content/10.1101/2020.01.23.20018549v1.full.pdf> (2020).
- 279 21. Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J. The reproductive number of COVID-19 is higher
280 compared to SARS coronavirus. *J Travel Med* (2020). doi: 10.1093/jtm/taaa021. PubMed PMID:
281 32052846.
- 282 22. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical Characteristics of Coronavirus
283 Disease 2019 in China. *The New England journal of medicine* (2020). Epub 2020/02/29. doi:
284 10.1056/NEJMoa2002032. PubMed PMID: 32109013.
- 285 23. Chan JF, Yuan S, Kok KH, To KK, Chu H, Yang J, et al. A familial cluster of pneumonia
286 associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a
287 family cluster. *Lancet* (2020) 395(10223):514-23. Epub 2020/01/28. doi: 10.1016/s0140-
288 6736(20)30154-9. PubMed PMID: 31986261.
- 289 24. Wang H, Wang Z, Dong Y, Chang R, Xu C, Yu X, et al. Phase-adjusted estimation of the number
290 of Coronavirus Disease 2019 cases in Wuhan, China. *Cell Discovery* (2020) 6(1):10. doi:
291 10.1038/s41421-020-0148-0.
- 292 25. Kraemer MUG, Yang C-H, Gutierrez B, Wu C-H, Klein B, Pigott DM, et al. The effect of
293 human mobility and control measures on the COVID-19 epidemic in China. *medRxiv*
294 (2020):2020.03.02.20026708. doi: 10.1101/2020.03.02.20026708.
- 295 26. Nishiura H, Kobayashi T, Miyama T, Suzuki A, Jung S, Hayashi K, et al. Estimation of the
296 asymptomatic ratio of novel coronavirus infections (COVID-19). *medRxiv*
297 (2020):2020.02.03.20020248. doi: 10.1101/2020.02.03.20020248.
- 298 27. Tian H, Liu Y, Li Y, Wu C-H, Chen B, Kraemer MUG, et al. The impact of transmission control
299 measures during the first 50 days of the COVID-19 epidemic in China. *medRxiv*
300 (2020):2020.01.30.20019844. doi: 10.1101/2020.01.30.20019844.
- 301 28. Aleta A, Moreno Y. Evaluation of the potential incidence of COVID-19 and effectiveness of
302 contention measures in Spain: a data-driven approach. *medRxiv* (2020):2020.03.01.20029801. doi:
303 10.1101/2020.03.01.20029801.
- 304

305 **Tables**

306

307 **Table 1. R0, incidence and endpoint in Hunan and Italy according to the eSIR model.**

308

Model	R0		Incidence (%)		Endpoint	
	Mean	95%CI	Mean	95%CI	Mean	95%CI
Hunan	3.15	1.71-5.21	0.0066	0.0013-0.0150	2020/2/23	2020/2/19-2020/3/23
Italy	4.10	2.15-6.77	0.0498	0.0131-0.1355	2020/4/25	2020/3/30-2020/8/07

309

Inf: The endpoint appears more than 200 days after t_0 .

310

311

312

313 **Table 2. R0, incidence and endpoint in Italy according to different beginning of country blockade.**

314

Begin time of country blockade	R0		Incidence (%)		Endpoint	
	Mean	95%CI	Mean	95%CI	Mean	95%CI
Mar 05	3.26	1.83-5.34	0.0176	0.0039-0.0386	2020/3/16	2020/3/9-2020/4/22
Mar 15	4.24	2.44-6.61	0.0745	0.0028-0.1601	2020/4/27	2020/4/1-2020/7/26

315

316

317

318 **Figures**

319

320 **Figure 1. Epidemic development of COVID-19 in Hunan, China and Italy.**

321

322 A, B: Daily new COVID-19 cases in Hunan, China and Italy.

323 C, D: Epidemic distribution of COVID-19 in Hunan, China and Italy.

324

325 **Figure 2. Epidemiological trend of COVID-19 under existing preventions in Hunan, China and**
326 **Italy according to eSIR model.**





327

328 A: Hunan, China; B: Italy.

329 The black dots left to the blue vertical line denote the observed proportions of the infected and removed
330 compartments on the last date of available observations or before. The blue vertical marks is the current
331 time up to which we have observed data (t_0). The green and purple vertical lines denote the first and
332 second turning points, respectively. The cyan and salmon color area denotes the 95% credible interval
333 of the predicted proportions of current infected cases before and after t_0 , respectively. The gray and red
334 curves are the posterior mean and median curves.

335 Epidemiological trend of COVID-19 under existing preventions of Hunan, China and Italy in eSIR
336 model

337 **Figure 1. Epidemic development of COVID-19 in Hunan, China and Italy.**




338

339

340

341

342 **Figure 2. Epidemiological trend of COVID-19 under existing preventions in Hunan, China and Italy according to eSIR model.**



343