Partial unlock model for COVID-19 or similar pandemic averts medical and economic disaster

Robert L. Shuler^{12*}

¹ShulerResearch.org

*Corresponding author

E-mail: robert@shulerresearch.org (RS)

March 30, 2020

1 **Abstract**

2 Data as of March 29, 2020 show that the “flattening” strategy for COVID-19 in the U.S. is
3 working so well that a clean removal of social distancing (aka “unlock”) at any time in 2020 will
4 produce a renewed catastrophe, overloading the healthcare system. Leaving the economy locked down
5 for a long time is its own catastrophe. An SIR-type model with clear parameters suitable for public
6 information, and both tracking and predictive capabilities which “learns” disease spread characteristics
7 rapidly as policy changes, suggests that a solution to the problem is a partial unlock. Case load can be
8 managed so as not to exceed critical resources such as ventilators, yet allow enough people to get sick
9 that herd immunity develops and a full unlock can be achieved in as little as five weeks from beginning
10 of implementation. The partial unlock could be for example 3 full working days per week. Given that
11 not all areas or individuals will respond, and travel and public gatherings are still unlikely, the partial
12 unlock might be 5 full working days per week. The model can be regionalized easily, and by
13 expediting the resolution of the pandemic in the U.S. medical equipment and volunteers, many of them
14 with already acquired immunity, can be made available to other countries.

15 **Introduction**

16 After failure to contain COVID-19 in China, the global response has largely been social isolation,
17 with effective cessation of global and local travel except by personal vehicle, and in some cities by bus
18 and subway with frequent sterilization protocols. The strategy is called “curve flattening” and is

19 intended to contain demand for special medical equipment, especially ventilators and related devices,
20 from exceeding supply, which would cause unnecessary deaths.

21 A vaccine reputed to be 18 months away, the remaining strategy is herd immunity. For COVID-
22 19 with an R_0 (R-naught, number of people to whom each victim transmits the virus) of 1.4 to 3.28 [1],
23 and seemingly everyone susceptible with no natural immunity, up to 70% of the population must have
24 the disease for herd immunity to come into play. With half a million known and likely 2.5 million
25 unknown cases worldwide, a high spread rate, many very mild cases, and a contagious incubation
26 period, containment is no longer a viable option. The disease is substantially different than MERS or
27 SARS. Although, many in the public and some officials seem to be behaving as if containment were
28 viable.

29 In fighting a disease with public policy that relies on compliance, it is important to avoid the
30 following sort of dilemma. Suppose we posit two policy choices, lockdown and unlock. Suppose the
31 implementation of a policy has two parts: a planned scientific and administrative decision, and the
32 public compliance with that decision. Suppose either one of those parts amounts to “If the disease is
33 spreading and there is horror in the hospitals, lockdown, otherwise unlock.” Then lockdown reduces
34 spread and produces a sense of relief, which results in unlock, which in turn restores the pandemic. A
35 Gödel undecidability ensues and the actual strategy flip flops ineffectively. In this paper we propose a
36 model which rapidly measures and uses public response to policy, rather than depending on how well a
37 particular policy is implemented or accepted, dodging the dilemma. This leaves individuals free to
38 self-isolate or risk working without overly compelling them. It leaves regional governments free to
39 modify the policy to fit their area, if for example the disease is spreading faster and threatening hospital

resources. To predict the effect of an “unlock” the model (or the person using it) can use a response calibration from a previous time, or estimate a new one. For this paper, we use the initial spread rate to model “unlock.” This may be high, as people will likely not go back to mass gatherings and flying on airplanes immediately, which simply means we model a worst case scenario which gives the model a margin of safety.

A principle objective of this paper is to share results and strategy possibilities before the conclusion of the COVID-19 outbreak. If studying frequently repeating phenomena, the scientific approach is to present complete data. However, we are studying a rare (roughly once in 100 years) phenomena, with the aim to intervene before it is concluded.

Approach

The goal of our approach then discards containment as an opportunity past (and perhaps not realistic from early on), and a vaccine as a prospect too far in the future to avoid economic catastrophe. Opinions differ as to the effect of severe and prolonged recession on mortality and health. For example, there are fewer motorway deaths due to less driving [2]. On the other hand the 2008 financial crisis resulted over the next few years at least 260,000 additional cancer deaths [3]. Economic losses from pandemics, even without a long term global shutdown, have been estimated at the low end of but within the range of impacts from climate change [4].

Therefore the goal we adopt is to use public policy supported by modeling to achieve herd immunity as rapidly as possible, without overloading the healthcare system’s critical resources (we use ventilators as a general proxy for resources) and causing unnecessary deaths. Given our assumptions

about vaccines, containment and R_0 , 70% of the population will get sick anyway. Our approach of voluntary compliance with unlocks allows individuals and regional governments to manage their health and economic risks as they see fit. As the U.S. has ventilators per capita among the highest in the world, and stands to suffer high economic damage, it is a logical place to attempt this strategy. If the U.S. develops herd immunity, then it might choose to make some of its ventilators, and its production capacity for more ventilators, available to other countries. At the current time the hoarding mentality which arises from uncertainty is impairing the sharing of medical equipment and New York and Italy plead for ventilators and masks, as other countries protest they need their own stockpiles. But China, feeling it has the spread under control, is more forthcoming.

Model parameters

The purpose of this paper is more to illustrate an approach than to propose a specific model. We suggest following Huppert and Katriel's guidance: *"To examine which of the predictions made by a model are trustworthy, it is essential to examine the outcomes of different models. Thus, if a highly simplified model makes a prediction, and if the same or a very similar prediction is made by a more elaborate model that includes some mechanisms or details that the first model did not, then we gain some confidence that the prediction is robust. An important benefit derived from mathematical modelling activity is that it demands transparency and accuracy regarding our assumptions, thus enabling us to test our understanding of the disease epidemiology by comparing model results and observed patterns. Models can also assist in decision-making by making projections regarding important issues such as intervention-induced changes in the spread of disease."* [5]

To that end we select parameters according to the following criteria:

1. Provide sufficient realism to assure relevance
2. Keep the model simple
3. Compatible with easily accessible public data for updating/tracking
4. Err toward the worst case rather than optimism

We expect numerous models to come into play in an actual policy implementation, and regional authorities to do their own modeling. In addition, we want our model to be comprehensible by the public and to aid in the important goal of informing the public and enlisting their rational choices in implementing the strategy. To this end our first parameter reformulates the traditional R_0 reproduction rate, which does not have a specific time base, as a time-based spreading rate “ R .” Data from the CDC <https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/cases-in-us.html#investigation> for number of cases for March 16-21 in the US (4226, 7038, 10442, 15219, 18747, 24583) represent a time period when local spread had begun to act exponentially. These data correspond to daily spread rates of (1.665, 1.48, 1.45, 1.23, 1.311). Previously a slow rate attributable to travel had been dominant. After March 21 the rate of spread decreases continuously, which we presume is due to increased government directed social distancing including widespread business closure, stay at home orders, and increased public compliance with these policies. For the model’s initial value we adopted a daily spread rate of 1.414. The fraction of *new* cases is then $R=0.414$. This number is immediately replaced in the model by the next day’s rate, and adjusted by the computed herd immunity factor when re-used following unlock. The unlock spread rate should be measured and adjusted, but there has yet been no unlock.

Our R can be compared with R_0 by selecting a time window for transmission. Several possibilities exist, and since our R is empirically measured we are not too concerned with determining the exact relation. But for example, suppose there is a window of 4 days before a person discovers

104 they have the disease and self-isolates. At our presumed R (cases doubling every two days) there
 105 would be 4 people with the disease at the end of this time, or 3 new cases. That corresponds to $R_0=3$,
 106 which is in line with the high end of estimates given above (3.28 maximum).

107 The number of ventilators in the U.S. including reserves, alternatives (anesthesia machines) and
 108 older equipment is taken at 200,000 [6].

109 The ratio of total likely cases to known reported cases is taken at 6 (16.6% known) with alternative
 110 scenarios checked at 5 (20% known). The lower numbers are more critical due to the way the model
 111 calculates ventilator requirements. Published numbers are typically around 14% [7], so our assumption
 112 has some margin of safety.

113 A precise number for how long a case of COVID-19 lasts is of course not obtainable due to the
 114 wide variation. However this number is not critical for our model. They could last forever and the
 115 herd immunity calculation would be unaffected since it includes both recovered and infected people.
 116 Somewhat more important is the length of time a ventilator is required, if one is required. The best
 117 estimate we could obtain was “up to weeks” since this too is highly variable. We used 14 days for both
 118 numbers.

119 The fraction of cases which require resources such as a ventilator is also important. We used 3.2%
 120 of known cases, or about half of critical cases, taken from Meng, et. al. [8] Lower estimates exist.
 121 This is an area in which additional data and input from more specialized simulations would be used
 122 before public policy formulated. Regionalization is also important as ventilators may not be
 123 distributed where needed. Publication daily of model predictions, assuming they predict or exceed the

124 actual data coming in, we believe would increase confidence and promote redistribution of ventilators
125 according to need.

126 Social networks, location tracking and other massive data mining efforts recommended in
127 research of more persistent (non-pandemic) diseases [9] are specifically not part of our approach. They
128 take time, where we require rapid feedback. They invite abuse for other applications later. But most
129 important, social networks change as soon as a pandemic is announced, change again when
130 government policy is announced, and keep changing. An aggregated tracking and feedback method
131 will work better.

132 **Model dynamics**

133 We use a standard SIR modeling approach [10] but with the time based reproductive factor as
134 described above. The number of new cases is our R factor times the existing cases which are only 4
135 days old or less.

136 During “lockdown” the R factor is adjusted according to (a) the ratio of new cases from the
137 previous day, and (b) the increase in herd immunity factor over the previous day. Only when the
138 caseload is rapidly peaking will day to day changes in herd immunity factor be relevant. When an
139 “unlock” policy is established in the predictive model, the R factor is re-established at the initial value
140 reduced by the herd immunity factor. When actual unlock data is available, this will need further
141 adjustment which we do not specify at this time.

142 This simple model was implemented in a spreadsheet with some facilities for setting days to be
143 unlocked, or they can be set manually. The spreadsheet will be available as supplementary materials

144 and in the <http://medRxiv.org> preprint repository, and is currently available at
 145 <http://shulerresearch.org/covid19.htm>. Each day actual data was used to replace predicted data based
 146 on CDC data at the link specified above. This affects the model's integration base and the effective
 147 reproduction rate R . The number of "new" cases, necessary for bookkeeping active and spreading
 148 cases and ventilator utilization, is deduced from the day to day change in number of cases. Total cases,
 149 used for the herd immunity calculation, is calculated by either 5 or 6 times the known cases as
 150 described.

151 Results

152 The progress toward "curve flattening" is shown in Figure 1, with predicted plots based on actual
 153 data from 3/21, 3/25 and 3/29. One can see that to say flattening was successful is putting it mildly.
 154 Thus arises the dilemma: not enough people are getting sick to develop herd immunity.

155 In the 3/25 and 3/29 cases in Figure 1, herd immunity is incomplete. In the 3/21 case it appears
 156 90% of the population is infected because of numerical overshoot during the rapid rise. One day
 157 quantization may not be enough resolution, the rise is so rapid. The extent to which such an overshoot
 158 might happen in reality (it might) is not of concern to us since this case will not be allowed. In the
 159 following cases it will be apparent that where a slower rise is allowed due to partial lockdown, the total
 160 cases peak around 70% just where it should for $R_0=3$.

161 Figure 2 shows two different dates for total unlock, beginning with the day after the current
 162 extension of lockdown through April 30th. The second scenario begins 6 months later on November 1.
 163 Though the peak is reduced somewhat, ventilator capacity is still wildly exceeded. More ventilators

164 may be produced by then, but 6 months have been wasted when people could have been building herd
 165 immunity, and the economic loss, the permanent loss of houses and business from loan default, and
 166 consequent loss of jobs, will be known is the worst “man-made” disaster of all time, since it was a
 167 reactionary policy decision which we will show is easily prevented.

168 Figure 3 shows a three day a week partial unlock strategy beginning as soon as April 9th.
 169 Actually it could begin April 2nd, but this paper is not likely to be in public view by then. Five weeks
 170 later, total unlock is possible. Critical resource (ventilator) usage expands smoothly toward the 70-
 171 80% range, then drops dramatically just at the time of full unlock. It is possible to unlock a week
 172 earlier, but doing so causes a rise in critical cases which might undermine public confidence.

173

Figure 1. Progress toward “flattening” from 3/21 to 3/29.

Figure 4 shows a sensitivity analysis for the multiple of unknown to known cases of 5 instead of 6.

The situation is worse, but capacity is not exceeded. Most expert opinion is that the multiple will be

179 higher rather than lower, however additional studies involving testing random samples would clarify
 180 this figure, which might vary by region.

181 It appears that the unlock date need not be pinned down in advance. There will be a precipitous
 182 drop in usage of critical resources, and unlock can then occur within a few days.

Figure 2. Alternatives for full unlock.

Figure 3. Three day a week unlock scenario.

Figure 4. Three day a week unlock scenario with 5x unknown to known cases.

Conclusion

It appears that options are available that are not now openly under consideration that would lessen economic impact of the COVID-19 worldwide response by six to ten times, while operating within existing critical medical resources and not causing unnecessary deaths.

194 The public desperately wants more concrete data and projections, projections which can be seen
195 to be valid as the data comes in. Current models involve obscure parameters that make investigators
196 hesitant to share them. The use of a time base reproductive factor would likely be more
197 comprehensible and easier for the public to verify for themselves.

198 It is likely the response to a “partial unlock” would be underwhelming. Many businesses would
199 be hesitant to reopen. However, within a few days this effect will be apparent in the case data, and the
200 unlock schedule can be adjusted higher or lower. Using days of the week is a customary and usual way
201 of managing work schedules that is easily understood.

202 The public will realize, and it must be made clear, that partial unlock is not a declaration of
203 safety from CORVID-19. It is a declaration of reasonable assurance of adequate medical capacity. We
204 recommend a flexible policy allowing those at risk or those simply afraid to continue to isolate, and
205 those who wish or need to work to accept the risk. The end result is a great reduction of the isolation
206 time required for those who choose to continue to isolate.

207 The final and perhaps most important benefit of this methodology and strategy is the rapid
208 availability of excess medical resources from the U.S. (once unlock is achieved, in about 5 weeks from
209 start) to assist the rest of the world which is not so fortunate in number of ventilators or comparable
210 devices. Further, healthcare workers who have acquired immunity may well want to volunteer to help
211 in other countries. Let’s do this.

212 Acknowledgements

213 The author appreciates feedback, comments, references and general moral support from Theodore
 214 Koukouvitis at every stage of this research, and for some time previously as the author expanded his
 215 modeling and theoretical work in social systems safety and reliability into cooperative and
 216 evolutionary/biological systems.

217 References

- 218 [1] Liu, Y., Gayle, A.A., Wilder-Smith, A., Rocklöv, J. (2020). The reproductive number of COVID-
 219 19 is higher compared to SARS coronavirus. *Journal of Travel Medicine*, **27**, 2, March, taaa021,
 220 <https://doi.org/10.1093/jtm/taaa021>.
- 221 [2] He, M.M. (2016). Driving through the great recession: why does motor vehicle fatality decrease
 222 when the economy slows down? *Soc. Sci. Med.*, **155**, 1-11, April,
 223 <https://doi.org/10.1016/j.socscimed.2016.02.016>
- 224 [3] Maruthappu, M., Watkins, J., Noor, A.M., Williams, C., Ali, R., Sullivan, R., et. al. (2016).
 225 Economic downturns, universal health coverage, and cancer mortality in high-income and middle-
 226 income countries, 1990–2010: a longitudinal analysis, *The Lancet*, **388**, 10045, 684-695, August,
 227 [https://doi.org/10.1016/S0140-6736\(16\)00577-8](https://doi.org/10.1016/S0140-6736(16)00577-8)
- 228 [4] Fan, V.Y., Jamison, D.T., Summers, L.H. (2018). Pandemic risk: how large are the expected
 229 losses? *Policy & Practice, Bulletin of the World Health Organization*, **96**, 129-134, December,
 230 <http://dx.doi.org/10.2471/BLT.17.199588>

- 231 [5] Huppert, A. and Katriel, G. (2013). Mathematical modeling and prediction in infectious disease
232 epidemiology, *Clin. Microbiol. Infect.*, **19**, 999-1005, June, [https://doi.org/10.1111/1469-](https://doi.org/10.1111/1469-0691.12308)
233 [0691.12308](https://doi.org/10.1111/1469-0691.12308)
- 234 [6] Halpern, N.A. and Tan, K.S., (2020). U.S. ICU Resource Availability for COVID-19, *Soc. Of Crit.*
235 *Care Med., SCCM Ventilator Taskforce*, ver. 2, March 25,
236 [https://www.sccm.org/getattachment/Blog/March-2020/United-States-Resource-Availability-for-](https://www.sccm.org/getattachment/Blog/March-2020/United-States-Resource-Availability-for-COVID-19/United-States-Resource-Availability-for-COVID-19.pdf?lang=en-US)
237 [COVID-19/United-States-Resource-Availability-for-COVID-19.pdf?lang=en-US](https://www.sccm.org/getattachment/Blog/March-2020/United-States-Resource-Availability-for-COVID-19/United-States-Resource-Availability-for-COVID-19.pdf?lang=en-US)
- 238 [7] Li, R., Pei, S., Chen, B., Song, Y., Zhang, T., Yang, W. and Shaman, J. (2020). Substantial
239 undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV2),
240 *Science*, eabb3221, March, <https://doi.org/10.1126/science.abb3221>
- 241 [8] Meng, L., Qiu, H., Wan, L., Ai, Y., Xue, Z., et. al. (2020). Intubation and Ventilation amid the
242 COVID-19 Outbreak: Wuhan's Experience, *Anesthesiology*, special section on COVID-19, no
243 pagination specified, March, <https://doi.org/10.1097/ALN.0000000000003296>
- 244 [9] Marathe, M. and Ramakrishnan, N. (2013). Recent Advances in Computational Epidemiology,
245 *IEEE Intell. Syst.*, **28**, 4, 96-101, July-August, <https://dx.doi.org/10.1109%2FMIS.2013.114>
- 246 [10] Marathe, M., Vullikanti, A.K.S. (2013). Computational epidemiology, *Communications of the*
247 *ACM*, **56**, 7, 88-96, July, <http://doi.acm.org/10.1145/2483852.2483871>