The Use of Adjuvant Therapy in Preventing Progression to Severe Pneumonia in Patients with Coronavirus Disease 2019: A Multicenter Data Analysis

Zhichao Feng, MD; Jennifer Li, MBBS; Shanhu Yao, MD; Qizhi Yu, MD; Wenming Zhou, MD;
Xiaowen Mao, MD; Huiling Li, BD; Wendi Kang, BD; Xin Ouyang, MD; Ji Mei, MD; Qiuhua
Zeng, MD; Jincai Liu, MD; Xiaoqian Ma, MD; Pengfei Rong, MD; Wei Wang, MD

Author Affiliations: Department of Radiology, Third Xiangya Hospital, Central South University,
Changsha, Hunan, China (Z. Feng, S. Yao, H. Li, W. Kang, X. Ma, P. Rong, W. Wang); Westmead
Clinical School, Faculty of Medicine and Health, University of Sydney, Sydney, Australia (J. Li);
Changsha Public Health Treatment Center, Changsha, Hunan, China (Q. Yu); Department of
Radiology, First Hospital of Changsha, Changsha, Hunan, China (Q. Yu); Department of Medical
Imaging, First Hospital of Yueyang, Yueyang, Hunan, China (W. Zhou); Department of Medical
Imaging, Central Hospital of Shaoyang, Shaoyang, Hunan, China (X. Mao); Department of
Radiology, Central Hospital of Xiangtan, Xiangtan, Hunan, China (X. Ouyang); Department of
Radiology, Second Hospital of Changde, Changde, Hunan, China (J. Mei); Department of
Radiology, Central Hospital of Loudi, Loudi, Hunan, China (Q. Zeng); Department of Radiology,
First Affiliated Hospital of University of South China (J. Liu).

Corresponding Authors: Pengfei Rong, MD, Department of Radiology, Third Xiangya Hospital,
Central South University, No. 138 Tongzipo Rd, Changsha, Hunan, 410013, China
(rongpengfei66@163.com); Wei Wang, MD, Department of Radiology, Third Xiangya Hospital,

Central South University, No. 138 Tongzipo Rd, Changsha, Hunan, 410013, China

(cjr.wangwei@vip.163.com).

Word count: 2992 words (main text, not including abstract, reference, and figure legends)

KEYPOINTS

Question: Does the use of adjuvant therapy reduce progression to severe pneumonia in patients with coronavirus disease 2019 (COVID-19)?

Findings: In this retrospective, observational cohort study involving 564 patients with confirmed COVID-19, hypertension was an independent risk factor for progression to severe pneumonia irrespective of age and those on angiotensin-converting enzyme inhibitor (ACEI) or angiotensin receptor blocker (ARB) therapy were less likely to develop severe COVID-19 pneumonia, while nonspecific antivirals or chloroquine did not have significant impact on clinical progression.

Meaning: Hypertensive patients with COVID-19 should not have ACEI or ARB ceased, unless there is a strong indication or further epidemiological evidence, given its potential protective effects.

ABSTRACT

IMPORTANCE Coronavirus disease 2019 (COVID-19) is a global pandemic associated with high mortality and effective treatment to prevent clinical deterioration to severe pneumonia has not yet been well clarified.

OBJECTIVE To investigate the role of several adjuvant treatments in preventing severe pneumonia in patients with COVID-19.

DESIGN, SETTING, AND PARTICIPANTS Multicenter, retrospective cohort study of 564 consecutively hospitalized patients with confirmed COVID-19 at Third Xiangya Hospital of Central South University, Changsha Public Health Treatment Center, First Hospital of Yueyang, Junshan People's Hospital of Yueyang, Central Hospital of Shaoyang, Central Hospital of Xiangtan, Second Hospital of Changde, Central Hospital of Loudi, and First Affiliated Hospital of University of South China in Hunan province from January 17, 2020 to February 28, 2020; The final date of follow-up was March 15, 2020.

EXPOSURES Nonspecific antivirals (arbidol, lopinavir/ritonavir, and interferon α), antihypertensives, and chloroquine.

MAIN OUTCOMES AND MEASURES The development of severe COVID-19 pneumonia; Demographic, epidemiological, clinical, laboratory, radiological, and treatment data were collected and analyzed.

RESULTS Of 564 patients, the median age was 47 years (interquartile range, 36-58 years), and 284 (50.4%) patients were men. Sixty-nine patients (12.2%) developed severe pneumonia. Patients who developed severe pneumonia were older (median age of 59 and 45 years, respectively), and more patients had comorbidities including hypertension (30.4% and 12.3%,

respectively), diabetes (17.4% and 6.7%, respectively), and cardiovascular disease (8.7% and 3.2%, respectively) and presented with fever (84.1% and 60.4%, respectively) and shortness of breath (10.1% and 3.8%, respectively) compared with those who did not. Nonspecific antiviral therapy did not prevent clinical progression to severe pneumonia, although fewer hypertensive patients on angiotensin-converting enzyme inhibitors or angiotensin-receptor blockers (ACEI/ARB) therapy developed severe pneumonia in contrast with those on non-ACEI/ARB antihypertensive therapy (1 of 16 [6.3%] patients and 16 of 49 [32.7%] patients, respectively [difference, 26.4%; 95% CI, 1.5% to 41.3%]). Multivariate logistic regression analysis showed that hypertension without receiving ACEI/ARB therapy was an independent risk factor (odds ratio [OR], 2.07; 95% CI, 1.07 to 4.00) for developing severe pneumonia irrespective of age. Besides, none of patients treated with chloroquine developed severe pneumonia, though without significance (difference, 12.0%; 95% CI, -3.5% to 30.0%) by propensity score matching.

CONCLUSIONS AND RELEVANCE Hypertensive patients on ACEI or ARB may be protective from severe pneumonia in COVID-19 and hence these therapies should not be ceased unless there is a strong indication or further epidemiological evidence. Though none of the current antiviral and immunoregulation therapy showed benefit in preventing COVID-19 progression, chloroquine deserved further investigation.

INTRODUCTION

The world has witnessed the rapid escalation of 2019 coronavirus disease (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), now becoming a global pandemic.¹ While many patients may be asymptomatic or experience only mild symptoms, reports so far have identified the elderly and patients with chronic comorbidities such as hypertension, diabetes, and cardiovascular disease to be at higher risk of progression to severe COVID-19 pneumonia, characterized by acute respiratory distress syndrome (ARDS) and/or multiorgan failure.²⁻⁴ These severe patients require more intensive medical resource utilization and have worse prognosis, with a case fatality rate about 20 times higher than that of non-severe patients.^{5,6} While the implementation of effective prevention measures is essential, there clearly exists an urgent need for interventions to modulate disease progression in the face of an increasing number of infected individuals, which have overwhelmed the capacity of hospital systems in some countries.

While there is no specific therapy for COVID-19, current progress with investigational therapies includes the development and trial of a coronavirus vaccine and interrogation of several orphan drugs in randomized control trials (RCTs).⁷ This includes remdesivir, which has garnered the most interest in an ongoing adaptive RCT and other nucleotide analogues including ritonavir, favipiravir and galidesvir.⁸ A recently published open-label RCT showed no benefit of the protease inhibitors lopinavir/ritonavir in reducing clinical severity or mortality associated with COVID-19, despite in vitro evidence of activity against SARS-CoV-2.⁹ Other drugs of interest include antiviral therapy (pegylated interferon-alpha, arbidol) and immunomodulators (hydroxychloroquine) are also under investigation and results are eagerly monitored.^{10,11}

Inhibition of the spike glycoprotein (S-protein) may also be a viable target and has recently garnered increasing interest - particularly the potential interactions with angiotensinogen converting enzyme 2 (ACE2). It is proposed that the binding of the SARS-CoV-2 S-protein to the functional receptor ACE2 results in proteolytic cleavage by type II transmembrane serine proteases (TMPRSS2).¹²⁻¹⁴ There has been considerable interest and debate regarding the risk of severe COVID-19 pneumonia in patients on angiotensinogen converting enzyme inhibitors (ACEI) or angiotensin receptor blocker (ARB) therapy, presumably through the modulation of ACE2 expression.^{4,15,16} The role of ACEI and ARB in COVID-19 infection remains controversial – as the data is not adjusted for confounding factors of age and other comorbidities to definitively show that ACEI or ARB are independently associated with worse outcomes – and of course, this does not show causality either.¹⁷ Hence, various independent medical societies have put out position statements that recognize the potential risks but given the current quality of evidence, recommend against inappropriate cessation or switch from ACEI/ARB therapy for hypertension, given its proven benefits.^{18,19}

Therefore, the objective of this study was to investigate the role of several adjuvant treatments in preventing severe pneumonia in COVID-19 patients from seven cities in Hunan province, China.

METHODS

Study Design and Participants

The Institutional Review Board of Third Xiangya Hospital of Central South University approved our study and waived informed consent for the retrospective nature of this study. All consecutive adult patients with confirmed COVID-19 who were treated at Third Xiangya Hospital of Central

South University, Changsha Public Health Treatment Center, First Hospital of Yueyang, Junshan People's Hospital of Yueyang, Central Hospital of Shaoyang, Central Hospital of Xiangtan, Second Hospital of Changde, Central Hospital of Loudi, and First Affiliated Hospital of University of South China from January 17, 2020 to February 28, 2020 were enrolled. The diagnosis of COVID-19 was established based on the World Health Organization (WHO) interim guidance, and a confirmed case was defined as a positive result to high-throughput sequencing or real-time reverse transcription-polymerase chain reaction (RT-PCR) assay for nasal and pharyngeal swab specimens. Patients without available clinical or imaging data were excluded. The clinical outcomes including severe pneumonia development, discharge, and death were monitored up to March 15, 2020.

Clinical and imaging data collection

We extracted the demographic, exposure history, comorbidities, clinical symptoms, and laboratory data on admission from electronic medical records. Medication history among individuals with hypertension was recorded, including ACEI/ARB, calcium-antagonists, β -blocker, and diuretic agents. All laboratory tests were performed according to the clinical care needs of the patient during hospitalization, including a complete blood count, assessment of liver and renal function, coagulation testing, and measures of electrolytes, creatine kinase, lactate dehydrogenase, procalcitonin, and C-reactive protein. The use of antiviral therapy (arbidol, lopinavir/ritonavir, and interferon α) and chloroquine were obtained from the medication administration record. The severity degree of COVID-19 at the time of admission is defined as follows: (1) mild type, with fever or respiratory tract symptoms but without radiological pneumonia. (2) moderate type, with fever, respiratory tract symptoms, and radiological evidence of pneumonia. (3) severe type, with

one of the following: a) respiratory distress (respiratory rate ≥ 30 beats/min); b) hypoxia (oxygen saturation $\leq 93\%$ in the resting state); c) hypoxemia (arterial blood oxygen partial pressure/oxygen concentration $\leq 300\text{mmHg}$). (4) critical type, with one of the following: a) respiratory failure requiring mechanical ventilation; b) shock; c) intensive care unit (ICU) admission is required for combined other organs failure. In this study, severe COVID-19 pneumonia broadly included severe and critical types.

The CT images were acquired on admission and at intervals of 3-5 days and were reviewed independently by two experienced radiologists. Typical CT findings for COVID-19 were defined as peripherally distributed multifocal ground-glass opacities (GGOs) with patchy consolidations and posterior part or lower lobe involvement predilection.²⁰ Each of the five lung lobes was reviewed for GGO and consolidation. The lesions extent within each lung lobe was semi-quantitatively evaluated by scoring from 0 to 5 based on the degree of involvement: score 0, none involvement; score 1, $\leq 5\%$ involvement; score 2, $6\sim 25\%$ involvement; score 3, $26\sim 50\%$ involvement; score 4, $51\sim 75\%$ involvement; score 5, $> 75\%$ involvement. The total score was calculated by summing up scores of all five lobes to provide a lung CT score ranging from 0 to 25.²¹ The mean value of lung CT scores evaluated by the two radiologists was used for analysis.

Statistical Analysis

Our primary outcome of analysis was progression to severe COVID-19 pneumonia. Continuous variables are presented as the median and interquartile range (IQR), and categorical variables are presented as frequency and percentage. Differences between groups were analyzed using Student's t-test or Mann-Whitney U test for quantitative variables according to the normal distribution and Chi-square test or Fisher's exact test for categorical variables. Meanwhile, the differences in

distributions were also reported using differences with 95% confidence intervals (CIs). Univariate and multivariate logistic regression with backward stepwise selection based on likelihood ratio was used to identify the risk factors for the development of severe pneumonia, and the corresponding odds ratios (ORs) and 95% CIs were determined. We also employed the propensity score matching (PSM) with a 1:1 ratio to minimize selection bias and adjust for the imbalance between patients with or without chloroquine therapy. A one-to-one nearest-neighbour matching algorithm with an optimal calliper of 0.2 without replacement was used to generate pairs of patients. The analyses regarding different factors were based on non-missing data, and missing data were not imputed. All statistical analyses were performed using SPSS statistics software (version 22.0, IBM SPSS Inc, Chicago, IL, USA). A two-sided *P* value of less than 0.05 considered to be statistically significant.

RESULTS

Patient Characteristics

The study population included 564 adult patients with confirmed COVID-19 from nine hospitals in seven cities (Figure 1), and their clinical characteristics on admission are presented in Table 1. The median age of patients was 47 years (IQR, 36-58; range, 19-84 years), and 284 (50.4%) were men. 132 (23.4%) patients had comorbidities, including hypertension (n = 82 [14.5%]), diabetes (n = 45 [8.0%]), cardiovascular disease (n = 22 [3.9%]), and chronic obstructive pulmonary disease (COPD, n = 16 [2.8%]). The most commonly self-reported symptoms at onset of illness were fever (n = 357 [63.3%]), cough (n = 323 [57.2%]), fatigue (n = 87 [15.4%]), sputum production (n = 51 [9.0%]), anorexia (n = 26 [4.6%]), and shortness of breath (n = 26 [4.6%]). A total of 516 (91.5%) patients had typical abnormal findings on chest CT (Supplementary Figure 1),

with a median lung CT score of 6 (IQR, 4-11; range, 0-25).

Differences between severe and non-severe patients

In total, 69 of 564 (12.2%) patients were identified to have severe pneumonia. As summarized in Table 1, these patients were older and more likely to have pre-existing comorbidities, including hypertension, diabetes, cardiovascular disease, COPD, and chronic renal disease compared with the non-severe patients. The clinical symptoms at illness onset also differed, with severe patients more likely to present with fever and shortness of breath. Furthermore, patients with severe pneumonia had elevated liver enzyme (aspartate aminotransferase [difference, -10.8; 95% CI, -14.2 to -7.6; $P < 0.001$]), renal dysfunction (blood urea nitrogen [difference, -1.0; 95% CI, -1.4 to -0.6; $P < 0.001$]), abnormal coagulation function (D-dimer ≥ 0.05 mg/L [difference, -18.3%; 95% CI, -29.5% to -5.7%; $P = 0.004$]), raised inflammation-related parameters (lactate dehydrogenase [difference, -112.2; 95% CI, -134.0 to -89.9; $P < 0.001$]), and worse lung CT score (difference, -8; 95% CI, -10 to -7; $P < 0.001$). However, lymphocyte count (difference, 0.4; 95% CI, 0.3 to 0.5; $P < 0.001$) was significantly lower in patients with severe pneumonia.

Antiviral therapy did not prevent severe pneumonia

Thirty patients were classified with severe pneumonia on admission and were excluded, to leave 534 patients with non-severe COVID-19 on admission for further analysis. Amongst this group, 39 (7.3%) patients developed severe pneumonia during hospitalization. Most ($n = 503$ [94.2%]) patients received single or combined antiviral therapy during their admission. After adjusting for age, gender, smoking history, hypertension, diabetes, cardiovascular disease, COPD, and lung CT score, none of these investigational antiviral therapies (all $P > 0.05$) were associated with reduced risk of progression to severe disease, as shown in Table 2.

ACEI/ARB therapy in patients with hypertension

Eighty-two individuals were recorded to have hypertension and only 16 patients were on ACEI/ARB therapy (including 6 cases receiving ARB monotherapy, 9 cases receiving combined ARB and calcium-antagonist, and 1 case receiving ACEI monotherapy). There were 49 patients on non-ACEI/ARB antihypertensive therapy (including 39 cases receiving calcium-antagonist monotherapy, 3 cases receiving β -blocker monotherapy, 2 cases receiving diuretic monotherapy, and 5 cases receiving combined calcium-antagonist and β -blocker) and all were maintained on their usual antihypertensive regimen during hospitalization. The remaining 17 cases were not on antihypertensive medications or limited by unavailable data and received amlodipine or nitrendipine to control blood pressure as necessary. As summarized in Table 3, hypertensive patients on ACEI/ARB therapy had similar baseline clinical characteristics and laboratory findings compared with those on non-ACEI/ARB antihypertensive therapy. However, patients on ACEI/ARB therapy were less likely to develop severe pneumonia, in contrast with those on non-ACEI/ARB therapy (difference, -26.4%; 95% CI, -41.3%~-1.5%; $P = 0.037$). Laboratory and imaging data from 65 patients with hypertension during hospitalization were analyzed (Supplementary Figure 2). Lactate dehydrogenase and CT lung scores were lower in patients on ACEI/ARB therapy than those on non-ACEI/ARB therapy at about 3 days after admission.

To further confirm the clinical risk factors for severe pneumonia in COVID-19 patients, univariate and multivariate logistic regression analyses were performed (Table 4). Univariate analyses identified age, hypertension, diabetes, and cardiovascular disease to be associated with the development of severe pneumonia (all $P < 0.05$), but only older age (OR and 95% CI, 1.05 [1.03 to 1.07]; $P < 0.001$) and hypertension without ACEI/ARB therapy (OR and 95% CI, 2.07 [1.07 to

4.00]; $P = 0.030$) were independent risk factors in multivariate analysis. Compared with patients without hypertension, hypertension on ACEI/ARB therapy was not associated with developing severe pneumonia (OR and 95% CI, 0.41 [0.05 to 3.19]; $P = 0.392$).

Chloroquine therapy in non-severe patients

Among 534 patients with non-severe COVID-19 on admission, only patients in Changsha received chloroquine therapy (500mg, bid). Thus, 192 non-severe COVID-19 patients without receiving ACEI/ARB therapy in Changsha (25 patients with chloroquine therapy and 167 patients without chloroquine therapy) were included and 25 well-matched pairs for comparison were identified for comparison using PSM, with matched covariates of age, gender, hypertension, and lung CT score (Supplementary Table 1). None of patients treated with chloroquine developed severe pneumonia, though without significance (difference, 12.0%; 95% CI, -3.5% to 30.0%; $P = 0.074$). In addition, there was no significant difference in reported adverse events and $\geq$ Grade 3 adverse events between the two groups ($P = 0.777$ and 0.440, respectively).

DISCUSSION

In this study, we described the use of several adjuvant treatments in preventing severe pneumonia development in COVID-19 patients. Hypertensive patients who received ACEI/ARB therapy had a lower risk of developing severe pneumonia compared with those on non-ACEI/ARB antihypertensive therapy, which suggested that ACEI/ARB therapy may be protective from severe pulmonary injury for hypertensive patients. However, these results should be interpreted with caution owing to potential bias and residual confounders in this observational study.

Consistent with earlier reports and clinical experience around the world, our cohort confirmed that

that older age and hypertension without ACEI/ARB therapy were independently associated with developing severe pneumonia in COVID-19.^{2,3,22} The association between age and severe pneumonia may be attributed to the accumulation of medical comorbidities with age or decline in immune function over time. Given the close collinearity between age and hypertension, our study adjusted for age by multivariate regression analysis and revealed that hypertension remained an independent risk factor for disease progression irrespective of age for patients without receiving ACEI/ARB therapy.

Our study examined the controversial topic of COVID-19 and ACEI/ARB therapy in hypertensive patients. We found that hypertensive patients on ACEI/ARB therapy were less likely to develop severe pneumonia, although we acknowledged that the analysis was conducted in limited sample size - with only 16 of 82 patients on ACEI/ ARB therapy. The dynamic profile of laboratory and radiological findings further showed that in patient on ACEI/ARB therapy, lactose dehydrogenase and CT lung score were lower in than those on non-ACEI/ARB therapy at about 3 days after admission. Together, these findings were suggestive that ACEI/ ARB therapy may be protective in hypertensive patients with COVID-19.

COVID-19 patients with hypertension may endure worse outcomes due to decreased ACE2 levels. ACE2, an important component of the renin-angiotensin-aldosterone system (RAAS), is highly expressed in both the lung and myocardium and converts angiotensin II (AII) into angiotensin 1-7, which can downregulate ACE and directly cause vasodilatation.²³⁻²⁵ Downregulation of ACE2 level has also been shown in hypertensive animal models, although there has been limited success in translational confirmation in human studies.^{26,27} None the less, this may be exacerbated by the further reduction in available ACE2 following coronavirus infection. Both SARS-CoV and

SARS-CoV-2 can gain entry into the host cell through binding of the S-protein to the membrane-bound ACE2 aminopeptidase, which then is cleaved by serine proteases to allow for membrane fusion and invasion.^{14,28} Following infection, there is a reduction in pulmonary ACE2, either through internalization with viral entry and/or downregulation of ACE2 enzyme during this process. Hence this may exacerbate the lower baseline ACE2 function in hypertensive patients to lead to the accumulation of AII and reduced angiotensin 1-7 and subsequently worse immune-related cytolysis and lung injury, which was validated by the linear association between elevated plasma AII level and viral load or lung injury in COVID-19 patients.²⁹

The use of ACEI or ARB likely promotes feedback upregulation of ACE2 expression in hypertension, although there are animal and human studies which have also found either no effect or downregulation of ACE2 level in the setting of diabetes, hypertension or cardiovascular disease, with or without ACE/ARB therapy.^{25,30-33} Nonetheless, this may explain our finding that ACEI/ARB therapy can be protective in SARS-CoV-2 infection. Further evidence that would support this come from earlier studies in which ACEI/statins were protective in severe pneumonia, recombinant ACE2 protected mice from lung injury following SARS-CoV infection, and RAAS blockade reduced acute lung injury in rats injected with SARS-CoV S-protein.^{34,35}

Currently, there is still no specific treatment for COVID-19 except for supportive care. In our cohort, 94.2% of non-severe patients received nonspecial antiviral therapy from just after admission. The combination protocol and medication dose varied in different hospitals. However, our results did not support any single or combined antiviral treatments to limit the progression of COVID-19. In addition, none of 25 patients treated with chloroquine developed severe pneumonia, though without statistical significance. This is unfortunately limited by the degree of heterogeneity

in our cohort and relatively small size. Chloroquine is known to block virus infection by increasing endosomal pH required for virus/cell fusion, as well as interfering with the glycosylation of cellular receptors of coronavirus, which has been confirmed to inhibit SARS-CoV-2 in vitro.³⁶⁻³⁸ A recent open-label non-randomized clinical trial reported that hydroxychloroquine combined with azithromycin is significantly associated with viral load reduction/disappearance in COVID-19 patients.³⁹ Even so, more RCTs to assess the therapeutic effects of chloroquine on COVID-19 need to be conducted.

LIMITATIONS

There were several limitations in this study. First, our study was conducted by retrospective data analysis. Second, some cases had incomplete laboratory test results, given the variation in the clinical management among different hospitals. All data generation was clinically driven and not systematic. Third, because most patients had recovered and discharged at the time of data cutoff and only 2 patients died during admission, the impact of adjuvant treatment on the clinical outcomes cannot be analyzed. Fourth, there were limited hypertensive patients who received ACEI or ARB therapy, so we cannot analyze the effect of ACEI and ARB respectively. A larger cohort study of COVID-19 patients with hypertension from other cities in China and other countries would help to further validate the potential protective effect of ACEI/ARB therapy.

CONCLUSIONS

COVID-19 is a fast-evolving, global pandemic with high mortality from severe pneumonia, ARDS, and multiorgan failure, with the highest risk in the elderly and patients with underlying hypertension. The current adjuvant antiviral and immunoregulation therapies have not yet proven

benefits in clinical settings. But importantly, our results show the potential protective effects of

ACEI or ARB therapy to limiting progression to severe lung disease in hypertensive patients.

ACKNOWLEDGEMENT

We thank Hongzhuan Tan, PhD, from the Department of Epidemiology, Xiangya School of Public Health, Central South University for his statistical analysis discussion. We also thank Wei Nie, MD, Qin Liu, MD, Jing Zhao, MD, Junhong Duan, MD, Zhimin Yan, MD, Min Yang, MD from Department of Radiology, Ying Li, MD from Health Management Center, Sujie Jia, MD from Department of Pharmacy, Third Xiangya Hospital, Central South University, for their data collection. They were not compensated for their contributions.

Conflict of Interest Disclosures: None reported.

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Author Contributions:

Drs W. Wang and P. Rong had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Drs P. Rong and W. Wang contributed equally to this article.

Concept and design: Z. Feng, J. Li, S. Yao, X. Ma, P. Rong, W. Wang; Acquisition, analysis, or interpretation of data: Z. Feng, Q. Yu, W. Zhou, X. Mao, H. Li, W. Kang, X. Ouyang, J. Mei, Q. Zeng, J. Liu; Drafting of the manuscript: Z. Feng, J. Li, S. Yao; Critical revision of the manuscript for important intellectual content: J. Li, X. Ma, P. Rong, W. Wang; Statistical analysis: Z. Feng, S. Yao, W. Kang; Obtained funding: P. Rong; Administrative, technical, or material support: S. Yao, Q. Yu, W. Zhou, X. Mao, X. Ma, P. Rong, W. Wang.

REFERENCES

1. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. 2020;10.1001/jama.2020.2648.
2. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA*. 2020.
3. Guan WJ, Ni ZY, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. 2020.
4. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med*. 2020.
5. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet (London, England)*. 2020;395(10223):507-513.
6. Yang Y, Lu Q, Liu M, et al. Epidemiological and clinical features of the 2019 novel coronavirus outbreak in China. *MedRxiv*. 2020:2020.2002.2010.20021675.
7. NIH clinical trial of investigational vaccine for COVID-19 begins. <https://www.nih.gov/news-events/news-releases/nih-clinical-trial-investigational-vaccine-covid-19-begins>. 2020.
8. Holshue ML, DeBolt C, Lindquist S, et al. First Case of 2019 Novel Coronavirus in the United States. *N Engl J Med*. 2020;382(10):929-936.
9. Cao B, Wang Y, Wen D, et al. A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-19. *N Engl J Med*. 2020.
10. Li G, De Clercq E. Therapeutic options for the 2019 novel coronavirus (2019-nCoV). *Nat Rev Drug Discov*. 2020;19(3):149-150.
11. Liu J, Cao R, Xu M, et al. Hydroxychloroquine, a less toxic derivative of chloroquine, is effective in inhibiting SARS-CoV-2 infection in vitro. *Cell Discov*. 2020;6(1):16.
12. Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020.
13. Wrapp D, Wang N, Corbett KS, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science*. 2020;367(6483):1260-1263.
14. Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell*. 2020.
15. Kuster GM, Pfister O, Burkard T, et al. SARS-CoV2: should inhibitors of the renin-angiotensin system be withdrawn in patients with COVID-19? *Eur Heart J*. 2020.
16. Peng YD, Meng K, Guan HQ, et al. [Clinical characteristics and outcomes of 112 cardiovascular disease patients infected by 2019-nCoV]. *Zhonghua xin xue guan bing za zhi*. 2020;48(0):E004.
17. Patel AB, Verma A. COVID-19 and Angiotensin-Converting Enzyme Inhibitors and Angiotensin Receptor Blockers: What Is the Evidence? *JAMA*. 2020.
18. HFSA/ACC/AHA Statement Addresses Concerns Re: Using RAAS Antagonists in COVID-19. <https://wwwacc.org/latest-in-cardiology/articles/2020/03/17/08/59/hfsa-acc-aha-statement-addresses-concerns-re-using-raas-antagonists-in-covid-19>. 2020.
19. The Coronavirus Conundrum: ACE2 and Hypertension Edition.

- <http://www.nephjccom/news/covidace2>. 2020.
20. Zu ZY, Jiang MD, Xu PP, et al. Coronavirus Disease 2019 (COVID-19): A Perspective from China. *Radiology*. 2020;200490.
 21. Pan F, Ye T, Sun P, et al. Time Course of Lung Changes On Chest CT During Recovery From 2019 Novel Coronavirus (COVID-19) Pneumonia. *Radiology*. 2020;0(0):200370.
 22. Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med*. 2020.
 23. Sodhi CP, Wohlford-Lenane C, Yamaguchi Y, et al. Attenuation of pulmonary ACE2 activity impairs inactivation of des-Arg(9) bradykinin/BKB1R axis and facilitates LPS-induced neutrophil infiltration. *Am J Physiol Lung Cell Mol Physiol*. 2018;314(1):L17-L31.
 24. Koka V, Huang XR, Chung AC, Wang W, Truong LD, Lan HY. Angiotensin II up-regulates angiotensin I-converting enzyme (ACE), but down-regulates ACE2 via the AT1-ERK/p38 MAP kinase pathway. *Am J Pathol*. 2008;172(5):1174-1183.
 25. Ocaranza MP, Godoy I, Jalil JE, et al. Enalapril attenuates downregulation of Angiotensin-converting enzyme 2 in the late phase of ventricular dysfunction in myocardial infarcted rat. *Hypertension*. 2006;48(4):572-578.
 26. Crackower MA, Sarao R, Oudit GY, et al. Angiotensin-converting enzyme 2 is an essential regulator of heart function. *Nature*. 2002;417(6891):822-828.
 27. Mendoza-Torres E, Oyarzun A, Mondaca-Ruff D, et al. ACE2 and vasoactive peptides: novel players in cardiovascular/renal remodeling and hypertension. *Ther Adv Cardiovasc Dis*. 2015;9(4):217-237.
 28. de Wit E, van Doremalen N, Falzarano D, Munster VJ. SARS and MERS: recent insights into emerging coronaviruses. *Nat Rev Microbiol*. 2016;14(8):523-534.
 29. Liu Y, Yang Y, Zhang C, et al. Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury. *Science China Life sciences*. 2020;63(3):364-374.
 30. Romero CA, Orias M, Weir MR. Novel RAAS agonists and antagonists: clinical applications and controversies. *Nat Rev Endocrinol*. 2015;11(4):242-252.
 31. Ferrario CM, Jessup J, Chappell MC, et al. Effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockers on cardiac angiotensin-converting enzyme 2. *Circulation*. 2005;111(20):2605-2610.
 32. Agata J, Ura N, Yoshida H, et al. Olmesartan is an angiotensin II receptor blocker with an inhibitory effect on angiotensin-converting enzyme. *Hypertens Res*. 2006;29(11):865-874.
 33. Burrell LM, Risvanis J, Kubota E, et al. Myocardial infarction increases ACE2 expression in rat and humans. *Eur Heart J*. 2005;26(4):369-375; discussion 322-364.
 34. Mortensen EM, Pugh MJ, Copeland LA, et al. Impact of statins and angiotensin-converting enzyme inhibitors on mortality of subjects hospitalised with pneumonia. *Eur Respir J*. 2008;31(3):611-617.
 35. Imai Y, Kuba K, Rao S, et al. Angiotensin-converting enzyme 2 protects from severe acute lung failure. *Nature*. 2005;436(7047):112-116.
 36. Yao X, Ye F, Zhang M, et al. In Vitro Antiviral Activity and Projection of Optimized Dosing Design of Hydroxychloroquine for the Treatment of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). *Clin Infect Dis*. 2020.

37. Wang M, Cao R, Zhang L, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res.* 2020;30(3):269-271.
38. Vincent MJ, Bergeron E, Benjannet S, et al. Chloroquine is a potent inhibitor of SARS coronavirus infection and spread. *Virology*. 2005;2(1):69.
39. Gautret P, Lagier J-C, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *International Journal of Antimicrobial Agents*. 2020:105949.

FIGURE LEGENDS

Figure 1 The study design flow chart of the included patients

Table 1 Clinical characteristics of 564 COVID-19 patients according to disease severity

Characteristics	Total (n =564)	Severe (n = 69)	Non-Severe (n = 495)	Difference (95% CI)	P
Age (years)	47 (36 to 58)	59 (51 to 66)	45 (35 to 56)	-13 (-16 to -9)	< 0.001
< 50	317 (56.2%)	16 (23.2%)	301 (60.8%)		
50~65	160 (28.4%)	31 (44.9%)	129 (26.1%)		
≥ 65	87 (15.4%)	22 (31.9%)	65 (13.1%)		
Male gender	284 (50.4%)	39 (56.5%)	245 (49.5%)	-7.0 (-18.9 to 5.5)	0.274
Exposure within 14 days				-7.1 (-19.3 to 5.2)	0.267
recently been to Wuhan	251 (44.5%)	35 (50.7%)	216 (43.6%)		
contacted with people from Wuhan or local infected patients	313 (55.5%)	34 (49.3%)	279 (56.4%)		
Smoking history	47 (8.3%)	9 (13.0%)	38 (7.7%)	-5.4 (-15.5 to 1.2)	0.131
Comorbidity					
Any	132 (23.4%)	37 (53.6%)	95 (19.2%)	-34.4 (-46.2 to -22.2)	< 0.001
Hypertension	82 (14.5%)	21 (30.4%)	61 (12.3%)	-18.1 (-30.1 to -8.0)	< 0.001
Diabetes	45 (8.0%)	12 (17.4%)	33 (6.7%)	-10.7 (-21.5 to -3.1)	0.002
Cardiovascular disease	22 (3.9%)	6 (8.7%)	16 (3.2%)	-5.5 (-14.6 to -0.4)	0.028
COPD	16 (2.8%)	9 (13.0%)	7 (1.4%)	-11.6 (-21.6 to -5.4)	< 0.001
Cerebrovascular disease	5 (0.9%)	0 (0)	5 (1.0%)	1.0 (-4.3 to 2.3)	0.402
Chronic renal disease	3 (0.5%)	2 (2.9%)	1 (0.2%)	-2.7 (-9.8 to 0.4)	0.004
Hepatitis B/C infection	9 (1.6%)	3 (4.4%)	6 (1.2%)	-3.1 (-10.8 to 0.1)	0.051
Malignancy	4 (0.7%)	1 (1.5%)	3 (0.6%)	-0.8 (-7.2 to 0.8)	0.434
Immunodeficiency	1 (0.2%)	0 (0)	1 (0.2%)	0.2 (-5.1 to 1.1)	0.709
Symptoms or signs					
Fever	357 (63.3%)	58 (84.1%)	299 (60.4%)	-23.7 (-31.7 to -12.4)	< 0.001
Cough	323 (57.2%)	37 (53.6%)	286 (57.8%)	4.2 (-7.9 to 16.6)	0.513
Fatigue	87 (15.4%)	8 (11.6%)	79 (16.0%)	4.4 (-5.7 to 11.0)	0.347
Myalgia	25 (4.4%)	3 (4.4%)	22 (4.4%)	0.1 (-7.7 to 3.7)	0.971

Sputum production	51 (9.0%)	6 (8.7%)	45 (9.1%)	0.4 (-8.9 to 5.9)	0.915
Anorexia	26 (4.6%)	6 (8.7%)	20 (4.0%)	-4.7 (-13.8 to 0.5)	0.084
Diarrhea	18 (3.2%)	4 (5.8%)	14 (2.8%)	-3.0 (-11.2 to 1.0)	0.189
Shortness of breath	26 (4.6%)	7 (10.1%)	19 (3.8%)	-6.3 (-15.8 to -0.8)	0.019
Hemoglobin, g/L	131 (121 to 143)	131 (119 to 142)	132 (121 to 144)	3 (-2 to 7)	0.258
Platelet count, $\times 10^9/L$	182 (143 to 236)	163 (138 to 234)	183 (144 to 237)	8 (-10 to 25)	0.390
Blood leukocyte count, $\times 10^9/L$	4.8 (3.7 to 6.1)	5.6 (3.9 to 8.3)	4.7 (3.6 to 5.9)	-0.9 (-1.6 to -0.3)	0.004
Neutrophil count, $\times 10^9/L$	3.0 (2.2 to 4.2)	4.4 (2.7 to 6.8)	2.9 (2.2 to 3.9)	-1.1 (-1.7 to -0.6)	< 0.001
Lymphocyte count, $\times 10^9/L$	1.1 (0.8 to 1.5)	0.7 (0.5 to 1.1)	1.2 (0.9 to 1.6)	0.4 (0.3 to 0.5)	< 0.001
Alanine aminotransferase, U/L	20.3 (15.0 to 30.4)	24.7 (16.8 to 41.0)	20.0 (15.0 to 29.5)	-3.9 (-7.2 to -0.9)	0.011
Aspartate aminotransferase, U/L	24.3 (19.5 to 31.5)	35.0 (25.6 to 47.0)	23.8 (19.0 to 29.0)	-10.8 (-14.2 to -7.6)	< 0.001
Total bilirubin, $\mu\text{mol/L}$	11.9 (8.7 to 17.6)	11.8 (9.0 to 17.4)	11.9 (8.7 to 17.6)	0.1 (-1.4 to 1.7)	0.887
Albumin, g/L	39.0 (35.7 to 42.4)	35.4 (31.7 to 38.3)	39.4 (36.4 to 42.7)	4.4 (3.2 to 5.7)	< 0.001
Blood urea nitrogen, mmol/L	4.0 (3.2 to 5.0)	4.9 (3.7 to 6.2)	3.9 (3.1 to 4.8)	-1.0 (-1.4 to -0.6)	< 0.001
Creatinine, $\mu\text{mol/L}$	59.0 (46.8 to 74.0)	65.1 (53.0 to 84.0)	58.2 (46.2 to 72.9)	-8.9 (-14.3 to -3.3)	0.002
Sodium, mmol/L	137.6 (135.4 to 140.1)	135.4 (133.3 to 137.2)	138.0 (135.9 to 140.3)	2.6 (1.7 to 3.5)	< 0.001
Potassium, mmol/L	4.0 (3.6 to 4.3)	3.9 (3.5 to 4.3)	4.0 (3.6 to 4.3)	0.1 (-0.1 to 0.2)	0.493
Creatine kinase, U/L	70.8 (48.0 to 112.8)	81.7 (48.0 to 181.2)	70.0 (48.0 to 109.3)	-11.4 (-27.0 to 1.7)	0.092
Lactose dehydrogenase, U/L	189.0 (152.0 to 244.0)	304.4 (236.8 to 378.5)	183.6 (148.3 to 230.3)	-112.2 (-134.0 to -89.9)	< 0.001
D-dimer ≥ 0.05 mg/L	269 (47.7%)	44 (63.8%)	225 (45.5%)	-18.3 (-29.5 to -5.7)	0.004
Procalcitonin ≥ 0.05 ng/L	207 (36.7%)	38 (55.1%)	169 (34.1%)	-20.9 (-32.8 to -8.5)	0.001
C-reactive protein level ≥ 10 mg/L	286 (50.7%)	61 (88.4%)	225 (45.5%)	-43.0 (-50.0 to -32.3)	< 0.001
Abnormal CT findings	516 (91.5%)	68 (98.6%)	448 (90.5%)	-8.1 (-11.2 to -1.3)	0.025
Lung CT score	6 (4 to 11)	15 (10 to 21)	6 (3 to 9)	-8 (-10 to -7)	< 0.001

Abbreviations: CI, confidence interval; COPD, chronic obstructive pulmonary disease; COVID-19; 2019 coronavirus disease; CT computed tomography.

Table 2 The impact of different treatment protocols on progression to severe pneumonia after admission (n = 534)

Protocols	Rude OR (95% CI)	<i>P</i> value	Adjusted OR (95% CI)	<i>P</i> value
No antiviral therapy (n = 31)	Reference	1	Reference	1
Arbidol (n = 43)	1.18 (0.18 to 7.51)	0.864	0.79 (0.11 to 6.01)	0.823
Lopinavir/ritonavir (n = 71)	0.91 (0.16 to 5.23)	0.912	0.66 (0.10 to 4.31)	0.661
Interferon α (n = 18)	1.81 (0.23 to 14.12)	0.570	1.33 (0.15 to 12.26)	0.800
Arbidol + Lopinavir/ritonavir (n = 34)	1.55 (0.24 to 10.01)	0.643	0.95 (0.12 to 7.38)	0.958
Arbidol + Interferon α (n = 16)	0.97 (0.08 to 11.54)	0.979	0.23 (0.01 to 4.43)	0.329
Lopinavir/ritonavir + Interferon α (n = 184)	1.04 (0.22 to 4.90)	0.959	0.62 (0.12 to 3.35)	0.579
Arbidol + Lopinavir/ritonavir + Interferon α (n = 137)	1.45 (0.31 to 6.84)	0.639	0.85 (0.16 to 4.60)	0.847

Abbreviations: CI, confidence interval; OR, odds ratio.

Adjusted OR was calculated by adjusting age, gender, smoking history, hypertension, diabetes, cardiovascular disease, chronic obstructive pulmonary disease, and lung CT score.

Table 3 Comparison between hypertensive patients on ACEI/ARB therapy and on Non-ACEI/ARB therapy (n = 65)

Characteristics	ACEI/ARB (n = 16)	Non-ACEI/ARB (n = 49)	Difference (95% CI)	P
Age (years)	57 (51 to 65)	63 (53 to 69)	5 (-1 to 12)	0.106
Male gender	10 (62.5%)	23 (46.9%)	15.6 (-11.9 to 38.7)	0.280
Smoking history	2 (12.5%)	5 (10.2%)	2.3 (-12.4 to 26.5)	0.797
Comorbidity				
Diabetes	2 (12.5%)	18 (36.7%)	24.2 (-2.2 to 40.9)	0.068
Cardiovascular disease	0 (0)	8 (16.3%)	16.3 (-4.6 to 29.0)	0.084
COPD	0 (0)	1 (2.0%)	2.0 (-17.4 to 10.7)	0.565
Cerebrovascular disease	0 (0)	3 (6.1%)	6.1 (-13.7 to 16.5)	0.311
Chronic renal disease	1 (6.3%)	1 (2.0%)	-4.2 (-26.4 to 5.9)	0.435
Hepatitis B infection	0 (0)	2 (4.1%)	4.1 (-15.5 to 13.7)	0.412
Medication history				
Aspirin	0 (0)	3 (6.1%)	6.1 (-13.7 to 16.5)	0.311
Statins	1 (6.3%)	2 (4.1%)	-2.2 (-24.4 to 8.8)	0.720
Metformin	1 (6.3%)	8 (16.3%)	10.1 (-13.3 to 23.8)	0.311
Other oral hypoglycemic agents	1 (6.3%)	10 (20.4%)	14.2 (-9.7 to 28.4)	0.190
Insulin	0 (0)	2 (4.1%)	4.1 (-15.5 to 13.7)	0.412
Systolic pressure, mmHg	132 (120 to 137)	136 (120 to 146)	7 (-2 to 15)	0.107
Diastolic pressure, mmHg	80 (71 to 89)	80 (78 to 87)	2 (-3 to 8)	0.547
Hemoglobin, g/L	133 (125 to 150)	133 (118 to 138)	-7 (-17 to 1)	0.060
Platelet count, $\times 10^9/L$	181 (135 to 241)	182 (147 to 239)	6 (-38 to 52)	0.837
Blood leukocyte count, $\times 10^9/L$	5.5 (4.6 to 6.5)	5.2 (3.9 to 7.0)	-0.2 (-1.4 to 0.9)	0.698
Neutrophil count, $\times 10^9/L$	3.7 (3.1 to 4.4)	3.7 (2.4 to 4.8)	-0.1 (-1.0 to 0.8)	0.733
Lymphocyte count, $\times 10^9/L$	1.1 (0.9 to 1.4)	1.0 (0.7 to 1.5)	-0.1 (-0.4 to 0.2)	0.322
Alanine aminotransferase, U/L	28.9 (16.2 to 40.7)	23.0 (17.2 to 31.5)	-2.3 (-12.6 to 5.3)	0.590
Aspartate aminotransferase, U/L	25.0 (20.0 to 30.9)	26.3 (20.0 to 36.3)	0.8 (-4.6 to 7.2)	0.800
Total bilirubin, $\mu\text{mol/L}$	12.6 (10.3 to 19.9)	12.5 (9.5 to 19.0)	-0.7 (-4.0 to 3.0)	0.634
Albumin, g/L	38.4 (34.6 to 44.5)	37.0 (32.2 to 40.4)	-2.5 (-6.2 to 1.4)	0.185
Blood urea nitrogen, mmol/L	4.8 (4.1 to 5.8)	4.5 (3.7 to 6.5)	-0.2 (-1.2 to 0.9)	0.611
Creatinine, $\mu\text{mol/L}$	64.7 (44.9 to 82.1)	63.3 (51.2 to 78.7)	1.3 (-14.5 to 14.7)	0.877
Sodium, mmol/L	136.3 (134.7 to 137.2)	137.5 (134.6 to 140.6)	1.3 (-0.8 to 3.4)	0.252
Potassium, mmol/L	4.0 (3.6 to 4.4)	3.9 (3.5 to 4.2)	-0.1 (-0.4 to 0.2)	0.438
Creatine kinase, U/L	118.5 (64.6 to 187.3)	64.5 (46.5 to 147.2)	-28.4 (-70.0 to 5.1)	0.083
Lactose dehydrogenase, U/L	184.0 (157.1 to 281.7)	238.0 (164.1 to 315.2)	33.4 (-19.7 to 89.2)	0.301
D-dimer ≥ 0.05 mg/L	10 (62.5%)	27 (55.1%)	-7.4 (-30.9 to 19.8)	0.604
Procalcitonin ≥ 0.05 ng/L	6 (37.5%)	27 (55.1%)	17.6 (-10.0 to 40.7)	0.221
C-reactive protein level ≥ 10 mg/L	11 (68.8%)	41 (83.7%)	14.9 (-6.4 to 40.5)	0.195
Abnormal CT findings	15 (93.8%)	46 (93.9%)	0.1 (-11.5 to 22.6)	0.985
CT severity score	7 (3 to 14)	10 (5 to 16)	2 (-3 to 8)	0.258
Combined treatments after admission				
Nonspecific antiviral therapy	15 (93.8%)	45 (91.8%)	-1.9 (-14.1 to 20.7)	0.803
Interferon α	8 (50.0%)	35 (71.4%)	21.4 (-4.6 to 45.9)	0.116
Intravenous antibiotics	12 (75.0%)	34 (69.4%)	-5.6 (-25.9 to 21.3)	0.668

Severe pneumonia	1 (6.3%)	16 (32.7%)	26.4 (1.5 to 41.3)	0.037
------------------	----------	------------	--------------------	--------------

Abbreviations: ACEI, angiotensin-converting enzyme inhibitors; ARB, angiotensin-receptor blockers; CI, confidence interval; COPD, chronic obstructive pulmonary disease; CT computed tomography.

Table 4 Clinical risk factors for severe pneumonia in COVID-19 patients

Variables	Univariate OR (95% CI)	<i>P</i> value	Multivariate OR (95% CI)	<i>P</i> value
Age (per year increase)	1.06 (1.04 to 1.08)	< 0.001	1.05 (1.03 to 1.07)	< 0.001
Male gender	1.33 (0.80 to 2.20)	0.275		
Hypertension		< 0.001		0.055
Non-hypertensive	Reference	1	Reference	1
Hypertensive on ACEI/ARB therapy	0.60 (0.08 to 4.66)	0.628	0.41 (0.05 to 3.19)	0.392
Hypertensive on other therapy	3.93 (2.15 to 7.19)	< 0.001	2.07 (1.07 to 4.00)	0.030
Diabetes	2.95 (1.44 to 6.03)	0.003		0.293
Cardiovascular disease	2.85 (1.08 to 7.56)	0.035		0.916

Abbreviations: ACEI, angiotensin-converting enzyme inhibitors; ARB, angiotensin-receptor blockers; CI, confidence interval; COVID-19; 2019 coronavirus disease; OR, odds ratio.