1
2 The proportion of deaths cases in confirmed patients of COVID-19 is still
3 increasing for cumulative cases reported up to 25 April 2020

4
5 Thomas Scheuerl¹

6
7 ¹Research Institute for Limnology, Mondsee, University of Innsbruck, Mondsee, Austria.
8 thomas.scheuerl@uibk.ac.at ORCID ID: <https://orcid.org/0000-0001-5216-5630>

9
10 **Abstract**

11 In this work I analyse how proportions of fatal cases after COVID-19 infection change since
12 outbreak of the disease. Using publicly available data I model the change in deaths probability
13 from day 30 of outbreak until 25 April 2020. The global trend is that the proportion of fatal cases is
14 still increasing and that many countries have not yet reached the maximum deaths proportion.
15 However, there are visual differences between countries and in some countries the proportions are
16 clearly below or above the global trend. A positive correlation between deaths cases and recorded
17 infections indicates that a higher infection number results in increased mortality numbers.

19 Introduction

20 The ongoing coronavirus (COVID-19) outbreak starting in China [1] expanded rapidly over the
 21 globe and many countries respond to the spread of the disease by a lock-down to limit contacts
 22 between people. Many countries are heavily affected by the outbreak and there is a high deaths rate
 23 [2]. The deaths probability will be an important parameter guiding how long lock-down and social
 24 distancing are necessary. Countries economies are under pressure and for older adults, who are the
 25 most vulnerable group, social distancing is predicted to come with serious negative effects [3], but
 26 obviously spread of the disease should be condemned as much as possible until the health system
 27 can cope with the disease. Here, I analyse the death proportion dynamics of COVID-19 infections
 28 since outbreak of the disease. I focus on the change of deaths probability over confirmed cases of a
 29 particular day and if this proportion changes over time. This indicates the probability of a fatal case
 30 for the current number of cases and reflects whether further increase can be expected, or if
 31 proportions have reached the maximum. The analysis reveals that at the current number of
 32 confirmed cases, the proportion of people dying is still increasing in most countries, indicating the
 33 situation may further worsen.

34

35 Methods

36 I analyse freely available data on the COVID-19 pandemic situation from Hopkins University
 37 <https://coronavirus.jhu.edu/data/mortality>. Data were downloaded from [https://github.com/datasets/](https://github.com/datasets/covid-19/tree/master/data)
 38 [covid-19/tree/master/data](https://github.com/datasets/covid-19/tree/master/data). The data frame contains 185 countries and collects *confirmed cases*,
 39 *recovered cases* and *deaths cases* each day starting from 22nd of January 2020. I limit the analysis to
 40 countries which have at least 1000 confirmed cases and countries that have at least 1 recovered and
 41 1 fatal case. Because the disease was spreading in China several days before it arrived in other
 42 countries, I also removed the first 30 days, during which deaths cases were limited to China.

43 Data were analysed in R statistical environment [4]. A time-series analysis was conducted to track
 44 changes over time using generalized additive mixed effects models (GAMMs) following ref. [5].
 45 This approach smoothes changes over time tracking non-linear trends. Change in proportion and
 46 count data were modelled using 'Days' as continuous variable building a global model for all
 47 countries applying a smooth term with family either *binomial* for proportion or *poisson* for count
 48 data. The change over time per country was introduced into the random effect. The models were
 49 build using the R packages *nlme* [6] and *mgcv* [7]. The change over time was smoothed using 10
 50 knots and a *cubic regression* spline. Model convergence was controlled using the *lmeControl*
 51 function, with the maximum number of iterations set to 500 and using optimisation *opt=optim*
 52 function. Temporal autocorrelation was introduced to account for violation of independence.
 53 Choosing based on the lowest AIC criterion, the best correlation structure was selected from a range
 54 of possibilities presented elsewhere [5]. All final models included a *corAR1* correlation structure
 55 including time and country to address temporal correlation. Data were *log10* transformed to
 56 improve model fit, inspected by validation plotting. To explore the correlation of deaths proportions
 57 over confirmed cases generalized least square models were used including the *varExp* variance
 58 structure to allow changing spread of residuals over changes in confirmed cases. Visualization was
 59 also done in R.

60

61 Results

62 There are >70 countries with more than 1000 confirmed cases and at least one 1 deaths and 1
 63 recovered case.

64 The global trend for new cases is still increasing but slightly slowing down (Supporting Fig. 1). The
 65 smoothed term across time for all countries indicates an increase with little decrease. Only in China
 66 and South Korea no new cases are confirmed. In all other countries the trend is still raising,
 67 however not exponential any more. Most cases are confirmed in the US, followed by Italy, Spain

68 and Germany, but in all countries the increase of new cases is slowing down. Japan is one of the
69 countries following the global trend. Thailand is one of the countries with an increase less than the
70 global trend in new cases.

71 There also is a positive trend for recovered cases. The smoothed term across all countries however
72 indicates that this trend is starting to slow down. Not surprisingly, those countries with highest
73 confirmed cases have most recovered cases.

74 The same seems true for deaths cases. In all countries, except China, the numbers of deaths cases
75 are increasing daily (Supporting Fig. 3).

76 I also looked into a measure like “Open cases” by subtracting daily deaths cases and daily recovered
77 cases from the total number confirmed (Supplemental Fig. 4). Using this kind of measure, global
78 open cases are still increasing as indicated by the global smooth term. The countries China, South-
79 Korea and Thailand seem to record a reduction of open cases, while in all other countries more open
80 cases are recorded daily.

81 Up to this point there is little new information more than in most countries more and more cases are
82 recorded. Next I was looking into how the temporal proportions change and if the likelihood to
83 either die or recover is changing since outbreak of the disease. I start on looking at the proportion of
84 deaths per confirmed cases. This measure may represent how likely deaths occurs and if the
85 maximum is reached already. The global trend is still positive, indicating the proportion of deaths
86 cases is still going to increase world wide based on total cases (Fig. 1). However, there is a decline
87 indicated and there is no further significant increase the last few days.

88

89

90

92

93 **Figure 1.** The proportion of fatal cases after COVID-19 infection since outbreak. The proportion is shown on a log
 94 scale. The global trend (black line) indicates an increasing trend (significant trend in red) with little reduction in daily
 95 new cases. Temporal tracks of individual countries are indicated by different colours as indicated in the legend.

96

97 In countries like Italy, Spain and the United Kingdom, the deaths proportions are particularly high,
 98 but now seem to level off and stabilise. In China this measure doesn't change because there are no
 99 new cases of infections, neither deaths cases recorded (Supporting Figures 1+3), which causes no

more temporal change. This indicates the maximum proportion was reached. Interestingly, in the country with most confirmed cases, the United States, the deaths proportions are lower than could be expected. Germany is reaching the global trend now, despite it has more than average confirmed cases. In Japan the deaths proportions are declining, which may be caused by the fact that there are more daily new infections than on average, but the deaths rates are increasing less than the global trend.

One likely explanation for the observed patterns of increased deaths proportions over time after the pandemic outbreak may be that more cases simply result in even more deaths cases because the countries health system experiences an overload. Because of this I looked into correlations between confirmed cases and deaths proportion, checking for positive trends. Indeed there seems to be a positive trend that more confirmed cases result in a higher probability of fatal recordings (Fig. 2).

Fig.2. The correlation between deaths proportions and number of confirmed cases for four different days within the last four weeks. The red line indicates the best fit line of a generalized least square model including a variance structure allowing for larger residual spread for low number of confirmed cases. Individual slopes are indicated per plot.

126

127 With more infected cases the probability to die from COVID-19 infections seems to increase. This
128 is true for 4 different days within the last month. In line with the global analysis the correlations are
129 all positive but slowly decrease over time.

130 The proportions of recovered cases is also increasing however this trend seems also to be slowing
131 down (Supporting Figure 5). Countries like the US, UK and Sweden seem to be below the global
132 trend.

133

134 **Discussion**

135 Coronavirus causes severe respiratory symptoms resulting in many fatal cases. The number of
136 infections is still increasing in many countries and so do the proportions of fatal cases. Giving hope,
137 there is a declining global trend that the proportions start to level off. However, there are clear
138 differences between countries and individual countries will have to follow individual actions to
139 condemn the spread of the disease. While in general it looks like more infections cause more fatal
140 cases, this is not true for all countries. Particularly in the US the deaths proportions are less
141 increasing than might be expected. This may be either that the healths system can still cope with the
142 cases, or confirmed cases are accumulating faster than fatal cases rise.

143 The deaths over confirmed analysis is a bit inaccurate as when there are no records neither in new
144 confirmed cases nor new deaths cases, like in China, there is no decline in the proportions. It only
145 tells if the maximum is yet reached. In case of China the measure indicates the country has reached
146 the maximum level in deaths probability. Another measure could be to calculate “Deaths cases”
147 over “Open cases”. This would require to calculate daily new cases of deaths only. This is a
148 promising future step and certainly worth following up. In this work I focus on whether countries
149 have already reached the maximum deaths proportion and if this trend seems stabilizing over time.

150

151

152 **Data Availability Statement:** All relevant data are within the paper and its Supporting
153 Information files.

154

155 **Competing interests:** The author declares no competing interests.

156

157 **Funding:** There is no funding body associated.

158

159

160 **References**

1. Chen S, Yang J, Yang W, Wang C, Bärnighausen T. 2020 COVID-19 control in China during mass population movements at New Year. *The Lancet* **395**, 764–766. (doi:10.1016/S0140-6736(20)30421-9)
2. Maitra S, Biswas M, Bhattacharjee S. 2020 Case- fatality rate in COVID- 19 patients: A meta-analysis of publicly accessible database. *medRxiv* , 2020.04.09.20059683. (doi:10.1101/2020.04.09.20059683)
3. Jawaid A. 2020 Protecting older adults during social distancing. *Science* **368**, 145. (doi:10.1126/science.abb7885)
4. R Development Core Team. 2010 The R project for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. <http://www.r-project.org/>. *Vienna, Austria: R Foundation for Statistical Computing*. See <http://www.r-project.org/> (accessed on 25 July 2012).
5. Zuur AF, Ieno EN, Walker N, Saveliev AA, Smith GM. 2009 *Mixed effects models and extensions in ecology with R*. 1st Edition. Springer, New York.
6. Pinheiro J, Bates D, DebRoy S, Sarkar D. 2012 Linear and nonlinear mixed effects models CRAN - Package nlme. <http://cran.r-project.org/web/packages/nlme/index.html>. See <http://cran.r-project.org/web/packages/nlme/index.html> (accessed on 12 September 2012).
7. Woods S. 2012 GAMs with GCV/AIC/REML smoothness estimation and GAMMs by PQL. R package version 1.7-9. <http://cran.r-project.org/web/packages/mgcv/index.html>. See <http://cran.r-project.org/web/packages/mgcv/index.html> (accessed on 12 September 2012).

161

162

163 Supplementary Figures

165

166 Supplementary Figure 1. Confirmed new cases of COVID-19 since outbreak. The confirmed
 167 number is shown on a log scale. The global trend (red line) indicates an increasing trend with little
 168 reduction in daily new cases. Temporal tracks of individual countries are indicated by different
 169 colours as indicated in the legend.

170

172

173 Supplementary Figure 2. Recovered cases since outbreak. The recovered cases are shown on a log
 174 scale. The global trend (red line) indicates an increasing trend with little reduction in daily new
 175 cases. Temporal tracks of individual countries are indicated by different colours as indicated in the
 176 legend.

177

179

180 Supporting Figure 3. Deaths cases since outbreak. The dearths cases are shown on a log scale. The
 181 global trend (red line) indicates an increasing trend with little reduction in daily new cases.
 182 Temporal tracks of individual countries are indicated by different colours as indicated in the legend.

183

184

186

187 Supporting Figure 4. Open cases since outbreak. The open cases are shown on a log scale. The

188 global trend (red line) indicates an increasing trend with little reduction in daily new cases.

189 Temporal tracks of individual countries are indicated by different colours as indicated in the legend.

190

191

193
 194 Supporting Figure 5. Proportion of recovered cases since outbreak. The proportions of recovered
 195 cases are shown on a log scale. The global trend (red line) indicates an increasing trend with little
 196 reduction in daily new cases. Temporal tracks of individual countries are indicated by different
 197 colours as indicated in the legend.

198

199