1 **Title: Prediction of the Epidemic Peak of Covid19 in Egypt, 2020**

2 **Author:** Eman D.El Desouky

3 Epidemiology and Biostatistics, National Cancer Institute, Cairo University,
4 eman.desouky@nci.cu.edu.eg

5

6 **Address:** Kasr Al Eini ‘Street ‘Fom El Khalig ‘Cairo Governorate,

7 **Postal code:** 11796

1 Prediction of the Epidemic Peak of Covid19 in Egypt, 2020

2 Abstract

3 **Objectives:** Since December 2019 a pandemic of new novel coronavirus has started from
4 Wuhan, China, in Egypt, the first case reported on February 14, 2020. In this study we
5 aimed to predict the time of possible peak and simulate the changes could be happen by
6 the social behavior of Egyptians during Ramadan (the holy month).

7
8 **Methods:** SIR and SEIR compartmental models were used to predict the peak time. We
9 simulated different expected scenarios based to examine their effects on the peak timing.

10 **Results:** We found that the peak most likely to be in middle of June 2020. Simulating
11 different transmission rate probability and R0 the earliest peak could to be in the May 20
12 and latest one could be in 18 July. The peak shifted much earlier to 11th April 2020
13 without lockdown and other mitigation strategies.

14 **Conclusion:** Social behaviors of citizens during the holy month will dramatically affect
15 the peak timing. Mitigations strategies and other lockdown measure helped to delay the
16 expected peak.

17 **Keywords:** COVID-19; compartmental models; Epidemic peak; Egypt

18

1 **Introduction**

2 The first case of respiratory disease caused by a novel coronavirus was identified in
3 Wuhan City, Hubei Province, China in In December 2019 and on 13 January 2020 the
4 first case outside of China was reported in Thailand.¹ On 11th March 2020 the World
5 Health Organization named it coronavirus disease 2019 (COVID-19) and considered as
6 pandemic.² The first case appeared in Egypt on 14th February 2020 for a foreigner and
7 fortunately was asymptomatic until 1st of March 2020 he was the only reported case after
8 that few cases started to appear till 14th March 2020 to reach 110 cumulatively, and then
9 the number of laboratory confirmed cases increased weekly (Table 1) to reach 3490 till
10 April 22, 2020.³

11 **Table 1. Number of cumulative COVID-19 cases in Egypt until 23th April 2020**

...	No of Cumulative cases
14 February -20 February	1
21 February -27 February	1
28 February -5 March	1
6 March -12 March	3
13 March -19 March	80
20 March -26 March	256
27 March -2 April	495
3 April - 9 April	865
10 April-16 April	1699
17 April-22 April	3490

12
13
14 As seen in table 1 the numbers started nearly to double weekly from the fifth week and
15 after closure of schools with partial lockdown the public concerned about the timing of
16 the cases surge. The aim of this study is to predict the time of possible peak and simulate
17 the changes that might happen by the social behavior of Egyptians during Ramadan (the
18 holy month) that will begins on Friday April 24 and ends on Saturday May 23. In
19 Ramadan families usually gather especially in first days , children have more activates
20 with each other's and everyone leaves work at the same time and need to be in home at
21 the same time ; currently with Covid19 pandemic and partial lock down it's challenging
22 to cope with social distancing and Ramadan habits .

23
24

1 **Methods:**

2 We tested two models SIR and SEIR models ⁴ to predict peak timing using data till 17th
3 April 2020 and simulated different scenarios based on hypothetical expected changes in
4 social behavior of Egyptians during Ramadan .

5 **1- SIR model**

9 **2- SEIR model**

14
15 Where $S(t)$, $E(t)$, $I(t)$ and $R(t)$ represent the susceptible, exposed, infective and removed
16 populations at time t , respectively. β , ε and γ represent the infection rate, the onset rate
17 and the removal rate, respectively. $1/\varepsilon$ and $1/\gamma$ imply the average incubation period and
18 the average infectious period, respectively. The force of infection (rate of infection per
19 unit time is βI and β denoted the rate of transmission from S to I or effective contact
20 between S and I .⁵

21 These models based on Euler`s Method: A numerical method for solving differential
22 equations and referred to as a compartment model since it is useful to refer to people
23 moving from one compartment to another.

24 Parameters used in the models based on previous studies assuming average Incubation
25 period = 5 days and so onset rate $\varepsilon = 0.2$ ⁶, assuming infectious period = 10 day and so
26 removal rate $\gamma = 0.1$.⁷ Egypt population (1×10^8) ⁸, no one infected at the time (t_0) time step
27 is one day and population remain constant so $S+E+I+R=1$.

Regarding transmission rate (β) and Basic reproduction number (R_0) researcher depended on the published data available on GitHub, midas-network⁹, R_0 for Egypt as published 2.3 and so $\beta=0.23$ by applying the formula;

$$R_0 = \frac{\beta}{\gamma}$$

For the purpose of simulation different scenario of effective contacts during Ramadan, different R_0 and β were used within the range of previous published data available on midas-network.⁹

Starting from model1 (M1) that represent less effective contact rate than the current situation ($\beta=0.22$, $R_0=2.2$) till model10 (M10) that represent higher effective contact rate with $\beta=0.32$ and $R_0=3.2$

The peak was identified by the maximum proportion of population infected in a year

12

13 **Results**

14

15 Researcher used SIR model with different simulation scenarios to predict peak timing, 16 illustrated in figure1. The earliest of peak could be on May 20, 2020 and latest peak could 17 be July 18, 2020, then applied Applying regression method to estimate the best fitted 18 models with the cumulative data released by MOHP till 17th April, 2020; peak on 15 19 June 2020 is more likely.

20

21 **Figure (1): Different models estimating different peaks with SIR model**

22

1 Researcher used SEIR Peak prediction models almost similar results obtained; earliest
2 peak could be May 17, 2020 and latest peak could be by July 15, 2020.

3
4 The 3rd scenario was estimating the peak time without lockdown and other strategies
5 based on mobility changes report released by google¹¹ that stated average 50% reduction
6 in Egypt since the time of lockdown and mitigation strategies. The peak was estimated to
7 occur on 11th April 2020 but with mitigation strategies and lockdown it was shifted about
8 69 days to reach June 15, 2020 (figure 2).

9

10

11

12 **Figure (2): Peak estimation with lockdown and other measures in comparison to the**
13 **situation without them.**

14

15 **Discussion**

16 In this study, by applying the SIR and SEIR compartmental model along with the daily
17 reported cases of Covid19 till 17th April in Egypt; we estimated the peak could be in the
18 middle of June. We hypothesized that the social and contact rate per individual changes
19 in Ramadan (the Holy month) that will start on April 24, 2020 will affect the timing of
20 peak as a result of Increase in β (transmission rate). We simulated different situations
21 with different rates of effective contact, transmission rate and R_0 consequently; the peak
22 timing changed to reach 20 May, 2020 as earliest one and latest peak could be July 18,
23 2020. These findings was in concordance with the study done by Pearsn et al¹² that

reported 10,000 cases could be in 20 May 2020. On the contrast a recent report released by **SUTD Data-Driven Innovation Lab**¹³ that reported the pandemic will end in Egypt on May 20, 2020. Researcher simulated another scenario based on current data to predict the timing of peak without lockdown the peak shifted earlier to be on April 11, 2020; the mitigation strategies helped to delay peak. The current study simulated different scenarios based on different values on transmission rate (calculated and estimated) in concordance with **Bootsma and Ferguson**¹⁴ who stated the most important factor affecting disease pandemic is the transmission rate which depend on crowding as it affects amount of contact per individual. The current study used SIR and SEIR models which are too idealistic for modeling COVID-19 through numerical simulation, but it gives an idea about when the COVID-19 cases can surge to.¹⁵

The study stated long range peak prediction to emphasis the role of social distancing to flatten the curve in concordance with **Anderson et al**¹⁶.

The result suggests that the epidemic of COVID-19 in Egypt would not end so quickly. This might be reliable with the WHO's statement on 6 March 2020 that it is a false hope that COVID-19 will disappear in the summer like the flu.¹⁷

Limitations

This is too idealistic for modeling COVID-19 as the reproduction number (R_0) changes overtime but the multiple simulation may overcome this hypothetically for certain time. The deterministic models are not accurate and used to give insight about what will happen to population on average. Study didn't estimate the hospitalized number at the surge however the timing of the peak didn't affected by the identified cases form the actual infective population.

Conclusions:

During COVID-19 pandemic it's hard to predict the real peak timing however the surge timing will depend mainly on the behavior of citizens towards social distancing and hygiene measures. The mitigation strategies and lockdown in Egypt has a positive effect on the delay of the epidemic peak, giving more time to the health sector to encompass the situation. There is an urgent need for Local and worldwide policy to deal with Covid19 that will extend to summer time. Egyptian government should monitor the reported cases daily along with the performance of citizens in the coming month to determine the proper

strategies to flatten the curve as much as possible. The optimistic view to the situation should be treated cautiously as COVID-19 is still unclear.

Declaration of interests: The author declares no competing interests.

Reference

- 1 CDC, Coronavirus Disease 2019 (COVID-19).
2 <https://www.cdc.gov/coronavirus/2019-ncov/index.html>. Accessed 20 Apr 2020
- 3 WHO, Coronavirus disease 2019.
4 <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>. Accessed 20 Apr
5 2020
- 6 Egypt cares. 2020. Available from:
7 <https://www.care.gov.eg/EgyptCare/Index.aspx> Accessed 22 Apr 2020
- 8 Inaba, H. Age-Structured Population Dynamics in Demography and Epidemiology;
9 Springer: Berlin/Heidelberg, Germany, 2017.
- 10 Abbey H. An examination of the Reed-Frost theory of epidemics. Human biology.
11 1952 Sep 1;24(3):201.
- 12 Linton NM, Kobayashi T, Yang Y, Hayashi K, Akhmetzhanov AR, Jung SM,
13 Yuan B, Kinoshita R, Nishiura H. Incubation period and other epidemiological
14 characteristics of 2019 novel coronavirus infections with right truncation: a statistical
15 analysis of publicly available case data. Journal of clinical medicine. 2020 Feb;9(2):538.
- 16 Sun H, Qiu Y, Yan H, Huang Y, Zhu Y, Chen SX. Tracking and Predicting
17 COVID-19 Epidemic in China Mainland. medRxiv. 2020 Jan 1.
- 18 CAPMAS. <https://www.capmas.gov.eg/Pages/populationClock.aspx>. Accessed 22
19 Apr 2020
- 20 COVID-19/README.md at master · midas-network/COVID-19.
21 https://github.com/midas-network/COVID-19/blob/master/parameter_estimates/2019_novel_coronavirus/README.md. Accessed
22 17 Apr 2020
- 23 Ridenhour B, Kowalik JM, Shay DK. Unraveling R 0: Considerations for Public
24 Health Applications. American journal of public health. 2018 Dec;108(S6):S445-54.

1 11 COVID-19 Community Mobility Report.
2 https://www.gstatic.com/covid19/mobility/2020-04-11_EG_Mobility_Report_en.pdf.
3 Accessed 20 Apr 2020
4
5
6 12 Pearson CA, Van Schalkwyk C, Foss AM, O'Reilly KM, Pulliam JR,. Projected
7 early spread of COVID-19 in Africa. CMMID COVID-19 working group 2020. 63:25–28

8 13 SUTD Data-Driven Innovation Lab. “Egypt.” *Data-Driven Prediction of COVID-*
9 *19 Pandemic End Dates*, ddi.sutd.edu.sg/portfolio/items/442705. Accessed 23 Apr 2020

10 14 Bootsma MC, Ferguson NM. The effect of public health measures on the 1918
11 influenza pandemic in US cities. Proceedings of the National Academy of Sciences. 2007
12 May 1;**104**(18):7588-93.
13

14 15 Wees J Van, Osinga S, Kuip M Van Der, Tanck M, Hanegraaf M, Pluymaekers
15 M, et al. Forecasting hospitalization and ICU rates of the COVID-19 outbreak: an
16 efficient SEIR model. [Submitted]. Bulletin World Health Organization. E-pub: 30
17 March 2020. doi: <http://dx.doi.org/10.2471/BLT.20.256743>
18

19 16 Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will
20 country-based mitigation measures influence the course of the COVID-19 epidemic? The
21 Lancet. 2020 Mar 21;395(10228):931-4.

22 17 “False hope” coronavirus will disappear in summer like flu, WHO says. Available
23 from: [https://www.cnn.com/2020/03/06/its-a-false-hope-coronavirus-will-disappear-in-](https://www.cnn.com/2020/03/06/its-a-false-hope-coronavirus-will-disappear-in-the-summer-like-the-flu-who-says.html)
24 [the-summer-like-the-flu-who-says.html](https://www.cnn.com/2020/03/06/its-a-false-hope-coronavirus-will-disappear-in-the-summer-like-the-flu-who-says.html). Accessed 18 Apr 2020.
25