

1 **Comprehensive Testing Highlights Racial, Ethnic, and Age Disparities in the COVID-19**
2 **Outbreak: Epidemiological and Clinical Characteristic of Cases in Utah**

3 Sharia M Ahmed¹ PhD, Rashmee U. Shah² MD, Margaret Bale³ MPH, Jordan B. Peacock³ BS,
4 Ben Berger³ BA, Alyssa Brown³ BS, Sara Mann³ BS, William West³ BS, Valerie Martin³ BS,
5 Valerie Fernandez⁴, Sara Grineski^{5,6} PhD, Benjamin Brintz^{1,7} PhD, Matthew H. Samore^{7,8} MD,
6 Matthew J. Ferrari⁹ PhD, Daniel T. Leung^{1†} MD, Lindsay T. Keegan^{7†*} PhD

7
8 ¹University of Utah School of Medicine, Department of Internal Medicine, Division of Infectious
9 Diseases, Salt Lake City UT

10 ²University of Utah School of Medicine, Department of Internal Medicine, Division of
11 Cardiovascular Medicine, Salt Lake City UT

12 ³University of Utah School of Medicine, Salt Lake City UT

13 ⁴University of Utah, Salt Lake City, UT

14 ⁵Center for Natural & Technological Hazards, University of Utah, Salt Lake City, UT

15 ⁶Department of Sociology, University of Utah, Salt Lake City, UT

16 ⁷University of Utah School of Medicine, Department of Internal Medicine, Division of
17 Epidemiology, Salt Lake City UT

18 ⁸Department of Veterans Affairs Salt Lake City Health Care System, Salt Lake City, Utah, USA

19 ⁹The Pennsylvania State University, Department of Biology, State College, PA

20
21 †These authors contributed equally

22 *Corresponding author:

23 Lindsay Keegan

24 lindsay.keegan@utah.edu

25

26

27 **Abstract**

28 The United States (US), which is currently the epicenter for the COVID-19 pandemic, is
29 a country whose demographic composition differs from that of other highly-impacted countries.
30 US-based descriptions of SARS-CoV-2 infections have, for the most part, focused on patient
31 populations with severe disease, captured in areas with limited testing capacity. The objective of
32 this study is to compare characteristics of positive and negative SARS-CoV-2 patients, in a
33 population primarily comprised of mild and moderate infections, identified from comprehensive
34 population-level testing. Here, we extracted demographics, comorbidities, and vital signs from
35 20,088 patients who were tested for SARS-CoV-2 at University of Utah Health clinics, in Salt
36 Lake County, Utah; and for a subset of tested patients, we performed manual chart review to
37 examine symptoms and exposure risks. To determine risk factors for testing positive, we used
38 logistic regression to calculate the odds of testing positive, adjusting for symptoms and prior
39 exposure. Of the 20,088 individuals, 1,229 (6.1%) tested positive for SARS-CoV-2. We found
40 that Non-White persons were more likely to test positive compared to non-Hispanic Whites
41 (adjOR=1.1, 95% CI: 0.8, 1.6), and that this increased risk is more pronounced among Hispanic
42 or Latino persons (adjOR=2.0, 95%CI: 1.3, 3.1). However, we did not find differences in the
43 duration of symptoms nor type of symptom presentation between non-Hispanic White and non-
44 White individuals. We found that risk of hospitalization increases with age (adjOR=6.9 95% CI:
45 2.1, 22.5 for age 60+ compared to 0-19), and additionally show that younger individuals (aged 0-
46 19), were underrepresented both in overall rates of testing as well as rates of testing positive. We
47 did not find major race/ethnic differences in hospitalization rates. In this analysis of
48 predominantly non-hospitalized individuals tested for SARS-CoV-2, enabled by expansive
49 testing capacity, we found disparities in both testing and SARS-CoV-2 infection status by
50 race/ethnicity and by age. Further work on addressing racial and ethnic disparities, particularly
51 among Hispanic/Latino communities (where SARS-CoV-2 may be spreading more rapidly due
52 to increased exposure and comparatively reduced testing), will be needed to effectively combat
53 COVID-19 in the US.

54

55 **Introduction**

56 Since its emergence in humans in late 2019, coronavirus disease 2019 (COVID-19) (caused by
57 severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection), has caused a global
58 pandemic with over 3 million infections and 230,000 deaths. Currently, the United States (US) is
59 the epicenter with approximately 30% of all global cases¹. Although the majority of case- and
60 disease-course descriptions have originated outside of the US²⁻⁵, emerging data suggests
61 symptoms, risk from comorbidities, as well as age and racial patterns of COVID-19 may differ in
62 the US compared to other highly-impacted countries such as China and Italy. The majority of
63 US-based clinical or epidemiological descriptions of COVID-19 have been among hospitalized
64 and ICU patients, due to limited testing capacity in much of the US⁶⁻¹¹. This bias in study
65 population is potentially masking differences between hospitalized and non-hospitalized SARS-
66 CoV-2 infections. Furthermore, due to the locations of most outbreaks, most US-based studies
67 have focused on highly urban, densely populated areas with high levels of SARS-CoV-2
68 infection and relatively low testing capacity¹⁰⁻¹³.

69

70 Widespread lack of SARS-CoV-2 testing capacity in the US has led to delays in case detection
71 and isolation, significant under-reporting of cases (particularly those with mild and moderate

72 symptoms), and gross underestimates of the true number of fatalities¹³⁻¹⁸. In contrast, Utah is
73 one of only two states (as well as Tennessee) that has maintained high per capita testing rates
74 (testing 188 per 100,000 persons per day) as well as a stable proportion of positive tests
75 (approximately 5.5% of all tests were positive since 12 March) (Figure 1)¹⁹. As such, testing in
76 Utah has not been restricted to the most critical patients, and the majority of SARS-CoV-2 tests
77 are administered in outpatient settings to community-dwelling, social distancing persons.
78

79 The objective of this study is to compare demographic characteristics of positive and negative
80 patients in a population for which testing is more readily available than previously published
81 cohorts. In this article, we present the clinical, epidemiological, and demographic characteristics
82 of all persons (hospitalized and community-dwelling) tested for SARS-CoV-2 infection at
83 University of Utah Health (UHealth) from 10 March to 24 April 2020. UHealth primarily serves
84 Salt Lake County, a diverse, medium-density, medium-sized metropolitan area of just over 1
85 million people. Due to the expanded testing capacity for SARS-CoV-2 in Utah, these data
86 provide an alternative picture of the full spectrum of SARS-CoV-2 epidemiology relative to
87 studies of hospitalized patients.
88

89 **Methods**

90 All patients tested for SARS-CoV-2 in the UHealth system were eligible for our study. We
91 developed an electronic, near real-time registry of all patients tested for SARS-CoV-2 at UHealth
92 prior to 30 April 2020, for a total sample size of 20,088, representing approximately 20% of all
93 tests conducted in the State of Utah during that time. The registry was built from the hospital
94 operations dashboards, which contain basic demographics and test results for all patients tested
95 for SARS-CoV-2. We linked this data with additional data using the Enterprise Data Warehouse
96 (EDW) at UHealth. The EDW aggregates data from the health system's disparate data collection
97 systems in support of operations and research. We extracted demographics, International
98 Classification of Diseases (ICD) billing codes, and vital signs, from the EDW to create the data
99 set for this project. Demographic characteristics, including age, sex, and race/ethnicity were
100 collected from all patients on intake for primary care or hospital visits, and via questionnaire for
101 mobile testing centers. The department that ordered SARS-CoV-2 test was recorded, and data on
102 comorbidities were collected via billing codes for all SARS-CoV-2 tested patients. While we
103 were unable to determine if individuals were hospitalized as a result of their SARS-CoV-2
104 infection, we approximated this by determining if each person tested was hospitalized for any
105 cause within 14 days of SARS-CoV-2 testing.
106

107 For each demographic variable, we calculated the odds ratios for testing SARS-COV-2 positive
108 for a series of variables including race or ethnicity, age, and sex, adjusted for cough, fever,
109 shortness of breath, and contact with a known SARS-COV-2 cases. Race/ethnicity was measured
110 by comparing non-Hispanic White persons to non-White persons. Non-white persons included
111 Hispanic/Latino, Asian, Black or African American, Native Hawaiian or other Pacific Islander,
112 American Indian/Native Alaskan and other race individuals, as well as those with unknown race
113 and those who chose not to disclose their race. We also compared Hispanic persons to non-
114 Hispanic White persons, as they are the largest minority group in Salt Lake County and comprise
115 19% of the population.
116

117 We reviewed clinical text collected between 24 hours before and after SARS-CoV-2 testing for a
118 subset of 2,043 persons tested prior to 31 March 2020 for additional manual chart review. Of the
119 20,088 total tests performed, 7,113 were performed prior to 31 March 2020. We selected a
120 random subset, stratified by date, such that we included at least 20% of patients presenting on
121 each day. This resulted in a sample size of 2043. Data on symptoms were extracted from the
122 subset of manually reviewed charts. Symptoms were selected based on previous literature of
123 SARS-CoV-2-positive patients²⁰. Epidemiologic risk factors collected from these charts
124 included history of travel if mentioned, exposure to a SARS-CoV-2-positive person, healthcare
125 worker status, and smoking status. Symptoms/exposure factors not mentioned in the medical
126 chart notes were recorded as ‘not mentioned.’ In analysis we assumed “not mentioned”
127 symptoms were absent. Percentage of present symptoms are seen below (Table 2).

128
129 In addition to collecting UHealth data, we extracted county-level and state-level data from the
130 Utah COVID-19 dashboard for comparison¹⁹.

131
132 This study was reviewed by the University of Utah Institutional Review Board (IRB) and
133 determined to be exempt.

134 135 **Results**

136 *Demographic Characteristics of Tested Individuals*

137 Of the 20,088 patients tested for SARS-CoV-2 between 10 March 2020 and 24 April 2020, 1,229
138 (6.1%) tested positive for SARS-CoV-2. Testing occurred at 10 different types of facilities,
139 including 18,399 (92%) in outpatient clinics, 279 (1%) at mobile testing sites, 958 (5%) in the
140 emergency department (ED), 219 (1%) in an acute care facility, and 107 (0.5%) tested in the
141 intensive care unit (ICU). The proportion of positive tests were similar between facility types
142 (Table 1).

143 144 *Race and Ethnicity*

145 While the racial/ethnic distribution of the population tested at UHealth was broadly similar to the
146 racial/ethnic distribution of Salt Lake County (Figure 2B), there were still statistical differences
147 ($p < 0.001$). Discrepancies were especially pronounced for persons identifying as Hispanic or
148 Latino; Hispanic/Latino persons comprise 19% of the county population but only 10% of all
149 tested at UHealth. Furthermore, we found that non-White persons were more likely to test
150 positive compared to non-Hispanic White/Caucasians (adjOR=1.1, 95% CI: 0.8, 1.6), and
151 Hispanic/Latino persons were more likely to test positive compared to non-Hispanic
152 White/Caucasian persons (adjOR=2.0, 95% CI: 1.3, 3.1; Table 1, Figure 2). Likewise, when
153 comparing the location where individuals presented for testing (e.g., ED, outpatient clinic,
154 mobile testing site), we found variation by race/ethnicity ($p < 0.001$), but Hispanic/Latino persons
155 were only slightly more likely to be tested in the ED compared to non-Hispanic White/Caucasian
156 (6.3% versus 4.9%). We did not find a difference in reported symptoms between
157 Hispanic/Latino and non-Hispanic White/Caucasian persons.

158 159 *Age Structure*

160 The population tested at UHealth is not similar to the age structure of Salt Lake County (Figure
161 3). However, the age distribution of those tested at UHealth is consistent between individuals
162 testing negative versus positive. Younger individuals (aged 0-19), were underrepresented in

163 overall rates of testing and rates of testing positive (Table 1, Figure 3). Without adjusting for the
164 underlying age distribution of the county population, we found that persons aged 20-59 had the
165 highest odds of testing positive compared to those 0-19 years old (adjOR=1.6, 95% CI: 0.7, 3.6,
166 Table 1).

167
168 *Sex*

169 Persons who tested positive for SARS-CoV-2 were more likely to be male. Among individuals
170 who tested positive for SARS-CoV-2, over 56% were male (44% were female), while 44% of
171 individuals who tested negative were male (56% of female). Excluding individuals for which
172 gender was unknown or non-binary (11 people), males had 1.4 times the odds of testing SARS-
173 COV-2 positive compared to females (95% CI: 1.0, 1.9).

174
175 ***Clinical Characteristics***

176 Among manually extracted charts, 136 (6.7%) individuals tested positive for SARS-CoV-2
177 between 10 March 2020 and 31 March 2020 (Table 2). This was similar to the proportion of
178 individuals testing positive for SARS-CoV-2 among all those tested 1229 (6.5%).

179
180 *Clinical symptoms*

181 Persons testing SARS-COV-2-positive and SARS-COV-2-negative had similar vital signs and
182 rates of cough (89%), but varied in presentation of other symptoms. Symptom profiles were
183 different in persons who tested SARS-COV-2 positive compared to those who tested SARS-
184 COV-2 negative (Table 2). The median duration between symptom onset and presentation for
185 testing was 4 days (IQR=2.1, 7.0) for both SARS-COV-2 positive and SARS-COV-2 negative
186 persons (Table 2).

187
188 *Hospitalization rates*

189 Using all-cause hospitalization occurrence within 14 days after testing, we found that 5.1% of
190 persons tested positive were hospitalized, and 5.1% of persons testing negative were
191 hospitalized. Rates of hospitalization were similar across racial ethnic groups, with 6% of people
192 identifying as Black/African American, 5% of people identifying as non-Hispanic
193 White/Caucasian, and 5% of people identifying as Hispanic/Latino being hospitalized within 14
194 days of SARS-CoV-2 testing. Those identifying as American Indian and Alaskan Native had
195 higher rates of 14-day hospitalization (9%), but the sample size was limited.

196
197 We found that hospitalization differed by age, and older adults had the highest odds of all-cause
198 hospitalization (adjOR=6.9, 95% CI: 2.1, 22.5 for adults 60+ compared to 0-19-year olds; Table
199 1, Figure 3). Additionally, we found all-cause hospitalization rates increased with age in both
200 SARS-COV-2-positive and SARS-COV-2-negative persons (Figure 3d).

201
202 *Epidemiological Risk Factors*

203 Persons testing SARS-CoV-2 positive were more likely to have known exposure to another
204 SARS-CoV-2 person or a history of travel (Table 2). Among SARS-COV-2-positive persons,
205 57% had been in contact with a confirmed SARS-CoV-2 case and 46% had travelled before their
206 own SARS-CoV-2 testing, compared to 29% and 24% in SARS-COV-2-negative persons,
207 respectively (Table 2). 9% of SARS-COV-2-positive persons and 11% of SARS-COV-2-
208 negative persons were healthcare workers (Table 2).

209

210 **Conclusions**

211 The high-level of SARS-CoV-2 testing availability in Utah has allowed for a higher proportion
212 of the outpatient, community dwelling (and usually social distancing) population to be tested.
213 This expands on previous studies that were limited to more severe or critically-ill patients, as a
214 result of limited test availability. These data provide insight into the epidemiological and clinical
215 characteristics of COVID-19 over a much broader range of disease severity, as well as highlight
216 disparities in COVID-19 response. While the data presented in this study represents only 20% of
217 the total number of people tested in Utah, based on publicly available state-level data, our
218 findings are largely representative of the overall state-level outbreak.

219

220 While the racial/ethnic makeup of the population tested for SARS-CoV-2 is broadly similar to
221 overall county demographics (Figure 2), important discrepancies remain. We found that persons
222 who identified as non-White had elevated odds of testing positive compared to non-Hispanic
223 White/Caucasians (adjOR=1.1, 95% CI: 0.8, 1.6). The discrepancy was especially pronounced
224 for persons identifying as Hispanic or Latino, who had 2.0 times the odds (95% CI: 1.3, 3.1) of
225 testing SARS-CoV-2 positive compared to those who identified as non-Hispanic White (Figure
226 2). Though our study is limited to those within a single health system, our results are
227 representative to those of the county and the state as a whole. For example, Hispanics make up
228 10% of our tested population, and 22% of those testing positive, which is broadly consistent with
229 state-level data in which Hispanics make up 14.2% of the population and 36.6% of those testing
230 positive ¹⁹.

231

232 There are likely several factors that contribute to the disparities in test-positive rate ²¹.
233 Hispanic/Latino persons are disproportionately employed as essential workers, which increases
234 exposure risk. While Hispanics make up 17.6% of the US workforce, they comprise 25% of
235 service occupations ²². This increased employment exposure is likely compounded by the lack of
236 paid sick leave, as Hispanic workers have lower rates of access to sick leave than do White
237 workers ²³. Workers without paid sick leave are likely to continue working even while sick,
238 leading to increased exposure to co-workers and family-members. Another compounding factor
239 is the decreased ability for public health personnel to conduct contact tracing within minority
240 communities due to historical and current political instances leading to lack of trust between
241 minority communities and government bodies ²⁴⁻²⁹. It is also the case that Hispanic workers may
242 be less able to stop working and stay home due to low incomes. Hispanic households in Salt
243 Lake County make \$56,498/year as compared to \$82,026 among non-Hispanic White households
244 ³⁰. Lower incomes also translate into more crowded housing situations, which facilitate virus
245 transmission. In the US, 10% of Hispanics live in overcrowded conditions ³¹ and that figure rises
246 to 33% among older Hispanics in the urban US ³².

247

248 Persons identifying as non-Hispanic White/Caucasian and Hispanic/Latino were tested at each
249 facility at approximately equal rates – approximately 5% were tested in the ED, and 90% in
250 outpatient clinics. Likewise, using symptom presentation or hypoxia (SpO₂ < 90%) as a proxy
251 for severity, we find no difference between persons identifying as non-Hispanic White/Caucasian
252 and as Hispanic/Latino. The duration of time from symptom onset to seeking testing was also
253 similar across races. This suggests racial disparities in SARS-COV-2 burden are not simply due
254 to Hispanic/Latino persons delaying care until symptoms are critical and presenting to the ED,

255 nor is it as straight forward as preferentially presenting to mobile testing centers. Identifying the
256 underlying cause of these health disparities is a critical step towards containing SARS-CoV-2.

257
258 Unlike previous studies⁶⁻¹¹ that focus only on severe or critical patients, we are also able to
259 identify differential patterns of testing by age. Promisingly, we find most adults (aged 20-79) in
260 our study have high testing rates (over 100 tests per 10,000 population), which indicates that we
261 are capturing large numbers of mild and moderate illness. On the other hand, we also show that
262 testing rates remain lower in children (under 20 years old), and of those under 20 years old who
263 are tested, rates of SARS-CoV-2 positivity are disproportionately lower. This supports previous
264 studies which highlight risk of asymptomatic SARS-CoV-2 in younger individuals, as inclusion
265 in our study was predicated by having at least one symptom (cough, fever, or shortness of breath)
266 ²⁰.

267
268 In addition to differences due to race, we also identified differences due to age. We found the
269 highest hospitalization rates were among middle-aged SARS-COV-2-positive patients (Figure 3).
270 This is broadly consistent with Myers et al.⁶ who found hospitalized and ICU admitted patients
271 in California tended to be middle-aged. Like similar studies⁸, we found hospitalization rates
272 were smallest among younger age individuals (aged 0-19). Indeed, we found persons under 20
273 years old who tested positive for SARS-CoV-2 were far less likely to be hospitalized than their
274 peers who were tested for SARS-CoV-2 but tested negative. Compared to other studies, we
275 found age-adjusted lower hospitalization rates than previously reported among those who test
276 positive.

277
278 Given that Utah has the youngest population in the country³³, a disease that is more likely to be
279 asymptomatic in younger people is especially salient for the overall state burden and the ability
280 to control the outbreak as social distancing measures are loosened. Given our finding that
281 younger individuals are underrepresented in testing due in part to testing criteria, state-level
282 strategies should be re-examined to consider strategies to target testing of younger age groups
283 based on geospatial and epidemiological risk factors, regardless of symptomatology.

284
285 A benefit of expanded testing locations in Salt Lake County was that we were able to assess the
286 demographic characteristics of testing locations. We found that those tested in outpatient
287 facilities tended to be younger and that the proportion of persons tested in the ED increased with
288 age. We found no difference in race/ethnicity by testing location. If testing were to be expanded
289 to lift symptom requirements in an effort to identify more mild and asymptomatic infections, our
290 study highlights the benefit of outpatient and mobile facilities to reach underrepresented
291 demographics.

292
293 This study offers an important contribution to existing COVID-19 literature, which has been
294 primarily focused on hospitalized patients. In contrast to previous work, over 90% of people
295 included in our study were tested in outpatient or community settings. However, our data is
296 limited to persons who met testing criteria; very few asymptomatic infections are captured in this
297 study. Further work understanding the demographic and epidemiological characteristics of
298 asymptomatic infections remains vital to understanding and controlling SARS-CoV-2. By
299 highlighting critical gaps in testing, particularly among Hispanic/Latino communities, where
300 SARS-CoV-2 may be spreading more rapidly due to increased exposure and comparatively

301 reduced testing, this study takes a critical first step towards reversing these health disparities. Our
 302 findings, based largely on those who received testing as outpatients, highlights the potential gaps
 303 in control of SARS-CoV-2 infection related to age, race, and ethnicity.

304
 305

Table 1: Characteristics of SARS-CoV-2-Tested Patients (n=20088)

	SARS-CoV-2 Positive (N=1229)	SARS-CoV-2 Negative (N=18859)	Adjusted¹ Odds Ratio (95% CI)
Age			
Median – (IQR)	38.4 (27.6 - 51)	39.2 (27.8 - 53.6)	
0-19 – no. (%)	91 (7.4%)	1745 (9.3%)	Reference group
20-59 – no. (%)	994 (81%)	13956 (74%)	1.6 (0.7, 3.6)
60+ – no. (%)	137 (11%)	3154 (17%)	1.4 (0.6, 3.4)
Missing – no. (%)	7 (0.57%)	4 (0.02%)	---
Sex			
Female – no. (%)	545 (44%)	10526 (56%)	Reference group
Male – no. (%)	683 (56%)	8323 (44%)	1.4 (1.0, 1.9)
Race/Ethnicity			
White non-Hispanic – no. (%)	489 (40%)	12662 (67%)	Reference group
Hispanic/Latino – no. (%)	402 (33%)	2402 (12%)	2.0 (1.3, 3.1)
Unknown – no. (%)	182 (15%)	1739 (9.2%)	0.8 (0.4, 1.6)
COMBINED ²			0.8 (0.4, 1.4)
Black/African American – no. (%)	31 (2.5%)	330 (1.8%)	---
Asian – no. (%)	30 (2.4%)	416 (2.2%)	---
American Indian and Alaska Native – no. (%)	6 (0.49%)	146 (0.77%)	---
Native Hawaiian and Other Pacific Islander – no. (%)	10 (0.8%)	213 (1.1%)	---
Other – no. (%)	44 (3.6%)	498 (2.6%)	---
Refuse – no. (%)	35 (2.9%)	453 (2.4%)	---
BMI			Not Calculated
Median – (IQR)	30.1 (26 - 34.9)	27.6 (23.2 - 32.9)	---
Missing – no. (%)	1161 (95%)	17433 (92%)	---
Comorbidities			Not Calculated
Any comorbidity	134 (11%)	3402 (18%)	---
fluid/electrolyte disorders – no. (%)	7 (0.57%)	221 (1.2%)	---
Renal Failure – no. (%)	8 (0.65%)	172 (0.91%)	---
Alcohol Abuse – no. (%)	5 (0.41%)	207 (1.1%)	---
Chronic blood loss anemia – no. (%)	5 (0.41%)	160 (0.85%)	---
Chronic pulmonary disease – no. (%)	10 (0.81%)	220 (1.2%)	---
Depression – no. (%)	5 (0.41%)	62 (0.33%)	---
Hypertensive renal disease without renal failure – no. (%)	14 (1.2%)	398 (2.1%)	---
Hypertension, uncomplicated – no. (%)	111 (9.0%)	2809 (15%)	---
Obesity – no. (%)	11 (0.90%)	340 (1.8%)	---
Other – no. (%)	12 (0.98%)	600 (3.2%)	---
Pulse Rate			Not Calculated
Median – (IQR)	87 (78 – 99)	85 (75 - 96)	---
Missing – no. (%)	138 (11%)	2068 (11%)	---
Respiratory Rate			Not Calculated
Median – (IQR)	18 (16 - 20)	18 (16 - 20)	---
Missing – no. (%)	1108 (90%)	16552 (88%)	---
SpO2			Not Calculated

Median – (IQR)	96 (94 – 97)	96 (95 - 98)	---
Missing – no. (%)	131 (10.7%)	2054 (11%)	---
Body Temperature (C)			Not Calculated
Median – (IQR)	37.1 (36.7 - 37.6)	37 (36.6 - 37.3)	---
Missing – no. (%)	762 (62%)	11118 (59%)	---
Facility that ordered SARS-CoV-2 test			Not Calculated
Acute care – no. (%)	7 (0.57%)	212 (1.12%)	---
ED – no. (%)	55 (4.48%)	903 (4.79%)	---
External – no. (%)	5 (0.41%)	0 (0%)	---
ICU – no. (%)	6 (0.49%)	101 (0.54%)	---
Inpatient rehab – no. (%)	0 (0%)	10 (0.05%)	---
LND/MNB – no. (%)	0 (0%)	34 (0.18%)	---
Mobile – no. (%)	230 (1.22%)	230 (1.2%)	---
Outpatient Clinics – no. (%)	1095 (89.1%)	17304 (92%)	---
Telemedicine – no. (%)	0 (0%)	25 (0.13%)	---
Other – no. (%)	12 (0.98%)	40 (0.21%)	---

306 ¹Adjusted for cough, fever, shortness of breath, and known contact with SARS-COV-2 case

307 ²Due to lack of model convergence due to small cell counts, the following race/ethnicity variables were combined
 308 for odds ratio calculations: Black/African American, Asian, American Indian and Alaska Native, Native Hawaiian
 309 and Other Pacific Islander, Other, Refuse


310

311 *Table 2: Symptoms in Subset of SARS-CoV-2-tested Patients (n=2043)*

Symptoms	SARS-CoV-2	SARS-CoV-2
	Positive	Negative
	(n=136)	(n=1907)
Cough – no. (%)	121 (89%)	1697 (89%)
Fever – no. (%)	103 (76%)	1229 (64%)
Shortness of breath – no. (%)	68 (50%)	1239 (65%)
Myalgia – no. (%)	57 (42%)	572 (30%)
Headache – no. (%)	50 (37%)	462 (24%)
Lethargy – no. (%)	43 (32%)	496 (26%)
Nasal – no. (%)	44 (32%)	562 (29%)
sore throat – no. (%)	41 (30%)	592 (31%)
Diarrhea – no. (%)	16 (12%)	188 (9.9%)
nausea/vomit – no. (%)	10 (7.4%)	163 (8.6%)
Duration of symptoms – median (IQR)	4.0 (2.3, 7.0)	4.0 (2.0, 7.0)
BEHAVIORS		
Prior exposure – no. (%)	77 (57%)	546 (29%)
Travel – no. (%)	63 (46%)	461 (24%)
Healthcare worker – no. (%)	12 (8.8%)	205 (11%)
Smoker – no. (%)		
never – no. (%)	78 (57%)	869 (46%)
former – no. (%)	5 (3.7%)	164 (8.6%)
current – no. (%)	4 (2.9%)	186 (9.8%)
missing – no. (%)	49 (36%)	688 (36%)


312

313




314
315
316
317
318
319
320

Figure 1: State-level SARS-CoV-2 testing data in Utah from 04 March to 28 April 2020. Grey bars represent the total number of tests performed each day in Utah and the red bars represent the total number of positive tests, based on the date they were ordered. The black line represents the percent of tests state wide that were positive each day.



321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340

Figure 2: Epidemiological characteristics of persons who were tested for SARS-CoV-2 at UHealth by race/ethnicity. A) Shows 7-day moving average of the percent of tests that were positive by race/ethnicity, each day over the course of the study, for select race and ethnicities. Red, teal, and yellow represent the percent positive for Black or African American, Hispanic, and White or Caucasian persons while black represents the 7-day moving average across all races. B) Shows the percent of persons by each race from the 2018 census estimates for Salt Lake County³⁴, the proportion of persons tested by race/ethnicity, as well as the proportion of persons who tested positive by race/ethnicity. C) The log-odds of testing positive by race, compared to White/Caucasian. Adjusted for cough, fever, shortness of breath, and contact with known SARS-COV-2 case. Due to lack of model convergence due to small cell counts, the following race/ethnicity variables were combined for odds ratio calculations: Black/African American, Asian, American Indian and Alaska Native, Native Hawaiian and Other Pacific Islander, Other, Refuse



341
 342 **Figure 3: Epidemiological characteristics of persons who were tested for SARS-CoV-2 at**
 343 **UHealth by age.** A) The log-odds of testing positive by age, compared to 0-9 year olds. Adjusted
 344 for cough, fever, shortness of breath, and contact with known SARS-COV-2 case. B) Shows the
 345 percent of persons in each age group from the 2018 census estimates for Salt Lake County³⁴. C)
 346 The rate of testing, per 10,000 population, by age for persons testing (light blue) SARS-CoV-2
 347 positive and (dark blue) SARS-CoV-2 negative. D) The rate of testing, per 10,000 population, by
 348 age all cause hospitalizations within 14 days of testing for (light blue) SARS-CoV-2 positive and
 349 (dark blue) SARS-CoV-2 negative.
 350
 351

352 **References**

- 353 1. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real
354 time. *Lancet Infect Dis*. 2020.
- 355 2. Young BE, Ong SWX, Kalimuddin S, et al. Epidemiologic features and clinical course of
356 patients infected with SARS-CoV-2 in Singapore. *Jama*. 2020.
- 357 3. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus
358 disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from
359 the Chinese Center for Disease Control and Prevention. *Jama*. 2020;323(13):1239-1242.
- 360 4. Huang SS, Datta R, Platt R. Risk of acquiring antibiotic-resistant bacteria from prior room
361 occupants. *Arch Intern Med*. 2006;166(18):1945-1951. doi:10.1001/archinte.166.18.1945
- 362 5. Riccardo F, Ajelli M, Andrianou X, et al. Epidemiological characteristics of COVID-19
363 cases in Italy and estimates of the reproductive numbers one month into the epidemic.
364 *medRxiv*. 2020.
- 365 6. Myers LC, Parodi SM, Escobar GJ, Liu VX. Characteristics of Hospitalized Adults With
366 COVID-19 in an Integrated Health Care System in California. *JAMA*. 2020.
- 367 7. Arentz M, Yim E, Klaff L, et al. Characteristics and outcomes of 21 critically ill patients
368 with COVID-19 in Washington State. *Jama*. 2020.
- 369 8. Garg S. Hospitalization rates and characteristics of patients hospitalized with laboratory-
370 confirmed coronavirus disease 2019—COVID-NET, 14 States, March 1--30, 2020.
371 *MMWR Morb Mortal Wkly Rep*. 2020;69.
- 372 9. Gold JAW. Characteristics and Clinical Outcomes of Adult Patients Hospitalized with
373 COVID-19—Georgia, March 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69.
- 374 10. Bhatraju PK, Ghassemieh BJ, Nichols M, et al. Covid-19 in critically ill patients in the
375 Seattle region—case series. *N Engl J Med*. 2020.
- 376 11. Vahidy FS, Nicolas JC, Meeks JR, et al. Racial and Ethnic Disparities in SARS-CoV-2
377 Pandemic: Analysis of a COVID-19 Observational Registry for a Diverse US
378 Metropolitan Population. *medRxiv*. 2020.
- 379 12. Project CT. Most recent data. *COVID Track Proj*. 2020. <https://covidtracking.com/data/>.
- 380 13. Bialek S, Bowen V, Chow N, et al. Geographic differences in COVID-19 cases, deaths,
381 and incidence—United States, February 12--April 7, 2020. 2020.
- 382 14. Padula W V. Why Only Test Symptomatic Patients? Consider Random Screening for
383 COVID-19. 2020.
- 384 15. Li R, Pei S, Chen B, et al. Substantial undocumented infection facilitates the rapid
385 dissemination of novel coronavirus (SARS-CoV2). *Science (80-)*. 2020.
- 386 16. Chow EJ, Schwartz NG, Tobolowsky FA, et al. Symptom screening at illness onset of
387 health care personnel with SARS-CoV-2 infection in King County, Washington. *JAMA*.
388 2020.
- 389 17. Sharfstein JM, Becker SJ, Mello MM. Diagnostic testing for the novel coronavirus. *Jama*.
390 2020.
- 391 18. Baggett TP, Keyes H, Sporn N, Gaeta JM. Prevalence of SARS-CoV-2 Infection in
392 Residents of a Large Homeless Shelter in Boston. *JAMA*. 2020.
- 393 19. coronavirus.utah.gov. Case Counts: coronavirus. Coronavirus.utah.gov.
394 <https://coronavirus.utah.gov/case-counts/>. Published 2020. Accessed May 1, 2020.
- 395 20. Bialek S, Gierke R, Hughes M, McNamara LA, Pilishvili T, Skoff T. Coronavirus Disease
396 2019 in Children—United States, February 12--April 2, 2020. *Morb Mortal Wkly Rep*.
397 2020;69(14):422.

- 398 21. for Disease Control TC, Prevention. COVID-19 in Racial and Ethnic Minority Groups.
399 2020. <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/racial-ethnic->
400 minorities.html. Accessed May 1, 2020.
- 401 22. U.S. Bureau of Labor Statistics. Labor Force Statistics from the Current Population
402 SurveyCPS CPS Program Links. U.S. Bureau of Labor Statistics.
403 <https://www.bls.gov/cps/cpsaat11.htm>. Published 2020. Accessed May 1, 2020.
- 404 23. of Labor Statistics USB. Report 1082, Labor force characteristics by race and ethnicity,
405 2018. 2019. <https://www.bls.gov/opub/reports/race-and-ethnicity/2018/home.html>.
406 Accessed May 1, 2020.
- 407 24. Munier N, Albarracin J, Boeckelman K. Determinants of rural Latino trust in the federal
408 government. *Hisp J Behav Sci*. 2015;37(3):420-438.
- 409 25. Armstrong K, Ravenell KL, McMurphy S, Putt M. Racial/ethnic differences in physician
410 distrust in the United States. *Am J Public Health*. 2007;97(7):1283-1289.
- 411 26. Shoff C, Yang T-C. Untangling the associations among distrust, race, and neighborhood
412 social environment: A social disorganization perspective. *Soc Sci Med*. 2012;74(9):1342-
413 1352.
- 414 27. Wilkes R, Wu C. Trust and minority groups. *Oxford Handb Soc Polit Trust*. 2018:231.
- 415 28. Thiede M. Information and access to health care: is there a role for trust? *Soc Sci Med*.
416 2005;61(7):1452-1462.
- 417 29. Stepanikova I, Mollborn S, Cook KS, Thom DH, Kramer RM. Patients' race, ethnicity,
418 language, and trust in a physician. *J Health Soc Behav*. 2006;47(4):390-405.
- 419 30. Department SLCH. Healthy Salt Lake, 2020 Demographics. 2020.
420 healthysaltlake.org/demographics. Accessed May 1, 2020.
- 421 31. on Aging NHC. Housing. 2020. <https://www.nhcoa.org/our-focus/housing/>. Accessed
422 May 1, 2020.
- 423 32. Burr JA, Mutchler JE, Gerst K. Patterns of residential crowding among Hispanics in later
424 life: Immigration, assimilation, and housing market factors. *Journals Gerontol Ser B*
425 *Psychol Sci Soc Sci*. 2010;65(6):772-782.
- 426 33. Census US. Median Age in 2010 and 2018 for the United States, Regions, and States.
427 United States Census Bureau. [https://www.census.gov/newsroom/press-](https://www.census.gov/newsroom/press-kits/2019/detailed-estimates.html)
428 kits/2019/detailed-estimates.html. Published 2020. Accessed May 1, 2020.
- 429 34. Census TUS. QuickFacts Salt Lake County, Utah; Utah. 2018.
430 <https://www.census.gov/quickfacts/fact/table/saltlakecountyutah,UT/PST045218>.
431 Accessed May 1, 2020.
- 432
- 433
- 434

