1 **Title:** Facemasks prevent influenza-like illness: implications for COVID-19

2

3 **Total word count:** 395

4

5 The coronavirus disease 2019 (COVID-19) pandemic is causing a huge toll on
6 individuals, families, communities and societies across the world. Currently, whether
7 wearing facemasks in public should be a measure to prevent transmission of severe
8 acute respiratory syndrome coronavirus-2 (SARS-CoV-2) remains controversial.¹ This
9 is largely because there have been no randomized controlled trials (RCTs) for
10 coronavirus to directly support this. However, lessons may be taken from published
11 RCTs examining influenza-like illness (ILI).^{2,3} Recent studies suggested that
12 SARS-CoV-2 shares similar transmission route with influenza virus,⁴ and the
13 incidence of community transmission of SARS-CoV-2 in individuals with ILI is
14 high.⁵ Therefore, we undertook this meta-analysis of RCTs examining the efficacy of
15 wearing facemasks to prevent ILI in community settings, irrespective of confirmatory
16 testing for the causative virus.

17

18 We undertook a systematic literature search for RCTs related to facemasks and
19 ILI between 1966 and April 2020 using PUBMED, EMBASE, and Cochrane library.
20 RCTs undertaken in community (not hospital) settings comparing wearing and not
21 wearing facemasks for ILI were included. Incidence of ILI (e.g., fever, cough,
22 headache, sore throat, aches or pains in muscles or joints) was estimated per group.

23 Relative risk (RR) and 95% confidence interval (CI) were calculated.

24

25 We screened 899 related abstracts and eventually included 8 RCTs (**Figure S1**).

26 Basic characteristics and quality of included RCTs are listed in **Supplement**.

27 Participants wearing facemasks had a significantly lower risk of developing ILI than
 28 those not wearing facemasks (pooled RR=0.81, 95% CI: 0.70-0.95) and there was no
 29 heterogeneity (**Figure 1**). The decreased risk of ILI was more pronounced if everyone
 30 wore facemask irrespective of whether they were infected or not (RR=0.77, 95% CI:
 31 0.65-0.91), compared to those wearing facemasks when infected (RR=0.95, 95% CI:
 32 0.58-1.56) or uninfected (RR=1.26, 95% CI: 0.69-2.31).

33

34 This study shows that wearing facemasks, irrespective of infection status, is
 35 effective in preventing ILI spread in the community. This situation mirrors what is
 36 happening now in public settings where we do not know who has been infected and
 37 who has not. Although there are no RCTs of facemasks for SARS-CoV-2, as with
 38 other simple measures such as social distancing and handwashing, these data support
 39 the recommendation to wear facemasks in public to further reduce transmission of
 40 SARS-CoV-2 and flatten the curve of this pandemic, especially when social
 41 distancing is impractical, such as shopping, or travelling with public transport for
 42 work that cannot be done from home.

43 We declare no competing interests.

44 **Jie Wei, Michael Doherty, Monica S.M. Persson, Aliya Sarmanova, Subhashisa**

45 **Swain, Changfu Kuo, Chao Zeng, Guanghua Lei, Weiya Zhang**

46 **lei_guanghua@csu.edu.cn**

47 **Weiya.zhang@nottingham.ac.uk**

48

49 Health Management Center, Xiangya Hospital, Central South University, Changsha,

50 Hunan, China (JW); Academic Rheumatology Department, Division of Rheumatology,

51 Orthopaedics and Dermatology, School of Medicine, University of Nottingham,

52 Nottingham, UK (MD, SS, WZ); Centre of Evidence Based Dermatology, Division of

53 Rheumatology, Orthopaedics and Dermatology, School of Medicine, University of

54 Nottingham, Nottingham, UK (MP); Musculoskeletal Research Unit, Bristol Medical

55 School (THS), University of Bristol, Bristol, UK (AS); Division of Rheumatology,

56 Allergy and Immunology, Chang Gung Memorial Hospital, Taoyuan, Taiwan (CK);

57 Department of Orthopaedics, Xiangya Hospital, Central South University, Changsha,

58 Hunan, China (CZ, GL)

59 **References**

60 1. Greenhalgh T, Schmid MB, Czypionka T, Bassler D, Gruer L. Face masks for the public

61 during the covid-19 crisis. *BMJ* 2020; **369**: m1435.

62 2. Aiello AE, Murray GF, Perez V, et al. Mask use, hand hygiene, and seasonal

63 influenza-like illness among young adults: a randomized intervention trial. *J Infect Dis* 2010;

64 **201**(4): 491-8.

- 65 3. Suess T, Remschmidt C, Schink SB, et al. The role of facemasks and hand hygiene in the
66 prevention of influenza transmission in households: results from a cluster randomised trial;
67 Berlin, Germany, 2009-2011. *BMC Infect Dis* 2012; **12**: 26.
- 68 4. Boelle PY, Souty C, Launay T, et al. Excess cases of influenza-like illnesses synchronous
69 with coronavirus disease (COVID-19) epidemic, France, March 2020. *Euro Surveill* 2020;
70 **25**(14).
- 71 5. Zwald ML, Lin W, Sondermeyer Cooksey GL, et al. Rapid Sentinel Surveillance for
72 COVID-19 - Santa Clara County, California, March 2020. *MMWR Morb Mortal Wkly Rep* 2020;
73 **69**(14): 419-21.
- 74

75 **Figure legends:**

76

77 **Figure 1. Relative risk (RR) and 95% confidence interval (CI) of developing**
78 **influenza-like symptoms between wearing versus not wearing facemasks in**
79 **general population.**