Rapid implementation of real-time SARS-CoV-2 sequencing to investigate healthcare-associated COVID-19 infections

Authors

Luke W. Meredith^{#*1}, William L. Hamilton^{#*2,3}, Ben Warne^{2,3}, Charlotte J. Houldcroft², Myra Hosmillo¹, Aminu S. Jahun¹, Martin D. Curran⁴, Surendra Parmar⁴, Laura G. Caller^{1,5}, Sarah L. Caddy², Fahad A. Khokhar², Anna Yakovleva¹, Grant Hall¹, Theresa Feltwell², Sally Forrest², Sushmita Sridhar², Michael P. Weekes², Stephen Baker², Nicholas Brown⁴, Elinor Moore³, Ashley Popay⁵, Iain Roddick⁵, Mark Reacher⁵, Theodore Gouliouris^{3,6}, Sharon J. Peacock^{2,7}, Gordon Dougan², M. Estée Török^{#\$2,6} and Ian Goodfellow^{#\$1}

These authors contributed equally to the work:

* Joint first author

\$ Joint last author

Affiliations

1. Department of Pathology, University of Cambridge, Cambridge, UK
2. Department of Medicine, University of Cambridge, Cambridge, UK
3. Cambridge University Hospitals NHS Foundation Trust, UK
4. Public Health England Clinical Microbiology and Public Health Laboratory, Cambridge, UK
5. Francis Crick Institute, London, United Kingdom
6. Field Epidemiology, Field Service, National Infection Service, Public Health England, Cambridge, United Kingdom
7. National Infection Service, London, Public Health England

Corresponding Authors:

Dr M. Estée Török et317@cam.ac.uk
 Professor Ian Goodfellow ig299@cam.ac.uk

Abstract

Background

The burden and impact of healthcare-associated COVID-19 infections is unknown. We aimed to examine the utility of rapid sequencing of SARS-CoV-2 combined with detailed epidemiological analysis to investigate healthcare-associated COVID-19 infections and to inform infection control measures.

Methods

We set up rapid viral sequencing of SARS-CoV-2 from PCR-positive diagnostic samples using nanopore sequencing, enabling sample-to-sequence in less than 24 hours. We established a rapid review and reporting system with integration of genomic and epidemiological data to investigate suspected cases of healthcare-associated COVID-19.

Results

Between 13 March and 24 April 2020 we collected clinical data and samples from 5191 COVID-19 patients in the East of England. We sequenced 1000 samples, producing 747 complete viral genomes. We conducted combined epidemiological and genomic analysis of 299 patients at our hospital and identified 26 genomic clusters involving 114 patients. 66 cases (57.9%) had a strong epidemiological link and 15 cases (13.2%) had a plausible epidemiological link. These results were fed back to clinical, infection control and hospital management teams, resulting in infection control interventions and informing patient safety reporting.

Conclusions

We established real-time genomic surveillance of SARS-CoV-2 in a UK hospital and demonstrated the benefit of combined genomic and epidemiological analysis for the investigation of healthcare-associated COVID-19 infections. This approach enabled us to detect cryptic transmission events and identify opportunities to target infection control interventions to reduce further healthcare-associated infections.

Introduction

SARS-CoV-2 emerged in the human population in December 2019¹, originating from an intermediate animal host². Owing to the error prone nature of the viral replication process, RNA viruses, such as SARS-CoV-2, accumulate mutations over time which results in sequence diversity. The current mutation rate of SARS-CoV-2 is estimated to be ~2.5 nucleotides per month^{3,4}. Sequencing of SARS-CoV-2 can provide valuable information on virus biology, transmission and population dynamics⁵⁻⁸. When linked with detailed epidemiological data and performed in a timescale of days, genomic data can support epidemiological investigations of potential hospital-acquired infections (HAI). On a larger population scale, genomic surveillance of SARS-CoV-2 can inform which lineages of the virus are currently circulating in the human population, how these change over time as an indicator of the success of control measures, how often new sources of virus are introduced from other geographic areas, and how the virus evolves in response to interventions.

Healthcare-associated infections (HCAI) can affect patients, resulting in increased morbidity and mortality, and healthcare workers (HCW), impacting on staff sickness and morale, to the detriment of patient care. It is crucial to rapidly detect and manage HCAI effectively to prevent both complications and onward transmission to susceptible patients and staff. The burden of nosocomial COVID-19 infections is unknown but one study from China has reported a 41% prevalence⁹. Worldwide, over 22,000 cases of COVID-19 infection in healthcare workers have been reported, but is likely to be an underestimate¹⁰. As the number of community-acquired COVID-19 cases reduces, healthcare settings are likely to act as reservoirs of infection. Identifying transmission events in these settings will therefore become increasingly important to manage outbreaks and monitor effective infection control.

We aimed to examine the utility of rapid sequencing of SARS-CoV-2, combined with detailed epidemiological analysis, to investigate healthcare-associated COVID-19 infections and to inform infection control measures in our hospital.

Methods

Study design, setting and participants

A prospective surveillance study of COVID-19 infections was conducted at Cambridge University Hospitals NHS Foundation Trust (CUH), a secondary care provider and tertiary referral centre in the East of England (EoE). Clinical specimens collected from patients presenting to 18 hospitals in EoE were submitted to the Public Health England Clinical Microbiology and Public Health Laboratory (CMPHL) at CUH for diagnostic testing. Samples underwent nucleic acid extraction and were tested for presence of SARS-CoV-2 using a validated in-house RT qPCR assay developed by CMPHL (Supplementary Methods). The test was reported as SARS-CoV-2 PCR positive if the cycle threshold (Ct) value was ≤36.

Study procedures

Demographic, clinical and laboratory data were extracted from the hospital information system (EPIC Systems Corporation, Verona, USA). All PCR-positive diagnostic samples were identified

and transferred from the CMPHL to the Department of Virology for nanopore sequencing. All CUH samples and a selection of samples from East of England hospitals were selected for sequencing.

Sequencing

Where Ct values were available prior to sample selection, positive samples with a Ct value ≤ 33 were sequenced using a multiplex PCR based approach according to the modified ARTIC v2 protocol v3 primer set^{11,12}. Amplicon libraries were sequenced using MinION flow cells v9.4.1 (Oxford Nanopore Technologies, Oxford, UK). Genomes were assembled using reference-based assembly and a bioinformatic pipeline¹³ using 20x minimum coverage cut-off for any region of the genome and 50.1% cut-off for calling single nucleotide polymorphisms.

Genomic analysis

All sequences underwent quality control (QC) filtering and de-duplication to remove repeat samples from the same patient. Multiple sequence alignment was performed using MAFFT¹⁴. Phylogenetic trees were produced using IQ-TREE¹⁵ and visualised in Microreact¹⁶ for weekly hospital reports and the R package ggtree¹⁷. A pairwise SNP distance matrix was produced from the alignment (Supplementary Methods). Viral lineages¹⁸ were assigned with the PANGOLIN package (<https://github.com/hCoV-2019/pangolin>) v1.07.

Epidemiological analysis

Patient movement data for all SARS-CoV-2 PCR positive samples were extracted from the hospital information system and transferred to the PHE Field Service (Epidemiology). Epidemiological analysis was performed using a cloud-based plotter application (Cluster Track, Camart Ltd, Cambridge, UK).

Review of healthcare-associated infections

Clusters of COVID-19 cases including HCWs were identified by the clinical and infection control teams and reviewed at a weekly meeting, co-ordinated by the Patient Safety team. The genomic and epidemiological analyses were presented, to help determine whether infections were healthcare-associated and to identify possible causes and interventions. A weekly report was fed back to the clinical, infection control and hospital management teams to inform changes in infection control practice and comply with patient safety procedures.

Ethical considerations

The study was conducted as part of surveillance for COVID-19 infections under the auspices of Section 251 of the NHS Act 2006. It therefore did not require individual patient consent or ethical approval. The COG-UK study protocol was approved by the Public Health England Research Ethics Governance Group.

Results

Baseline characteristics and COVID-19 epidemic curve

368 PCR-confirmed COVID-19 patients were admitted to CUH between 10 March and 24 April 2020 (Figure 1, Table 1). The median age was 66 years (range 0 to 98 years) and 63% were male. 19.6% of patients were admitted to critical care units and 20.1% died. Between 13 March

to 24 April 2020 1000 samples were selected for sequencing and, after QC and de-duplication, 747 were used for downstream analysis (Supplementary Figure 1). 256/368 (69.6%) of CUH COVID-19 admissions had sequencing data available (Supplementary Table 1). 52/368 (14.1%) of admissions were suspected or highly likely hospital-acquired, of which 47 (90.4%) had genome sequences available. A further 10.3% of admissions were community-acquired but likely healthcare-associated, and 2.2% were healthcare workers (Supplementary Table 1). The CUH epidemic curve showed that weekly admissions peaked in week 4 (commencing 30 March 2020) and then declined (Figure 2). The UK went into full lockdown on 23 March 2020. In the early stages of the epidemic community-onset community-acquired infections predominated but the frequency of healthcare-associated infections increased over time.

Genomic analysis

Each week a sample set was locked and underwent bioinformatic analysis. Out of 1,000 sequenced genomes, 253 were excluded from downstream analysis because they failed to pass QC thresholds, metadata were missing or they were repeat samples from the same patient (Figure 1). The median genome depth of coverage was 6,612x. We compared Ct value versus depth of coverage and found that the latter declined at Ct values >30 (Supplementary Figure 2). We also examined the location and frequency of SNPs across the genomes (Supplementary Figure 2). Genomes were assigned to a lineage based on the combination of mutations that have accumulated since the virus emerged. As of 23 March 2020, 12 lineages had been described in the UK¹⁹. The vast majority of samples in both the EoE and CUH belonged to lineage B.1. There were no lineage A samples, which have mainly been identified in China, USA, South Korea and Australia¹⁸ (Figure 3, Supplementary Figure 3).

Phylogenetic trees were used to explore potential genetic clustering and correlation with sampling ward location and/or cases of suspected HAI (Figure 4). Samples collected from the Emergency Department (ED) were phylogenetically dispersed, likely reflecting unconnected recent transmission events in community-acquired infections (Supplementary Figure 4). In contrast, samples collected from several wards in CUH and an outpatient dialysis unit were genetically clustered (Figure 4). The putative ward clusters include a high proportion of suspected HAI, suggestive of linked transmission chains in the hospital.

Due to its recent introduction into the human population, SARS-CoV-2 has low genetic diversity; here, a median of 8 single nucleotide polymorphisms (SNPs) separated any two samples at CUH (Supplementary Figure 5). This low sequence diversity makes interpretation of putative clusters challenging, as samples may be identical “by chance” rather than because of a recent connected transmission chain. To investigate genomic clustering further we adopted a combined genetic, clinical and epidemiological approach. Samples with zero SNP differences were identified and clusters named numerically by decreasing sample size (Supplementary Table 2).

Overall 114 genomes in 26 clusters shared at least one identical sequence. The two largest clusters each had 15 identical genomes. Patients’ medical records (including address, social setting, clinical details and ward movements) were reviewed for all putative genomic clusters to assess whether cases had plausible or likely linked recent transmission (Supplementary

Methods). Of 114 cases from 26 putative clusters, 63 cases (55.3%) had strong epidemiological evidence to support recent transmission, 27 cases (23.7%) had intermediate evidence, and 24 (21.1%) no evidence of connected transmission. Clusters with strong evidence of linked recent transmission include cases where a connection was already suspected, such as groups of likely HAI seen on several CUH hospital wards (Figure 4), and clusters that were previously not recognised as being linked, such as a care home outbreak involving HCWs based in hospital and community settings. These are described further below.

Application and impact of combined epidemiological and genomic analyses

Directed by clinical and infection control teams we established a process to conduct focused genomic and epidemiological analyses of suspected HAI cases (Supplementary Figure 6). These were discussed in weekly meetings with an accompanying written report submitted to the hospital. Anonymised examples are described here.

Hospital cluster 1. Six surgical patients on ward A (a non-COVID area) were diagnosed with COVID-19 infection (Figure 4B). All cases had been on the ward for >14 days prior to their specimen date and were classified as probable HAI. Genomic data were available for all cases; five differed by zero SNPs and one by a single SNP, consistent with recent ward-based transmission events. In response to this, universal mask use was introduced.

Hospital cluster 2. Four transplant patients on ward B (a non-COVID area) were diagnosed with COVID-19 infection between 3 and 20 April 2020 (Supplementary Figure 7A). A fifth patient, who had been recently discharged from the ward, presented to the ED with COVID-19 infection. Genomic analysis revealed that all 5 cases had identical genomes. Three HCWs were found to have identical genomes in the same cluster as the ward B cases; two had worked on ward B, one of whom had professional contact with the other HCW. These findings led to a review of infection control and PPE procedures for staff and patients in the transplant service.

Dialysis unit cluster. Renal dialysis patients are among the most vulnerable to COVID-19 (up to 19% mortality²⁰), with challenging infection control arrangements as most units consist of large open rooms with no barriers between patients. Six patients with end-stage renal failure were diagnosed with COVID-19 between 1 and 20 April 2020, testing positive in several locations including ED and an acute admissions ward (Supplementary Figure 7B). Their viral genomes were identical, and epidemiological investigation revealed they dialysed at the same outpatient dialysis unit on the same days of the week. This suggests linked recent transmission of community-onset healthcare-associated infections. These findings led to a review of infection control procedures in dialysis patients and identified shared patient transportation as a possible risk factor. Genomics can also be used to “rule out” linked transmission. The renal ward (which shares patients with the outpatient dialysis unit) also had a group of COVID-19 cases at around the same time. However, the dialysis unit genomes belonged to lineage B.2 (relatively rare in EoE), whereas the renal ward genomes were the common B.1 lineage, making it very unlikely that infections between the two patient groups were related.

Community cluster. Fifteen patients were admitted to CUH with COVID-19 between 7 and 21 April 2020 with genetically identical viruses. Eight patients were residents at a community care home (care home A). A review of the medical records revealed that another patient in this genetic cluster worked in care home A and one was a retired nurse who worked in an unknown care home. Two cases were paramedics and two were nurses (who worked in different wards at CUH but lived with paramedics). The final case did not have any discernible epidemiological links with the others. In summary, this investigation revealed a cluster of cases with evidence of linked transmission coming from the same care home with potential links between HCW and the care home, none of which had been detected by clinicians or infection control.

The information from these combined epidemiological / genomic investigations was fed back to the clinical, infection control, and hospital management teams. This triggered further investigations of patient isolation, ward cleaning procedures, use of personal protective equipment (PPE) and staff social distancing behaviour. HCAI were also assessed in relation to potential harm caused to patients and recorded in hospital's patient safety reporting system for follow-up and further action.

Discussion

The value of real-time viral genome sequencing has been demonstrated in previous epidemics^{21–26}. We sought to embed genomic surveillance as part of an active SARS-CoV-2 infection control process in a large UK hospital. A rapid sequencing workflow was established on 23 March 2020 using multiplex PCR-based nanopore sequencing, which has been proven to be effective in a wide range of clinical samples and viral loads²⁷. We aimed to sequence all available positive samples from CUH and a selection from each of the EoE regional hospitals submitted to the diagnostic microbiology laboratory, linking with clinical metadata pulled from the hospital electronic patient records system. We also included 44 samples collected as part of a local HCW screening programme (Rivett et al, unpublished). In 5 weeks, over 1,000 SARS-CoV-2 genomes had been sequenced including the majority of CUH samples from this phase of the epidemic. We applied this system to investigate nosocomial and HCW COVID-19 infections at CUH, integrating genomics with epidemiological and clinical data.

We examined the diversity in SARS-CoV-2 at CUH and found that overall genetic diversity was low and reflected the pattern seen in the EoE as a whole, with most viruses belonging to the B.1 lineage. We identified a median of 8 (range 0 to 24) SNP differences between viruses, and 4.5% of pairwise comparisons between CUH genomes had 0 to 1 SNP differences. Given the virus's mutation rate and infectious timeframe, cases may share linked recent transmission if there are fewer than ~2 SNP differences. However, unravelling potential transmission networks requires rapid integration with in-depth epidemiological data.

We investigated clusters of patients and HCW cases at CUH in response to queries from the clinical and infection control teams. Using ward location data for patients and HCWs we performed epidemiological analyses to determine if there had been ward-based contact. We compared the genomes of patients and HCWs in the suspected clusters with those of other patients at CUH and in EoE hospitals to examine relatedness. This approach enabled us to confirm or refute evidence

for transmission between patients and / or HCWs within these clusters. Genomic analysis also enabled us to identify cryptic transmission i.e. additional cases that were not initially suspected to be linked to the original ward or HCW clusters. We were also able to use genomic data to refute suspected transmission events (e.g. different lineages between two isolates that were epidemiologically linked), also important as a mechanism to monitor infection control measures. These cases illustrate the power of combining rapid genomic and epidemiological analyses in near real time. In contrast to previous studies²⁸⁻³¹, we reported results of our investigations to the clinical, infection control and management teams on a weekly basis, thus enabling them to respond to this information and act accordingly within the timeframe of ongoing ward outbreaks. The genomic data informed reviews of patient placement and isolation procedures, assessment of PPE use and staff break arrangements, supporting us to better focus efforts at a time of unprecedented demand on infection control teams. Finally, these analyses are currently being used to inform existing patient safety review processes within our hospital, including investigations related to hospital-onset COVID-19 where the patient has come to harm.

Our study highlights the importance of understanding SARS-CoV-2 transmission within healthcare settings in managing the pandemic. The possible transmission networks that we have identified were complex, involving patients and HCW in both community and hospital settings. However, we have also identified potential transmission networks in other environments including care homes, outpatient units and ambulance services, that have been poorly studied. Although there are strong epidemiological and genomic associations between cases, the mechanism and direction of transmissions remain unclear. The role of asymptomatic intermediates, fomites (including PPE) and the environment are not well understood and require further investigation. During the time frame of this study a number of infection control interventions were implemented across the hospital, as well as national public health measures to reduce community spread. Understanding the interaction between such interventions and nosocomial transmission are complex (especially in the context of the comparably long incubation period for SARS-CoV-2 relative to other respiratory viruses), but critical in enabling healthcare providers to safely deliver existing services in the context of a pandemic.

We acknowledge several limitations to our study. Firstly, we were unable to sequence all of the genomes from samples that were collected during the study period. We may therefore have missed the opportunity to investigate all potential transmission events. Furthermore, we only had main ward location data for most of the HCW and could have overlooked potential epidemiological links with patients with whom they had contact on other wards. Similarly, we were unable to track HCW movements outside their main ward location to identify potential contact with patients on other wards or HCW within shared communal areas. Finally, due to the recent introduction of SARS-CoV-2 into the human population, highly similar genomes are not sufficient evidence to infer a recent transmission, in the absence of confirmatory epidemiological data. However, our experience indicates that further investigation of genomic clusters with highly similar genomes can uncover previously unknown epidemiological links. Furthermore, we were able to use rapidly generated genomic data to investigate HCAI within the hospital setting. The next initiative is to expand this platform into a more detailed analysis of infections in HCW and in community settings, such as residential homes. As the practical challenges associated with implementing real-time

genome sequencing during epidemics are overcome, unlocking the real power of genomic epidemiology will require its integration with clinical and public health systems to support decision-making on local, national and international scales.

Acknowledgments

We acknowledge the assistance of the laboratory staff of CMPHL for processing the diagnostic samples and the clinical teams (infectious diseases, microbiology, virology, infection control) at CUH for their assistance with the investigation of healthcare-associated infections. We are grateful to Richard Smith (CUH Patient Safety Lead), Lucy Rivett, Dominic Sparkes, Nick K. Jones and Matthew Routledge (for providing some HCW data), and Anthony Underwood (Centre for Genomic Pathogen Surveillance) for helpful discussions and advice.

Funding

This work was funded COG-UK, which is supported by funding from the Medical Research Council (MRC) part of UK Research & Innovation (UKRI), the National Institute of Health Research (NIHR) and Genome Research Limited, operating as the Wellcome Sanger Institute. It was also supported by the Wellcome Trust (Senior Fellowship 207498/Z/17/Z and ARTIC Network Collaborative Award 206298/B/17/Z to IG, Collaborative Award 204870/Z/16/Z supporting CJH, Senior Research Fellowship to SGB 215515/Z/19/Z, Senior Clinical Research Fellowship 108070/Z/15/Z to MPW), the Academy of Medical Sciences and the Health Foundation (Clinician Scientist Fellowship to MET), and the National Institute for Health Research Cambridge Biomedical Research Centre at the Cambridge University Hospitals NHS Foundation Trust (BW, GD, MET). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

References

1. Zhou P, Yang X Lou, Wang XG, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;579(7798):270–3.
2. Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. The proximal origin of SARS-CoV-2. *Nat. Med.* 2020;26(4):450–2.
3. Duchene S, Featherstone L, Haritopoulou-Sinanidou M, Rambaut A, Lemey P, Baele G. Temporal signal and the phylodynamic threshold of SARS-CoV-2. *bioRxiv* 2020;2020.05.04.077735.
4. Bedford T, Greninger AL, Roychoudhury P, et al. Cryptic transmission of SARS-CoV-2 in Washington State. *medRxiv* 2020;2020.04.02.20051417.
5. Grubaugh ND, Ladner JT, Lemey P, et al. Tracking virus outbreaks in the twenty-first century. *Nat. Microbiol.* 2019;4(1):10–9.
6. Gardy JL, Loman NJ. Towards a genomics-informed, real-time, global pathogen surveillance system. *Nat Rev Genet* 2018;19(1):9–20.
7. Houldcroft CJ, Beale MA, Breuer J. Clinical and biological insights from viral genome sequencing. *Nat Rev Microbiol* 2017;15(3):183–92.
8. Grubaugh ND, Ladner JT, Kraemer MUG, et al. Genomic epidemiology reveals multiple introductions of Zika virus into the United States. *Nature* 2017;546(7658):401–5.
9. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients with 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA - J Am Med Assoc*

- 2020;323(11):1061–9.
10. WHO. Coronavirus disease 2019 (COVID-19) Situation Report – 82 [Internet]. 2020 [cited 2020 May 8]; Available from: <https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200411-sitrep-82-covid-19.pdf>
11. Quick J. artic-ncov2019/primer_schemes/nCoV-2019/V3 at master · artic-network/artic-ncov2019 [Internet]. 2020 [cited 2020 May 6]; Available from: https://github.com/artic-network/artic-ncov2019/tree/master/primer_schemes/nCoV-2019/V3
12. Quick J. nCoV-2019 sequencing protocol [Internet]. 2020 [cited 2020 May 6]; Available from: <https://www.protocols.io/view/ncov-2019-sequencing-protocol-bbmuk6w>
13. Loman N, Rowe W, Rambaut A. nCoV-2019 novel coronavirus bioinformatics protocol [Internet]. 2020 [cited 2020 May 6]; Available from: <https://artic.network/ncov-2019/ncov2019-bioinformatics-sop.html>
14. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Mol Biol Evol* 2013;30(4):772–80.
15. Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 2015;32(1):268–74.
16. Argimón S, Abudahab K, Goater RJE, et al. Microreact: visualizing and sharing data for genomic epidemiology and phylogeography. *Microb genomics* 2016;2(11):e000093.
17. Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. ggtree: an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol Evol* 2017;8(1):28–36.
18. Rambaut A, Holmes EC, Hill V, et al. A dynamic nomenclature proposal for SARS-CoV-2 to assist genomic epidemiology. *bioRxiv* 2020;2020.04.17.046086.
19. Consortium C-U. COVID-19 Genomics UK (COG-UK) Consortium Weekly Report #1 23rd March 2020 [Internet]. 2020 [cited 2020 May 7]; Available from: <https://www.cogconsortium.uk/wp-content/uploads/2020/04/14-COG-UK-Weekly-Report-23rd-March-2020.pdf>
20. Association TR. COVID-19 data [Internet]. 2020 [cited 2020 May 8]; Available from: <https://renal.org/covid-19/data/>
21. Arias A, Watson SJ, Asogun D, et al. Rapid outbreak sequencing of Ebola virus in Sierra Leone identifies transmission chains linked to sporadic cases. *Virus Evol* 2016;2(1).
22. Quick J, Loman NJ, Duraffour S, et al. Real-time, portable genome sequencing for Ebola surveillance. *Nature* 2016;530(7589):228–32.
23. Dudas G, Carvalho LM, Bedford T, et al. Virus genomes reveal factors that spread and sustained the Ebola epidemic. *Nature* 2017;544(7650):309–15.
24. Gardy JL, Naus M, Amlani A, et al. Whole-genome sequencing of measles virus genotypes H1 and D8 during outbreaks of infection following the 2010 Olympic Winter Games reveals viral transmission routes. *J Infect Dis* 2015;212(10):1574–8.
25. MacFadden DR, McGeer A, Athey T, et al. Use of genome sequencing to define institutional influenza outbreaks, Toronto, Ontario, Canada, 2014–15. *Emerg Infect Dis* 2018;24(3):492–7.
26. Andersen KG, Shapiro BJ, Matranga CB, et al. Clinical Sequencing Uncovers Origins and Evolution of Lassa Virus. *Cell* 2015;162(4):738–50.
27. Quick J, Grubaugh ND, Pullan ST, et al. Multiplex PCR method for MinION and Illumina sequencing of Zika and other virus genomes directly from clinical samples. *Nat Protoc* 2017;12(6):1261–6.
28. Köser CU, Holden MTG, Ellington MJ, et al. Rapid Whole-Genome Sequencing for Investigation of a Neonatal MRSA Outbreak. *N Engl J Med* 2012;366(24):2267–75.
29. Harris SR, Cartwright EJP, Török ME, et al. Whole-genome sequencing for analysis of an outbreak of meticillin-resistant *Staphylococcus aureus*: A descriptive study. *Lancet Infect*

- Dis 2013;13(2):130–6.
30. Coll F, Harrison EM, Toleman MS, et al. Longitudinal genomic surveillance of MRSA in the UK reveals transmission patterns in hospitals and the community. *Sci Transl Med* 2017;9(413).
31. Houldcroft CJ, Roy S, Morfopoulou S, et al. Use of Whole-Genome Sequencing of Adenovirus in Immunocompromised Pediatric Patients to Identify Nosocomial Transmission and Mixed-Genotype Infection. *J Infect Dis* 2018;218(8):1261–71.
32. Kalyaanamoorthy S, Minh BQ, Wong TKF, Von Haeseler A, Jermiin LS. ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nat Methods* 2017;14(6):587–9.
33. Paradis E, Schliep K. Ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* 2019;35(3):526–8.

Figure 1. Study flow diagram.

*Possible healthcare-associated infections. This category includes (i) Community onset suspected healthcare-associated; (ii) Hospital onset indeterminate healthcare-associated; (iii) Hospital onset suspected healthcare-associated; (iv) healthcare workers tested as part of routine clinical care. Out of 1,000 sequenced genomes, 166 failed QC (2 were removed as <29Kb, 164 removed as >2,990N)

Table 1. Baseline characteristics of COVID-19 patients at CUH

Baseline characteristics	
Number of patients	374
Age in years, mean (range) Age in years, median (IQR)	64 (0-98) 67 (51-79)
Male sex	233 (62.3%)
Female sex	141 (37.7%)
Ethnicity – White	267 (71.4%)
Ethnicity – Black, Asian and minority ethnic	28 (7.5%)
Ethnicity – not stated/missing	79 (21.1%)
Co-morbidities	
Hypertension	115 (31.3%)
Ischaemic heart disease	53 (14.4%)
Cardiac failure	25 (6.8%)
Asthma	45 (12.8%)
Chronic obstructive pulmonary disease	30 (8.2%)
Diabetes mellitus	78 (21.2%)
Chronic kidney disease	37 (9.9%)
Hepatic cirrhosis	16 (4.4%)
Dementia	32 (8.7%)

Obesity	51 (13.9%)
Classification of infection*	
Community onset, community associated	261 (69.9%)
Community onset, suspected healthcare-associated	40 (10.7%)
Hospital onset, indeterminate healthcare-associated	11 (2.9%)
Hospital onset, suspected healthcare-associated	13 (3.5%)
Hospital onset, healthcare-associated	39 (10.5%)
Healthcare worker	9 (2.4%)
Unknown	1 (0.3%)
Outcome	
Admission to hospital	347 (92.5%)
Admission to critical care	72 (19.3%)
Died as an inpatient	75 (20.1%)

Definitions of COVID-19 infections

1. Community onset, community associated = diagnostic sample positive within 48 hours of admission and no healthcare contact during the previous 14 days
2. Community onset, suspected healthcare-associated = diagnostic sample positive within 48 hours of admission with healthcare contact during the previous 14 days
3. Hospital onset, indeterminate healthcare-associated = diagnostic sample positive after 48 hours but less than 7 days post-admission
4. Hospital onset, suspected healthcare-associated = diagnostic sample positive 7 to 14 days post admission
5. Hospital onset, healthcare-associated = diagnostic sample positive >14 days post-admission
6. Healthcare worker

Figure 2. Epidemic curve of COVID-19 infections at CUH

Classification of infection: 1. Community onset, community associated; 2. Community onset, suspected healthcare-associated; 3. Hospital onset, indeterminate healthcare-associated; 4. Hospital onset, suspected healthcare-associated; 5. Hospital onset, healthcare-associated; 6. Healthcare worker

Figure 3. SARS-CoV-2 lineages identified in the East of England and CUH.

3A. Map of the EoE showing the distribution of SARS CoV-2 lineages in different sites

Map of East of England showing the different hospital sites (in circles) and the breakdown on SARS-CoV-2 lineages by site.

3B. Comparison of the frequency of SARS-CoV-2 lineages in CUH versus other sites in the EoE

Frequency distribution of SARS-CoV-2 lineages in CUH (blue) versus other sites in the East of England (shown in red). One sample was excluded due to missing location metadata.

Figure 4. Phylogenetic tree and epidemiological timeline from CUH

4A. Phylogenetic tree of SARS-CoV-2 genomes from CUH

Phylogenetic tree of 299 SARS-CoV-2 genomes from CUH and 30 international genomes (Supplementary Methods, Supplementary Figure 4). The tree is rooted on a December 2019 genome from Wuhan, China. The left hand column highlights the hospital and community clusters in colours. Wards A, B and C all had clusters of HAI cases with viruses <2 SNPs different. Eight Ward C cases are contained within one of the largest clusters of identical viruses (n=15). Genomic clusters containing the cases for Wards B and C, the dialysis unit and Care home A all include HCWs. The right hand column shows the classification of infection: 1. Community onset, community associated; 2. Community onset, suspected healthcare-associated; 3. Hospital onset, indeterminate healthcare-associated; 4. Hospital onset, suspected healthcare-associated; 5. Hospital onset, healthcare-associated; 6. Healthcare worker; 7. Unable to determine/ data missing.

4B. Epidemiological timeline of ward A cluster

Six patients on ward A were diagnosed with COVID-19 infection. All had been on the ward for >14 days prior to their specimen date and were considered likely to be hospital-acquired infections. The date of the first positive sample collection is shown with a red box. Five of the viral genomes (Patients A to E) had zero SNP differences between them, and one of the viral genomes (Patient F) differed by 1 SNP.

Supplementary Methods

SARS-CoV-2 molecular testing

Nucleic acid extraction was undertaken using the NUCLEISENS easyMAG platform (BioMerieux, Marcy L'Etoile), in accordance with manufacturers' instructions. Nucleic acids were extracted from 500µL of sample, with a dilution of MS2 bacteriophage added pre-extraction to act as an internal extraction and inhibition control. The presence of SARS-CoV-2 was assessed using an in-house generated and validated one-step RT q-PCR assay that detects a 222 base-pair region of the SARS-CoV-2 RdRp genes, along with an MS2 bacteriophage internal extraction control. The RdRp gene was detected using the RdRp For primer (ATGGGTTGGGATTATCCTAAATGTGA) and the RdRp Rev primer (AGCAGTTGTGGCATCTCCTGATGAG) with a FAM labelled MGB RdRp Probe 3 (ATGCTTAGAATTATGGCCTCAC). The internal extraction control was detected using the MS2 For primer (TGGCACTACCCCTCTCCGTATTCACG), the MS2 Rev primer (GTACGGGCGACCCACGATGAC) and a ROX-BHQ2 labelled MS2 probe (CACATCGATAGATCAAGGTGCCTACAAGC). Amplification reactions and detection of PCR products were performed using the Rotorgene™ PCR instrument. A typical reaction contained 400nM of For and Rev primers for the RdRp genes and 200nM of the the MS2 internal control For and Rev primer pair, along with 120nM of the RdRp and MS2 probes. Reactions typically contained 25% extracted nucleic acid and were cycled through the following conditions 25°C 2min, 50°C 15min, 95°C 2min followed by 45 cycles of 95°C and 60°C. Samples that generated a Ct value ≤36 were considered positive. Samples and negative control (molecular grade water) were individually spiked with MS2 bacteriophage internal control (4600 pfu per extraction) prior to nucleic acid extraction to identify any inhibitors or extraction issues. Positive control material, BetaCoV/England/02/2020, was obtained from PHE Colindale and was essentially purified virus RNA diluted down to give a cycle threshold value of 26-28. Negative controls included extracted molecular grade water.

Sample and data collection

Sample collection, metadata curation and linkage to sequencing IDs involved coordination between multiple teams and critically depended on good relations between the sequencing and diagnostic laboratory staff (Supplementary Figure 1). Each day samples with positive SARS-CoV-2 PCR results from the last 24 to 72 hours were identified from the hospital information system and clinical metadata was extracted, formatted and integrated into a master metadata file. A 15 microlitre aliquot of the RNA extract of all CUH samples and a random selection of EoE samples were collected from the diagnostic microbiology laboratory for local sequencing. These were assigned COG-UK sequencing codes which were integrated back into the master metadata file. For samples that were not sequenced locally, the remaining RNA extract was collected from the diagnostic microbiology laboratory and sent to the Wellcome Sanger Institute for sequencing as part of the COG-UK consortium. Each week, clinical metadata and sequencing data were combined and formatted for upload to the MRC CLIMB system. Data manipulations were performed in R (v 3.6.2) using the *tidyverse* packages (v 1.3.0) installed onto computers within the Trust network.

Sequencing details

Samples were sequenced using Nanopore technology following the ARTICnetwork V3 protocol (<https://dx.doi.org/10.17504/protocols.io.bbmuik6w>) and assembled using the ARTICnetwork assembly pipeline (<https://artic.network/ncov-2019/ncov2019-bioinformatics-sop.html>). Median genome depth of coverage was 6,612x across all 747 genomes. 14 samples in our dataset were also sequenced with Illumina technology at the Wellcome Sanger Institute as part of COG-UK. There was 100% concordance in called nucleotides between sample pairs. Four genomes differed because of base pairs called in the Illumina data that were missing in the Nanopore sequences. The accession numbers of the samples included in this study are available in Supplementary Table 4.

Bioinformatic analysis

Consensus fasta sequence quality control cutoffs were: size >29Kb, N count <2990 (~10%). After QC filtering, de-duplication and matching with metadata, the first sample set analysed comprised 197 genomes collected up to 10th April 2020; set 2 had 444 genomes up to 15th April, and set 3 (presented here) had 747 genomes up to 24th April. 30 reference genomes were added to the sample sets downloaded from GISAID (<https://www.gisaid.org/>; Supplementary Table 3). The reference genomes were chosen to represent the major branches of the global phylogenetic tree as visualised in Nextstrain (<https://nextstrain.org/>) to provide broader context, including a sample from December 2019 collected in Wuhan, China, used to root the tree. Multiple sequence alignment was performed using MAFFT (v 7.458) with default settings, command:

```
/PATH/mafft" --retree 2 --inputorder "multi_fasta.fasta" > "aligned_multi_fasta"
```

The alignment was manually inspected using AliView. Maximum likelihood trees were produced using IQ-TREE software¹⁵ for all samples passing QC filters and the subset of samples from CUH (n=299 for this dataset). Initial tests with the ModelFinder Plus option³², which selects the optimal nucleotide substitution model out of over 200 options

(<http://www.iqtree.org/doc/Substitution-Models>), consistently identified GTR+F+I as the best model. Therefore from 24th April (including analysis presented here) we specified GTR+F+I.

Command using ModelFinder Plus:

```
/PATH/iqtree -s aligned_filtered_multi_fasta -m MFP
```

Command with GTR+F+I model specified:

```
/PATH/iqtree -s aligned_filtered_multi_fasta -m GTR+F+I
```

Trees were manually inspected in FigTree, rooted on the 2019 Wuhan sample (EPI_ISL_402123), ordered by descending node and exported as Newick files. Trees were visualised in online software Microreact¹⁶ in a private account to explore relationships between wards and clinico-epidemiological questions for our weekly reports. Further visualisations were produced in R using the packages *Ape*³³ (v 5.3) and *ggtree*¹⁷ (v 2.0.4).

A SNP difference matrix was produced from the multiple sequence alignment using the *snp-dists* package (v 0.7.0; <https://github.com/tseemann/snp-dists>; installed into a conda environment), command:

```
snp-dists -c aligned_filtered_multi_fasta.aln > snp_dist_matrix.csv
```


The matrix was exported as .csv and manipulated in R using the *Matrix* and *tidyverse* packages for ward and pairwise SNP comparisons and plotted using the *ggplot2* (v 3.3.0) package. A heatmap was produced in python using the seaborn (v 0.10.0) clustermap function. To identify clusters with zero SNP differences we used the scipy.cluster.hierarchy functions *linkage* and *fcluster* setting a maximum distance of zero (scipy v 1.4.1). Clusters were named in descending size order and linked with sample metadata and lineage data.

Investigation of genetic clusters with zero SNP differences

Patient records from each case within a putative genomic cluster were manually reviewed in detail by authors BW, WLH and MET and assigned a score of 1 (strong evidence supporting a recent linked transmission chain, e.g. patients co-located on the same ward becoming positive within the incubation period of the virus), 2 (a plausible transmission chain is present e.g. patients becoming positive while located in nearby wards within the hospital but who did not appear to be in direct contact), and 3 (no evidence of any transmission connections between cases).

Epidemiological analysis methods

Timeline plotting

Space time relationships between patients were plotted using patient specific time lines by exporting the bed and ward admission dates, dates of transfer, dates of discharge or date of death obtained from hospital information system (EPIC Systems Corporation, Verona, USA) and importing them into a cloud-based timeline plotter application (Cluster Track, Camart Ltd, Cambridge available at Clusterack.com). Earliest positive specimen date for COVID19 was obtained from the laboratory records and date of onset of symptoms from the clinical records and uploaded.

The application aided visualisation of ward and time relations by assigning unique colours to wards and then ward presence by date along the x axis shown in days, such that a solid timeline bar by colour and by date permitted the visualisation of the location of each patient over time. The positive specimen date for COVID 19, genomic cluster, death and discharge were each overlaid on the patient timeline using standard visual representation built into the application. Visualisation was aided by using the sort command within the application on admission date, earliest positive specimen date, or first ward to which admitted. Separate plots of subset of the total cases were created to provide clearer visualisation when needed

Ward time and genomic cluster plots

A clustering and network analysis function was used in the Cluster track application in which an algorithm links patients with admission days to the same ward on the same date and displays a network diagram to indicate these overlapping cases.

More advanced space time clustering was undertaken by exporting these timeline data sets into an SQL database running a more advanced clustering algorithm in which time parameters were set for the presumed susceptible, infectious and non-infectious/ recovered intervals counting from the earliest positive specimen date. The algorithm identified and linked cases in which two or more patients had an overlap on the same ward of the time interval of infectiousness of an earlier

case with the interval of susceptibility in a later case or cases. Links continue to be made until no further overlaps of the infectious interval in an earlier case occurred with the interval of susceptibility in a later case on the same ward: this ended the space time cluster.

The cluster diagrams of the space time clustered cases were reviewed. Cases within the same space time cluster that belonged to the same genomic cluster were deemed to be supportive of a recent transmission event.

Supplementary Figures

Supplementary Figure 1. Data and Sample Processing at CUH

HAI = Hospital Acquired Infection
HCW = Healthcare Worker

Supplementary Figure 2.

2A. Ct value plotted against percentage coverage for 947 SARS-CoV-2 genomes

The Ct value plotted against the percent of SARS-CoV-2 genome sequenced prior to internal screening and for which a Ct value is available (N=947)

2B. Location and frequency of SNPs across sequenced genomes.

Cumulative number and location of SNPs compared to the original Wuhan strain (Accession No. MN908947) observed across 747 genomes sequenced in East of England. This shows the total occurrence of SNPs, 10,536nt across 22,337,541nt sequenced (0.005%) occurring at 1,196 positions. Of the 1,196 positions, 1,192 SNPs were found to be single SNPs, while 4 sites had 2 SNPs. 5 common SNPs were found in the majority of sequenced genomes (A23403G, C14408T, C241T, C3037T, T deletion at 24981) while G28881A, G28882A, G28883 were also found in ~50% of samples. These are not unique mutations and have been observed in other cases in the global NextStrain analysis.

Supplementary Figure 3. SARS-CoV-2 lineages over time

In the weeks commencing 9 and 13 March 2020 lineages (and descendent lineages) of B, B.1, B.2, B.3 and B.8 were present in the EoE (amalgamated here as there were only 2 samples for week commencing 9 March). Diversification of lineages already present from earlier weeks was seen over time, with the detection of descendants of lineages B.1 and B.2, but no new lineages emerged during this period, likely an impact of lockdown measures preventing new viruses being introduced from other regions. Changes in lineage frequency may be stochastic due to changes in the available sample size during each week of the sampling period. Lineage B.8 was only detected in the week commencing 16 March 2020. Lineage B.4 viruses (associated with export from Iran) were not seen in our sample set. Lineage A viruses (or A descendants), most commonly reported in China, USA, South Korea and Australia, were not detected in our EoE samples.

Supplementary Figure 4. Phylogenetic trees of East of England and CUH genomes

4A. Phylogenetic tree of East of England SARS-CoV-2 genomes

Phylogenetic tree of 747 East of England SARS-CoV-2 genomes and 30 reference genomes used to provide further genetic and geographic context. The reference genomes are highlighted with coloured tips and are the same used in Figure 4A. As with Figure 4A, the tree is rooted on a December 2019 sample from Wuhan, China, with older samples from Asia represented at the base of the tree as expected. The lineages are indicated by the colour bar.

4B. Phylogenetic tree of CUH SARS-CoV-2 genomes

Phylogenetic tree of 299 CUH SARS-CoV-2 genomes and 30 reference genomes. The inner ring shows emergency department (ED) samples in blue and samples collected from all other sites in grey. The outer ring shows the different classifications of infection: 1. Community onset, community associated; 2. Community onset, suspected healthcare-associated; 3. Hospital onset, indeterminate healthcare-associated; 4. Hospital onset, suspected healthcare-associated; 5. Hospital onset, healthcare-associated; 6. Healthcare worker; 7. Unable to determine/ data missing.

Supplementary Figure 5. SNP difference matrix, distribution and box plots

5A. SNP difference matrix

SNP difference matrix for 299 SARS-CoV-2 genomes from CUH. Darker colouring indicates more similar genomes. Several clusters of identical viruses are present in the dataset, as discussed in main text. The left hand bar shows wards A, B, C and the dialysis unit highlighted in colour and other wards in grey

5B. Distribution of SNP differences among CUH samples

Distribution of pairwise SNP differences for 299 CUH samples. The total number of pairwise comparisons is 44,551. The median difference was 8 SNPs (range 0 to 24 SNPs). 4.5% of genomes were identical or 1-SNP different.

5C. Box plot of SNP differences between SARS-CoV-2 genomes

Box plot of SNP differences between SARS-CoV-2 genomes at CUH within different sampling locations. The number of SNP differences was very low on certain wards (0 to 1 SNPs) compared with the emergency department (ED), the Trust (CUH) and the East of England (EoE) as a whole, consistent with shared recent transmission events on these wards (discussed in main text).

Supplementary Figure 6. Process for investigating healthcare associated COVID-19 infections at Cambridge University Hospitals NHS Foundation Trust

Supplementary Figure 7. Epidemiological timelines of hospital clusters

7A. Hospital cluster 2 (Ward B)

Four transplant patients on ward B (shown in khaki) were diagnosed with COVID-19 infection between 3 and 20 April 2020. A fifth patient, who had been recently discharged from the ward, presented to the ED with COVID-19 infection. The sample dates are shown in yellow circles (patients) and diamonds (HCW). Genomic analysis revealed that all 5 patients had identical genomes. Two HCW were found to have identical genomes in the same cluster as the ward B cases; one had worked on ward B and had professional contact with the other HCW.

7B. Dialysis unit cluster

Six patients with end-stage renal failure were diagnosed with COVID-19 between 1 and 20 April 2020, testing positive in several locations including ED and an acute admissions ward. The sample dates are shown in yellow circles. Their viral genomes were identical, and epidemiological investigation revealed they dialysed at the same outpatient dialysis unit. This suggests linked recent transmission of community-onset healthcare-associated infections.

Supplementary Tables

Supplementary Table 1. CUH COVID-19 infections and sequence data availability

Classification of infection	No.	No. with available sequence (%)
Community onset, community associated	261	174 (66.7%)
Community onset, suspected healthcare-associated	40	27 (67.5%)
Hospital onset, indeterminate healthcare-associated	11	7 (63.6%)
Hospital onset, suspected healthcare-associated	13	11 (84.6%)
Hospital onset, healthcare-associated	39	36 (92.3%)
Healthcare worker	9	6 (66.7%)
Unknown	1	1 (100%)
Total	374	262 (70.1%)

Supplementary Table 2. Transmission cluster table

No.	Size	CAIs	Possible HAIs*	HAIs	HCWs	First Case	Last Case	Cases with good epi links	Cases with possible epi links	Cases with no epi links	Notes
1	15	4	4	5	2	20/3/20	16/4/20	14	0	1	Eight patients swabbed during a suspected outbreak on ward 1. The first patient to be swabbed was recently discharged from ward 2 and then re-admitted to ward 3. The two HCWs worked on ward 3; one had direct contact with the first patient to be swabbed in this cluster. Two patients, distinct from the larger outbreak, co-habit (spouses). The final patient has no identifiable association with any other case in this cluster.
2	15	10	1	0	4	27/3/20	20/4/20	9	5	1	8 patients were resident in the care home A. One patient works as a carer in the same home. Another patient also works as a carer in an unspecified home. Two of the HCWs were paramedics. The other two HCWs work in different clinical areas, but both live with paramedics. One patient had no identifiable contact with the other cases in this cluster.

3	10	1	2	3	4	1/4/20	20/4/20	7	1	2	Four patients swabbed during a suspected outbreak on ward 4 (ward B); a fifth patient was discharged from ward 4 and re-admitted with COVID within 2 weeks of discharge. Two HCWs work on ward 4 and a third works in the same department as one of these HCWs. There are no identified associations between the remaining patient or the fourth HCW.
4	8	7	0	0	1	20/3/20	17/4/20	1	0	7	No identified epidemiological link between the seven patients. The HCW had direct contact with two of these patients.
5	5	0	3	2	0	27/3/20	11/4/20	5	0	0	Five patients all swabbed on ward 5 (ward A)
6	5	4	0	1	0	27/3/20	19/4/20	0	0	5	No identified epidemiological link between the 5 patients.
7	5	0	4	0	1	20/3/20	16/4/20	4	1	0	Three patients swabbed during a suspected outbreak on ward 2. HCW works on ward 2. The first patient in this cluster was recently discharged from ward 8 and readmitted to ward 8. Wards 2 and 8 are in the same department and share some staff
8	5	2	0	0	3	7/4/20	17/4/20	5	0	0	Two patients resident in care home B; one of these patients admitted to ward 6. Two HCWs work on

											ward 6, including one with direct contact with this patient. The third HCW works on ward 8.
9	4	2	0	0	2	4/4/20	16/4/20	4	0	0	One patient resident in care home B, admitted to ward 7. Second patient is a carer in care home B. One HCW works on wards 7 and 8. The second HCW works on ward 8. The isolates from clusters 8 and 9 diverge by 1 SNP, but share epidemiological links of both patients and HCWs.
10	6	0	6	0	0	1/4/20	20/4/20	6	0	0	All six patients receive dialysis at the same unit
11	3	2	1	0	0	9/4/20	17/4/20	0	0	3	Two patients live in separate care homes. The third patient is a community carer but has no other identified association with these patients
12	3	3	0	0	0	11/4/20	20/4/20	0	2	1	One patient is a resident in a specialist dementia care home. A second patient works in a specialist dementia care home, but it is unclear whether this is the same home as the first patient. The third patient has no other identified associations with the other patients.
13	3	0	0	1	2	31/3/20	17/4/20	0	0	3	The first patient acquired infection in hospital. The HCWs have no known association with each other or the patient.

14	3	0	0	2	1	12/4/20	20/4/20	3	0	0	The two patients were co-located on ward 10. The HCW worked on the same ward, but there is no documented direct contact with either patient.
15	2	1	0	0	1	2/4/20	13/4/20	2	0	0	The patient was admitted to ward 6 with documented direct contact with the HCW.
16	2	1	1	0	0	4/4/20	5/4/20	2	0	0	Two patients co-located on ward 9.
17	2	0	0	0	2	16/4/20	17/4/20	0	2	0	Two HCWs, working in separate wards in the same department.
18	2	1	0	0	1	15/4/20	18/4/20	0	0	2	The HCW lives with a care assistant in an unspecified care home. The patient was admitted from a care home. No other identified association
19	2	1	0	0	1	19/4/20	20/4/20	0	0	2	The patient works in a local community hospital. The HCW works in a rehabilitation unit. There are no other known epidemiological associations.
20	2	0	0	1	1	9/4/20	12/4/20	0	0	2	The HCW and patient have moved between multiple ward areas. No direct contact documented.
21	2	1	0	1	0	3/4/20	3/4/20	0	2	0	Two patients briefly co-located within 24 hrs of testing, likely insufficient duration for transmission. One patient was resident with community acquired

											infection in a care home; the second patient was on a rehabilitation ward prior to swabbing
22	2	1	0	1	0	22/3/20	27/3/20	0	0	2	No identified associations between the two patients
23	2	1	1	0	0	30/3/20	19/4/20	0	2	0	No direct associations between two patients, but they were co-located on neighbouring wards (wards 1, 4 and 10), which share some HCWs
24	2	2	0	0	0	29/3/20	7/4/20	0	0	2	No identified associations between the two patients
25	2	0	0	2	0	29/3/20	7/4/20	2	0	0	Two patients co-located on ward 11
26	2	2	0	0	0	1/4/20	2/4/20	2	0	0	Two patients co-habiting (siblings)
Total	114		23	19	25			66	15	33	

Supplementary Table 3. GISAID genomes included in phylogenetic tree

GISAID ID	Country of origin
EPI_ISL_402123*	China
EPI_ISL_406716	China
EPI_ISL_406801	China
EPI_ISL_414598	Spain
EPI_ISL_416396	China
EPI_ISL_416757	France
EPI_ISL_417969	Spain
EPI_ISL_418034	USA
EPI_ISL_419228	China
EPI_ISL_419232	China
EPI_ISL_420064	France
EPI_ISL_421905	UK
EPI_ISL_422024	UK
EPI_ISL_423034	Netherlands
EPI_ISL_424657	Belgium
EPI_ISL_426019	UK

EPI_ISL_427119	Australia
EPI_ISL_427144	Australia
EPI_ISL_427322	Russia
EPI_ISL_427391	Turkey
EPI_ISL_427441	USA
EPI_ISL_428482	India
EPI_ISL_428848	Singapore
EPI_ISL_428857	The Gambia
EPI_ISL_429061	USA
EPI_ISL_429175	Thailand
EPI_ISL_429177	Thailand
EPI_ISL_429206	Switzerland
EPI_ISL_429259	DRC
EPI_ISL_429773	Luxembourg

*Sample from China used to root the tree

Supplementary Table 4. Table of sequences / accession numbers

GISAID Accession ID	Virus name	COG-UK	GISAID Accession ID	Virus name	COG-UK
EPI_ISL_425231	hCoV-19/England/CAMB-7374F/2020	CAMB-7374F	EPI_ISL_433840	hCoV-19/England/CAMB-7AB00/2020	CAMB-7AB00
EPI_ISL_425232	hCoV-19/England/CAMB-7375E/2020	CAMB-7375E	EPI_ISL_433841	hCoV-19/England/CAMB-7AB1F/2020	CAMB-7AB1F
EPI_ISL_425233	hCoV-19/England/CAMB-7376D/2020	CAMB-7376D	EPI_ISL_433842	hCoV-19/England/CAMB-7AB2E/2020	CAMB-7AB2E
EPI_ISL_425234	hCoV-19/England/CAMB-7377C/2020	CAMB-7377C	EPI_ISL_433843	hCoV-19/England/CAMB-7AB3D/2020	CAMB-7AB3D
EPI_ISL_425235	hCoV-19/England/CAMB-7378B/2020	CAMB-7378B	EPI_ISL_433844	hCoV-19/England/CAMB-7AB79/2020	CAMB-7AB79
EPI_ISL_425236	hCoV-19/England/CAMB-7379A/2020	CAMB-7379A	EPI_ISL_433845	hCoV-19/England/CAMB-7AB88/2020	CAMB-7AB88
EPI_ISL_425237	hCoV-19/England/CAMB-737A9/2020	CAMB-737A9	EPI_ISL_433846	hCoV-19/England/CAMB-7AB97/2020	CAMB-7AB97
EPI_ISL_425238	hCoV-19/England/CAMB-737B8/2020	CAMB-737B8	EPI_ISL_433847	hCoV-19/England/CAMB-7ABA6/2020	CAMB-7ABA6
EPI_ISL_425239	hCoV-19/England/CAMB-737C7/2020	CAMB-737C7	EPI_ISL_433848	hCoV-19/England/CAMB-7ABB5/2020	CAMB-7ABB5
EPI_ISL_425240	hCoV-19/England/CAMB-737D6/2020	CAMB-737D6	EPI_ISL_433849	hCoV-19/England/CAMB-7ABC4/2020	CAMB-7ABC4
EPI_ISL_425241	hCoV-19/England/CAMB-737E5/2020	CAMB-737E5	EPI_ISL_433850	hCoV-19/England/CAMB-7ABE2/2020	CAMB-7ABE2
EPI_ISL_425242	hCoV-19/England/CAMB-737F4/2020	CAMB-737F4	EPI_ISL_433851	hCoV-19/England/CAMB-7ABF1/2020	CAMB-7ABF1
EPI_ISL_425243	hCoV-19/England/CAMB-73800/2020	CAMB-73800	EPI_ISL_433852	hCoV-19/England/CAMB-7AC0D/2020	CAMB-7AC0D
EPI_ISL_425244	hCoV-19/England/CAMB-7381F/2020	CAMB-7381F	EPI_ISL_433853	hCoV-19/England/CAMB-7AC1C/2020	CAMB-7AC1C
EPI_ISL_425245	hCoV-19/England/CAMB-7382E/2020	CAMB-7382E	EPI_ISL_433854	hCoV-19/England/CAMB-7AC2B/2020	CAMB-7AC2B
EPI_ISL_425246	hCoV-19/England/CAMB-7386A/2020	CAMB-7386A	EPI_ISL_433855	hCoV-19/England/CAMB-7AC3A/2020	CAMB-7AC3A
EPI_ISL_425247	hCoV-19/England/CAMB-73879/2020	CAMB-73879	EPI_ISL_433856	hCoV-19/England/CAMB-7AC49/2020	CAMB-7AC49
EPI_ISL_425248	hCoV-19/England/CAMB-73888/2020	CAMB-73888	EPI_ISL_433857	hCoV-19/England/CAMB-7AC58/2020	CAMB-7AC58
EPI_ISL_425249	hCoV-19/England/CAMB-738A6/2020	CAMB-738A6	EPI_ISL_433858	hCoV-19/England/CAMB-7AC67/2020	CAMB-7AC67
EPI_ISL_425250	hCoV-19/England/CAMB-738B5/2020	CAMB-738B5	EPI_ISL_433859	hCoV-19/England/CAMB-7AC76/2020	CAMB-7AC76
EPI_ISL_425251	hCoV-19/England/CAMB-738C4/2020	CAMB-738C4	EPI_ISL_433860	hCoV-19/England/CAMB-7AC85/2020	CAMB-7AC85
EPI_ISL_425252	hCoV-19/England/CAMB-738D3/2020	CAMB-738D3	EPI_ISL_433861	hCoV-19/England/CAMB-7AC94/2020	CAMB-7AC94
EPI_ISL_425253	hCoV-19/England/CAMB-738E2/2020	CAMB-738E2	EPI_ISL_433862	hCoV-19/England/CAMB-7ACA3/2020	CAMB-7ACA3
EPI_ISL_425254	hCoV-19/England/CAMB-738F1/2020	CAMB-738F1	EPI_ISL_433863	hCoV-19/England/CAMB-7ACB2/2020	CAMB-7ACB2
EPI_ISL_425255	hCoV-19/England/CAMB-7391C/2020	CAMB-7391C	EPI_ISL_433864	hCoV-19/England/CAMB-7ACC1/2020	CAMB-7ACC1
EPI_ISL_425256	hCoV-19/England/CAMB-7393A/2020	CAMB-7393A	EPI_ISL_433865	hCoV-19/England/CAMB-7ACD0/2020	CAMB-7ACD0
EPI_ISL_425257	hCoV-19/England/CAMB-73949/2020	CAMB-73949	EPI_ISL_433866	hCoV-19/England/CAMB-7ACFE/2020	CAMB-7ACFE
EPI_ISL_425258	hCoV-19/England/CAMB-73976/2020	CAMB-73976	EPI_ISL_433867	hCoV-19/England/CAMB-7AD0A/2020	CAMB-7AD0A
EPI_ISL_425259	hCoV-19/England/CAMB-73985/2020	CAMB-73985	EPI_ISL_433868	hCoV-19/England/CAMB-7AD28/2020	CAMB-7AD28
EPI_ISL_425260	hCoV-19/England/CAMB-73994/2020	CAMB-73994	EPI_ISL_433869	hCoV-19/England/CAMB-7AD37/2020	CAMB-7AD37
EPI_ISL_425261	hCoV-19/England/CAMB-739A3/2020	CAMB-739A3	EPI_ISL_433870	hCoV-19/England/CAMB-7AD46/2020	CAMB-7AD46
EPI_ISL_425262	hCoV-19/England/CAMB-739B2/2020	CAMB-739B2	EPI_ISL_433871	hCoV-19/England/CAMB-7AD55/2020	CAMB-7AD55
EPI_ISL_425263	hCoV-19/England/CAMB-739D0/2020	CAMB-739D0	EPI_ISL_433872	hCoV-19/England/CAMB-7AD64/2020	CAMB-7AD64
EPI_ISL_425264	hCoV-19/England/CAMB-739EF/2020	CAMB-739EF	EPI_ISL_433873	hCoV-19/England/CAMB-7AD82/2020	CAMB-7AD82
EPI_ISL_425265	hCoV-19/England/CAMB-739FE/2020	CAMB-739FE	EPI_ISL_433874	hCoV-19/England/CAMB-7AD91/2020	CAMB-7AD91
EPI_ISL_425266	hCoV-19/England/CAMB-73A0A/2020	CAMB-73A0A	EPI_ISL_433875	hCoV-19/England/CAMB-7ADA0/2020	CAMB-7ADA0
EPI_ISL_425267	hCoV-19/England/CAMB-73A19/2020	CAMB-73A19	EPI_ISL_433876	hCoV-19/England/CAMB-7ADBF/2020	CAMB-7ADBF
EPI_ISL_425268	hCoV-19/England/CAMB-73A28/2020	CAMB-73A28	EPI_ISL_433877	hCoV-19/England/CAMB-7ADDD/2020	CAMB-7ADDD
EPI_ISL_425269	hCoV-19/England/CAMB-73A37/2020	CAMB-73A37	EPI_ISL_433878	hCoV-19/England/CAMB-7ADEC/2020	CAMB-7ADEC
EPI_ISL_425270	hCoV-19/England/CAMB-73A46/2020	CAMB-73A46	EPI_ISL_433879	hCoV-19/England/CAMB-7ADF8/2020	CAMB-7ADF8
EPI_ISL_425271	hCoV-19/England/CAMB-73A64/2020	CAMB-73A64	EPI_ISL_433880	hCoV-19/England/CAMB-7AE07/2020	CAMB-7AE07
EPI_ISL_425272	hCoV-19/England/CAMB-73A73/2020	CAMB-73A73	EPI_ISL_433881	hCoV-19/England/CAMB-7AE25/2020	CAMB-7AE25
EPI_ISL_425273	hCoV-19/England/CAMB-73A82/2020	CAMB-73A82	EPI_ISL_433882	hCoV-19/England/CAMB-7AE34/2020	CAMB-7AE34
EPI_ISL_425274	hCoV-19/England/CAMB-73AA0/2020	CAMB-73AA0	EPI_ISL_433883	hCoV-19/England/CAMB-7AE43/2020	CAMB-7AE43
EPI_ISL_425275	hCoV-19/England/CAMB-73ABF/2020	CAMB-73ABF	EPI_ISL_433884	hCoV-19/England/CAMB-7AE52/2020	CAMB-7AE52
EPI_ISL_425276	hCoV-19/England/CAMB-73ACE/2020	CAMB-73ACE	EPI_ISL_433885	hCoV-19/England/CAMB-7AE61/2020	CAMB-7AE61
EPI_ISL_425277	hCoV-19/England/CAMB-73ADD/2020	CAMB-73ADD	EPI_ISL_433886	hCoV-19/England/CAMB-7AE70/2020	CAMB-7AE70
EPI_ISL_425278	hCoV-19/England/CAMB-73AEC/2020	CAMB-73AEC	EPI_ISL_433887	hCoV-19/England/CAMB-7AE9E/2020	CAMB-7AE9E
EPI_ISL_425279	hCoV-19/England/CAMB-73AFB/2020	CAMB-73AFB	EPI_ISL_433888	hCoV-19/England/CAMB-7AEA9/2020	CAMB-7AEA9
EPI_ISL_433666	hCoV-19/England/CAMB-73B16/2020	CAMB-73B16	EPI_ISL_433889	hCoV-19/England/CAMB-7AECB/2020	CAMB-7AECB
EPI_ISL_433667	hCoV-19/England/CAMB-73B61/2020	CAMB-73B61	EPI_ISL_433890	hCoV-19/England/CAMB-7AEDA/2020	CAMB-7AEDA
EPI_ISL_433668	hCoV-19/England/CAMB-73B70/2020	CAMB-73B70	EPI_ISL_433891	hCoV-19/England/CAMB-7AEE9/2020	CAMB-7AEE9
EPI_ISL_433669	hCoV-19/England/CAMB-73B8F/2020	CAMB-73B8F	EPI_ISL_433892	hCoV-19/England/CAMB-7AEF8/2020	CAMB-7AEF8
EPI_ISL_433670	hCoV-19/England/CAMB-73B9E/2020	CAMB-73B9E	EPI_ISL_433893	hCoV-19/England/CAMB-7AF04/2020	CAMB-7AF04
EPI_ISL_433671	hCoV-19/England/CAMB-73BAD/2020	CAMB-73BAD	EPI_ISL_433894	hCoV-19/England/CAMB-7AF13/2020	CAMB-7AF13
EPI_ISL_433672	hCoV-19/England/CAMB-73BBC/2020	CAMB-73BBC	EPI_ISL_433895	hCoV-19/England/CAMB-7AF22/2020	CAMB-7AF22
EPI_ISL_433673	hCoV-19/England/CAMB-73BCB/2020	CAMB-73BCB	EPI_ISL_433896	hCoV-19/England/CAMB-7AF31/2020	CAMB-7AF31
EPI_ISL_433674	hCoV-19/England/CAMB-73BDA/2020	CAMB-73BDA	EPI_ISL_433897	hCoV-19/England/CAMB-7AF40/2020	CAMB-7AF40
EPI_ISL_433675	hCoV-19/England/CAMB-73BF8/2020	CAMB-73BF8	EPI_ISL_433898	hCoV-19/England/CAMB-7AF5F/2020	CAMB-7AF5F
EPI_ISL_433676	hCoV-19/England/CAMB-73C04/2020	CAMB-73C04	EPI_ISL_433899	hCoV-19/England/CAMB-7AF6E/2020	CAMB-7AF6E
EPI_ISL_433677	hCoV-19/England/CAMB-73C13/2020	CAMB-73C13	EPI_ISL_433900	hCoV-19/England/CAMB-7AF7D/2020	CAMB-7AF7D
EPI_ISL_433678	hCoV-19/England/CAMB-73C31/2020	CAMB-73C31	EPI_ISL_433901	hCoV-19/England/CAMB-7AF8C/2020	CAMB-7AF8C
EPI_ISL_433679	hCoV-19/England/CAMB-73C40/2020	CAMB-73C40	EPI_ISL_433902	hCoV-19/England/CAMB-7AF9A/2020	CAMB-7AF9A
EPI_ISL_433680	hCoV-19/England/CAMB-73C5F/2020	CAMB-73C5F	EPI_ISL_433903	hCoV-19/England/CAMB-7AFB9/2020	CAMB-7AFB9
EPI_ISL_433681	hCoV-19/England/CAMB-73C7D/2020	CAMB-73C7D	EPI_ISL_433904	hCoV-19/England/CAMB-7AFF5/2020	CAMB-7AFF5
EPI_ISL_433682	hCoV-19/England/CAMB-73C9B/2020	CAMB-73C9B	EPI_ISL_433905	hCoV-19/England/CAMB-7B000/2020	CAMB-7B000
EPI_ISL_433683	hCoV-19/England/CAMB-73CAA/2020	CAMB-73CAA	EPI_ISL_433906	hCoV-19/England/CAMB-7B01F/2020	CAMB-7B01F
EPI_ISL_433684	hCoV-19/England/CAMB-73CC8/2020	CAMB-73CC8	EPI_ISL_433907	hCoV-19/England/CAMB-7B02E/2020	CAMB-7B02E
EPI_ISL_433685	hCoV-19/England/CAMB-73CD7/2020	CAMB-73CD7	EPI_ISL_433908	hCoV-19/England/CAMB-7B05B/2020	CAMB-7B05B
EPI_ISL_433686	hCoV-19/England/CAMB-73D01/2020	CAMB-73D01	EPI_ISL_433909	hCoV-19/England/CAMB-7B06A/2020	CAMB-7B06A
EPI_ISL_433687	hCoV-19/England/CAMB-73D10/2020	CAMB-73D10	EPI_ISL_433910	hCoV-19/England/CAMB-7B079/2020	CAMB-7B079
EPI_ISL_433688	hCoV-19/England/CAMB-73D2F/2020	CAMB-73D2F	EPI_ISL_433911	hCoV-19/England/CAMB-7B088/2020	CAMB-7B088
EPI_ISL_433689	hCoV-19/England/CAMB-73D3E/2020	CAMB-73D3E	EPI_ISL_433912	hCoV-19/England/CAMB-7B097/2020	CAMB-7B097
EPI_ISL_433690	hCoV-19/England/CAMB-73D4D/2020	CAMB-73D4D	EPI_ISL_433913	hCoV-19/England/CAMB-7B0A6/2020	CAMB-7B0A6
EPI_ISL_433691	hCoV-19/England/CAMB-73D5C/2020	CAMB-73D5C	EPI_ISL_433914	hCoV-19/England/CAMB-7B0B5/2020	CAMB-7B0B5
EPI_ISL_433692	hCoV-19/England/CAMB-73D6B/2020	CAMB-73D6B	EPI_ISL_433915	hCoV-19/England/CAMB-7B0C4/2020	CAMB-7B0C4
EPI_ISL_433693	hCoV-19/England/CAMB-73D7A/2020	CAMB-73D7A	EPI_ISL_433916	hCoV-19/England/CAMB-7B0D3/2020	CAMB-7B0D3
EPI_ISL_433694	hCoV-19/England/CAMB-73D89/2020	CAMB-73D89	EPI_ISL_433917	hCoV-19/England/CAMB-7B0E2/2020	CAMB-7B0E2
EPI_ISL_433695	hCoV-19/England/CAMB-73D98/2020	CAMB-73D98	EPI_ISL_433918	hCoV-19/England/CAMB-7B0F1/2020	CAMB-7B0F1
EPI_ISL_433696	hCoV-19/England/CAMB-73DA7/2020	CAMB-73DA7	EPI_ISL_433919	hCoV-19/England/CAMB-7B10D/2020	CAMB-7B10D
EPI_ISL_433697	hCoV-19/England/CAMB-73DB6/2020	CAMB-73DB6	EPI_ISL_433920	hCoV-19/England/CAMB-7B11C/2020	CAMB-7B11C
EPI_ISL_433698	hCoV-19/England/CAMB-73DC5/2020	CAMB-73DC5	EPI_ISL_433921	hCoV-19/England/CAMB-7B12B/2020	CAMB-7B12B
EPI_ISL_433699	hCoV-19/England/CAMB-73DD4/2020	CAMB-73DD4	EPI_ISL_433922	hCoV-19/England/CAMB-7B13A/2020	CAMB-7B13A
EPI_ISL_425280	hCoV-19/England/CAMB-74359/2020	CAMB-74359	EPI_ISL_433923	hCoV-19/England/CAMB-7B149/2020	CAMB-7B149

GISAID Accession ID	Virus name	COG-UK	GISAID Accession ID	Virus name	COG-UK
EPI_ISL_425281	hCoV-19/England/CAMB-74368/2020	CAMB-74368	EPI_ISL_433924	hCoV-19/England/CAMB-7B158/2020	CAMB-7B158
EPI_ISL_425282	hCoV-19/England/CAMB-74377/2020	CAMB-74377	EPI_ISL_433925	hCoV-19/England/CAMB-7B167/2020	CAMB-7B167
EPI_ISL_425283	hCoV-19/England/CAMB-74386/2020	CAMB-74386	EPI_ISL_433926	hCoV-19/England/CAMB-7B194/2020	CAMB-7B194
EPI_ISL_425284	hCoV-19/England/CAMB-74395/2020	CAMB-74395	EPI_ISL_433927	hCoV-19/England/CAMB-7B1A3/2020	CAMB-7B1A3
EPI_ISL_425285	hCoV-19/England/CAMB-743D1/2020	CAMB-743D1	EPI_ISL_433928	hCoV-19/England/CAMB-7B1B2/2020	CAMB-7B1B2
EPI_ISL_425286	hCoV-19/England/CAMB-743E0/2020	CAMB-743E0	EPI_ISL_433929	hCoV-19/England/CAMB-7B1C1/2020	CAMB-7B1C1
EPI_ISL_425287	hCoV-19/England/CAMB-743FF/2020	CAMB-743FF	EPI_ISL_433930	hCoV-19/England/CAMB-7B1D0/2020	CAMB-7B1D0
EPI_ISL_425288	hCoV-19/England/CAMB-7440B/2020	CAMB-7440B	EPI_ISL_433931	hCoV-19/England/CAMB-7B1EF/2020	CAMB-7B1EF
EPI_ISL_425289	hCoV-19/England/CAMB-7441A/2020	CAMB-7441A	EPI_ISL_433932	hCoV-19/England/CAMB-7B20A/2020	CAMB-7B20A
EPI_ISL_425290	hCoV-19/England/CAMB-74438/2020	CAMB-74438	EPI_ISL_433933	hCoV-19/England/CAMB-7B237/2020	CAMB-7B237
EPI_ISL_425291	hCoV-19/England/CAMB-74456/2020	CAMB-74456	EPI_ISL_433934	hCoV-19/England/CAMB-7B273/2020	CAMB-7B273
EPI_ISL_425292	hCoV-19/England/CAMB-74465/2020	CAMB-74465	EPI_ISL_433935	hCoV-19/England/CAMB-7B291/2020	CAMB-7B291
EPI_ISL_425293	hCoV-19/England/CAMB-74474/2020	CAMB-74474	EPI_ISL_433936	hCoV-19/England/CAMB-7B2A0/2020	CAMB-7B2A0
EPI_ISL_425294	hCoV-19/England/CAMB-74483/2020	CAMB-74483	EPI_ISL_433937	hCoV-19/England/CAMB-7B2BF/2020	CAMB-7B2BF
EPI_ISL_425295	hCoV-19/England/CAMB-74492/2020	CAMB-74492	EPI_ISL_433938	hCoV-19/England/CAMB-7B2CE/2020	CAMB-7B2CE
EPI_ISL_425296	hCoV-19/England/CAMB-744A1/2020	CAMB-744A1	EPI_ISL_433939	hCoV-19/England/CAMB-7B2EC/2020	CAMB-7B2EC
EPI_ISL_425297	hCoV-19/England/CAMB-744B0/2020	CAMB-744B0	EPI_ISL_433940	hCoV-19/England/CAMB-7B307/2020	CAMB-7B307
EPI_ISL_425298	hCoV-19/England/CAMB-744CF/2020	CAMB-744CF	EPI_ISL_433941	hCoV-19/England/CAMB-7B316/2020	CAMB-7B316
EPI_ISL_425299	hCoV-19/England/CAMB-744DE/2020	CAMB-744DE	EPI_ISL_433942	hCoV-19/England/CAMB-7B325/2020	CAMB-7B325
EPI_ISL_425300	hCoV-19/England/CAMB-74508/2020	CAMB-74508	EPI_ISL_433943	hCoV-19/England/CAMB-7B334/2020	CAMB-7B334
EPI_ISL_425301	hCoV-19/England/CAMB-74517/2020	CAMB-74517	EPI_ISL_433944	hCoV-19/England/CAMB-7B343/2020	CAMB-7B343
EPI_ISL_425302	hCoV-19/England/CAMB-74526/2020	CAMB-74526	EPI_ISL_433945	hCoV-19/England/CAMB-7B361/2020	CAMB-7B361
EPI_ISL_425303	hCoV-19/England/CAMB-74535/2020	CAMB-74535	EPI_ISL_433946	hCoV-19/England/CAMB-7B370/2020	CAMB-7B370
EPI_ISL_425304	hCoV-19/England/CAMB-74544/2020	CAMB-74544	EPI_ISL_433947	hCoV-19/England/CAMB-7B38F/2020	CAMB-7B38F
EPI_ISL_425305	hCoV-19/England/CAMB-74553/2020	CAMB-74553	EPI_ISL_433948	hCoV-19/England/CAMB-7B39E/2020	CAMB-7B39E
EPI_ISL_425306	hCoV-19/England/CAMB-74562/2020	CAMB-74562	EPI_ISL_433949	hCoV-19/England/CAMB-7B3BC/2020	CAMB-7B3BC
EPI_ISL_425307	hCoV-19/England/CAMB-74571/2020	CAMB-74571	EPI_ISL_433950	hCoV-19/England/CAMB-7B3CB/2020	CAMB-7B3CB
EPI_ISL_425308	hCoV-19/England/CAMB-74580/2020	CAMB-74580	EPI_ISL_433951	hCoV-19/England/CAMB-7B3DA/2020	CAMB-7B3DA
EPI_ISL_425309	hCoV-19/England/CAMB-7459F/2020	CAMB-7459F	EPI_ISL_433952	hCoV-19/England/CAMB-7B3E9/2020	CAMB-7B3E9
EPI_ISL_425310	hCoV-19/England/CAMB-745AE/2020	CAMB-745AE	EPI_ISL_433953	hCoV-19/England/CAMB-7B3F8/2020	CAMB-7B3F8
EPI_ISL_425311	hCoV-19/England/CAMB-745BD/2020	CAMB-745BD	EPI_ISL_433954	hCoV-19/England/CAMB-7B404/2020	CAMB-7B404
EPI_ISL_425312	hCoV-19/England/CAMB-745CC/2020	CAMB-745CC	EPI_ISL_433955	hCoV-19/England/CAMB-7B413/2020	CAMB-7B413
EPI_ISL_425313	hCoV-19/England/CAMB-745DB/2020	CAMB-745DB	EPI_ISL_433956	hCoV-19/England/CAMB-7B422/2020	CAMB-7B422
EPI_ISL_425314	hCoV-19/England/CAMB-745EA/2020	CAMB-745EA	EPI_ISL_433957	hCoV-19/England/CAMB-7B431/2020	CAMB-7B431
EPI_ISL_425315	hCoV-19/England/CAMB-74605/2020	CAMB-74605	EPI_ISL_433958	hCoV-19/England/CAMB-7B440/2020	CAMB-7B440
EPI_ISL_425316	hCoV-19/England/CAMB-74614/2020	CAMB-74614	EPI_ISL_433959	hCoV-19/England/CAMB-7B45F/2020	CAMB-7B45F
EPI_ISL_425317	hCoV-19/England/CAMB-74623/2020	CAMB-74623	EPI_ISL_433960	hCoV-19/England/CAMB-7B47D/2020	CAMB-7B47D
EPI_ISL_425318	hCoV-19/England/CAMB-74641/2020	CAMB-74641	EPI_ISL_433961	hCoV-19/England/CAMB-7B48C/2020	CAMB-7B48C
EPI_ISL_425319	hCoV-19/England/CAMB-74650/2020	CAMB-74650	EPI_ISL_433962	hCoV-19/England/CAMB-7B49B/2020	CAMB-7B49B
EPI_ISL_425320	hCoV-19/England/CAMB-7467E/2020	CAMB-7467E	EPI_ISL_433963	hCoV-19/England/CAMB-7B4AA/2020	CAMB-7B4AA
EPI_ISL_425321	hCoV-19/England/CAMB-7468D/2020	CAMB-7468D	EPI_ISL_433964	hCoV-19/England/CAMB-7B4B9/2020	CAMB-7B4B9
EPI_ISL_425322	hCoV-19/England/CAMB-7469C/2020	CAMB-7469C	EPI_ISL_433965	hCoV-19/England/CAMB-7B4C8/2020	CAMB-7B4C8
EPI_ISL_425323	hCoV-19/England/CAMB-746AB/2020	CAMB-746AB	EPI_ISL_433966	hCoV-19/England/CAMB-7B4D7/2020	CAMB-7B4D7
EPI_ISL_425324	hCoV-19/England/CAMB-746BA/2020	CAMB-746BA	EPI_ISL_433967	hCoV-19/England/CAMB-7B4E6/2020	CAMB-7B4E6
EPI_ISL_425325	hCoV-19/England/CAMB-746D8/2020	CAMB-746D8	EPI_ISL_433968	hCoV-19/England/CAMB-7B4F5/2020	CAMB-7B4F5
EPI_ISL_425326	hCoV-19/England/CAMB-74702/2020	CAMB-74702	EPI_ISL_433969	hCoV-19/England/CAMB-7B501/2020	CAMB-7B501
EPI_ISL_425327	hCoV-19/England/CAMB-74711/2020	CAMB-74711	EPI_ISL_433970	hCoV-19/England/CAMB-7B510/2020	CAMB-7B510
EPI_ISL_425328	hCoV-19/England/CAMB-74720/2020	CAMB-74720	EPI_ISL_433971	hCoV-19/England/CAMB-7B52F/2020	CAMB-7B52F
EPI_ISL_425329	hCoV-19/England/CAMB-7473F/2020	CAMB-7473F	EPI_ISL_433972	hCoV-19/England/CAMB-7B53E/2020	CAMB-7B53E
EPI_ISL_425330	hCoV-19/England/CAMB-7474E/2020	CAMB-7474E	EPI_ISL_433973	hCoV-19/England/CAMB-7B54D/2020	CAMB-7B54D
EPI_ISL_425331	hCoV-19/England/CAMB-7475D/2020	CAMB-7475D	EPI_ISL_433974	hCoV-19/England/CAMB-7B55C/2020	CAMB-7B55C
EPI_ISL_425332	hCoV-19/England/CAMB-7476C/2020	CAMB-7476C	EPI_ISL_433975	hCoV-19/England/CAMB-7B56B/2020	CAMB-7B56B
EPI_ISL_425333	hCoV-19/England/CAMB-7477B/2020	CAMB-7477B	EPI_ISL_433976	hCoV-19/England/CAMB-7B5F6/2020	CAMB-7B5F6
EPI_ISL_425334	hCoV-19/England/CAMB-748D2/2020	CAMB-748D2	EPI_ISL_433977	hCoV-19/England/CAMB-7B5FE/2020	CAMB-7B5FE
EPI_ISL_425335	hCoV-19/England/CAMB-748E1/2020	CAMB-748E1	EPI_ISL_433978	hCoV-19/England/CAMB-7B5F4/2020	CAMB-7B5F4
EPI_ISL_425336	hCoV-19/England/CAMB-748F0/2020	CAMB-748F0	EPI_ISL_433979	hCoV-19/England/CAMB-7C00F/2020	CAMB-7C00F
EPI_ISL_425337	hCoV-19/England/CAMB-7490C/2020	CAMB-7490C	EPI_ISL_433980	hCoV-19/England/CAMB-7C01E/2020	CAMB-7C01E
EPI_ISL_425338	hCoV-19/England/CAMB-7491B/2020	CAMB-7491B	EPI_ISL_433981	hCoV-19/England/CAMB-7C02D/2020	CAMB-7C02D
EPI_ISL_425339	hCoV-19/England/CAMB-7492A/2020	CAMB-7492A	EPI_ISL_433982	hCoV-19/England/CAMB-7C03C/2020	CAMB-7C03C
EPI_ISL_425340	hCoV-19/England/CAMB-7498A/2020	CAMB-7498A	EPI_ISL_433983	hCoV-19/England/CAMB-7C04B/2020	CAMB-7C04B
EPI_ISL_425341	hCoV-19/England/CAMB-749A2/2020	CAMB-749A2	EPI_ISL_433984	hCoV-19/England/CAMB-7C05A/2020	CAMB-7C05A
EPI_ISL_425342	hCoV-19/England/CAMB-74A09/2020	CAMB-74A09	EPI_ISL_433985	hCoV-19/England/CAMB-7C069/2020	CAMB-7C069
EPI_ISL_425343	hCoV-19/England/CAMB-74A18/2020	CAMB-74A18	EPI_ISL_433986	hCoV-19/England/CAMB-7C078/2020	CAMB-7C078
EPI_ISL_425344	hCoV-19/England/CAMB-74A72/2020	CAMB-74A72	EPI_ISL_433987	hCoV-19/England/CAMB-7C087/2020	CAMB-7C087
EPI_ISL_425345	hCoV-19/England/CAMB-74ABE/2020	CAMB-74ABE	EPI_ISL_433988	hCoV-19/England/CAMB-7C096/2020	CAMB-7C096
EPI_ISL_425346	hCoV-19/England/CAMB-74B24/2020	CAMB-74B24	EPI_ISL_433989	hCoV-19/England/CAMB-7C0A5/2020	CAMB-7C0A5
EPI_ISL_425347	hCoV-19/England/CAMB-74B33/2020	CAMB-74B33	EPI_ISL_433990	hCoV-19/England/CAMB-7C0B4/2020	CAMB-7C0B4
EPI_ISL_425348	hCoV-19/England/CAMB-74B42/2020	CAMB-74B42	EPI_ISL_433991	hCoV-19/England/CAMB-7C0C3/2020	CAMB-7C0C3
EPI_ISL_425349	hCoV-19/England/CAMB-74B51/2020	CAMB-74B51	EPI_ISL_433992	hCoV-19/England/CAMB-7C0D2/2020	CAMB-7C0D2
EPI_ISL_425350	hCoV-19/England/CAMB-74B60/2020	CAMB-74B60	EPI_ISL_433993	hCoV-19/England/CAMB-7C0E1/2020	CAMB-7C0E1
EPI_ISL_425351	hCoV-19/England/CAMB-74B9D/2020	CAMB-74B9D	EPI_ISL_433994	hCoV-19/England/CAMB-7C0F0/2020	CAMB-7C0F0
EPI_ISL_425352	hCoV-19/England/CAMB-74BAC/2020	CAMB-74BAC	EPI_ISL_433995	hCoV-19/England/CAMB-7C10C/2020	CAMB-7C10C
EPI_ISL_425353	hCoV-19/England/CAMB-74BBB/2020	CAMB-74BBB	EPI_ISL_433996	hCoV-19/England/CAMB-7C11B/2020	CAMB-7C11B
EPI_ISL_425354	hCoV-19/England/CAMB-74BCA/2020	CAMB-74BCA	EPI_ISL_433997	hCoV-19/England/CAMB-7C12A/2020	CAMB-7C12A
EPI_ISL_425355	hCoV-19/England/CAMB-74BF7/2020	CAMB-74BF7	EPI_ISL_433998	hCoV-19/England/CAMB-7C139/2020	CAMB-7C139
EPI_ISL_425356	hCoV-19/England/CAMB-74C03/2020	CAMB-74C03	EPI_ISL_433999	hCoV-19/England/CAMB-7C148/2020	CAMB-7C148
EPI_ISL_425357	hCoV-19/England/CAMB-74C12/2020	CAMB-74C12	EPI_ISL_434000	hCoV-19/England/CAMB-7C157/2020	CAMB-7C157
EPI_ISL_425358	hCoV-19/England/CAMB-74C21/2020	CAMB-74C21	EPI_ISL_434001	hCoV-19/England/CAMB-7C166/2020	CAMB-7C166
EPI_ISL_425359	hCoV-19/England/CAMB-74C30/2020	CAMB-74C30	EPI_ISL_434002	hCoV-19/England/CAMB-7C175/2020	CAMB-7C175
EPI_ISL_425360	hCoV-19/England/CAMB-74C4F/2020	CAMB-74C4F	EPI_ISL_434003	hCoV-19/England/CAMB-7C184/2020	CAMB-7C184
EPI_ISL_425361	hCoV-19/England/CAMB-74C5E/2020	CAMB-74C5E	EPI_ISL_434004	hCoV-19/England/CAMB-7C193/2020	CAMB-7C193
EPI_ISL_425362	hCoV-19/England/CAMB-74C8B/2020	CAMB-74C8B	EPI_ISL_434005	hCoV-19/England/CAMB-7C1A2/2020	CAMB-7C1A2
EPI_ISL_425363	hCoV-19/England/CAMB-74CA9/2020	CAMB-74CA9	EPI_ISL_434006	hCoV-19/England/CAMB-7C1B1/2020	CAMB-7C1B1
EPI_ISL_425364	hCoV-19/England/CAMB-74CC7/2020	CAMB-74CC7	EPI_ISL_434007	hCoV-19/England/CAMB-7C1C0/2020	CAMB-7C1C0

GISAID Accession ID	Virus name	COG-UK	GISAID Accession ID	Virus name	COG-UK
EPI_ISL_425365	hCoV-19/England/CAMB-74CD6/2020	CAMB-74CD6	EPI_ISL_434008	hCoV-19/England/CAMB-7C1DF/2020	CAMB-7C1DF
EPI_ISL_425366	hCoV-19/England/CAMB-74CF4/2020	CAMB-74CF4	EPI_ISL_434009	hCoV-19/England/CAMB-7C1EE/2020	CAMB-7C1EE
EPI_ISL_425367	hCoV-19/England/CAMB-74D00/2020	CAMB-74D00	EPI_ISL_434010	hCoV-19/England/CAMB-7C1FD/2020	CAMB-7C1FD
EPI_ISL_425368	hCoV-19/England/CAMB-74D1F/2020	CAMB-74D1F	EPI_ISL_434011	hCoV-19/England/CAMB-7C209/2020	CAMB-7C209
EPI_ISL_425369	hCoV-19/England/CAMB-74D2E/2020	CAMB-74D2E	EPI_ISL_434012	hCoV-19/England/CAMB-7C218/2020	CAMB-7C218
EPI_ISL_425370	hCoV-19/England/CAMB-74D3D/2020	CAMB-74D3D	EPI_ISL_434013	hCoV-19/England/CAMB-7C227/2020	CAMB-7C227
EPI_ISL_425371	hCoV-19/England/CAMB-74D4C/2020	CAMB-74D4C	EPI_ISL_434014	hCoV-19/England/CAMB-7C236/2020	CAMB-7C236
EPI_ISL_425373	hCoV-19/England/CAMB-74D5B/2020	CAMB-74D5B	EPI_ISL_434015	hCoV-19/England/CAMB-7C245/2020	CAMB-7C245
EPI_ISL_425374	hCoV-19/England/CAMB-74D6A/2020	CAMB-74D6A	EPI_ISL_434016	hCoV-19/England/CAMB-7C254/2020	CAMB-7C254
EPI_ISL_425375	hCoV-19/England/CAMB-74DE2/2020	CAMB-74DE2	EPI_ISL_434017	hCoV-19/England/CAMB-7C263/2020	CAMB-7C263
EPI_ISL_425376	hCoV-19/England/CAMB-74DF1/2020	CAMB-74DF1	EPI_ISL_434018	hCoV-19/England/CAMB-7C272/2020	CAMB-7C272
EPI_ISL_425377	hCoV-19/England/CAMB-74E0D/2020	CAMB-74E0D	EPI_ISL_434019	hCoV-19/England/CAMB-7C281/2020	CAMB-7C281
EPI_ISL_425378	hCoV-19/England/CAMB-74E49/2020	CAMB-74E49	EPI_ISL_434020	hCoV-19/England/CAMB-7C290/2020	CAMB-7C290
EPI_ISL_425379	hCoV-19/England/CAMB-74E58/2020	CAMB-74E58	EPI_ISL_434021	hCoV-19/England/CAMB-7C2AF/2020	CAMB-7C2AF
EPI_ISL_425380	hCoV-19/England/CAMB-74E67/2020	CAMB-74E67	EPI_ISL_434022	hCoV-19/England/CAMB-7C2BE/2020	CAMB-7C2BE
EPI_ISL_425381	hCoV-19/England/CAMB-74E76/2020	CAMB-74E76	EPI_ISL_434023	hCoV-19/England/CAMB-7C2CD/2020	CAMB-7C2CD
EPI_ISL_425382	hCoV-19/England/CAMB-74E85/2020	CAMB-74E85	EPI_ISL_434024	hCoV-19/England/CAMB-7C2DC/2020	CAMB-7C2DC
EPI_ISL_425383	hCoV-19/England/CAMB-74EA3/2020	CAMB-74EA3	EPI_ISL_434025	hCoV-19/England/CAMB-7C2FA/2020	CAMB-7C2FA
EPI_ISL_425384	hCoV-19/England/CAMB-74EB2/2020	CAMB-74EB2	EPI_ISL_434026	hCoV-19/England/CAMB-7C306/2020	CAMB-7C306
EPI_ISL_425385	hCoV-19/England/CAMB-74EC1/2020	CAMB-74EC1	EPI_ISL_434027	hCoV-19/England/CAMB-7C315/2020	CAMB-7C315
EPI_ISL_425386	hCoV-19/England/CAMB-7537E/2020	CAMB-7537E	EPI_ISL_434028	hCoV-19/England/CAMB-7C324/2020	CAMB-7C324
EPI_ISL_425387	hCoV-19/England/CAMB-75419/2020	CAMB-75419	EPI_ISL_434029	hCoV-19/England/CAMB-7C333/2020	CAMB-7C333
EPI_ISL_425388	hCoV-19/England/CAMB-75437/2020	CAMB-75437	EPI_ISL_434030	hCoV-19/England/CAMB-7C342/2020	CAMB-7C342
EPI_ISL_425389	hCoV-19/England/CAMB-75455/2020	CAMB-75455	EPI_ISL_434031	hCoV-19/England/CAMB-7C351/2020	CAMB-7C351
EPI_ISL_425390	hCoV-19/England/CAMB-75464/2020	CAMB-75464	EPI_ISL_434032	hCoV-19/England/CAMB-7C360/2020	CAMB-7C360
EPI_ISL_425391	hCoV-19/England/CAMB-75473/2020	CAMB-75473	EPI_ISL_434033	hCoV-19/England/CAMB-7C37F/2020	CAMB-7C37F
EPI_ISL_425392	hCoV-19/England/CAMB-75482/2020	CAMB-75482	EPI_ISL_434034	hCoV-19/England/CAMB-7C38E/2020	CAMB-7C38E
EPI_ISL_425393	hCoV-19/England/CAMB-75491/2020	CAMB-75491	EPI_ISL_434035	hCoV-19/England/CAMB-7C39D/2020	CAMB-7C39D
EPI_ISL_425394	hCoV-19/England/CAMB-754A0/2020	CAMB-754A0	EPI_ISL_434036	hCoV-19/England/CAMB-7C3AC/2020	CAMB-7C3AC
EPI_ISL_425395	hCoV-19/England/CAMB-754B7/2020	CAMB-754B7	EPI_ISL_434037	hCoV-19/England/CAMB-7C3BB/2020	CAMB-7C3BB
EPI_ISL_425396	hCoV-19/England/CAMB-754CE/2020	CAMB-754CE	EPI_ISL_434038	hCoV-19/England/CAMB-7C3CA/2020	CAMB-7C3CA
EPI_ISL_425397	hCoV-19/England/CAMB-754DD/2020	CAMB-754DD	EPI_ISL_434039	hCoV-19/England/CAMB-7C3E8/2020	CAMB-7C3E8
EPI_ISL_425398	hCoV-19/England/CAMB-754EC/2020	CAMB-754EC	EPI_ISL_434040	hCoV-19/England/CAMB-7E10A/2020	CAMB-7E10A
EPI_ISL_425399	hCoV-19/England/CAMB-754FB/2020	CAMB-754FB	EPI_ISL_434041	hCoV-19/England/CAMB-7E119/2020	CAMB-7E119
EPI_ISL_425400	hCoV-19/England/CAMB-75507/2020	CAMB-75507	EPI_ISL_434042	hCoV-19/England/CAMB-7E128/2020	CAMB-7E128
EPI_ISL_425401	hCoV-19/England/CAMB-75516/2020	CAMB-75516	EPI_ISL_434043	hCoV-19/England/CAMB-7E137/2020	CAMB-7E137
EPI_ISL_425402	hCoV-19/England/CAMB-75525/2020	CAMB-75525	EPI_ISL_434044	hCoV-19/England/CAMB-7E146/2020	CAMB-7E146
EPI_ISL_425403	hCoV-19/England/CAMB-75534/2020	CAMB-75534	EPI_ISL_434045	hCoV-19/England/CAMB-7E155/2020	CAMB-7E155
EPI_ISL_425404	hCoV-19/England/CAMB-75543/2020	CAMB-75543	EPI_ISL_434046	hCoV-19/England/CAMB-7E164/2020	CAMB-7E164
EPI_ISL_425405	hCoV-19/England/CAMB-75552/2020	CAMB-75552	EPI_ISL_434047	hCoV-19/England/CAMB-7E1BF/2020	CAMB-7E1BF
EPI_ISL_425406	hCoV-19/England/CAMB-75570/2020	CAMB-75570	EPI_ISL_434048	hCoV-19/England/CAMB-7E1CE/2020	CAMB-7E1CE
EPI_ISL_425407	hCoV-19/England/CAMB-7559E/2020	CAMB-7559E	EPI_ISL_434049	hCoV-19/England/CAMB-7E1DD/2020	CAMB-7E1DD
EPI_ISL_433700	hCoV-19/England/CAMB-755AD/2020	CAMB-755AD	EPI_ISL_434050	hCoV-19/England/CAMB-7E1EC/2020	CAMB-7E1EC
EPI_ISL_425408	hCoV-19/England/CAMB-755BC/2020	CAMB-755BC	EPI_ISL_434051	hCoV-19/England/CAMB-7E1FB/2020	CAMB-7E1FB
EPI_ISL_425409	hCoV-19/England/CAMB-755CB/2020	CAMB-755CB	EPI_ISL_434052	hCoV-19/England/CAMB-7E207/2020	CAMB-7E207
EPI_ISL_425410	hCoV-19/England/CAMB-755DA/2020	CAMB-755DA	EPI_ISL_434053	hCoV-19/England/CAMB-7E225/2020	CAMB-7E225
EPI_ISL_425411	hCoV-19/England/CAMB-755E9/2020	CAMB-755E9	EPI_ISL_434054	hCoV-19/England/CAMB-7E234/2020	CAMB-7E234
EPI_ISL_425412	hCoV-19/England/CAMB-755F8/2020	CAMB-755F8	EPI_ISL_434055	hCoV-19/England/CAMB-7E243/2020	CAMB-7E243
EPI_ISL_425413	hCoV-19/England/CAMB-75604/2020	CAMB-75604	EPI_ISL_434056	hCoV-19/England/CAMB-7E252/2020	CAMB-7E252
EPI_ISL_425414	hCoV-19/England/CAMB-75613/2020	CAMB-75613	EPI_ISL_434057	hCoV-19/England/CAMB-7E261/2020	CAMB-7E261
EPI_ISL_425415	hCoV-19/England/CAMB-75622/2020	CAMB-75622	EPI_ISL_434058	hCoV-19/England/CAMB-7E270/2020	CAMB-7E270
EPI_ISL_425416	hCoV-19/England/CAMB-75631/2020	CAMB-75631	EPI_ISL_434059	hCoV-19/England/CAMB-7E29E/2020	CAMB-7E29E
EPI_ISL_425417	hCoV-19/England/CAMB-75640/2020	CAMB-75640	EPI_ISL_434060	hCoV-19/England/CAMB-7E2AD/2020	CAMB-7E2AD
EPI_ISL_425418	hCoV-19/England/CAMB-7565F/2020	CAMB-7565F	EPI_ISL_434061	hCoV-19/England/CAMB-7E2BC/2020	CAMB-7E2BC
EPI_ISL_425419	hCoV-19/England/CAMB-7566E/2020	CAMB-7566E	EPI_ISL_434062	hCoV-19/England/CAMB-7E2CB/2020	CAMB-7E2CB
EPI_ISL_425420	hCoV-19/England/CAMB-7567D/2020	CAMB-7567D	EPI_ISL_433466	hCoV-19/England/CAMB-7E2DA/2020	CAMB-7E2DA
EPI_ISL_425421	hCoV-19/England/CAMB-7569B/2020	CAMB-7569B	EPI_ISL_433467	hCoV-19/England/CAMB-7E2E9/2020	CAMB-7E2E9
EPI_ISL_425422	hCoV-19/England/CAMB-756AA/2020	CAMB-756AA	EPI_ISL_433468	hCoV-19/England/CAMB-7E2F8/2020	CAMB-7E2F8
EPI_ISL_425423	hCoV-19/England/CAMB-76D1D/2020	CAMB-76D1D	EPI_ISL_433469	hCoV-19/England/CAMB-7E304/2020	CAMB-7E304
EPI_ISL_425424	hCoV-19/England/CAMB-76D3B/2020	CAMB-76D3B	EPI_ISL_433470	hCoV-19/England/CAMB-7E313/2020	CAMB-7E313
EPI_ISL_425425	hCoV-19/England/CAMB-76D59/2020	CAMB-76D59	EPI_ISL_433471	hCoV-19/England/CAMB-7E322/2020	CAMB-7E322
EPI_ISL_425426	hCoV-19/England/CAMB-76D77/2020	CAMB-76D77	EPI_ISL_433472	hCoV-19/England/CAMB-7E331/2020	CAMB-7E331
EPI_ISL_425427	hCoV-19/England/CAMB-76D95/2020	CAMB-76D95	EPI_ISL_433473	hCoV-19/England/CAMB-7E340/2020	CAMB-7E340
EPI_ISL_425428	hCoV-19/England/CAMB-76DB3/2020	CAMB-76DB3	EPI_ISL_433474	hCoV-19/England/CAMB-7F7E9/2020	CAMB-7F7E9
EPI_ISL_425429	hCoV-19/England/CAMB-76DD1/2020	CAMB-76DD1	EPI_ISL_433475	hCoV-19/England/CAMB-7F804/2020	CAMB-7F804
EPI_ISL_425430	hCoV-19/England/CAMB-76DFF/2020	CAMB-76DFF	EPI_ISL_433476	hCoV-19/England/CAMB-7F822/2020	CAMB-7F822
EPI_ISL_425431	hCoV-19/England/CAMB-76E1A/2020	CAMB-76E1A	EPI_ISL_433477	hCoV-19/England/CAMB-7F88C/2020	CAMB-7F88C
EPI_ISL_425432	hCoV-19/England/CAMB-76E38/2020	CAMB-76E38	EPI_ISL_433478	hCoV-19/England/CAMB-7F8E6/2020	CAMB-7F8E6
EPI_ISL_425433	hCoV-19/England/CAMB-76E56/2020	CAMB-76E56	EPI_ISL_433479	hCoV-19/England/CAMB-7FB29/2020	CAMB-7FB29
EPI_ISL_425434	hCoV-19/England/CAMB-76E74/2020	CAMB-76E74	EPI_ISL_433480	hCoV-19/England/CAMB-7FB47/2020	CAMB-7FB47
EPI_ISL_425435	hCoV-19/England/CAMB-76E92/2020	CAMB-76E92	EPI_ISL_433481	hCoV-19/England/CAMB-7FB56/2020	CAMB-7FB56
EPI_ISL_425436	hCoV-19/England/CAMB-76EB0/2020	CAMB-76EB0	EPI_ISL_433482	hCoV-19/England/CAMB-7FB65/2020	CAMB-7FB65
EPI_ISL_425437	hCoV-19/England/CAMB-76EDE/2020	CAMB-76EDE	EPI_ISL_433483	hCoV-19/England/CAMB-7FB74/2020	CAMB-7FB74
EPI_ISL_425438	hCoV-19/England/CAMB-76EFC/2020	CAMB-76EFC	EPI_ISL_433484	hCoV-19/England/CAMB-7FB83/2020	CAMB-7FB83
EPI_ISL_425439	hCoV-19/England/CAMB-76F17/2020	CAMB-76F17	EPI_ISL_433485	hCoV-19/England/CAMB-7FB92/2020	CAMB-7FB92
EPI_ISL_425440	hCoV-19/England/CAMB-76F35/2020	CAMB-76F35	EPI_ISL_433486	hCoV-19/England/CAMB-7FBA1/2020	CAMB-7FBA1
EPI_ISL_425441	hCoV-19/England/CAMB-76F62/2020	CAMB-76F62	EPI_ISL_433487	hCoV-19/England/CAMB-7FBB0/2020	CAMB-7FBB0
EPI_ISL_425442	hCoV-19/England/CAMB-76F71/2020	CAMB-76F71	EPI_ISL_433488	hCoV-19/England/CAMB-7FBCF/2020	CAMB-7FBCF
EPI_ISL_425443	hCoV-19/England/CAMB-76F9F/2020	CAMB-76F9F	EPI_ISL_433489	hCoV-19/England/CAMB-7FBDE/2020	CAMB-7FBDE
EPI_ISL_425444	hCoV-19/England/CAMB-76FAE/2020	CAMB-76FAE	EPI_ISL_433490	hCoV-19/England/CAMB-7FBFC/2020	CAMB-7FBFC
EPI_ISL_425445	hCoV-19/England/CAMB-76FBD/2020	CAMB-76FBD	EPI_ISL_433491	hCoV-19/England/CAMB-7FC08/2020	CAMB-7FC08
EPI_ISL_425446	hCoV-19/England/CAMB-76FCC/2020	CAMB-76FCC	EPI_ISL_433492	hCoV-19/England/CAMB-7FC17/2020	CAMB-7FC17
EPI_ISL_425447	hCoV-19/England/CAMB-76FDB/2020	CAMB-76FDB	EPI_ISL_433493	hCoV-19/England/CAMB-7FC26/2020	CAMB-7FC26
EPI_ISL_425448	hCoV-19/England/CAMB-76FEA/2020	CAMB-76FEA	EPI_ISL_433494	hCoV-19/England/CAMB-7FC35/2020	CAMB-7FC35

GISAIID Accession ID	Virus name	COG-UK	GISAIID Accession ID	Virus name	COG-UK
EPI_ISL_425449	hCoV-19/England/CAMB-76FF9/2020	CAMB-76FF9	EPI_ISL_433495	hCoV-19/England/CAMB-7FC44/2020	CAMB-7FC44
EPI_ISL_425450	hCoV-19/England/CAMB-77004/2020	CAMB-77004	EPI_ISL_433496	hCoV-19/England/CAMB-7FC53/2020	CAMB-7FC53
EPI_ISL_425451	hCoV-19/England/CAMB-77013/2020	CAMB-77013	EPI_ISL_433497	hCoV-19/England/CAMB-7FC62/2020	CAMB-7FC62
EPI_ISL_425452	hCoV-19/England/CAMB-77022/2020	CAMB-77022	EPI_ISL_433856	hCoV-19/England/CAMB-7AC49/2020	CAMB-7AC49
EPI_ISL_425453	hCoV-19/England/CAMB-77031/2020	CAMB-77031	EPI_ISL_433857	hCoV-19/England/CAMB-7AC58/2020	CAMB-7AC58
EPI_ISL_425454	hCoV-19/England/CAMB-77040/2020	CAMB-77040	EPI_ISL_433858	hCoV-19/England/CAMB-7AC67/2020	CAMB-7AC67
EPI_ISL_425455	hCoV-19/England/CAMB-7705F/2020	CAMB-7705F	EPI_ISL_433859	hCoV-19/England/CAMB-7AC76/2020	CAMB-7AC76
EPI_ISL_425456	hCoV-19/England/CAMB-7706E/2020	CAMB-7706E	EPI_ISL_433855	hCoV-19/England/CAMB-7AC3A/2020	CAMB-7AC3A
EPI_ISL_425457	hCoV-19/England/CAMB-7707D/2020	CAMB-7707D	Pending	hCoV-19/England/CAMB-7FC71/2020	CAMB-7FC71
EPI_ISL_425458	hCoV-19/England/CAMB-7708C/2020	CAMB-7708C	Pending	hCoV-19/England/CAMB-7FC80/2020	CAMB-7FC80
EPI_ISL_425459	hCoV-19/England/CAMB-7709B/2020	CAMB-7709B	Pending	hCoV-19/England/CAMB-7FC9F/2020	CAMB-7FC9F
EPI_ISL_425460	hCoV-19/England/CAMB-770B9/2020	CAMB-770B9	Pending	hCoV-19/England/CAMB-7FCBD/2020	CAMB-7FCBD
EPI_ISL_425461	hCoV-19/England/CAMB-770C8/2020	CAMB-770C8	Pending	hCoV-19/England/CAMB-7FCF9/2020	CAMB-7FCF9
EPI_ISL_433701	hCoV-19/England/CAMB-779FA/2020	CAMB-779FA	Pending	hCoV-19/England/CAMB-7FD05/2020	CAMB-7FD05
EPI_ISL_433702	hCoV-19/England/CAMB-78197/2020	CAMB-78197	Pending	hCoV-19/England/CAMB-7FD23/2020	CAMB-7FD23
EPI_ISL_433703	hCoV-19/England/CAMB-781A6/2020	CAMB-781A6	Pending	hCoV-19/England/CAMB-7FD32/2020	CAMB-7FD32
EPI_ISL_433704	hCoV-19/England/CAMB-781B5/2020	CAMB-781B5	Pending	hCoV-19/England/CAMB-7FD6F/2020	CAMB-7FD6F
EPI_ISL_433705	hCoV-19/England/CAMB-781E2/2020	CAMB-781E2	Pending	hCoV-19/England/CAMB-7FD7E/2020	CAMB-7FD7E
EPI_ISL_433706	hCoV-19/England/CAMB-7820D/2020	CAMB-7820D	Pending	hCoV-19/England/CAMB-7FDBD/2020	CAMB-7FDBD
EPI_ISL_433707	hCoV-19/England/CAMB-7822B/2020	CAMB-7822B	Pending	hCoV-19/England/CAMB-7FDBA/2020	CAMB-7FDBA
EPI_ISL_433708	hCoV-19/England/CAMB-78276/2020	CAMB-78276	Pending	hCoV-19/England/CAMB-7FDC9/2020	CAMB-7FDC9
EPI_ISL_433709	hCoV-19/England/CAMB-78285/2020	CAMB-78285	Pending	hCoV-19/England/CAMB-7FE20/2020	CAMB-7FE20
EPI_ISL_433710	hCoV-19/England/CAMB-78294/2020	CAMB-78294	Pending	hCoV-19/England/CAMB-7FE3F/2020	CAMB-7FE3F
EPI_ISL_433711	hCoV-19/England/CAMB-782A3/2020	CAMB-782A3	Pending	hCoV-19/England/CAMB-7FE4E/2020	CAMB-7FE4E
EPI_ISL_433712	hCoV-19/England/CAMB-782B2/2020	CAMB-782B2	Pending	hCoV-19/England/CAMB-7FE5D/2020	CAMB-7FE5D
EPI_ISL_433713	hCoV-19/England/CAMB-782D0/2020	CAMB-782D0	Pending	hCoV-19/England/CAMB-7FETB/2020	CAMB-7FETB
EPI_ISL_433714	hCoV-19/England/CAMB-782EF/2020	CAMB-782EF	Pending	hCoV-19/England/CAMB-7FE87/2020	CAMB-7FE87
EPI_ISL_433715	hCoV-19/England/CAMB-782FE/2020	CAMB-782FE	Pending	hCoV-19/England/CAMB-7FEC6/2020	CAMB-7FEC6
EPI_ISL_433716	hCoV-19/England/CAMB-7830A/2020	CAMB-7830A	Pending	hCoV-19/England/CAMB-7FEE4/2020	CAMB-7FEE4
EPI_ISL_433717	hCoV-19/England/CAMB-78319/2020	CAMB-78319	Pending	hCoV-19/England/CAMB-80AFB/2020	CAMB-80AFB
EPI_ISL_433718	hCoV-19/England/CAMB-78328/2020	CAMB-78328	Pending	hCoV-19/England/CAMB-80E3B/2020	CAMB-80E3B
EPI_ISL_433719	hCoV-19/England/CAMB-78337/2020	CAMB-78337	Pending	hCoV-19/England/CAMB-80E4A/2020	CAMB-80E4A
EPI_ISL_433720	hCoV-19/England/CAMB-78346/2020	CAMB-78346	Pending	hCoV-19/England/CAMB-80E59/2020	CAMB-80E59
EPI_ISL_433721	hCoV-19/England/CAMB-78355/2020	CAMB-78355	Pending	hCoV-19/England/CAMB-80E68/2020	CAMB-80E68
EPI_ISL_433722	hCoV-19/England/CAMB-78364/2020	CAMB-78364	Pending	hCoV-19/England/CAMB-80E77/2020	CAMB-80E77
EPI_ISL_433723	hCoV-19/England/CAMB-78373/2020	CAMB-78373	Pending	hCoV-19/England/CAMB-80E86/2020	CAMB-80E86
EPI_ISL_433724	hCoV-19/England/CAMB-78382/2020	CAMB-78382	Pending	hCoV-19/England/CAMB-80E95/2020	CAMB-80E95
EPI_ISL_433725	hCoV-19/England/CAMB-78391/2020	CAMB-78391	Pending	hCoV-19/England/CAMB-80EA4/2020	CAMB-80EA4
EPI_ISL_433726	hCoV-19/England/CAMB-783BF/2020	CAMB-783BF	Pending	hCoV-19/England/CAMB-80EB3/2020	CAMB-80EB3
EPI_ISL_433727	hCoV-19/England/CAMB-783CE/2020	CAMB-783CE	Pending	hCoV-19/England/CAMB-80EC2/2020	CAMB-80EC2
EPI_ISL_433728	hCoV-19/England/CAMB-783DD/2020	CAMB-783DD	Pending	hCoV-19/England/CAMB-80ED1/2020	CAMB-80ED1
EPI_ISL_433729	hCoV-19/England/CAMB-783EC/2020	CAMB-783EC	Pending	hCoV-19/England/CAMB-80EE0/2020	CAMB-80EE0
EPI_ISL_433730	hCoV-19/England/CAMB-783FB/2020	CAMB-783FB	Pending	hCoV-19/England/CAMB-80EFF/2020	CAMB-80EFF
EPI_ISL_433731	hCoV-19/England/CAMB-78407/2020	CAMB-78407	Pending	hCoV-19/England/CAMB-80F0B/2020	CAMB-80F0B
EPI_ISL_433732	hCoV-19/England/CAMB-78416/2020	CAMB-78416	Pending	hCoV-19/England/CAMB-80F1A/2020	CAMB-80F1A
EPI_ISL_433733	hCoV-19/England/CAMB-78425/2020	CAMB-78425	Pending	hCoV-19/England/CAMB-80F29/2020	CAMB-80F29
EPI_ISL_433734	hCoV-19/England/CAMB-78452/2020	CAMB-78452	Pending	hCoV-19/England/CAMB-80F38/2020	CAMB-80F38
EPI_ISL_433735	hCoV-19/England/CAMB-78461/2020	CAMB-78461	Pending	hCoV-19/England/CAMB-80F47/2020	CAMB-80F47
EPI_ISL_433736	hCoV-19/England/CAMB-78470/2020	CAMB-78470	Pending	hCoV-19/England/CAMB-80F56/2020	CAMB-80F56
EPI_ISL_433737	hCoV-19/England/CAMB-7848F/2020	CAMB-7848F	Pending	hCoV-19/England/CAMB-80F65/2020	CAMB-80F65
EPI_ISL_433738	hCoV-19/England/CAMB-7849E/2020	CAMB-7849E	Pending	hCoV-19/England/CAMB-80F74/2020	CAMB-80F74
EPI_ISL_433739	hCoV-19/England/CAMB-784AD/2020	CAMB-784AD	Pending	hCoV-19/England/CAMB-80F83/2020	CAMB-80F83
EPI_ISL_433740	hCoV-19/England/CAMB-784BC/2020	CAMB-784BC	Pending	hCoV-19/England/CAMB-80F80/2020	CAMB-80F80
EPI_ISL_433741	hCoV-19/England/CAMB-784CB/2020	CAMB-784CB	Pending	hCoV-19/England/CAMB-80FCF/2020	CAMB-80FCF
EPI_ISL_433742	hCoV-19/England/CAMB-784DA/2020	CAMB-784DA	Pending	hCoV-19/England/CAMB-80FDE/2020	CAMB-80FDE
EPI_ISL_433743	hCoV-19/England/CAMB-784E9/2020	CAMB-784E9	Pending	hCoV-19/England/CAMB-80FFC/2020	CAMB-80FFC
EPI_ISL_433744	hCoV-19/England/CAMB-78504/2020	CAMB-78504	Pending	hCoV-19/England/CAMB-81007/2020	CAMB-81007
EPI_ISL_433745	hCoV-19/England/CAMB-78531/2020	CAMB-78531	Pending	hCoV-19/England/CAMB-81016/2020	CAMB-81016
EPI_ISL_433746	hCoV-19/England/CAMB-78540/2020	CAMB-78540	Pending	hCoV-19/England/CAMB-81025/2020	CAMB-81025
EPI_ISL_433747	hCoV-19/England/CAMB-7855F/2020	CAMB-7855F	Pending	hCoV-19/England/CAMB-81034/2020	CAMB-81034
EPI_ISL_433748	hCoV-19/England/CAMB-7856E/2020	CAMB-7856E	Pending	hCoV-19/England/CAMB-81043/2020	CAMB-81043
EPI_ISL_433749	hCoV-19/England/CAMB-7857D/2020	CAMB-7857D	Pending	hCoV-19/England/CAMB-81061/2020	CAMB-81061
EPI_ISL_433750	hCoV-19/England/CAMB-7858C/2020	CAMB-7858C	Pending	hCoV-19/England/CAMB-81070/2020	CAMB-81070
EPI_ISL_433751	hCoV-19/England/CAMB-7859B/2020	CAMB-7859B	Pending	hCoV-19/England/CAMB-8108F/2020	CAMB-8108F
EPI_ISL_433752	hCoV-19/England/CAMB-785AA/2020	CAMB-785AA	Pending	hCoV-19/England/CAMB-8109E/2020	CAMB-8109E
EPI_ISL_433753	hCoV-19/England/CAMB-785C8/2020	CAMB-785C8	Pending	hCoV-19/England/CAMB-810AD/2020	CAMB-810AD
EPI_ISL_433754	hCoV-19/England/CAMB-785E6/2020	CAMB-785E6	Pending	hCoV-19/England/CAMB-810BC/2020	CAMB-810BC
EPI_ISL_433755	hCoV-19/England/CAMB-785F5/2020	CAMB-785F5	Pending	hCoV-19/England/CAMB-810CB/2020	CAMB-810CB
EPI_ISL_433756	hCoV-19/England/CAMB-78601/2020	CAMB-78601	Pending	hCoV-19/England/CAMB-810E9/2020	CAMB-810E9
EPI_ISL_433757	hCoV-19/England/CAMB-7862F/2020	CAMB-7862F	Pending	hCoV-19/England/CAMB-81104/2020	CAMB-81104
EPI_ISL_433758	hCoV-19/England/CAMB-7864D/2020	CAMB-7864D	Pending	hCoV-19/England/CAMB-81113/2020	CAMB-81113
EPI_ISL_433759	hCoV-19/England/CAMB-7866B/2020	CAMB-7866B	Pending	hCoV-19/England/CAMB-81122/2020	CAMB-81122
EPI_ISL_433760	hCoV-19/England/CAMB-7867A/2020	CAMB-7867A	Pending	hCoV-19/England/CAMB-81131/2020	CAMB-81131
EPI_ISL_433761	hCoV-19/England/CAMB-78689/2020	CAMB-78689	Pending	hCoV-19/England/CAMB-81140/2020	CAMB-81140
EPI_ISL_433762	hCoV-19/England/CAMB-786A7/2020	CAMB-786A7	Pending	hCoV-19/England/CAMB-8115F/2020	CAMB-8115F
EPI_ISL_433763	hCoV-19/England/CAMB-786C5/2020	CAMB-786C5	Pending	hCoV-19/England/CAMB-8117D/2020	CAMB-8117D
EPI_ISL_433764	hCoV-19/England/CAMB-786E3/2020	CAMB-786E3	Pending	hCoV-19/England/CAMB-8118C/2020	CAMB-8118C
EPI_ISL_433765	hCoV-19/England/CAMB-7870E/2020	CAMB-7870E	Pending	hCoV-19/England/CAMB-8119B/2020	CAMB-8119B
EPI_ISL_433766	hCoV-19/England/CAMB-7872C/2020	CAMB-7872C	Pending	hCoV-19/England/CAMB-811AA/2020	CAMB-811AA
EPI_ISL_433767	hCoV-19/England/CAMB-7873B/2020	CAMB-7873B	Pending	hCoV-19/England/CAMB-811C8/2020	CAMB-811C8
EPI_ISL_433768	hCoV-19/England/CAMB-7874A/2020	CAMB-7874A	Pending	hCoV-19/England/CAMB-81F91/2020	CAMB-81F91
EPI_ISL_433769	hCoV-19/England/CAMB-78777/2020	CAMB-78777	Pending	hCoV-19/England/CAMB-81FA0/2020	CAMB-81FA0
EPI_ISL_433770	hCoV-19/England/CAMB-78786/2020	CAMB-78786	Pending	hCoV-19/England/CAMB-81FBF/2020	CAMB-81FBF
EPI_ISL_433771	hCoV-19/England/CAMB-787A4/2020	CAMB-787A4	Pending	hCoV-19/England/CAMB-81FCE/2020	CAMB-81FCE

GISAID Accession ID	Virus name	COG-UK	GISAID Accession ID	Virus name	COG-UK
EPI_ISL_433772	hCoV-19/England/CAMB-787B3/2020	CAMB-787B3	Pending	hCoV-19/England/CAMB-81FDD/2020	CAMB-81FDD
EPI_ISL_433773	hCoV-19/England/CAMB-787C2/2020	CAMB-787C2	Pending	hCoV-19/England/CAMB-81FEC/2020	CAMB-81FEC
EPI_ISL_433774	hCoV-19/England/CAMB-787D1/2020	CAMB-787D1	Pending	hCoV-19/England/CAMB-81FFB/2020	CAMB-81FFB
EPI_ISL_433775	hCoV-19/England/CAMB-787E0/2020	CAMB-787E0	Pending	hCoV-19/England/CAMB-82006/2020	CAMB-82006
EPI_ISL_433776	hCoV-19/England/CAMB-7880B/2020	CAMB-7880B	Pending	hCoV-19/England/CAMB-82015/2020	CAMB-82015
EPI_ISL_433777	hCoV-19/England/CAMB-7881A/2020	CAMB-7881A	Pending	hCoV-19/England/CAMB-82024/2020	CAMB-82024
EPI_ISL_433778	hCoV-19/England/CAMB-78829/2020	CAMB-78829	Pending	hCoV-19/England/CAMB-82033/2020	CAMB-82033
EPI_ISL_433779	hCoV-19/England/CAMB-78838/2020	CAMB-78838	Pending	hCoV-19/England/CAMB-82042/2020	CAMB-82042
EPI_ISL_433780	hCoV-19/England/CAMB-78847/2020	CAMB-78847	Pending	hCoV-19/England/CAMB-82051/2020	CAMB-82051
EPI_ISL_433781	hCoV-19/England/CAMB-78865/2020	CAMB-78865	Pending	hCoV-19/England/CAMB-82060/2020	CAMB-82060
EPI_ISL_433782	hCoV-19/England/CAMB-78874/2020	CAMB-78874	Pending	hCoV-19/England/CAMB-8207F/2020	CAMB-8207F
EPI_ISL_433783	hCoV-19/England/CAMB-78883/2020	CAMB-78883	Pending	hCoV-19/England/CAMB-8208E/2020	CAMB-8208E
EPI_ISL_433784	hCoV-19/England/CAMB-78892/2020	CAMB-78892	Pending	hCoV-19/England/CAMB-8209D/2020	CAMB-8209D
EPI_ISL_433785	hCoV-19/England/CAMB-788A1/2020	CAMB-788A1	Pending	hCoV-19/England/CAMB-820AC/2020	CAMB-820AC
EPI_ISL_433786	hCoV-19/England/CAMB-788B0/2020	CAMB-788B0	Pending	hCoV-19/England/CAMB-820CA/2020	CAMB-820CA
EPI_ISL_433787	hCoV-19/England/CAMB-788CF/2020	CAMB-788CF	Pending	hCoV-19/England/CAMB-82103/2020	CAMB-82103
EPI_ISL_433788	hCoV-19/England/CAMB-788DE/2020	CAMB-788DE	Pending	hCoV-19/England/CAMB-82121/2020	CAMB-82121
EPI_ISL_433789	hCoV-19/England/CAMB-788ED/2020	CAMB-788ED	Pending	hCoV-19/England/CAMB-82130/2020	CAMB-82130
EPI_ISL_433790	hCoV-19/England/CAMB-788FC/2020	CAMB-788FC	Pending	hCoV-19/England/CAMB-8214F/2020	CAMB-8214F
EPI_ISL_433791	hCoV-19/England/CAMB-78917/2020	CAMB-78917	Pending	hCoV-19/England/CAMB-8216D/2020	CAMB-8216D
EPI_ISL_433792	hCoV-19/England/CAMB-78926/2020	CAMB-78926	Pending	hCoV-19/England/CAMB-8217C/2020	CAMB-8217C
EPI_ISL_433793	hCoV-19/England/CAMB-78935/2020	CAMB-78935	Pending	hCoV-19/England/CAMB-8218B/2020	CAMB-8218B
EPI_ISL_433794	hCoV-19/England/CAMB-78944/2020	CAMB-78944	Pending	hCoV-19/England/CAMB-821A9/2020	CAMB-821A9
EPI_ISL_433795	hCoV-19/England/CAMB-78953/2020	CAMB-78953	Pending	hCoV-19/England/CAMB-8218B/2020	CAMB-8218B
EPI_ISL_433796	hCoV-19/England/CAMB-78962/2020	CAMB-78962	Pending	hCoV-19/England/CAMB-821C7/2020	CAMB-821C7
EPI_ISL_433797	hCoV-19/England/CAMB-78971/2020	CAMB-78971	Pending	hCoV-19/England/CAMB-82200/2020	CAMB-82200
EPI_ISL_433798	hCoV-19/England/CAMB-78980/2020	CAMB-78980	Pending	hCoV-19/England/CAMB-8222E/2020	CAMB-8222E
EPI_ISL_433799	hCoV-19/England/CAMB-7899F/2020	CAMB-7899F	Pending	hCoV-19/England/CAMB-7540A/2020	CAMB-7540A
EPI_ISL_433800	hCoV-19/England/CAMB-789AE/2020	CAMB-789AE	Pending	hCoV-19/England/CAMB-756D7/2020	CAMB-756D7
EPI_ISL_433801	hCoV-19/England/CAMB-789BD/2020	CAMB-789BD	Pending	hCoV-19/England/CAMB-756F5/2020	CAMB-756F5
EPI_ISL_433802	hCoV-19/England/CAMB-789CC/2020	CAMB-789CC	Pending	hCoV-19/England/CAMB-75701/2020	CAMB-75701
EPI_ISL_433803	hCoV-19/England/CAMB-789DB/2020	CAMB-789DB	Pending	hCoV-19/England/CAMB-7572F/2020	CAMB-7572F
EPI_ISL_433804	hCoV-19/England/CAMB-789EA/2020	CAMB-789EA	Pending	hCoV-19/England/CAMB-7576B/2020	CAMB-7576B
EPI_ISL_433805	hCoV-19/England/CAMB-789F9/2020	CAMB-789F9	Pending	hCoV-19/England/CAMB-7577A/2020	CAMB-7577A
EPI_ISL_433806	hCoV-19/England/CAMB-78A05/2020	CAMB-78A05	Pending	hCoV-19/England/CAMB-75798/2020	CAMB-75798
EPI_ISL_433807	hCoV-19/England/CAMB-78A14/2020	CAMB-78A14	Pending	hCoV-19/England/CAMB-757A7/2020	CAMB-757A7
EPI_ISL_433808	hCoV-19/England/CAMB-78A32/2020	CAMB-78A32	Pending	hCoV-19/England/CAMB-757B6/2020	CAMB-757B6
EPI_ISL_433809	hCoV-19/England/CAMB-78A50/2020	CAMB-78A50	Pending	hCoV-19/England/CAMB-757C5/2020	CAMB-757C5
EPI_ISL_433810	hCoV-19/England/CAMB-78A7E/2020	CAMB-78A7E	Pending	hCoV-19/England/CAMB-757E3/2020	CAMB-757E3
EPI_ISL_433811	hCoV-19/England/CAMB-78A9C/2020	CAMB-78A9C	Pending	hCoV-19/England/CAMB-757F2/2020	CAMB-757F2
EPI_ISL_433812	hCoV-19/England/CAMB-78ABA/2020	CAMB-78ABA	Pending	hCoV-19/England/CAMB-7580E/2020	CAMB-7580E
EPI_ISL_433813	hCoV-19/England/CAMB-79A04/2020	CAMB-79A04	Pending	hCoV-19/England/CAMB-7582C/2020	CAMB-7582C
EPI_ISL_433814	hCoV-19/England/CAMB-79A22/2020	CAMB-79A22	Pending	hCoV-19/England/CAMB-7584A/2020	CAMB-7584A
EPI_ISL_433815	hCoV-19/England/CAMB-79A40/2020	CAMB-79A40	Pending	hCoV-19/England/CAMB-75859/2020	CAMB-75859
EPI_ISL_433816	hCoV-19/England/CAMB-79A5F/2020	CAMB-79A5F	Pending	hCoV-19/England/CAMB-75868/2020	CAMB-75868
EPI_ISL_433817	hCoV-19/England/CAMB-79A8C/2020	CAMB-79A8C	Pending	hCoV-19/England/CAMB-75886/2020	CAMB-75886
EPI_ISL_433818	hCoV-19/England/CAMB-79A9B/2020	CAMB-79A9B	Pending	hCoV-19/England/CAMB-75895/2020	CAMB-75895
EPI_ISL_433819	hCoV-19/England/CAMB-79AB9/2020	CAMB-79AB9	Pending	hCoV-19/England/CAMB-758A4/2020	CAMB-758A4
EPI_ISL_433820	hCoV-19/England/CAMB-79AC8/2020	CAMB-79AC8	Pending	hCoV-19/England/CAMB-758B3/2020	CAMB-758B3
EPI_ISL_433821	hCoV-19/England/CAMB-79AD7/2020	CAMB-79AD7	Pending	hCoV-19/England/CAMB-758C2/2020	CAMB-758C2
EPI_ISL_433822	hCoV-19/England/CAMB-79AE6/2020	CAMB-79AE6	Pending	hCoV-19/England/CAMB-758D1/2020	CAMB-758D1
EPI_ISL_433823	hCoV-19/England/CAMB-79AF5/2020	CAMB-79AF5	Pending	hCoV-19/England/CAMB-758E0/2020	CAMB-758E0
EPI_ISL_433824	hCoV-19/England/CAMB-79B01/2020	CAMB-79B01	Pending	hCoV-19/England/CAMB-758FF/2020	CAMB-758FF
EPI_ISL_433825	hCoV-19/England/CAMB-79B10/2020	CAMB-79B10	Pending	hCoV-19/England/CAMB-78522/2020	CAMB-78522
EPI_ISL_433826	hCoV-19/England/CAMB-7A8BE/2020	CAMB-7A8BE	Pending	hCoV-19/England/CAMB-78768/2020	CAMB-78768
EPI_ISL_433827	hCoV-19/England/CAMB-7A8DC/2020	CAMB-7A8DC	Pending	hCoV-19/England/CAMB-78759/2020	CAMB-78759
EPI_ISL_433828	hCoV-19/England/CAMB-7A8FA/2020	CAMB-7A8FA	Pending	hCoV-19/England/CAMB-79934/2020	CAMB-79934
EPI_ISL_433829	hCoV-19/England/CAMB-7AA30/2020	CAMB-7AA30	Pending	hCoV-19/England/CAMB-79943/2020	CAMB-79943
EPI_ISL_433830	hCoV-19/England/CAMB-7AA4F/2020	CAMB-7AA4F	Pending	hCoV-19/England/CAMB-79961/2020	CAMB-79961
EPI_ISL_433831	hCoV-19/England/CAMB-7AA6D/2020	CAMB-7AA6D	Pending	hCoV-19/England/CAMB-79970/2020	CAMB-79970
EPI_ISL_433832	hCoV-19/England/CAMB-7AA7C/2020	CAMB-7AA7C	Pending	hCoV-19/England/CAMB-7999E/2020	CAMB-7999E
EPI_ISL_433833	hCoV-19/England/CAMB-7AA8B/2020	CAMB-7AA8B	Pending	hCoV-19/England/CAMB-799BC/2020	CAMB-799BC
EPI_ISL_433834	hCoV-19/England/CAMB-7AAA9/2020	CAMB-7AAA9			
EPI_ISL_433835	hCoV-19/England/CAMB-7AAB8/2020	CAMB-7AAB8			
EPI_ISL_433836	hCoV-19/England/CAMB-7AAC7/2020	CAMB-7AAC7			
EPI_ISL_433837	hCoV-19/England/CAMB-7AAD6/2020	CAMB-7AAD6			
EPI_ISL_433838	hCoV-19/England/CAMB-7AAE5/2020	CAMB-7AAE5			
EPI_ISL_433839	hCoV-19/England/CAMB-7AAF4/2020	CAMB-7AAF4			
EPI_ISL_433840	hCoV-19/England/CAMB-7AB00/2020	CAMB-7AB00			
EPI_ISL_433841	hCoV-19/England/CAMB-7AB1F/2020	CAMB-7AB1F			
EPI_ISL_433842	hCoV-19/England/CAMB-7AB2E/2020	CAMB-7AB2E			
EPI_ISL_433843	hCoV-19/England/CAMB-7AB3D/2020	CAMB-7AB3D			
EPI_ISL_433844	hCoV-19/England/CAMB-7AB79/2020	CAMB-7AB79			
EPI_ISL_433845	hCoV-19/England/CAMB-7AB88/2020	CAMB-7AB88			
EPI_ISL_433846	hCoV-19/England/CAMB-7AB97/2020	CAMB-7AB97			
EPI_ISL_433847	hCoV-19/England/CAMB-7ABA6/2020	CAMB-7ABA6			
EPI_ISL_433848	hCoV-19/England/CAMB-7ABB5/2020	CAMB-7ABB5			
EPI_ISL_433849	hCoV-19/England/CAMB-7ABC4/2020	CAMB-7ABC4			
EPI_ISL_433850	hCoV-19/England/CAMB-7ABE2/2020	CAMB-7ABE2			
EPI_ISL_433851	hCoV-19/England/CAMB-7ABF1/2020	CAMB-7ABF1			
EPI_ISL_433852	hCoV-19/England/CAMB-7AC0D/2020	CAMB-7AC0D			
EPI_ISL_433853	hCoV-19/England/CAMB-7AC1C/2020	CAMB-7AC1C			
EPI_ISL_433854	hCoV-19/England/CAMB-7AC2B/2020	CAMB-7AC2B			