

COVID-19: Easing the coronavirus lockdowns with caution

Akpojoto Siemuri M.Sc^{a*}, Rasheed Omobolaji Alabi M.Sc^{a*+}, Mohammed Elmusrati D.Sc^a

^aDepartment of Industrial Digitalization, School of Technology and Innovations, University of Vaasa, Vaasa, Finland.

Abstract count: 244

Total Text count: 1955

***The last two authors have equal contributions and first authorship.**

+ Corresponding Author: Rasheed Omobolaji Alabi

Department of Industrial Digitalization, School of Technology and Innovations, University of Vaasa, Vaasa, Finland. **E-mail address:** rasheed.alabi@student.uwasa.fi

Disclosure: The authors declare no conflicts of interest.

Abstract

Background: The spread of the novel severe acute respiratory syndrome coronavirus (SARS-CoV-2) has reached a global level, creating a pandemic. The government of various countries, their citizens, politicians, and business owners are worried about the unavoidable economic impacts of this pandemic. Therefore, there is an eagerness for the pandemic peaking. **Objectives:** This study uses an objective approach to emphasize the need to be pragmatic with easing of lockdowns measures worldwide through the forecast of the possible trend of COVID-19. This is necessary to ensure that the enthusiasm about SARS-CoV-2 peaking is properly examined, easing of lockdown is done systematically to avoid second-wave of the pandemic. **Methods:** We used the Facebook prophet on the World Health Organization data for COVID-19 to forecast the spread of SARS-CoV-2 for the 7th April until 3rd May 2020. The forecast model was further used to forecast the trend of the virus for the 8th until 14th May 2020. We presented the forecast of the confirmed and death cases. **Results:** Our findings from the forecast showed an increase in the number of new cases for this period. Therefore, the need for easing the lockdown with caution becomes imperative. Our model showed good performance when compared to the official report from the World Health Organization. The average forecasting accuracy of our model was 79.6%. **Conclusion:** Although, the global and economic impact of COVID-19 is daunting. However, excessive optimism about easing the lockdown should be appropriately weighed against the risk of underestimating its spread. As seen globally, the risks appeared far from being symmetric. Therefore, the forecasting provided in this study offers an insight into the spread of the virus for effective planning and decision-making in terms of easing the lockdowns in various countries.

KEYWORDS: COVID; Lockdowns, Ease, Forecasting; SARS-COV-2

1. Introduction

The year 2019 ended on a remarkable note over the novel severe acute respiratory syndrome coronavirus (SARS-CoV-2) [1]. Patients with SARS-CoV-2 infection can develop coronavirus disease 2019 (COVID-19) [2] with an incubation period ranging from 2 to 14 days [1]. The first case of the coronavirus disease 2019 (COVID-19) was reported in Wuhan, Hubei province, China in December 2019 [3]. It was reported to have originated in bats and transmitted to humans through yet unknown animals within Wuhan, China. From the primary place where it first emerged, it has spread globally. Thus, the World Health Organization (WHO) declared the outbreak a pandemic [4].

The mode of transmission in humans includes inhalation or contact with infected droplets where the symptoms include fever, cough, sore throats, difficulty in breathing properly, fatigue among others [1]. Of note, many people are asymptomatic (showing no symptoms of the virus), mild symptoms in some people, and severe symptoms in others especially the elderly and those with comorbidities. These severe symptoms may progress to multi-organ dysfunction, acute respiratory distress syndrome, respiratory failure, and pneumonia [1]. The severe symptoms have usually resulted in high rates of contacts to the healthcare providers, hospitalization, and admission to the intensive care units (ICU) [5]

Around 96, 000 cases were reported in early March 2020 (05/03/2020) [1]. By early April, the number of cases globally has increased to 1,603,330 (10/03/2020) [6]. The rise in the number of cases had resulted in various measures to reduce the spread of this virus. The example of such measures includes lockdown by countries [7]. Other measures include social distancing, contact tracing, quarantine, increased handwashing and sanitizing, temporary closedown of schools, businesses, and international trades. Therefore, the global economic impact of this pandemic is of great concern.

Entrepreneurs have been rendered temporarily unemployed. Governments have either introduced laws, update social welfare, or provides palliatives to ensure that the overall impacts of this pandemic on her citizens are relatively reduced. Though, the long term sustainability of these approaches remains doubtful. Thus, Government, politicians, and business owners are eager to get back to their normal daily routine. This eagerness has led to optimism about the pandemic peaking.

Despite the significant economic impacts of the COVID-19 pandemic for the past 100 days (since the virus had been identified), reasonable caution against excessive optimism about the pandemic peaking must be exercised to minimize the overall impacts (economic, social, and death) [8]. Therefore, this paper introduces an objective approach to forecasting the global spread of COVID-19 crisis. Accurate forecasting of the associated risks and spread of the virus is necessary to ensure that the implications can be properly analyzed, planning by the government and business owners improved, and informed decisions about the pandemic can be made in terms of easing the lockdown.

2. Methods

2.1. World Health Organization (WHO) data

The study data was obtained from the World Health Organization (WHO). The WHO COVID-19 data was used to assess the worldwide and country-specific trends of COVID-19 new confirmed and death cases around the world. The data contained cases from 09 January 2020 until 6 April 2020. The input variables contained in the dataset were day, country, country name, region, death, cumulative deaths, confirmed, cumulative confirmed. As the data is publicly available, informed consent was not needed.

2.2. Trend of the virus

The forecasting of the trend of the virus was done using Facebook's Prophet in Google Colaboratory (a free Jupyter notebook environment). The dataset and the corresponding libraries were imported to Jupyter Notebook. The data was further preprocessed to remove missing values and ensure that the data is in the proper format. Additionally, the columns were renamed for convenience in the analyses. The visualization of the top 11 countries, Finland, and Nigeria (most populous African country) as of 7th May 2020 is given in Figure 1. The actual decomposition of the time series data was done using an additive model. The additive model has been implemented because it is easy to develop, fast to train, and provides interpretable patterns ⁹. We used the last 3 months of the COVID-19 cases to forecast for the next 5 days (8th until 14th May 2020) using the Facebook Prophet, with a 95% prediction interval by creating a base model using seasonality-related parameters and additional regressors. Note that the yearly and daily seasonality was not set since COVID-19 is not a seasonal disease. The forecast given by our model was compared with the official data reported on the World Health Organization (WHO) official website for COVID-19 updates for the same date. The dashboard of WHO's official website for comparison is available at <https://covid19.who.int/>. The result of the comparison for 07.04.2020 – 03.05.2020 is presented in Table 1. We also presented the model forecast for the 8th until 14th May 2020 to further validate our model prior to the official release of the cases by the World Health Organization.

3. Results

The study cohort included 210 countries at the time of data extraction from WHO official website (06.04.2020). After building the forecast model, it showed good forecasting accuracy

in predicting the confirmed and death cases for the date of data retrieval (6.04.2020) as shown in Table 1, respectively. As shown in Table 1, the prediction accuracy of the model for the confirmed cases was 79.6% and 80.3% (average of 79.9%) for the death cases when its performance for forecasting for the period 07.04.2020 – 03.05.2020 was compared with the reported cases on WHO official website for confirmed and death cases, respectively. The details of the forecast for this period is given in Table 1. As shown in Figures 2-3, there was increase in the number of cases, although not spike increase in the number of confirmed and death cases as seen in previous months. Additionally, the next 7 days (8th – 14th May 2020) prediction of the pandemic COVID-19 showed a rise in the number of confirmed cases and deaths from COVID-19 globally (Figure 2-3). As shown in Figure 2, few cases were confirmed between Mondays until Wednesdays. Likewise, most deaths occurred between Tuesdays and Thursdays (Figure 3). Towards easing the lockdown, the World Health Organization published six (6) important criteria that includes evidence that the transmission of COVID-19 is controlled, effective mechanism of contract tracing, quarantine, and testing, minimized outbreaks amongst the vulnerable in care homes, continuous adherence to preventive measures (physical distancing, handwashing, etc.), importation risks can be managed, and proper information and engagement of the communities concerning the pandemic [9]. Most countries have published tentative plans to ease the lockdown measures to address the spread of the COVID-19 pandemic. As shown by our predictive model, restarting the economies that had been significantly affected by the sudden shutdowns to curb the devastating effects of COVID-19 this should be done with caution [10]. This is necessary to avoid a possible second-wave of the virus which could have more damaging consequences [10].

4. Discussion

In this study, we used a time series forecasting approach to forecast the trend of COVID-19 from 07.04.2020 until 03.05.2020 (Table 1) and forecast the trend for the next 7 days (08-14.05.2020). This approach was aimed at emphasizing that the easing of lockdowns in various countries should be done systematically to avoid second-wave of the pandemic. We produced a forecast model using a Facebook prophet that used additive models [11]. A Facebook prophet was used as it is accurate, fast, fully automatic, offers tunable forecast, and compatible with Python or R [12]. Of note, as this is not a seasonal occurrence and the fact that there is no known vaccine yet, we assume that the trend will continue indefinitely in the modelling. The average forecasting accuracy of our model was 79.9%. The differences between our forecast model and the WHO official reported cases may be due to the possible errors, delay or inadequate reporting by various countries. Despite this variation, our forecast model still showed reasonable accuracy to forecast the confirmed and death cases for the target dates. The fact that the confirmed cases increases significantly between Wednesdays and Saturdays may account for the time taken to get and record the results from these tests. Similarly, the curve decreased on the weekend as most death cases were usually reported and recorded during the weekdays.

Our model produced a comparable performance in terms of the trend of the pandemic as shown in Table 1. It was observed that, there was no spike increase in the number of confirmed and death cases within this period. This may suggest that the pandemic may have started to peak in various countries. Despite this, for the period of 8th- 14th May 2022, our model predicted a rise in the number of confirmed cases globally (Figures 2-3). Therefore, excessive optimism about the easing the lockdown in various countries should be met with caution. This is because, at present, the life-and-death pressure on our health facilities is enormous [13–15]. The testing places and hospitals currently have limited or no resources to treat COVID-19 patients [14]. Thus, change of policies to relax the lockdown and social

distancing should be done systematically, otherwise, an increase in the number of hospitalized patients may be inevitable. With the current states of the various hospitals globally, it may be difficult to properly manage COVID-19 patients.

The global infection is expected to rise to 40-70% in the coming year [16]. The concern about the rise in the number of confirmed cases was also echoed by the Governor of Massachusetts echoed when he projected that between 0.7 and 2.5 % of the state's population are likely to contract the coronavirus [17]. The main concern is for how long do we continue to flatten the curve with the lockdown, quarantine, contract tracing, and social distancing measures? Admittedly, the economic and global impact as a result of these measures is massive. Therefore, the concerned authorities (government, politicians and other stakeholders) have started to relax the lockdown policies to minimize the economic impacts.

However, as predicted by our forecast model, a possible rise in the number of confirmed and death cases emphasize the need to have widespread testing in place and hospitals have the resources needed to treat COVID-19 patients prior to switching of gears to allow for controlled exposures [17]. Therefore, the government and policymakers need to get the trend of the spread of the virus right. The understanding of the trajectory of COVID-19 spread ensures that the right decision regarding public policies is made [17]. As such, with the forecast of the possible rise in the number of confirmed and death cases as presented by our model, and the suggestion that only possible COVID-19 vaccine may bring back normalcy [18,19], informing the citizens of the possibility of spending a year or more at home prior to the development of a possible vaccine for COVID-19, or get immune in a hard way becomes pertinent [17].

As pharmaceutical treatments for COVID-19 are yet to be found, the modality of exposure of citizens to the virus needs to be controlled. With herd immunity (exposure to the

virus), the vulnerable citizens fall to save the less vulnerable. The main question with this strategy is how this can be done ethically and responsibly. As predicted by our model, the rise in the cases meant that exposure of the world population should be done reasonably, sensibly and controllably for the citizens to acquire herd immunity while the world awaits the breakthrough in the vaccine for COVID-19 [19]. Thus, the government needs to strike a balance between the health and economic impact of this pandemic. If this is not properly considered, there is a high risk of viral reintroduction (those tested negative after initial exposure) [19], intense pressure on health facilities, and increased deaths from COVID-19. As various countries such as France starts to ease the lockdown, important measures such as making face masks compulsory, extra vigilance in the capital regions, reducing the number of public gatherings and events, and temporary closing of place of schools and places worships may still be in effect to avoid a possible second-wave of the pandemic [20].

In conclusion, despite the economic impact of COVID-19, the government and policy-makers in various countries should be very pragmatic in the easing the lockdown measures. Systematic easing of the lockdowns ensures that curve is not only flattened but crushed with controlled exposure to the virus pending the breakthrough of possible pharmaceutical treatment. The fact that our model predicted a spike in the rise should not lead to overly panicking [21] but to understand the trend for the proper decision-making process, especially in high-risk cases. This ensures that we are not only flattening the curve but crushing the curve through controlled herd immunity of the world's population.

Figure Legend

Figure 1. The number of confirmed cases and death in top 11 countries in the world, Finland and Nigeria.

Figure 2. Forecast plot for new confirmed NCOVID-19 cases

Figure 3. Forecast plot for new death in NCOVID-19 cases.

Table Legend


Table 1. The comparison of the forecast model with the reported cases on WHO's official website

REFERENCES




- 1 Singhal T. A Review of Coronavirus Disease-2019 (COVID-19). *The Indian Journal of Pediatrics* 2020;**87**:281–6. doi:10.1007/s12098-020-03263-6
- 2 Grasselli G, Zangrillo A, Zanella A, *et al.* Baseline Characteristics and Outcomes of 1591 Patients Infected With SARS-CoV-2 Admitted to ICUs of the Lombardy Region, Italy. *JAMA* Published Online First: 6 April 2020. doi:10.1001/jama.2020.5394
- 3 Wang C, Horby PW, Hayden FG, *et al.* A novel coronavirus outbreak of global health concern. *The Lancet* 2020;**395**:470–3. doi:10.1016/S0140-6736(20)30185-9
- 4 World Health Organization. WHO Director-General's opening remarks at the media briefing on COVID-19. 2020.0. <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020> (accessed 10 Apr 2020).
- 5 Grasselli G, Pesenti A, Cecconi M. Critical Care Utilization for the COVID-19 Outbreak in Lombardy, Italy: Early Experience and Forecast During an Emergency Response. *JAMA* Published Online First: 13 March 2020. doi:10.1001/jama.2020.4031
- 6 Center for Systems Science and Engineering (CSSE). Coronavirus COVID-19 global cases by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins. Updated April 10, 2020. <https://gisanddata.maps.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467b48e9ecf6> (accessed 10 Apr 2020).
- 7 Juliana K, Lauren F, Morgan MJ. A third of the global population is on coronavirus lockdown - here's our constantly updated list of countries and restrictions.

- 2020.<https://www.businessinsider.com/countries-on-lockdown-coronavirus-italy-2020-3?r=US&IR=T>
- 8 Wang V. Coronavirus epidemic keeps growing but spread in China slows. 2020. <https://www.nytimes.com/2020/02/18/world/asia/china-coronavirus-cases.html?referringSource=articleShare> (accessed 10 Apr 2020).
- 9 World Health Organization. WHO/Europe published considerations for gradual easing of COVID-19 measures. 2020.<http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/news/news/2020/4/whoeurope-publishes-considerations-for-gradual-easing-of-covid-19-measures> (accessed 7 May 2020).
- 10 Alasdair S. Coronavirus lockdowns: How and when do European countries plan to ease restrictions? Euro News. 2020.<https://www.euronews.com/2020/03/19/coronavirus-which-countries-are-under-lockdown-and-who-s-next>
- 11 Facebook Prophet. Facebook Open Source. Prophet: Forecasting at Scale. Documentation. <https://facebook.github.io/prophet/>; <https://facebook.github.io/prophet/docs/>; <https://github.com/facebook/prophet>; https://facebook.github.io/prophet/docs/quick_start.html (accessed 6 Apr 2020).
- 12 Taylor SJ, Letham B. Forecasting at scale. PeerJ Preprints 2017. doi:10.7287/peerj.preprints.3190v2
- 13 Li R, Rivers C, Tan Q, *et al.* The demand for inpatient and ICU beds for COVID-19 in the US: lessons from Chinese cities. Infectious Diseases (except HIV/AIDS) 2020. doi:10.1101/2020.03.09.20033241
- 14 Nicole Winfield. Not a wave, a tsunami. Italy hospitals at virus limit (AP News). 2020. <https://apnews.com/a4497f31bf5dbc1ff263e4263fc9f69e> (accessed 6 Apr 2020).
- 15 Marina V, Michael H. COVID-19 infections rise in New York with peak weeks away (AP News). 2020. <https://apnews.com/7c7563cb82626a4042797c6aa6da260a> (accessed 6 Apr 2020).
- 16 Nash CM. Harvard Professor Sounds Alarm on 'Likely Coronavirus Pandemic: 40% to 70% of world could be infected this year. <https://www.mediaite.com/news/harvard-professor-sounds-alarmon-likely-coronavirus-pandemic-40-to-70-of-world-could-be-infected-this-year/> (accessed 15 Apr 2020).
- 17 Jeff H. The only way this ends: herd immunity. It has a bad rap, but in the long run it's our best hope. 2020. <https://www.bostonglobe.com/2020/04/10/opinion/its-possible-flatten-curve-too-long/> (accessed 15 Apr 2020).
- 18 The Guardian News. Only a COVID-19 vaccine will allow return to "normalcy": UN chief. 2020. <https://guardian.ng/news/world/only-a-covid-19-vaccine-will-allow-return-to-normalcy-un-chief/> (accessed 16 Apr 2020).
- 19 Leung K, Wu JT, Liu D, *et al.* First-wave COVID-19 transmissibility and severity in China outside Hubei after control measures, and second-wave scenario planning: a modelling impact assessment. *The Lancet* Published Online First: April 2020. doi:10.1016/S0140-6736(20)30746-7




- 20 France 24 News. France confirms plan to start easing COVID-19 lockdown on May 11. 2020.<https://www.france24.com/en/20200507-live-france-unveils-final-plan-on-easing-covid-19-lockdown> (accessed 7 May 2020).
- 21 Salzberg F. There are better things to do than panic.
<https://www.forbes.com/sites/stevensalzberg/2020/02/29/coronavirus-time-to-panic-yet/> (accessed 15 Apr 2020).



Forecast Plot for New Confirmed NCOVID-19 cases



Forecast Plot for New Deaths in NCOVID-19 cases



Validated Forecasted data						
Date	Model Forecast for confirmed cases	WHO Reported cases for confirmed cases	Model Accuracy	Model Forecast for death cases	WHO Reported cases for death cases	Model Accuracy
7.4.2020	84833	68465	76,1	5623	5009	87,7
8.4.2020	88895	73849	79,6	5975	6685	89,4
9.4.2020	93295	82736	87,2	6278	6325	99,3
10.4.2020	95575	84635	87,1	6467	7231	89,4
11.4.2020	100173	90778	89,7	6891	6946	99,2
15.4.2020	92791	69728	66,9	7154	5909	78,9
16.4.2020	98110	77846	74,0	7394	7911	93,5
17.4.2020	86878	82622	94,8	7597	8473	89,7
18.4.2020	88480	85501	96,5	6926	6681	96,3
19.4.2020	87380	82180	93,7	6860	6507	94,6
20.4.2020	90228	72397	75,4	6950	5261	67,9
21.4.2020	91276	83596	90,8	7083	5118	61,6
22.4.2020	92340	73448	74,3	7430	6060	77,4
23.4.2020	97801	73911	67,7	7747	6675	83,9
24.4.2020	95541	82089	83,6	7562	6290	79,8
28.4.2020	94744	73759	71,5	6925	3894	22,2
29.4.2020	95283	66321	56,3	7297	5378	64,3
30.4.2020	99878	72955	63,1	7598	9785	77,6
1.5.2020	101002	84983	81,2	7719	6405	79,5
2.5.2020	104273	90520	84,8	7808	5803	65,4
3.5.2020	102431	83985	78,0	7721	8626	89,5
		Average	79,6		Average	80,3