Type of article: Original

Title of the article: Knowledge and awareness of Dengue and Chikungunya amidst recurrent outbreaks amongst urban slum community members of Pune, India

Running title: Knowledge and awareness of Dengue and Chikungunya amidst outbreaks

Contributors:

1. Godbharle Swapnil R., M.Sc. Health Sciences, Interdisciplinary School of Health Sciences, Savitribai Phule Pune University, Pune
2. Giri Bibek R., Master of Public Health, Interdisciplinary School of Health Sciences, Savitribai Phule Pune University, Pune
3. Kudale Abhay M., Ph.D., Interdisciplinary School of Health Sciences, Savitribai Phule Pune University, Pune

Department(s) and institution(s):

Interdisciplinary School of Health Sciences, Savitribai Phule Pune University, Ganeshkhind, Pune – 411007

Corresponding Author:

Name: Abhay M. Kudale

Address: Interdisciplinary School of Health Sciences

Phone number: +91-9881435808

E-mail address: amkudale@unipune.ac.in, abhay.kudale@gmail.com

○ **ORCID (Open Researcher and Contributor ID)**

Swapnil Rajendra Godbharle: ORCID ID: <https://orcid.org/0000-0001-5793-1285>

Bibek Raj Giri: ORCID ID: <https://orcid.org/0000-0001-7829-0878>

Abhay Machindra Kudale: ORCID ID: <https://orcid.org/0000-0003-2887-1636>

Title of the article: Knowledge and awareness of Dengue and Chikungunya amidst recurrent outbreaks amongst urban slum community members of Pune, India

Abstract:

Context: World health organization's (WHO) new strategy on Global Vector Control Response (GVCR) considered engaging and mobilizing communities and emphasized on local adaptation of communication strategies for achieving sustainable impact in vector control. Public awareness studies amidst and post epidemic/ outbreak situations can fast-track useful information to local programmes to set risk communication agenda.

Aims: To assess knowledge and awareness regarding dengue and chikungunya among general community members during and post-epidemic/outbreak situations.

Settings and Design: Cross-sectional study in slums of Pune city wherein several recurrent outbreaks of Dengue and Chikungunya were reported.

Methods and Material: In total 309 general community members residing in urban slums of Pune in Maharashtra were randomly chosen using population proportionate sampling. Semi-structured interview schedule was administered.

Statistical analysis used: Knowledge and awareness scores for dengue and chikungunya were calculated and analytical statistical tests were performed.

Results: It was found that most of the respondents were in the 18-39 years age group, ever married, literate, and had regular income. The study showed relatively good knowledge among respondents about symptoms, causes, mode of transmission and prevention methods for dengue and chikungunya. Respondents scored well with regard to awareness indicators concerning availability of supportive and symptomatic treatments for dengue and chikungunya. Average knowledge and awareness score for dengue and chikungunya together were 5.4 ± 1.96 (SD) and 1.68 ± 0.63 (SD) respectively. Better knowledge and awareness levels were reported among literate respondents.

Conclusions: Findings indicate need for sustaining these better knowledge and awareness levels into efficacious and community-owned prevention practices to fast-track appropriate public health actions during and post-epidemic situations leading to sustained community mobilization and engagement to promote successful implementation of WHO recommended GVCR strategy.

Key-words: Dengue; Chikungunya; Knowledge; Public Awareness; WHO GVCR

Key Messages: Findings indicate capitalizing on better knowledge and awareness levels achieved due to public health campaigns amidst recurrent epidemic/outbreak situations to facilitate appropriate public health actions on continual basis for strengthening vector control response and suggests need for aggressive and sustained health education campaigning to promote implementation of WHO recommended GVCR strategy.

Background: Worldwide social, demographic and environmental factors strongly influence transmission patterns of vector-borne pathogens, with major outbreaks of vector borne diseases such as Dengue, Chikungunya and Zika virus disease since 2014^{1,2}. In view of these ongoing VBD outbreaks in many countries in 2017, the World Health Organization has developed a new strategy to strengthen vector control globally, widely regarded as the Global vector control response 2017-2030 (GVCR). The 70th World Health Assembly adopted resolution of GVCR and encouraged Member States to develop or adapt national vector control strategies and operational plans in consonance with this global strategy. Engaging and mobilizing communities has been considered as one of the important pillars of action in GVCR strategy. And in spelt out research needs GVCR document emphasized on adaptation of information, education and communication strategies to ensure acceptability, participation and appropriate use of vector control tools^{1,2}.

Vector control has been considered worldwide as one of the best ways to control any VBDs. The success of vector control methods depends on public engagement and pro-active social mobilization. It is important to assess public awareness regarding the VBDs, along with its mode of transmission and prevention measures³. Public awareness studies can serve as a useful tool to strengthen vector control response and necessary public health actions⁴. This information can help the programmes to set communication agenda linked to increased public engagement and mobilizing demand for necessary services. It can also help in developing tailored strategies which can be appropriate for the socio-cultural and politico-economic contexts of different communities². Pune city in Maharashtra India had been hit by several Dengue and Chikungunya outbreaks in recent years³. The poor environmental and sanitation conditions in the slums not only contribute to the spread of the disease but also make it difficult for local municipal corporation health services to control the vector population effectively in these areas. Hence, a cross-sectional study was carried out to assess the knowledge and awareness regarding Dengue and Chikungunya among general community members residing in slums of Pune city.

Subjects and Methods:

Study setting: Pune is the second largest city in the Indian state of Maharashtra, after Mumbai. It is the ninth most populous city in the country with an estimated population of 3.13 million⁵. Along with its extended city limits Pimpri Chinchwad and the three cantonment towns of Pune, Khadki and Dehu Road, Pune forms the urban core of the Pune Metropolitan Region (PMR)⁶. According to the 2011 census, the urban area has a combined population of 5.05 million while the population of the metropolitan region is estimated at 7.27 million⁷.

A community-based cross-sectional study was carried out during 2017-18 in urban slums which were low-income settings in Pune Municipal Corporation (PMC), Maharashtra, India. The sampling method

was driven by the need to include a representative number of participants for the study purposes. The results of previous knowledge and awareness studies on Dengue and Chikungunya in India were reviewed taking into consideration key areas of enquiry and indicators to be examined in the present study⁸⁻¹⁰. The review provided the expected minimum value of knowledge and awareness proportion as 0.25. The formula for calculating sample size for proportions of large populations was used. One-third of the total 15 administrative wards in Pune city (5 administrative wards) were randomly selected for the survey. Population proportionate sampling was used to calculate a sample size of 309 respondents from 5 administrative wards. From each of the selected wards one low-income setting officially regarded as slum having highest number of slum structures was selected. Altogether a total of five slums (low-income settings) were selected for the study. Among the general community member participants from each slum a 1:1 sex ratio was maintained. Based on the number of slum structures of five selected slums, the population proportionate sampling for each respective slum was calculated. The households from every selected slum were randomly selected and one adult member from each household was interviewed if fitting into the pre-decided inclusion criteria. Inclusion criteria were community members who were residents of PMC (Pune Municipal Corporation) area and aged between 18-60 years.

A semi-structured pre-tested interview schedule was administered through tablet version of “EpiCollect 5” for data collection by trained investigators¹¹. All the questions asked in the interview schedule were multiple choice questions. Pilot testing was done before conducting the main survey study. For validation of tool, the interview schedule was checked for the flow and consistency of questions, and then it was revised which mainly comprised of formatting changes. The survey helped in documenting public knowledge and awareness about Dengue and Chikungunya symptoms, causes, modes of transmission, and prevention methods, as well as awareness about availability of the treatment. The complete data collection process was closely monitored by research supervisor to ensure overall quality control. Written informed consent was obtained from individual general community member prior to the interview. The entire study protocol was approved by institutional ethics committee of Savitribai Phule Pune University (Ref. SPPU/ISHS/117/2018).

Survey data were analysed into SPSS (Statistical Package for the Social Sciences, Chicago, IL, USA), version 19. Descriptive statistics were initially performed to summarize data, and the data were categorized into knowledge and awareness indicators to calculate the scores for each category. Knowledge indicators were those providing biomedical information or facts about Dengue and Chikungunya, while awareness indicators were related to the availability of treatment services and hearing information about Dengue and Chikungunya. The Chi-square test was performed to determine the factors associated with socio-demographic indicators such as gender, age, education and marital status. The t-test was used to study association between knowledge and awareness indicators across

education. For all analysis purposes, p value less than 0.05 (two-tailed) was considered for statistical significance.

Development of Dengue and Chikungunya Composite knowledge and awareness scores:

Descriptive statistics were initially performed to summarize data, and the data were categorized into knowledge and awareness indicators to calculate the scores for each category. After testing correlation between items for each knowledge topic, 9 knowledge items were retained as valid and included in the calculation of knowledge score as follows: for each individual item, correct answer recoded into '1' and 'incorrect or don't know' answers into '0'. A single summary score for each knowledge topic was obtained by summing the number of correct responses and dividing the total by the number of items in the topic. An aggregate measure was then computed, summing the values of each topic yielding a knowledge score for each participant. The theoretical range of knowledge was 0-9, with a higher score indicating a higher level of knowledge. The average knowledge-scores for the group were obtained by summing each participant's score and dividing it by the total number of survey participants.

Calculation of maximal mean knowledge and awareness score: Mean knowledge and awareness score by topic as percentages of maximal scores were calculated and considered as maximal mean knowledge and awareness scores. For mean knowledge score maximal theoretical possible score was 9 and for mean awareness score maximal theoretical possible score was 2.

Results:

Out of the total respondents 50% (154) were men and 50% (155) were women. The mean age of respondents was 32 ± 10.77 (SD). More than half ($n = 231$, 75%) of the total 309 respondents were in the younger age group (18-39). Of total, almost three-fourth of the respondents ($n=229$, 74%) were under the category of ever married and rest were in never married category. Among the respondents, 91% ($n=280$) respondents were literate, that means they have at least attended formal education. Table 1 shows the socio-demographic characteristics of the respondents.

Table 1: Socio-demographic profile of the respondents (n = 309)

Socio-demographic characteristics	Total n (%)	Men n (%)	Women n (%)	P-Value
Age-group in years				
18 -39	231(75)	114 (74)	117 (76)	0.869
40-60	078 (25)	040 (26)	038 (24)	
Marital Status				
Never Married	080 (26)	58 (38)	022 (14)	0.0000**
Ever Married	229 (74)	96 (62)	133 (86)	
Education				
Illiterate	029 (09)	07 (5)	22 (14)	0.00369*
Literate	280 (91)	147(95)	133 (86)	
Occupation				
Unemployed	04 (1)	04 (3)	00 (0.0)	Not-significant
Service (Govt.+Pvt.)	112 (36)	91(59)	21 (13)	
Petty Business	35 (11)	27 (17)	08(5)	
Students	51 (17)	32(21)	19 (13)	
Homemaker	107 (35)	0	107 (69)	
Income regular and dependable				
Yes	206 (67)	106 (69)	100 (64)	0.840
No	103 (33)	55 (35)	48 (31)	

Chi-square test applied, *P value= <0.05: Significant and **P value=<0.001: Highly significant

Table 2 shows the respondents knowledge about Dengue and Chikungunya. For Dengue high fever (76%) was the most commonly reported symptom, followed by nausea and vomiting (34%), while for Chikungunya joint swelling (65%) was the most commonly reported symptom, followed by high fever (32%). More than one-third of the respondents (44%) reported virus as a cause for Dengue and two-third of the respondents (66%) were able to report virus as a cause for Chikungunya. More than two-thirds (70%) reported mosquito bite as a mode of transmission of Dengue whereas more than three-fourth of the respondents (78%) were able to report mosquito bite as a mode of transmission for Chikungunya. Prevent standing/ stagnant water near house was reported as a common way of preventing Dengue by 81% (n = 250) of the respondents, followed by Indoor residual spray (71%). Almost 90% of the respondents were able to report that prevention methods of Chikungunya are similar to those of Dengue. There was no statistically significant gender-differences across knowledge indicators of Dengue and Chikungunya. Respondents knowledge about Dengue and Chikungunya was found to be relatively good. The average knowledge score for Dengue and Chikungunya together was 5.4 ± 1.96 (SD), with a theoretical maximum of 9.

Table 2: Distribution of knowledge indicators by gender

Items accurately identified as correct	Total (n=309) (Correct answers)	Men (n=154) n (%)	Women (n=155) n (%)
Symptoms related to Dengue			
High Fever	235 (76)	114 (74)	121 (78)
Headache	64 (21)	36 (23)	28 (18)
Pain behind eyes	55 (18)	28 (18)	27 (17)
Joint pain	88 (29)	55 (36)	33 (21)
Muscle and/or bone pain	53 (17)	31 (20)	22 (14)
Rash	57 (18)	18 (12)	39 (25)
Nausea and vomiting	104 (34)	49 (32)	55 (35)
Symptoms related to Chikungunya			
High Fever	100 (32)	49 (32)	51 (33)
Headache	31 (10)	15 (10)	16 (10)
Nausea and vomiting	51 (17)	27 (17)	24 (15)
Joint swelling	200 (65)	96 (62)	104 (67)
Muscle pain	97 (31)	53 (34)	44 (28)
Cause of Dengue			
Virus	136 (44)	72 (47)	64 (41)
Cause of Chikungunya			
Virus	193 (66)	101 (66)	92 (59)
Mode of transmission of Dengue			
Through mosquito bite	215 (70)	101 (66)	114 (74)
Mode of transmission of Chikungunya			
Through mosquito bite	240 (78)	117 (76)	123 (79)
Prevention methods for Dengue			
Bed nets	96 (31)	45 (29)	51 (33)
Indoor residual spray	219 (71)	112 (73)	107 (69)
Prevent standing water near house	250 (81)	128 (83)	122 (79)
Applying mosquito repellent on body	98 (32)	46 (30)	52 (34)
General Prevention methods for Chikungunya			
Similar to Dengue	278 (90)	143 (93)	135 (87)

Table 3 shows the respondents awareness about Dengue and Chikungunya. More than three-fourth of the respondents (90%) reported that they had previously heard about Dengue. When asked about availability of supportive and symptomatic treatment, 89% and 80% reported that such supportive and symptomatic treatment is available for Dengue and Chikungunya respectively. There was no statistically significant gender-differences across awareness indicators of Dengue and Chikungunya. Respondents awareness about Dengue and Chikungunya was found to be relatively good. The average awareness score for Dengue and Chikungunya together was 1.68 ± 0.63 (SD), with a maximum of 2.

Table 3: Distribution of awareness indicators by gender			
Items accurately identified as correct	Total (n=309) (Correct answers)	Men (n=154) n (%)	Women (n=155) n (%)
Heard about Dengue or not			
Yes	293 (90)	151 (98)	142 (92)
Availability of supportive and symptomatic treatment for Dengue			
Yes	275 (89)	139 (90)	136 (88)
Availability of supportive and symptomatic treatment for Chikungunya			
Yes	246 (80)	123 (80)	123 (79)

The following figures 1 and 2 shows gender differences across the respondents correct knowledge and awareness for Dengue and Chikungunya. Although not statistically significant, in case of Dengue women have had better knowledge while men have had slightly better awareness with regard to Dengue. In case of Chikungunya, men respondents have had better knowledge and awareness as compared to women respondents, however, difference was not statistically significant.

Figure 1: Proportion of maximal mean knowledge score (%) by topic as percentage of maximal score for Dengue by gender

Figure 2: Proportion of maximal mean knowledge score (%) by topic as percentage of maximal score for Chikungunya by gender

Table 4 shows literacy status-wise differences about Dengue and Chikungunya across knowledge and awareness indicators. The t-test results suggest that there is a significant statistical difference in knowledge and awareness about symptoms, cause, prevention methods, mode of transmission and availability of treatment. Table 4 clearly shows that literate respondents had more knowledge and awareness about Dengue and Chikungunya than their illiterate counterparts.

Table 4: Literacy status-wise differences across knowledge and awareness indicators of Dengue and Chikungunya						
Parameters	Illiterate n (%)	Literate n (%)	t-test	df	Significance (2-tailed)	P value
Knowledge						
Symptoms of Dengue and Chikungunya	22 (76)	246 (88)	22.79	308	0.00	< 0.001*
Causes of Dengue and Chikungunya	9 (31)	207 (74)	26.74	308	0.00	< 0.001*
Prevention & transmission of Dengue and Chikungunya	28 (96)	272 (97)	101.32	308	0.00	< 0.001*
Awareness						
Availability of supportive and symptomatic treatment for Dengue and Chikungunya	18 (62)	263 (94)	55.59	308	0.00	< 0.001*

* Highly Significant
df = degree of freedom

Discussion: Against the background of the occurrence of recurrent outbreaks of Dengue and Chikungunya in Pune city in recent years¹², present cross-sectional study documented the knowledge and awareness of general community members residing in low-income settings regarded as slums in Pune city with regard to Dengue and Chikungunya. Public awareness is considered a key to successful vector control programmes⁴. Such awareness is crucial as it improves people's understanding of risks and of prevention strategies and is considered a key to success in vector-borne diseases control programmes^{2,4}.

The study findings documented relatively better Dengue-related knowledge and awareness among the general population residing in slum areas of study amidst recurrent outbreaks. Similar findings related to awareness about Dengue was also documented in a study conducted in urban low-income areas of south Delhi⁸. Knowledge about Dengue was comparatively more in women respondents while awareness was found to be comparatively more among the men respondents. The literate respondents had better knowledge and awareness about Dengue as compared to illiterate respondents. Similar findings have been reported by a study conducted in Karachi, Pakistan¹³. High fever was regarded as the most commonly reported symptom of Dengue. Similar findings have been reported by other studies conducted in different cities of India^{9,14}. Knowledge about causes of Dengue was relatively low, as only 44% of respondents could report virus as a cause of Dengue. Most of the respondents reported mosquito bite as the mode of transmission of Dengue. Similar findings have been reported by a study conducted in Myanmar¹⁵. Stagnant water management was most commonly reported preventive method for Dengue. Most of the respondents in this study had previously heard information about Dengue. Similar findings have been reported by a study conducted in urban areas of south Delhi⁸. Most of the respondents in the study were aware about the availability of supportive and or symptomatic treatment for Dengue. The study findings also indicated that there was relatively better Chikungunya-related knowledge and awareness. Knowledge about Chikungunya was more among men respondents while awareness was almost identical in both men and women respondents. The literate respondents had better knowledge

and awareness about Chikungunya as compared to illiterate respondents. Joint swelling was the most commonly reported symptom of Chikungunya. Knowledge about cause of Chikungunya was relatively high, and majority respondents could report virus as a cause of Chikungunya. Most of the respondents reported mosquito bite as the mode of transmission of Chikungunya. Similar findings have been reported by a study conducted in Dhaka city of Bangladesh¹⁶. Most of the respondents reported that prevention methods for both Dengue and Chikungunya are similar.

Most of the respondents were unable to differentiate and distinguish symptoms of VBDs. They mentioned few general symptoms of febrile illnesses. Also, respondents had comparatively less information and knowledge about causative agents of the Dengue and Chikungunya. These finding would feed and enable strengthening of the to be prepared National plan for effective community engagement and mobilization in vector control.

The World Health Organization GVCR strategy envisage that for achieving sustainable impact in vector control, increased intersectoral and interdisciplinary action is required, thereby linking and converging efforts in environmental management, health education, and reorienting relevant governments programmes around proactive strategies to control new and emerging threats¹. Further the GVCR strategy already acknowledged the importance of engaging local authorities and communities wherein broad-based and inclusive intersectoral collaboration has been considered as important element for improved vector control delivery, through catering of interventions to specific local needs and scenarios as informed by locally generated epidemiological data¹. In that regard the present study findings would help in planning and implementation of community mobilization strategies amidst recurrent outbreaks to facilitate community engagement and collaboration leading to development of sustainable control programme efforts.

Conclusion: The findings from the study revealed that there were better knowledge and awareness levels about Dengue and Chikungunya in the urban slum population. These could be due to public health campaigns held during outbreak situations. The knowledge and awareness levels of VBDs were found to be more in literate respondents as compared to illiterates. This indicates urgent need of sustaining these knowledge and awareness into efficacious prevention practices to facilitate appropriate and sustained public health actions and individual risk protection measures.

Programmatic recommendations: As revealed through the study, most of the respondents were unable to differentiate and distinguish symptoms of VBDs resulting into reporting of general symptoms of febrile illnesses. In view of the overlap of symptoms which have implications on early help-seeking for diagnosis and treatment specifically during the outbreak situations; differentiating features of Dengue and Chikungunya should be mentioned in the currently used NVBDCP health education materials so that the general community will be able to accurately differentiate, distinguish and could take necessary actions to prevent and control the VBDs and thereby recurrent epidemics. Further, it was also found that respondents had comparatively less information and knowledge about causative agent of the Dengue and Chikungunya. This suggests the need to incorporate more scientific aspects and information about the causative agents of VBDs in the health education materials used by the NVBDCP local programme authorities. There is overall need for aggressive health campaigning and rigorous health education activities in the low-income settings of urban communities particularly amidst outbreak situations, through effective health education strategies to promote successful and sustained implementation of WHO recommended GVCR strategy.

○ **Conflicts of Interest**

None.

○ **Authorship declaration**

Swapnil Godbharle and Bibek Raj Giri conceptualized the study, prepared the study tools and participated in the data collection, analysed the data and wrote the manuscript. Abhay Kudale participated in conceptualization of the study, guided the design and coordination of the study, help analysed the data, and critically revised and reviewed the paper. All authors read and approved the final paper.

○ **Funding Source**

The study was self-funded and hence there is no role played and influence of any funding agency on the conduct and preparation of this paper.

○ **Ethical approval**

The study was approved by institutional ethics committee of Savitribai Phule Pune University.

○ **Acknowledgement**

We gratefully acknowledge our institute the Interdisciplinary School of Health Sciences at Savitribai Phule Pune University for giving us an opportunity to conduct this study. Furthermore, we would like to thank Dr. Pradeep Awate, State Surveillance Programme Officer In-charge, Integrated Disease Surveillance Project, Maharashtra State for their facilitation and guidance in the initial planning stages of the study. Last but not least, we acknowledge the general community members from low-income settings of Pune city who participated in this study.

○ **ORCID (Open Researcher and Contributor ID)**

Swapnil Rajendra Godbharle: ORCID ID: <https://orcid.org/0000-0001-5793-1285>

Bibek Raj Giri: ORCID ID: <https://orcid.org/0000-0001-7829-0878>

Abhay Machindra Kudale: ORCID ID: <https://orcid.org/0000-0003-2887-1636>

References:

1. World Health Organization. A global brief on vector-borne diseases [Internet]. Geneva, Switzerland: World Health organization; 2014. Available from: https://apps.who.int/iris/bitstream/handle/10665/111008/WHO_DCO_WHD_2014.1_eng.pdf;jsessionid=A106CD56AEFA0EE22D6CAE3F959B070F?sequence=1
2. World Health Organization, UNICEF. Global vector control response 2017-2030. 2017;
3. Directorate of National Vector Borne Disease Control Programme. National Guidelines for Clinical Management of Chikungunya. [Internet]. India; 2016. Available from: <http://nvbdcp.gov.in/Doc/National-Guidelines-Clinical-Management-Chikungunya-2016.pdf>.
4. Fournet F, Jourdain F, Bonnet E, Degroote S, Ridde V. Effective surveillance systems for vector-borne diseases in urban settings and translation of the data into action: a scoping review. *Infect Dis Poverty* [Internet]. 2018 Dec [cited 2019 Sep 16];7(1). Available from: <https://idpjournal.biomedcentral.com/articles/10.1186/s40249-018-0473-9>
5. Pune Municipal Corporation. City Census Department Population Data 2011 [Internet]. 2019. Available from: <https://www.pmc.gov.in/en/census>
6. Registrar General & Census Commissioner India. “Constituents of Urban Agglomerations having Population 1 lakh and above – Census of India 2011” [Internet]. India; 2011. Available from: http://censusindia.gov.in/2011-prov-results/paper2/data_files/India2/Table_4_PR_UAs_1Lakh_and_Above_Appendix.pdf

7. PMRDA. Pune Metropolitan Region Development Authority (PMRDA) [Internet]. Background. 2018. Available from: www.pmrda.gov.in/background.html
8. Acharya A., Goswami K., Srinath S., Goswami A. Awareness about dengue syndrome and related preventive practices amongst residents of an urban resettlement colony of south Delhi. *J Vector Borne Dis.* 2005;3(42):122–7.
9. Malhotra G, Yadav A, Dudeja P. Knowledge, Awareness and Practices Regarding Dengue among Rural and Slum Communities in North Indian city, India. *Int J Med Sci Public Health.* 2014;3(3):295.
10. Jeelani S, Sabesan S, Subramanian S. Community knowledge, awareness and preventive practices regarding dengue fever in Puducherry – South India. *Public Health.* 2015 Jun;129(6):790–6.
11. Imperial College London, Wellcome Trust. Epicollect5 [Internet]. London; 2017. Available from: <https://five.epicollect.net/>
12. Dayaraj Cecilia. Current status of dengue and chikungunya in India. *WHO South-East Asia J Public Heal.* 2014;
13. Itrat A, Khan A, Javaid S, Kamal M, Khan H, Javed S, et al. Knowledge, Awareness and Practices Regarding Dengue Fever among the Adult Population of Dengue Hit Cosmopolitan. Carr JK, editor. *PLoS ONE.* 2008 Jul 9;3(7):e2620.
14. Gupta P., Kumar P., Aggarwal OP. Knowledge, attitude and practices related to dengue in rural and slum areas of Delhi after the dengue epidemic of 1996. *J Commun Dis.* 1998;2(30):107–12.
15. Win KT, Nang SZ, Min A. Community-based assessment of dengue-related knowledge among caregivers. 2004;
16. Rashid MH, Sultana H, Zzaman MT. Knowledge and Awareness Regarding Chikungunya among Urban Community People of Selected Area of Dhaka City Bangladesh. *J Infect Dis Ther* [Internet]. 2018 [cited 2019 Mar 30];6(1). Available from: <https://www.omicsonline.org/open-access/knowledge-and-awareness-regarding-chikungunya-among-urban-community-people-of-selected-area-of-dhaka-city-bangladesh-2332-0877-1000355-99317.html>