Mental health in the UK during the COVID-19 pandemic: early observations

Brief title: Mental Health in the UK and COVID-19

¹Ru Jia MSc, ¹Kieran Ayling PhD, ²Trudie Chalder PhD, ¹Adam Massey PhD, ³Elizabeth Broadbent PhD,
¹Carol Coupland PhD, ^{1*}Kavita Vedhara PhD

¹Division of Primary Care
 University of Nottingham
 University Park
 Nottingham, NG7 2RD

²Department of Psychological Medicine,
 Institute of Psychiatry, Psychology & Neuroscience,
 King's College London,
 16, De Crespigny Park,
 London, SE5 8AF,
 United Kingdom.

³Department of Psychological Medicine,
 University of Auckland,
 Private bag 92019, Auckland,
 New Zealand.

* Author to whom all correspondence should be addressed: Professor Kavita Vedhara, Division of Primary Care, University of Nottingham, University Park, Nottingham, NG7 2RD, UK.
Kavita.vedhara@nottingham.ac.uk; Tel +44 115 8466931

Abstract

Background:

Previous pandemics have resulted in significant consequences for mental health. Here we report the mental health sequela of the COVID-19 pandemic on the UK population and examine modifiable and non-modifiable explanatory factors associated with mental health outcomes. We focus on the short-term consequences for mental health, as reported during the first four-six weeks of social distancing measures being introduced.

Methods:

A community cohort study was conducted with adults aged ≥ 18 years recruited through a mainstream and social media campaign between 3/4/20-30/4/20. Consenting participants completed an online survey measuring depression, anxiety and stress and explanatory variables hypothesised to be related to these mental health outcomes.

Outcomes:

N=3097 eligible individuals participated. The cohort was predominantly female (85%); mean age forty-four years; 10% from minority ethnic groups; 50% described themselves as key-workers and 20% identified as having clinical risk factors putting them at increased risk of COVID-19. Mean scores for depression, stress and anxiety significantly exceeded population norms. Analysis of non-modifiable factors indicated that being younger and female were associated with all outcomes, with the final multivariable models accounting for 7-13% of variance. When adding modifiable factors, significant independent effects emerged for positive mood, perceived loneliness and worry about getting COVID-19 for all outcomes, with the final multivariable models accounting for 54-57% of variance.

Interpretation:

Increased psychological morbidity was evident in this UK cohort, with younger people and women at particular risk. Interventions targeting perceptions of: loneliness, risk of COVID-19, worry about COVID-19, and positive mood may be effective.

Acknowledgements:

KA was supported by funding from the National Institute for Health Research School for Primary Care Research (NIHR SPCR). The views expressed are those of the author(s) and not necessarily those of the NIHR, the NHS or the Department of Health.

TC acknowledges the financial support of the Department of Health via the National Institute for Health Research (NIHR) Specialist Biomedical Research Centre for Mental Health award to the South London and Maudsley NHS Foundation Trust (SLaM) and the Institute of Psychiatry at King's College London. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Background

The COVID-19 (Coronavirus, 2019) pandemic has resulted in unprecedented disruption to the fabric of society, our health service and economy. However, the multitude of challenges presented by the pandemic may also pose a significant threat to our psychological health.¹ Individuals are facing a panoply of stressors including: serious illness, bereavement, social distancing, and unemployment. The consequences of these stressors for mental health will not be uniform, rather they will be influenced by a range of modifiable and non-modifiable factors. Understanding these factors will be critical in determining who is at greatest risk of mental health difficulties and potential approaches to intervention. We report here cross-sectional findings from a community cohort study designed to capture both the mental health sequela of the COVID-19 pandemic, as well as the modifiable and non-modifiable explanatory factors associated with adverse mental health outcomes. Our focus is on the immediate consequences for mental health, as reported during the first 4-6 weeks of social distancing measures being introduced in the UK.

In keeping with its recent emergence, much remains unknown about COVID-19 and its consequences. However, the expectation is that the consequences for mental health will be profound and far reaching.¹ Evidence on the impact of the pandemic on people living in China attests to this possibility. For example, one study reported the prevalence of depression and anxiety in medical staff to be over 50% and 44% respectively.² Similarly, a study with a general population sample during Level 1 of the epidemic (classified as a 'significant' emergency) revealed that greater than 70% of participants reported moderate or high psychological symptoms of obsessive compulsion, interpersonal sensitivity, phobic anxiety, and psychoticism.³

The experience of previous pandemics also suggests that significant mental health consequences are probable. During the Middle East Respiratory Syndrome outbreak, for example, untoward psychological effects occurred rapidly: with symptoms of anxiety and anger evident after brief isolation periods of 2 weeks.⁴ Similarly, the Severe Acute Respiratory Syndrome epidemic was associated with a 31% increase in suicide amongst the over 65s in Hong Kong.⁵

Preliminary evidence from the UK suggests that these experiences may be replicated here. For example, two online surveys conducted in the first week after social distancing measures were introduced, highlighted some early concerns.¹ One survey was conducted with people with lived experience of mental health problems and stakeholders (n=2198), and the other a nationally representative sample of the general population (n=1099). Both cohorts were asked about their concerns regarding the mental health impact of the COVID-19 pandemic. Although no validated assessments of mental health were undertaken, a common theme for both cohorts was concern about increasing levels of anxiety, and the role of social distancing measures and well-being of family members in contributing to such anxiety. Similarly, a qualitative study examining the mental health impact of social distancing measures suggested that deleterious effects were evident within 5-12 days of such measures being in place.⁶

So it would seem likely that we should brace ourselves for a significant increase in mental health difficulties associated with the pandemic. But who might be at greatest risk? Individuals at increased risk of the disease and/or adverse outcomes might be expected to experience greater mental health difficulties. Several such groups have now been identified. For example, the death rate is known to be higher in men and older individuals.^{7,8} The latter also being more likely to have co-existing conditions and be socially-isolated through shielding. The ethnic diversity of countries such as the US and UK has also highlighted that individuals from Black, Asian and Minority Ethnic (BAME) backgrounds appear to be affected disproportionately by the disease.⁹ Finally, recent UK data suggest that some key-workers, in particular those in social care, are at greater risk of COVID-19 related mortality.⁷ This combined with evidence from China of increased anxiety and depression among health care workers,³ suggests that key-workers may also be at greater risk of adverse mental health outcomes.

The aforementioned factors are, however, largely non-modifiable. Do modifiable risk factors exist which could be targets for intervention? Stress and coping theory.¹⁰ attests that emotional responses to challenging situations vary according to both our appraisal of stressors and the availability of psychological and social resources. Cognitions are central to our appraisal process and, in the current context, one might expect that individuals who appraise that they are at greatest risk of disease may have poorer mental health outcomes. Evidence from the Netherlands during the H1N1 pandemic supports this hypothesis, with prospective data indicating that levels of anxiety declined as perceptions of the risk of acquiring severe disease declined.¹¹ Another relevant feature of our appraisal process concerns the extent to which we attend to and worry about threats to our health. This is

particularly pertinent in contexts where individuals receive health threat information from the media, as in the current pandemic.¹² Early evidence of this mechanism has emerged in the COVID-19 pandemic: with frequency of exposure to information about COVID-19 on social media found to be positively associated with mental health outcomes.¹³

In terms of resources, social support, and its corollary loneliness, are among the best established determinants of our emotional responses to stressors. Successive systematic reviews demonstrate not only poorer mental health outcomes, but also increased morbidity and mortality in individuals who perceive themselves to be more lonely and lacking in support.^{14,15} A determinant also attracting attention is positive mood. It is now recognised that positive mood is not just the opposite of negative mood, but may in fact confer direct effects on well-being as well as protective effects in challenging situations.^{10,16} In terms of mental health outcomes, evidence suggests that the existence of positive mood reduces the risk of mood disorders by 28% and anxiety disorders by 53%, and also influences recovery from some mental health conditions.^{17,18}

Taken together there is an urgent need to report evidence on the prevalence of mental health problems during the COVID-19 pandemic, to understand who may be at greatest risk, and to explore the psychological and social resources that may mitigate this risk. To that end, we report cross sectional findings from a community cohort survey conducted between 3rd and 30th April 2020 which coincided with the first 4-6 weeks of social distancing measures being introduced in the UK.

Methods

Recruitment and Eligibility

Ethical approval was granted from the University of Nottingham Faculty of Medicine and Health Sciences (ref: 506-2003) and the NHS Health Research Authority (ref: 20/HRA/1858). The study was launched on 3/4/20 with participants recruited in the community through a social and mainstream media campaign involving, but not limited to, Facebook and Twitter. In addition, HRA regulatory approval enabled us to approach NHS organisations and request they advertise the research through their routine communications. Recruitment continued until 30/4/20. All media directed potential participants to the study website (www.covidstressstudy.co.uk) through which they accessed the information sheet, consent form and online survey.

Eligibility criteria specified that participants should be: aged 18 and over; able to give informed consent; able to read English; residing in the UK at the time of completing the survey and able to provide a sample of hair at least 1 cm long. The latter was collected for the determination of the stress biomarker cortisol which will be the subject of future manuscripts.

Procedures

Consenting participants completed an online survey implemented through JISC Online Survey (<https://www.onlinesurveys.ac.uk/>). This included validated measures capturing the mental health outcomes: anxiety ($\alpha=0.88$), depression ($\alpha=0.92$) and stress ($\alpha=0.76$).¹⁹⁻²² We also measured modifiable and non-modifiable variables we hypothesised would be related to these mental health outcomes due to being (i) associated with an increased risk of contracting COVID-19 and/or adverse disease outcomes; or (ii) known to be directly associated with adverse mental health outcomes. These were: age, gender, ethnicity, key-worker status, living alone, positive mood, worry about contracting COVID-19 and perceived loneliness and risk of COVID-19 (see supplementary appendix).

Statistical analysis

We first summarised the outcome variables (depression, anxiety and stress) and participant characteristics with appropriate summary statistics and examined histograms and scatterplots. To explore the associations between

non-modifiable and modifiable explanatory factors on outcome variables we conducted univariable linear regression analyses (see supplementary appendix). Multivariable linear regression analyses were then used to explore the independent relationships of non-modifiable factors (age, gender, ethnicity, keyworker status, living alone) on outcome variables. Then, in subsequent models, modifiable explanatory factors (perceived loneliness, perceived risk of COVID-19, positive mood, worry about contracting COVID-19) were added to examine the additional and independent contribution of these factors to explaining variation in the outcome variables. The variable assessing COVID-19 worry was treated as a categorical variable in all models, with “occasional worry” treated as the reference value as this was the most common response. Assumptions of linear regression (normality and homoscedasticity of residuals, linearity with continuous variables) and presence of outliers were assessed graphically. Square root transformations were used for depression and anxiety scores to satisfy assumptions. Robustness of the models were examined by removing data points with large residuals (<-3 or >3) and comparing results to the original models. In the vast majority of models, this had no substantive effect on interpretation. Thus these results are only mentioned where interpretation may be affected. Additionally, as perceived risk of getting COVID-19 was not assessed in those who thought they had had it ($n=519$) these participants are not represented in final multivariable models. As a sensitivity analysis, models were additionally re-specified excluding this explanatory variable (see supplementary appendix).

For depression and anxiety we carried out additional analyses dichotomising according to established cut-offs (scores of 10 or greater indicating moderate or severe levels). We used multiple logistic regression to estimate odds ratios with 95% confidence intervals for their associations with non-modifiable and modifiable variables.

Statistical analyses were performed using STATA (version 16).

Role of funding source

The study sponsor did not play a role in the study design, collection; analysis, and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

Results

Cohort characteristics

The final number of participants recruited was $n=3102$. Of these, five were ineligible due to being less than 18 years old. Thus, yielding $n=3097$ eligible participants. The largest proportion of visitors to the website came direct to the URL (62%/ $n=15,218$), followed by 25% ($n=6068$) via Facebook (the remainder through other websites). The vast majority of respondents accessed the website via a mobile phone (70%/ $n=17045$). The survey was completed in full by 100% of those who started it, consequently there were no missing data, with the exception of age, for which 2 participants entered non-numeric values.

Table 1 summarises the main characteristics of the participants and reveals that our sample was predominantly female; with a mean age of 44 years (standard deviation=15); with participation across the UK (albeit primarily from England) and 10%/ $n=296$ from minority ethnic backgrounds. Fifty percent ($n=1559$) described themselves as key-workers (39%/ $n=1198$ identifying as working in health and social care). Twenty percent ($n=649$) identified themselves as having clinical risk factors which would put them at increased or greatest risk of COVID-19.

Table 1: Participant Demographics ($n=3097$)

	Participants
	n (%)
Gender	

Male	476 (15.4%)
Female	2618 (84.5%)
Prefer not to say	3 (0.1%)
Age groups (years)	
18-24	363 (11.7%)
25-34	528 (17.1%)
35-44	635 (20.5%)
45-54	687 (22.2%)
55-64	568 (18.3%)
65-74	254 (8.2%)
≥75	62 (2.0%)
Ethnicity	
White – British, Irish, other	2796 (90.3%)
Asian/Asian British – Indian, Pakistani, Bangladeshi, other	119 (3.8%)
Black/Black British – Caribbean, African, other	42 (1.4%)
Chinese/Chinese British	28 (0.9%)
Mixed race – White and Black/Black British	19 (0.6%)
Middle Eastern/Middle Eastern British – Arab, Turkish, other	23 (0.7%)
Mixed race – other	40 (1.3%)
Other ethnic group	25 (0.8%)
Prefer not to say	5 (0.2%)
Relationship status	
Single, never married	574 (18.5%)
Single, divorced or widowed	263 (8.5%)
In a relationship/married but living apart	254 (8.2%)
In a relationship/married and cohabiting	1981 (64.0%)
Prefer not to say	25 (0.8%)
Education (highest level of attainment)	
No qualifications	33 (1.1%)
Completed GSCE/CSE/O-levels or equivalent	252 (8.1%)
Completed post-16 vocational course	101 (3.3%)
A-levels or equivalent (at school until aged 18)	403 (13.0%)
Undergraduate degree or professional qualification	1306 (42.2%)
Postgraduate degree	976 (31.5%)
Prefer not to say	26 (0.8%)
Place of residence	
South West England	241 (7.8%)
East Midlands	762 (24.6%)
Yorkshire and Humber	293 (9.5%)
North East	147 (4.8%)
East of England	153 (4.9%)
North West	357 (11.5%)
South East England	415 (13.4%)
Greater London	329 (10.6%)

West Midlands	165 (5.3%)
Northern Ireland	8 (0.3%)
Wales	73 (2.4%)
Scotland	154 (5.0%)
Key-worker status	
Health, social care or relevant related support worker	1198 (38.7%)
Teacher or childcare worker still travelling in to work	70 (2.3%)
Transport worker still travelling in to work	1 (0.03%)
Food chain worker (e.g. production, sale, delivery)	33 (1.1%)
Key public services worker (e.g. justice staff, religious staff, public service journalist or mortuary worker)	22 (0.7%)
Local or national government worker delivering essential public services	41 (1.3%)
Utility worker (e.g. energy, sewerage, postal service)	5 (0.2%)
Public safety or national security worker	11 (0.4%)
Worker involved in medicines or protective equipment production or distribution	10 (0.3%)
Other key worker role not listed	168 (5.4%)
Not a key worker	1538 (49.7%)
Living alone (or with others)	
Living alone	406 (13.1%)
Living with others	2691 (86.9%)
COVID-19 risk status	
Most at risk (e.g. suffering from advanced cancer, severe asthma/COPD, etc.)	121 (3.9%)
At increased risk (e.g., being pregnant, aged over 70, etc.)	528 (17.1%)
Not at-risk	2448 (79.0%)

Mental health outcomes

Table 2 summarises findings in relation to levels of stress, anxiety and depression in the cohort. The mean values for all measures indicate levels that are higher in women than men and decrease with age. Overall mean values are significantly higher than previously reported population norms²³⁻²⁵. For both anxiety and depression the means for the cohort were higher for both genders compared with their respective population norms, and also for all age ranges between 25-64 years. In contrast, both men and women aged over 65 years had anxiety and depression scores consistent with previous population norms. The data suggested no significant differences in stress scores by gender, despite the combined mean score exceeding the population norm.

Table 3 shows the categorisation of participants in line with established cut-offs for anxiety and depression. This shows 64% of participants reported symptoms of depression and 57% reported symptoms of anxiety. When considering the thresholds at which someone would qualify for high intensity psychological support (score of 10 or greater) in the NHS,²⁶ we observe that 31.6% reported moderate to severe depression and 26% moderate to severe anxiety.

Table 2: Depression (PHQ-9), anxiety (GAD-7) and stress (PSS-4) scores and published population normative data[†]

	PHQ-9 score			GAD-7 score			PSS-4 score		
	Participants		t	Participants		t	Participants		t
	Mean (SD)	Norms Mean (SD)		Mean (SD)	Norms Mean (SD)		Mean (SD)	Norms Mean (SD)	
Total Score	7.69 (6.0)	2.91 (3.5)	45.31****	6.59 (5.6)	2.95 (3.4)	36.52****	6.48 (3.3)	6.11 (3.1)	3.80****
Gender									
Male	6.49 (6.1)	2.7 (3.5)	18.56****	5.22 (5.4)	2.66 (3.2)	13.77****	5.88 (3.3)	5.56 (3.0)	1.57 (<i>p</i> =0.12)
Female	7.91 (6.0)	3.1 (3.5)	35.80****	6.84 (5.5)	3.20 (3.5)	28.83****	6.59 (3.3)	6.38 (3.2)	1.73 (<i>p</i> =0.084)
Age groups (years)									
18-24	11.26 (6.4)	..	..	9.04 (5.9)	..	..	7.64 (3.3)	..	..
25-34	8.74 (5.9)	2.3 (3.2)	22.46****	7.73 (5.6)	2.81 (3.3)	13.85****	6.97 (3.3)	..	..
35-44	8.24 (6.0)	2.6 (3.5)	23.48****	7.25 (5.7)	2.82 (3.3)	14.09****	6.40 (3.1)	..	..
45-54	7.34 (5.7)	2.8 (3.5)	19.31****	6.28 (5.3)	3.14 (3.4)	10.71****	6.06 (3.1)	..	..
55-64	6.32 (5.6)	3.2 (3.5)	12.90****	5.43 (5.1)	3.25 (3.6)	7.36****	5.40 (3.1)	..	..
65-74	3.81 (4.2)	3.3 (3.6)	1.88 (<i>p</i> =0.060)	3.33 (3.8)	2.79 (3.2)	1.95 (<i>p</i> =0.052)	4.83 (2.9)	..	..
≥75	4.69 (5.7)	4.4 (3.9)	0.29 (<i>p</i> =0.61)	3.26 (4.4)	3.05 (3.4)	0.37 (<i>p</i> =0.71)	5.22 (3.1)	..	..

[†] PHQ-9, the 9-item Patient Health Questionnaire;¹⁹ GAD-7, the 7-item Generalized Anxiety Disorder Scale;²⁰ PSS-4, the 4-item Perceived Stress Scale.²¹ Published population normative data for PHQ-9²³, GAD-7²⁵, PSS-4²⁴.

**** *p*<0.0001, *** *p*<0.001, ** *p*<0.01, * *p*<0.05

Table 3: Prevalence of depressive and anxiety cases[†]

		Whole sample		Male		Female	
Categories		n	%	n	%	n	%
Depression (PHQ-9 [‡])	No-Minimal Depression (0-4)	1125	36.3	230	48.3	894	34.1
	Mild Depression (5-9)	994	32.1	125	26.3	868	33.2
	Moderate Depression (10-14)	525	17.0	64	13.4	461	17.6
	Moderately Severe Depression (15-19)	276	8.9	35	7.4	241	9.2
	Severe Depression (20-27)	177	5.7	22	4.6	154	5.9
Anxiety (GAD-7 [‡])	No-Minimal Anxiety (0-4)	1344	43.4	276	58.0	1066	40.7
	Mild Anxiety (5-9)	947	30.6	108	22.7	839	32.0
	Moderate Anxiety (10-14)	430	13.9	44	9.2	386	14.7
	Severe Anxiety (15-21)	376	12.1	48	10.1	327	12.5

[†] Cut-offs for categories in line with published guidelines for PHQ-9²³ and GAD-7.²⁵

[‡] PHQ-9, the 9-item Patient Health Questionnaire;¹⁹ GAD-7, the 7-item Generalized Anxiety Disorder Scale.²⁰

Individuals at greatest risk of mental health problems: associations with age, gender, ethnicity, living alone and key-worker status

When non-modifiable explanatory variables were included in a multivariable model (Table 4), we observed that for depression (square-root transformed scores), being younger ($B=-0.28$, 95% CI: -0.31, -0.25 per decade), female ($B=0.36$, 95% CI: 0.25, 0.47) and living alone ($B=0.34$, 95% CI: 0.25, 0.47) were all independently significantly associated with greater levels of depression. This model accounted for approximately 13% of the variance in depression scores. These results were replicated when considering depression as a binary outcome (i.e., cases requiring high intensity intervention versus not) with females having a 49% increased odds of depression and living alone associated with a 55% increase.

Table 4: Regression models showing associations between non-modifiable explanatory variables and depression scores

	B	95% CI Lower	95% CI Upper	β	p
PHQ-9 Total Score ^a					
Age (per decade)	-0.28	-0.31	-0.25	-0.35	<.0001****
Female	0.36	0.25	0.47	0.11	<.0001****
Live alone	0.34	0.22	0.46	0.09	<.0001****
BAME background	0.03	-0.11	0.17	0.01	.64
Key-worker	0.06	-0.02	0.14	0.02	.15
Adjusted R²=0.13, n=3090					
	Odds Ratio	95% CI Lower	95% CI Upper	β	p
PHQ-9 “Cases” ^b					
Age (per decade)	0.67	0.63	0.71	-1.30	<.0001****
Female	1.49	1.18	1.89	0.31	.00079***
Live alone	1.55	1.23	1.97	0.32	.00025***
BAME background	1.14	0.88	1.48	0.08	.32
Key-worker	1.14	0.97	1.33	0.14	.11
Pseudo R²=0.06, n=3090					

**** $p<0.0001$, *** $p<0.001$, ** $p<0.01$, * $p<0.05$

^a A square-root transformation was applied to the dependent variable.

^b a “case” is defined as a PHQ-9 score greater than or equal to 10, at which level someone would qualify for high intensity psychological support in the NHS.

For anxiety (square-root transformed scores) being younger ($B=-0.24$, 95% CI: -0.27, -0.22 per decade) and female ($B=0.43$, 95% CI: 0.32, 0.55) were independently significantly associated with greater levels of anxiety (Table 5). This model accounted for approximately 11% of the variance and these results were replicated when considering anxiety as a binary outcome (i.e., cases requiring high intensity intervention versus not).

Table 5: Regression models showing associations between non-modifiable explanatory variables and anxiety scores

	B	95% CI Lower	95% CI Upper	β	p
GAD-7 Total Score^a					
Age (per decade)	-0.24	-0.27	-0.22	-0.30	<.0001****
Female	0.43	0.32	0.55	0.13	<.0001****
Live alone	-0.03	-0.15	0.09	-0.01	.64
BAME background	0.02	-0.12	0.16	0.01	.77
Key-worker	0.08	-0.00	0.16	0.03	.06
Adjusted R²=0.11, n=3090					
	Odds Ratio	95% CI Lower	95% CI Upper	β	p
GAD-7 “Cases”^b					
Age (per decade)	0.70	0.66	0.74	-1.23	<.0001****
Female	1.61	1.25	2.08	0.39	0.00020***
Live alone	1.02	0.78	1.32	0.01	.91
BAME background	1.15	0.88	1.51	0.10	.30
Key-worker	1.13	0.96	1.34	0.14	.15
Pseudo R²=0.05, n=3090					

**** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$

^a A square-root transformation was applied to the dependent variable.

^b a “case” is defined as a GAD-7 score greater than or equal to 10, at which level someone would qualify for high intensity psychological support in the NHS.

For stress, being younger ($B = -0.54$, 95% CI: -0.61, -0.46 per decade), female ($B = 0.78$, 95% CI: 0.46, 1.09), living alone ($B = 0.48$, 95% CI: 0.15, 0.82), being from a BAME background ($B = 0.45$, 95% CI: 0.07, 0.84), were all independently significantly associated with greater stress; while being a key-worker was independently significantly associated with a lower stress ($B = -0.24$, 95% CI: -0.47, -0.02). Together the model accounted for approximately 7% of the variance in stress scores (Table 6).

Table 6: Regression model showing associations between non-modifiable explanatory variables and stress scores

	B	95% CI Lower	95% CI Upper	β	p
PSS-4 Total Score					
Age (per decade)	-0.54	-0.61	-0.46	-0.25	<.0001****
Female	0.78	0.46	1.09	0.09	<.0001****
Live alone	0.48	0.15	0.82	0.05	0.0049**
BAME background	0.45	0.07	0.84	0.04	0.022*
Key-worker	-0.24	-0.47	-0.02	-0.04	0.033*
Adjusted R²=0.07, n=3090					

**** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$

Individuals at greatest risk of mental health problems: associations with perceived risk of COVID-19, perceived loneliness, COVID-19 worry and positive mood

Table 7 shows levels of modifiable explanatory variables (perceived risk, perceived loneliness, COVID-19 worry, and positive mood) across the whole sample, as well as by gender and age-groups.

Table 7: Loneliness, worry about COVID-19, perceived risk of COVID-19, and positive mood

		Gender		Age groups (years)						
		Whole sample	Male	Female	18-24	25-34	35-44	45-54	55-64	65-74
Loneliness										
Mean (SD)	3.86 (2.7)	3.56 (2.7)	3.91 (2.7)	5.36 (2.7)	4.36 (2.7)	3.76 (2.7)	3.62 (2.8)	3.47 (2.7)	2.70 (2.1)	2.71 (2.4)
Positive mood										
Mean (SD)	18.99 (5.1)	19.76 (5.1)	18.85 (5.0)	17.67 (4.9)	18.82 (5.1)	18.67 (5.0)	18.92 (5.1)	19.38 (5.0)	20.72 (4.6)	21.56 (5.2)
Perceived risk of COVID-19										
Mean (SD)	4.75 (2.2)	4.46 (2.2)	4.80 (2.2)	4.10 (2.0)	4.92 (2.2)	5.15 (2.2)	5 (2.2)	4.78 (2.3)	4.21 (2.1)	3.31 (1.9)
Worry about COVID-19										
No worry (n, %)	512 (16.5%)	105 (22.1%)	406 (15.5%)	105 (28.9%)	108 (20.5%)	91 (14.3%)	91 (13.3%)	65 (11.4%)	39 (15.4%)	13 (21.0%)
Occasional worry (n, %)	2050 (66.2%)	318 (66.8%)	1731 (66.1%)	208 (57.3%)	320 (60.6%)	427 (67.2%)	466 (67.8%)	396 (69.7%)	188 (74.0%)	45 (72.6%)
Much worry (n, %)	413 (13.3%)	40 (8.4%)	373 (14.3%)	39 (10.7%)	77 (14.6%)	91 (14.3%)	94 (13.7%)	85 (15.0%)	24 (9.5%)	3 (4.8%)
Most worry (n, %)	122 (3.9%)	13 (2.7%)	108 (4.1%)	11 (3.0%)	23 (4.4%)	26 (4.1%)	36 (5.2%)	22 (3.9%)	3 (1.2%)	1 (1.6%)

When modifiable explanatory variables were added into the multivariable model for depression: this revealed that greater perceived loneliness (B=0.10, 95% CI: 0.09, 0.12), lower positive mood (B=-0.12, 95% CI: -0.12, -0.11) and greater than occasional worry about getting COVID-19 (much of time: B=0.28, 95% CI: 0.18, 0.38; most of time: B=0.32, 95% CI: 0.13, 0.50), were all independently and significantly associated with greater levels of depression, in addition to age and gender. The model accounted for approximately 56% of the variance in depression scores. While perceived risk of COVID-19 was not statistically significant, in sensitivity analyses where large residuals were excluded (<-3/>3) this became statistically significant (B=0.02, 95% CI: 0.00, 0.03). These results were largely replicated when considering depression as a binary outcome although gender was no longer statistically significant (Table 8).

Table 8: Regression models showing associations between modifiable explanatory variables and depression scores

	B	95% CI Lower	95% CI Upper	β	p
PHQ-9 Total Score^a					
Age (per decade)	-0.18	-0.20	-0.15	-0.22	<0001****
Female	0.18	0.09	0.27	0.05	<0001****
Live alone	0.02	-0.08	0.12	0.01	0.71
BAME background	-0.03	-0.14	0.08	-0.01	0.61
Key-worker	0.01	-0.06	0.08	0.00	0.84
Perceived loneliness (per unit)	0.10	0.09	0.12	0.23	<0001****
Positive mood (per unit)	-0.12	-0.12	-0.11	-0.48	<0001****
COVID-19 worry ^b					
No worry	-0.01	-0.10	0.09	-0.00	0.89
Much of time	0.28	0.18	0.38	0.08	<0001****
Most of time	0.32	0.13	0.50	0.05	0.00067***
Perceived risk of COVID-19 (per unit)	0.01	-0.00	0.03	0.02	0.13
Adjusted R²=0.56, n=2494					
	Odds Ratio	95% CI Lower	95% CI Upper	β	p
PHQ-9 “Cases”^c					
Age (per decade)	0.68	0.63	0.74	-1.30	<0001****
Female	1.06	0.76	1.47	0.04	0.75
Live alone	0.88	0.61	1.25	-0.10	0.46
BAME background	0.95	0.65	1.39	-0.03	0.79
Key-worker	1.07	0.85	1.36	0.08	0.57
Perceived loneliness (per unit)	1.22	1.17	1.28	1.20	<0001****
Positive mood (per unit)	0.76	0.74	0.79	-3.02	<0001****
COVID-19 worry ^b					
No worry	1.00	0.71	1.41	0.00	0.98
Much of time	1.74	1.28	2.36	0.41	0.00037***
Most of time	2.08	1.17	3.72	0.30	0.013*
Perceived risk of COVID-19 (per unit)	1.04	0.98	1.09	0.17	0.23
Pseudo R²=0.35, n=2494					

**** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$

^a A square-root transformation was applied to the dependent variable.

^b Comparison reference group “I occasionally worry about getting COVID-19”.

^c a “case” is defined as a PHQ-9 score greater than or equal to 10, at which level someone would qualify for high intensity psychological support in the NHS.

For anxiety, the model revealed that greater perceived loneliness (B=0.06, 95% CI: 0.04, 0.07), lower positive mood (B=-0.12, 95% CI: -0.13, -0.11) and greater perceived risk of COVID-19 (B=0.04, 95% CI: 0.02, 0.05) were all independently and significantly associated with greater anxiety, in addition to the non-modifiable factors of being younger, female and living alone. Further, those participants who experienced greater than occasional worry about getting COVID-19 were significantly more likely to have higher levels of anxiety (much of time: B=0.58, 95% CI: 0.47, 0.68; most of time: B=0.87, 95% CI: 0.68, 1.06); with those who did not worry at all about getting COVID-19 being likely to have lower anxiety (B=-0.19, 95% CI: -0.28, -0.09). The model accounted for approximately 54% of the variance in anxiety scores. These results were largely replicated when considering anxiety as a binary outcome, although gender and not worrying at all about getting COVID-19 were no longer statistically significant (Table 9).

Table 9: Regression models showing associations between modifiable explanatory variables and anxiety

	B	95% CI Lower	95% CI Upper	β	p
GAD-7 Total Score^a					
Age (per decade)	-0.16	-0.18	-0.13	-0.19	<.0001****
Female	0.25	0.15	0.34	0.07	<.0001****
Live alone	-0.25	-0.35	-0.15	-0.07	<.0001****
BAME background	-0.08	-0.20	0.03	-0.02	0.17
Key-worker	-0.04	-0.11	0.03	-0.02	0.27
Perceived loneliness (per unit)	0.06	0.04	0.07	0.12	<.0001****
Positive mood (per unit)	-0.12	-0.13	-0.11	-0.48	<.0001****
COVID-19 worry ^b					
No worry	-0.19	-0.28	-0.09	-0.05	0.00015***
Much of time	0.58	0.47	0.68	0.16	<.0001****
Most of time	0.87	0.68	1.06	0.13	<.0001****
Perceived risk of COVID-19 (per unit)	0.04	0.02	0.05	0.06	<.0001****
Adjusted R²=0.54, n=2494					
	Odds Ratio	95% CI Lower	95% CI Upper	β	p
GAD-7 “Cases”^c					
Age (per decade)	0.69	0.63	0.75	-1.34	<.0001****
Female	1.17	0.82	1.67	0.14	0.37
Live alone	0.67	0.46	0.99	-0.32	0.044*
BAME background	0.96	0.65	1.43	-0.03	0.85
Key-worker	0.90	0.70	1.15	-0.13	0.40
Perceived loneliness (per unit)	1.11	1.06	1.17	0.67	<.0001****
Positive mood (per unit)	0.77	0.75	0.80	-3.07	<.0001****
COVID-19 worry ^b					
No worry	0.75	0.52	1.09	-0.24	0.14
Much of time	3.86	2.86	5.22	1.06	<.0001****
Most of time	11.57	5.88	22.77	1.06	<.0001****
Perceived risk of COVID-19 (per unit)	1.07	1.01	1.14	0.35	0.024*
Pseudo R²=0.36, n=2494					

**** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$

^a A square-root transformation was applied to the dependent variable.

^b Comparison reference group “I occasionally worry about getting COVID-19”.

^c a “case” is defined as a GAD-7 score greater than or equal to 10, at which level someone would qualify for high intensity psychological support in the NHS.

The multivariable model for stress scores showed that greater perceived loneliness (B=0.19, 95% CI: 0.15, 0.23), lower positive mood (B=-0.38, 95% CI: -0.40, -0.36), greater than occasional worry about getting COVID-19 (much of time: B=0.37, 95% CI: 0.10, 0.63; most of time: B=1.02, 95% CI: 0.54, 1.50), and greater perceived risk of getting COVID-19 (B=0.06, 95% CI: 0.02, 0.11) were all independently and significantly associated with greater stress, in addition to being younger, female, living alone and not being a key-worker. In robustness analyses, when removing large residuals (<-3 or >3) having a BAME background was also a statistically significant independent predictor (B=0.29, 95% CI: 0.00, 0.58). This model accounted for approximately 57% of the variance in stress scores (Table 10).

Table 10: Regression model showing associations between modifiable explanatory variables and stress scores

	B	95% CI Lower	95% CI Upper	β	p
PSS-4 Total Score					
Age (per decade)	-0.24	-0.30	-0.18	-0.11	<.0001****
Female	0.35	0.12	0.59	0.04	0.0035**
Live alone	-0.41	-0.67	-0.14	-0.04	0.0025**
BAME background	0.26	-0.04	0.55	0.02	0.088
Key-worker	-0.40	-0.58	-0.21	-0.06	<.0001****
Perceived loneliness (per unit)	0.19	0.15	0.23	0.15	<.0001****
Positive mood (per unit)	-0.38	-0.40	-0.36	-0.60	<.0001****
COVID-19 worry ^a					
No worry	-0.05	-0.30	0.19	-0.01	0.67
Much of time	0.37	0.10	0.63	0.04	0.0068**
Most of time	1.02	0.54	1.50	0.06	<.0001****
Perceived risk of COVID-19 (per unit)	0.06	0.02	0.11	0.04	0.0037**

Adjusted R²=57, n=2494

**** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$

^a Comparison reference group "I occasionally worry about getting COVID-19".

Discussion

We report findings from a community cohort study established in the UK to examine the mental health consequences of the COVID-19 pandemic. Our results pertain to the experiences of people within the first four to six weeks of social distancing measures being introduced, and focus on self-reported depression, anxiety and stress scores. The findings indicated that mean levels of depression, anxiety and stress significantly exceeded recent population norms.²³⁻²⁵ Models examining the relationship between these mental health outcomes and non-modifiable explanatory factors accounted for only a modest proportion of the variance (7-13%). Increased depression was associated with being younger, female and living alone; increased anxiety was associated with being younger and female; and increased stress was associated with being younger, female, living alone, being from a BAME background and not being a keyworker. In contrast, when we added the hypothesised modifiable variables into our multivariable models we observed that they accounted for a much larger proportion of the variance (54-57%) with significant independent effects emerging for positive mood, perceived loneliness and worry about getting COVID-19 for all three outcomes, as well as perceived risk of COVID-19 emerging as significant for anxiety and stress.

These findings highlight a number of issues worthy of discussion. First, both mean scores and measures of case-ness indicate that the COVID-19 pandemic is having widespread and deleterious effects on the emotional well-being of people in the UK. This is true for depression, generalised anxiety disorder and stress and is in keeping with observations from other countries.^{2,3} Indeed, the proportion of participants who would require intensive support for depression and anxiety in the NHS does not compare favourably with recent historical estimates of the prevalence of mental health problems in the UK. For example, the 2014 ONS report on adult psychiatric morbidity reported a prevalence of 17% for six different common mental disorders.²⁷ The prevalence of depression alone in the context of this pandemic is almost double this.

Second, the non-modifiable explanatory variables significantly associated with all three of our mental health outcomes were being younger and being female. These findings are consistent with unpublished data from another UK community cohort recruited during the COVID-19 pandemic with a similar gender profile to our own,²⁸ suggesting that these groups may be the most in need of intervention. Although this runs counter to our hypothesis that the greatest psychological morbidity would be observed in individuals at greatest risk of COVID-19, it is consistent with previous work which has shown that individual's perceptions of disease risk are often poorly related to actual risk.²⁹ Alternatively, the results may reflect the fact that the pandemic has resulted in a panoply of challenges to mental health that go beyond the disease itself. It could be hypothesised, for example, that some of the more immediate consequences such as unemployment, financial concerns and increased domestic violence would disproportionately affect younger people and women and this may explain our findings.

A third, and related issue, is that although being younger and female were consistently associated with poorer mental health, the relationship was modest, accounting for, at best, 13% of the variance. In contrast, the

modifiable explanatory measures when added to the multivariable models accounted for 52-57% of the variance. These findings are encouraging as they suggest that there is considerable potential for us to develop interventions to mitigate the mental health effects of the pandemic. The measures of perceived loneliness, positive mood and worry about getting COVID-19 were strongly associated with all three outcomes and thus would be appropriate cognitions to be targeted in future interventions.³⁰

A further issue concerns the effects of the pandemic beyond mental health. It is well known that when negative mood states persist over time they result in the dysregulation of physiological systems involved in the regulation of the immune system.³¹ Thus, there exists significant potential for the psychological harm inflicted by the pandemic to translate into physical harm. This could include an increased susceptibility to the virus, worse outcomes if infected, or indeed poorer responses to vaccinations in the future.³² Studies providing longitudinal data on the prevalence of psychological morbidity and appropriate biomarkers (e.g., cortisol) will be required to determine whether the risks to physical health go beyond the hypothetical.

Finally, we would like to acknowledge several limitations. These include the cross-sectional design of the work which impedes an analysis of cause and effect; the limited generalisability of our cohort inflicted by the self-selected community cohort design and the absence of information on pre-existing mental health conditions which are likely to impact on the severity and prevalence of psychological morbidity.¹ Nonetheless, we are among the first to provide evidence from a large cohort on the mental health impact of the COVID-19 pandemic on people in the UK; to identify groups who may be at particular risk, as well as potential targets for therapeutic intervention.

Author contributions

Ru Jia: study design, coordination and management of recruitment, preparation, analysis and interpretation of data and preparation of manuscript.

Kieran Ayling: study design, coordination and management of recruitment, preparation, analysis and interpretation of data and preparation of manuscript.

Trudie Chalder: study design, analysis and interpretation of data and preparation of manuscript

Adam Massey: study design, coordination and management of recruitment, preparation, analysis and interpretation of data and preparation of manuscript.

Elizabeth Broadbent: study design, interpretation of data and preparation of manuscript

Carol Coupland: study design, analysis and interpretation of data and preparation of manuscript

Kavita Vedhara: research lead, study design, coordination and management of recruitment, preparation, analysis and interpretation of data and preparation of manuscript.

As corresponding author, KV had access to all the data in the study and had final responsibility for the decision to submit for publication

References

1. Holmes EA, O'Connor RC, Perry VH, et al. Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science [published online ahead of print, 2020 Apr 15]. *Lancet Psychiatry*. 2020;S2215-0366(20)30168-1. doi:10.1016/S2215-0366(20)30168-1
2. Liu S, Yang L, Zhang C *et al.* Online mental health services in China during the COVID-19 outbreak. *Lancet Psychiatry* 2020; **18**:e17–e18 doi.org/10.1016/S2215-0366(20)30077-8
3. Tian F, Li H, Tian S, Yang J, Shao J, Tian C. Psychological symptoms of ordinary Chinese citizens based on SCL-90 during the level I emergency response to COVID-19 [published online ahead of print, 2020 Apr 11]. *Psychiatry Res*. 2020; 288-112992. doi:10.1016/j.psychres.2020.112992
4. Jeong H, Yim HW, Song YJ, et al. Mental health status of people isolated due to Middle East Respiratory Syndrome. *Epidemiol Health*. 2016; 38:e2016048. Published 2016 Nov 5. doi:10.4178/epih.e2016048
5. Chan SM, Chiu FK, Lam CW, Leung PY, Conwell Y. Elderly suicide and the 2003 SARS epidemic in Hong Kong. *Int J Geriatr Psychiatry*. 2006; **21** (2):113–118. doi:10.1002/gps.1432
6. Williams SN, Armitage CJ, Tampe T, Dienes K. Public perceptions and experiences of social distancing and social isolation during the COVID-19 pandemic: A UK-based focus group study. medRxiv 2020 doi: <https://doi.org/10.1101/2020.04.10.20061267> (preprint)
7. <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/causesofdeath/bulletins/coronaviruscovid19relateddeathsbyoccupationenglandandwales/deathsregistereduptoandincluding20april2020> [accessed 11/05/20]
8. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study; [published correction appears in *Lancet*. 2020 Mar 28;395(10229):1038]. *Lancet*. 2020; **395** (10229):1054–1062. doi:10.1016/S0140-6736(20)30566-3
9. Khunti K, Singh AK, Pareek M, Hanif W. Is ethnicity linked to incidence or outcomes of covid-19? *BMJ*. 2020; 369:m1548. Published 2020 Apr 20. doi:10.1136/bmj.m1548
10. Folkman S. 1997. Positive psychological states and coping with severe stress. *Soc. Sci. Med.* **45**:1207–21
11. Bults M, Beaujean DJ, de Zwart O, et al. Perceived risk, anxiety, and behavioural responses of the general public during the early phase of the Influenza A (H1N1) pandemic in the Netherlands: results of three consecutive online surveys. *BMC Public Health*. 2011; 11:2. Published 2011 Jan 3. doi:10.1186/1471-2458-11-2.
12. Salkovskis PM, Rimes KA, Warwick HM, Clark DM. The Health Anxiety Inventory: development and validation of scales for the measurement of health anxiety and hypochondriasis. *Psychol Med*. 2002; **32**(5):843–853. doi:10.1017/s0033291702005822

13. Gao J, Zheng P, Jia Y, et al. Mental health problems and social media exposure during COVID-19 outbreak. *PLoS One*. 2020; **15**(4):e0231924. Published 2020 Apr 16. doi:10.1371/journal.pone.0231924
14. Wang J, Mann F, Lloyd-Evans B, Ma R, Johnson S. Associations between loneliness and perceived social support and outcomes of mental health problems: a systematic review. *BMC Psychiatry*. 2018; **18**(1):156. Published 2018 May 29. doi:10.1186/s12888-018-1736-5
15. Leigh-Hunt N, Baggeley D, Bash K, Turner V, Turnbull S, Valtorta N, Caan W. An overview of systematic reviews on the public health consequences of social isolation and loneliness. *Public Health*. 2017 Nov; **152**:157-171. doi: 10.1016/j.puhe.2017.07.035.
16. Pressman SD, Jenkins BN, Moskowitz JT. Positive Affect and Health: What Do We Know and Where Next Should We Go? *Ann Rev Psychol*. 2019; **70**:627-650. doi:10.1146/annurev-psych-010418-102955
17. Schotanus-Dijkstra M, ten Have M, Lamers SMA, de Graaf R, Bohlmeijer ET. The longitudinal relationship between flourishing mental health and incident mood, anxiety and substance use disorders. *Eur J Public Health*. 2017; **27**(3):563-568. doi:10.1093/eurpub/ckw202
18. Schotanus-Dijkstra M, Keyes CLM, de Graaf R, ten Have M. Recovery from mood and anxiety disorders: The influence of positive mental health. *J Affect Disord*. 2019; **252**:107-113. doi:10.1016/j.jad.2019.04.051
19. Kroenke K, Spitzer RL, Williams JB, Lowe B. The Patient Health Questionnaire Somatic, Anxiety, and Depressive Symptom Scales: a systematic review. *Gen Hosp Psychiatry*. 2010;**32**(4):345-59
20. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med*. 2006; **166**(10):1092-1097. doi:10.1001/archinte.166.10.1092
21. Cohen S, Williamson G. (1988). Perceived Stress in a Probability Sample of the United States. In S. Spacapan & S. Oskamp (Eds.), *The social psychology of health: Claremont Symposium on Applied Social Psychology* (pp. 31-67). Newbury Park, CA: Sage.
22. Diener E., Wirtz D, Tov W, et al. (2010). New well-being measures: Short scales to assess flourishing, positive, and negative feelings. *Social Indicators Research*, **97**(2), 143-156. <https://doi.org/10.1007/s11205-009-9493-y>
23. Kocalevent RD, Hinze A, Brähler E. Standardization of the depression screener patient health questionnaire (PHQ-9) in the general population. *Gen Hosp Psychiatry*. 2013; **35** (5):551-555. doi:10.1016/j.genhosppsych.2013.04.006
24. Warrtig SL, Forshaw MJ, South J, White AK. New, normative, English-sample data for the Short Form Perceived Stress Scale (PSS-4). *J Health Psychol*. 2013; **18**(12):1617-1628. doi:10.1177/1359105313508346
25. Löwe B, Decker O, Müller S, et al. Validation and standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the general population. *Med Care*. 2008; **46** (3):266-274. doi:10.1097/MLR.0b013e318160d093.
26. https://www.rcpsych.ac.uk/docs/default-source/improving-care/nccmh/iapt/the-iapt-manual-final-republished-7-3-18.pdf?sfvrsn=a192d1af_0 [accessed 5/5/20]
27. https://files.digital.nhs.uk/pdf/q/3/mental_health_and_wellbeing_in_england_full_report.pdf [accessed 03/05/20]
28. <https://www.marchnetwork.org/research> [accessed 04/05/20]
29. Westhoff CL. Breast cancer risk: perception versus reality. *Contraception*. 1999; **59** (1 Suppl):25S-28S. doi:10.1016/s0010-7824(98)00134-6
30. Masi CM, Chen HY, Hawkey LC, Cacioppo JT. A meta-analysis of interventions to reduce loneliness. *Per Soc Psychol Rev*. 2011; **15** (3):219-266. doi:10.1177/1088868310377394
31. O'Connor D, Thayer J, Vedhara K. Stress and Health: A Review of Psychobiological Processes. *Annual Review of Psychology* (in press).
32. <https://thepsychologist.bps.org.uk/we-will-have-live-risk-covid-19-psychology-has-much-say-about-risk> [accessed 06/05/2020]