Title: Depression and loneliness during COVID-19 restrictions in the United States, and their associations with frequency of social and sexual connections

Authors: Molly Rosenberg^{1,2}, PhD; Maya Luetke¹, MS-MPH; Devon Hensel^{2,3,4}, PhD; Sina Kianersi¹, DVM; Tsung-chieh Fu², PhD; Debby Herbenick^{2,5}, PhD

Author affiliations:

¹Department of Epidemiology, Indiana University School of Public Health-Bloomington

²Center for Sexual Health Promotion, Indiana University School of Public Health-Bloomington

³Department of Pediatrics, Indiana University School of Medicine

⁴Department of Sociology, Indiana University Purdue University-Indianapolis

⁵Department of Applied Health Science, Indiana University School of Public Health-Bloomington

Corresponding author:

Molly Rosenberg
1025 E. 7th Street, Office C032
Bloomington, IN 47405
ORCID: 0000-0001-6679-6791
Email: rosenmol@indiana.edu
Phone: +1 (812) 856-2509
Fax: +1 (812) 855-9107

DECLARATIONS

Funding: The 2020 National Survey of Sexual and Reproductive Health During COVID-19 (NSRHDC) was generously supported by grants from Pure Romance as well the Indiana University Office of the Vice Provost for Research.

Ethics approval: Ethical approval for the study protocol was provided by the Indiana University Human Subjects Office (#2004194314).

Authors' contributions: M. Rosenberg and D. Herbenick conceived the study. D. Herbenick, M. Rosenberg, M. Luetke, and D. Hensel designed the survey. M. Rosenberg conducted the analysis and wrote the first draft of the manuscript with key drafting contributions from ML and scientific contributions from D. Hensel, T. Fu, and D. Herbenick. S. Kianersi created the data visualization. M. Rosenberg, M. Luetke, D. Hensel, T. Fu, S. Kianersi, and D. Herbenick contributed to the interpretation of the findings, critical review of the manuscript, and approval of the final manuscript as submitted.

ABSTRACT

Purpose: To estimate the prevalence of depression and loneliness during the US COVID-19 response, and examine their associations with frequency of social and sexual connections.

Methods: We conducted an online cross-sectional survey of a nationally representative sample of American adults (n=1010), aged 18-94, running from April 10-20, 2020. We assessed depressive symptoms (CES-D-10 scale), loneliness (UCLA 3-Item Loneliness scale), and frequency of in-person and remote social connections (4 items, e.g. hugging family member, video chats) and sexual connections (4 items, e.g. partnered sexual activity, dating app use).

Results: One-third of participants (32%) reported depressive symptoms, and loneliness was high [mean (SD): 4.4 (1.7)]. Those with depressive symptoms were more likely to be women, age 20-29, unmarried, and low-income. Frequent in-person connections were associated with lower depression and loneliness; frequent remote connections were not.

Conclusions: Depression and loneliness were elevated during the early US COVID-19 response. Those who maintained frequent in-person, but not remote, social and sexual connections had better mental health outcomes. While COVID-19 social restrictions remain necessary, it will be critical to expand mental health services to serve those most at-risk and identify effective ways of maintaining social and sexual connections from a distance.

Key words: Depression; Loneliness; COVID-19; Social isolation; American adults

INTRODUCTION

The spread of coronavirus disease 2019 (COVID-19) has led to unprecedented changes in the daily lives of Americans. Measures to mitigate the spread of COVID-19, including social distancing and sheltering in place, limit people's ability to maintain social, relational, and sexual connections and, accordingly, may have dramatic consequences for people's psychological well-being.[1, 2] Social isolation, which is essentially mandated to reduce transmission of coronavirus, has been previously linked to loneliness and negative mental health outcomes in numerous populations.[3, 4] Further, recent research has shown that the spread of COVID-19 and the related public health response have had negative effects on people's emotional well-being,[5-8] with more than one-third of American adults reporting serious changes in their mental health due to the COVID-19 spread and response.[9, 10]

Under normal conditions, depression is common worldwide and has a significant health burden, affecting some 264 million people each year, according to the World Health Organization.[11] Loneliness is also prevalent and strongly correlated with adverse mental and physical health outcomes, including depression.[12] In the United States, depression has increased among adults over the last decade,[13] largely among young adults and women.[14] Recent estimates from the National Institute of Mental Health indicate that 17.3 million American adults (7.1%) experienced a major depressive episode in 2017.[15] Loneliness tends to follow similar demographic trends as depression, with elevated levels observed in women and young adults, but also among low-income and unmarried people. [12, 16]

While previous studies have evaluated the psychological consequences of other respiratory disease epidemics,[17-20] few studies have been conducted to assess these outcomes related to the current pandemic of coronavirus (COVID-19).[5-7] To our knowledge, no COVID-19-era study has yet estimated the population-representative prevalence of depression and loneliness

in American adults. Further, there is limited understanding of how social and sexual connections are associated with mental well-being amidst a global pandemic and the associated containment measures such as quarantine, social distancing, and self-isolation.

This study fills these knowledge gaps by estimating the population-representative prevalence of depression and loneliness in American adults during the early stages of the COVID-19 spread and restrictions, and by examining the frequency of social and sexual connections as potential predictors of these mental health outcomes.

METHODS

Study setting, participants, and data collection

We conducted a nationally representative on-line survey of American adults (age 18+ years) from April 10-20, 2020 - the 2020 National Survey of Sexual and Reproductive Health During COVID-19. Participants were recruited into the study using the Ipsos KnowledgePanel, a nationally representative, probability-based sample established using address-based sampling via the U.S. Postal Service's Delivery Sequence File, of which other surveys have used to underpin high-quality and generalizable results.[21-23] Using an equal probability selection method, members of the panel were sampled and invited to participate in our study. Sampled participants were emailed an invitation and link to an on-line survey querying information on mental health, relationships, sexual behaviors. Ipsos maintains an incentive program for participation in individual surveys, including sweepstakes/raffles for prizes and cash rewards. Ethical approval for the study protocol was provided by the Indiana University Human Subjects Office (#2004194314).

Key variables

We assessed two key outcomes: depression and loneliness. We measured depression with the 10-item Center for Epidemiologic Studies Depression Scale.[24] As in other studies,[25, 26] we used a scale cutpoint of greater than or equal to 10 to identify those likely experiencing significant depressive symptoms. We measured loneliness with the UCLA 3-item Loneliness scale.[27] We calculated the continuous scale score (possible range 3-9), and used a score of 4 or greater to define a dichotomous loneliness category for comparison with prior studies.[28, 29] We used a more conservative loneliness definition (score of 6 or greater) for categorization for subsequent analyses. This categorization decision was made to address concerns about the likely lower discriminatory value of one of the scale items ('How often do you feel isolated from others?').

We assessed the frequency of eight social and sexual connection exposures with reference to the time frame of the last month, each with response categories: not at all, once or a few times, 1-3 times a week, and almost every day. We grouped the social connections into in-person (hug/kiss a family member, visit a non-household friend or family member) and remote (video chat with friend/family, send/receive a letter in the mail). Similarly, we grouped the sexual connections into in-person (hug/kiss a partner, partnered sexual activities: mutual masturbation, oral sex, or intercourse) and remote (sex over phone, video chat or texting; use of dating or hookup app). Few endorsed these final two remote sexual exposures at high frequency, so for those we combined the '1-3 times a week' and 'almost every day' responses. We also assessed relationship tension among participants reporting romantic relationship. We measured romantic tension with an item that queried whether participants noted increased tension or arguments with their partner due to COVID-19 spread and restrictions. We categorized responses into 'yes' (often, sometimes, or rarely) and 'no' (no significant change, or feeling more connected).

We assessed other key demographic variables with data maintained in the Ipsos KnowledgePanel: gender; age, in years, categorized in 10-year increments; race/ethnicity; marital status; region: northeast, south, Midwest, west; household size, and household makeup (living alone, children under 13 years in home). To assess the potential importance of living with a romantic partner, we created a dichotomous *relationship status* variable separating participants who lived with a romantic partner from participants who did not live with a romantic partner, either because their romantic partner lived elsewhere or because they did not have a romantic partner.

Statistical analysis

To account for any differential nonresponse that may have produced different distributions between our survey sample and the overall Ipsos KnowledgePanel, all analyses were conducted on a sample weighted to the overall to the geo-demographic distribution of the United States based on US Census data. The magnitude of missing data was <5% for all variables, so we conducted complete case analyses. We calculated descriptive estimates of the prevalence of depressive symptoms and loneliness, and used chi-square and t-tests to assess which categorical and continuous demographic variables differed significantly by depression status. We used log-binomial regression to estimate the associations between each of the nine exposures (frequency of social and sexual connections and relationship tension) and each of the two mental health outcomes (depression and loneliness). We estimated these associations in unadjusted models, and in models adjusted for gender, age, marital status, and income. To examine whether the associations differed by relationship status, we added an interaction term between relationship status and each of the nine exposures, dichotomizing the frequency variables at 1-3 times per week (at or above/below). We estimated stratified results and used the magnitude of differences in point estimates and Wald p-values to inform our conclusions

around effect measure modification. All analyses were conducted using SAS statistical software, version 9.4.

RESULTS

Of the 1632 KnowledgePanel members invited to participate in the study, 1010 completed the survey for a 61.9% response rate. The weighted sample was 52% female, 63% non-Hispanic White, 12% non-Hispanic Black, 16% Hispanic, and 9% other race or multiple races. Most (62%) were married, and one-third (33%) had a college education or higher (Table 1). The demographic differences between the weighted and the unweighted sample were minor. The ages of those in the sample ranged from 18 to 94 years and the weighted median age was 48 years (IQR 32-62).

The weighted prevalence of major depressive symptoms in the last week was 31.7% (95% CI: 28.9, 34.8%). People reporting major depressive symptoms were more likely to be women, age 20-29, unmarried, in the lowest income bracket, and living alone. Loneliness scale scores were also significantly higher among those with major depressive symptoms [mean (SD): 5.5 (2.1)], than among those without [mean (SD) 3.9 (1.3)]. The weighted prevalence of loneliness was 54.0% (95% CI: 51.0, 57.2%) using our first definition, and 23.8% (95% CI: 21.3, 26.6%) using our second definition.

In general, frequent in-person social connections were associated with lower depression and loneliness prevalence, but frequent remote social connections were not (Table 2, Figure 1). Compared to those who reported ‘not at all,’ people who hugged or kissed a family member almost every day in last month were 26% less likely to report major depressive symptoms [aPR (95% CI): 0.74 (0.56, 0.97)], and 28% less likely to report loneliness [aPR (95% CI): 0.72 (0.51, 1.02)]. Though not statistically significant, we observed similar trends for the exposure of in-

person visits with non-household members [$aPR_{\text{depression}}$ (95% CI): 0.79 (0.48, 1.31) and $aPR_{\text{loneliness}}$ (95% CI): 0.86 (0.51, 1.02)]. This trend was not observed for remote social connections (video chats and sending/receiving letters in the mail). Relative to 'not at all' responses, high frequency of video chats were not associated with lower depression or loneliness prevalence [aPR (95% CI): 1.02 (0.78, 1.35) and aPR (95% CI): 1.12 (0.79, 1.59), respectively], whereas those with high frequency of sending/receiving letters tended towards higher depression and loneliness [aPR (95% CI): 1.57 (0.95, 2.59) and aPR (95% CI): 2.49 (1.33, 4.65), respectively].

Similarly, frequent in-person sexual connections were associated with lower prevalence of depression and loneliness in ways that remote sexual connections were not (Table 2, Figure 1). For the hug/kiss and partnered sex exposures, a dose response relationship was observed where, relative to 'not at all' responses, increasing frequencies were associated with decreased depression and loneliness prevalence. Those who endorsed the most frequent hug/kiss exposure were 31% less likely to report depressive symptoms and 67% less likely to report loneliness [aPR (95% CI): 0.69 (0.53, 0.90) and aPR (95% CI): 0.33 (0.23, 0.48), respectively]. Those who endorsed the most frequent partnered sex exposure were 57% less likely to report depressive symptoms [aPR (95% CI): 0.43 (0.19, 0.97)]. No participant with partnered sex at this high frequency had loneliness scores above 6. Relatively few participants endorsed frequent remote sex or dating app usage, but those who did tended to have slightly higher rates of depression and loneliness. However, these estimates were measured imprecisely with wide confidence limits spanning the null. Participants who endorsed experiencing increased tension with their romantic partner due to COVID-19 spread and restrictions were more likely to report major depressive symptoms [aPR (95% CI): 2.79 (2.19, 3.55)] and loneliness [aPR (95% CI): 3.23 (2.37, 4.39)].

The patterns for the associations between social and sexual connections and mental health outcomes generally did not differ depending on if the participant lived with a romantic partner (Table 3). One possible exception to this trend emerged for the remote sex over phone, video chat, or texting outcome, though the low numbers of participants reporting this exposure frequently limited the precision of our estimates. For participants who lived with a partner, more frequent remote sex was associated with lower depression and loneliness [aPR (95% CI): 0.89 (0.38, 2.07) and aPR (95% CI): 0.34 (0.04, 2.70), respectively], while the opposite was true for participants who did not live with a partner [aPR (95% CI): 1.66 (0.97, 2.84) and aPR (95% CI): 1.69 (0.97, 2.84), respectively, Wald p-values for interaction both=0.2].

DISCUSSION

In a nationally representative probability survey of American adults taking place during the early phases of the COVID-19 response (April 2020), we found high levels of significant depressive symptoms and loneliness. Frequent in-person social and sexual contacts during the time of COVID-19 restrictions in the last month were generally associated with lower prevalence of depression and loneliness, while remote contacts were not similarly protective. We also found that relationship tension due to COVID-19 spread and restrictions was strongly predictive of depression and loneliness, and that these associations were significantly stronger among people who lived with their romantic partners. Our findings suggest that the COVID-19 spread and response has had a tremendous mental health impact on Americans.

We found that nearly one-third of Americans reported depressive symptoms in April 2020, notably much higher than previous estimates among American adults. From 2013-2016, the prevalence of major depression in a given 2 week period in US adults age 20+ years was 8%, nearly 4 times lower than our estimate.[30] Similarly, the mean loneliness score we found (4.4) was higher than prior estimates in three western European countries (3.5-3.7),[31] and the

prevalence of loneliness we found in our general US population (Definition 1: 54%), is a similar magnitude as previously observed in older and elderly Americans (43-56%).[28, 29] Patterns of higher depression prevalence among women, young adults, and people with lower incomes have been observed in previous studies.[14, 30] We also observed these gender, age, and socioeconomic patterns in our study, suggesting that while the magnitude of depression may be expanding during the time of COVID-19 spread and restrictions, the disproportionate burden continues to be felt in these populations.

The observed relationships between social and sexual connections and the outcomes of depression and loneliness are largely consistent with our understanding of the importance of human connection for mental health and well-being. Close touching in family interactions and among relationship partners has been associated with decreased heart rate, higher levels of oxytocin, and lower levels of cortisol;[32, 33] which, in turn could provide important mental health protections. These kinds of connections cannot easily be recreated with remote technology where direct touch is not possible, though some researchers have explored this through “huggable” and other similar communication devices.[34] We also know that common technical difficulties with video calls can cause misattributed negative feelings towards people on the call,[35] perhaps contributing to the null effects we observed for remote connections and mental health outcomes. As the COVID-19 spread and response have dramatically limited access to many previously routine and familiar options for human connection, our findings are consistent with an explanation that the decreased frequency of social and sexual connections is contributing to poor mental health outcomes. In addition to limiting people’s social interactions, restrictions have also likely resulted in many people missing counselling or therapy appointments or other activities (e.g., exercise, massage, support group meetings, etc.) that may have been supportive of their mental health.

There are several other plausible explanations for the observed relationship between social and sexual connections and mental health outcomes. First, people who are struggling with poor mental health and feelings of isolation during COVID-19 restrictions may be reaching out more frequently for remote connections. As this was a cross-sectional study, our ability to infer temporality is limited. Future studies with a longitudinal design would be better able to disentangle the temporal relationships between social/sexual connections and mental health outcomes. Second, we used self-reported data for our key exposure and outcome assessments, with the potential for biased results. However, web-based surveys have shown to be effective modes to elicit sensitive sexual behaviors from study participants, alleviating some of the bias concern.[36] Similarly, our depression and loneliness scales are widely used in web-based surveys, are validated, and have good psychometric properties.[24, 27] However, their use could be complemented with clinical assessments in future studies. Relatedly, it is not yet known how these scales perform during times of social restrictions when daily lives have shifted so dramatically. It is possible that the wording of certain items (e.g. ‘How often do you feel isolated from others?’) may lose their discriminatory ability in times where isolation is essentially mandated. For this reason, we used a more conservative categorical definition of loneliness (scale score of 6 or greater) for our analyses than has been commonly used in prior studies (scale score of 4 or greater). We also did not explore whether participants had run out of prescription medication supportive of their mental health or had missed counselling or therapy visits.

A key strength of this study is its external validity. Our findings are broadly representative of the US adult population during the early stages of the US COVID-19 response. However, these results may not hold during later phases of the COVID-19 response, or for countries outside of the US. Surveys should be conducted at frequent intervals and in other countries affected by

COVID-19 to update our understanding of the changing mental health landscape across time and space.

Conclusions

Depressive symptoms and loneliness are being experienced at very high levels in Americans during the COVID-19 spread and response. At this time, social restrictions are a critical tool to mitigate the spread of COVID-19 and its associated morbidity and mortality. Our findings should not be interpreted in support of prematurely lifting these necessary restrictions. Rather, public health and mental health professionals should be aware of the increasing mental health needs that are likely co-occurring with the COVID-19 response, and funding should be allocated to expand mental health services to those most likely to be experiencing poor mental health. Women, young adults, and low-income Americans seem to be particularly at-risk for these outcomes. We also found that Americans who could maintain frequent in-person social and sexual connections appeared to have better mental health outcomes, though remote connections did not appear to confer the same benefits. Given that extra-household connections are not advised at this time and possibly will be contraindicated intermittently in the future, a mental health priority should be to identify innovative and effective ways of supporting interpersonal connections from a distance.

TABLES AND FIGURES

Figure 1. Associations between frequencies of social and sexual connections and outcomes of depressive symptoms (top panel) and loneliness (bottom panel) in 1010 American adults, April 10-20, 2020.

Table 1. Characteristics of the unweighted and weighted sample, overall and by depressive symptoms in 1010 American adults, April 10-20, 2020

Characteristic	Unweighted Overall N=1010	Weighted Overall N=1010	Major depressive symptoms ¹ N=304	Below threshold for major depressive symptoms ¹ N=657	p-value
	N (%)	N (%)	N (%)	N (%)	
Sex					0.002
Female	516 (51.1)	521 (51.6)	179 (58.9)	315 (47.9)	
Male	494 (48.9)	489 (48.4)	125 (41.1)	342 (52.1)	
Age group					<0.0001
18-19	15 (1.5)	27 (2.7)	10 (3.4)	14 (2.2)	
20-29	114 (11.3)	184 (18.2)	76 (24.9)	99 (15.1)	
30-39	156 (15.5)	175 (17.4)	55 (17.9)	106 (16.1)	
40-49	138 (13.7)	142 (14.1)	45 (14.6)	92 (14.0)	
50-59	215 (21.3)	185 (18.3)	59 (19.4)	119 (18.1)	
60-69	206 (20.4)	165 (16.3)	33 (10.9)	126 (19.3)	
70-79	129 (12.8)	103 (10.2)	23 (7.6)	76 (11.6)	
80+	37 (3.7)	28 (2.8)	3 (1.1)	24 (3.7)	
Race/ethnicity					0.3
White, non-Hispanic	721 (71.4)	639 (63.2)	182 (59.7)	434 (66.1)	
Black, non-Hispanic	85 (8.4)	119 (11.8)	41 (13.6)	71 (10.8)	
Other or 2+ races, non-Hispanic	81 (8.0)	87 (8.6)	29 (9.6)	53 (8.1)	
Hispanic	123 (12.2)	165 (16.4)	52 (17.2)	99 (15.1)	
Marital status					<0.0001
Married or living with partner	668 (66.1)	622 (61.6)	152 (50.0)	444 (67.5)	
Separated/divorced	129 (12.8)	124 (12.3)	40 (13.1)	76 (11.6)	
Widowed	44 (4.4)	40 (3.6)	14 (4.6)	23 (3.6)	
Never married	169 (16.7)	224 (22.2)	98 (32.3)	114 (17.3)	
Region					0.8
Northeast	189 (18.7)	177 (17.5)	52 (17.1)	111 (17.0)	
Midwest	228 (22.6)	210 (20.8)	66 (21.6)	134 (20.4)	
South	341 (33.8)	383 (37.9)	110 (36.1)	258 (39.2)	
West	252 (25.0)	240 (23.8)	77 (25.2)	154 (23.4)	
Household income					0.0005
<\$25k	95 (9.4)	136 (13.5)	61 (20.2)	69 (10.6)	
\$25-49k	174 (17.2)	184 (18.2)	54 (17.7)	120 (18.3)	
\$50-74k	183 (18.1)	174 (17.2)	57 (18.6)	105 (15.9)	
\$75-99k	147 (14.6)	141 (14.0)	32 (10.5)	105 (16.0)	
\$100-124k	108 (10.7)	103 (10.3)	33 (10.8)	67 (10.2)	
\$125-149k	73 (7.2)	66 (6.5)	112 (3.8)	52 (7.9)	
\$150-174k	90 (8.9)	82 (8.1)	25 (8.1)	53 (8.0)	
\$175k+	140 (13.9)	124 (12.3)	32 (10.4)	86 (13.1)	
Education					0.1
Less than high school	75 (7.4)	98 (9.7)	39 (12.7)	52 (7.9)	
High school	278 (27.5)	295 (29.2)	83 (27.4)	186 (28.4)	
Some college	273 (27.0)	281 (27.8)	85 (28.1)	189 (28.7)	

Bachelor's degree or higher	384 (38.0)	336 (33.3)	97 (31.9)	230 (35.0)	
Children in home (<13 years)					1.0
Yes	194 (19.2)	223 (22.0)	67 (22.0)	145 (22.0)	
No	816 (80.8)	787 (78.0)	238 (78.0)	512 (78.0)	
Living alone					0.05
Yes	190 (18.8)	184 (18.2)	66 (21.5)	107 (16.3)	
No	820 (81.2)	826 (81.8)	239 (78.5)	550 (83.7)	
Household size					0.4
Mean (SD)	2.65 (1.50)	2.77 (1.60)	2.72 (1.66)	2.81 (1.58)	
Loneliness scale²					<0.0001
Mean (SD)	4.31 (1.69)	4.37 (1.74)	5.48 (2.08)	3.90 (1.33)	

¹Major depressive symptoms measured with the CES-D-10 scale. Possible scores range 0-30. The frequency of missing data for this variable was n=44 (4.4%).

²Loneliness measured with the UCLA 3-Item Loneliness scale. Possible scores ranged 3-9. The frequency of missing observations for this variable was n=19 (1.9%).

Table 2. Unadjusted and adjusted log-binomial regression for the associations between frequency of social and sexual connections and outcomes of depression and loneliness

Exposure variables	Weighted N (%)	Depressive symptoms		Loneliness	
		PR (95%CI)	aPR ¹ (95% CI)	PR (95% CI)	aPR ¹ (95% CI)
<i>In-person social connection in last month</i>					
Hug or kiss a family member (not partner)					
Not at all	519 (51.6)	1	1	1	1
Once or a few times	196 (19.5)	1.14 (0.91, 1.43)	1.00 (0.81, 1.24)	1.01 (0.77, 1.34)	0.98 (0.75, 1.28)
1-3 times a week	84 (8.4)	0.95 (0.67, 1.36)	0.85 (0.60, 1.19)	0.65 (0.40, 1.07)	0.65 (0.40, 1.06)
Almost every day	206 (20.5)	0.77 (0.58, 1.00)	0.74 (0.56, 0.97)	0.65 (0.46, 0.90)	0.72 (0.51, 1.02)
Missing	n=6				
In-person visit with non-household member					
Not at all	499 (49.8)	1	1	1	1
Once or a few times	353 (35.3)	1.13 (0.93, 1.38)	1.03 (0.84, 1.23)	1.26 (1.00, 1.60)	1.14 (0.91, 1.44)
1-3 times a week	110 (10.9)	0.98 (0.71, 1.36)	0.91 (0.67, 1.24)	0.98 (0.66, 1.45)	0.92 (0.63, 1.35)
Almost every day	40 (4.0)	0.94 (0.56, 1.58)	0.79 (0.48, 1.31)	0.91 (0.47, 1.76)	0.86 (0.45, 1.63)
Missing	n=9				
<i>Remote social connection in last month</i>					
Video chat with friend/family member					
Not at all	328 (32.8)	1	1	1	1
Once or a few times	321 (32.0)	0.83 (0.66, 1.05)	0.78 (0.62, 0.98)	0.86 (0.65, 1.15)	0.85, 0.64, 1.12)
1-3 times a week	239 (23.8)	0.91 (0.71, 1.16)	0.83 (0.65, 1.05)	0.95 (0.71, 1.28)	0.95 (0.71, 1.28)
Almost every day	114 (11.4)	1.12 (0.85, 1.49)	1.02 (0.78, 1.35)	1.08 (0.76, 1.55)	1.12 (0.79, 1.59)
Missing	n=8				
Send or receive a card or letter in the mail					
Not at all	614 (61.2)	1	1	1	1
Once or a few times	322 (32.2)	0.82 (0.66, 0.99)	0.92 (0.79, 1.08)	0.94 (0.74, 1.21)	1.10 (0.86, 1.40)
1-3 times a week	52 (5.2)	0.79 (0.50, 1.27)	0.91 (0.65, 1.26)	0.73 (0.40, 1.33)	0.83, 0.46, 1.50)
Almost every day	14 (1.4)	2.09 (1.45, 3.00)	1.57 (0.95, 2.59)	1.88 (1.01, 3.50)	2.49 (1.33, 4.65)

Missing	n=7				
<i>In-person sexual connection in last month</i>					
Hug or kiss romantic partner					
Not at all	333 (33.4)	1	1	1	1
Once or a few times	179 (18.0)	0.93 (0.73, 1.19)	0.93 (0.73, 1.17)	0.74 (0.56, 0.98)	0.77 (0.58, 1.02)
1-3 times a week	116 (11.7)	0.86 (0.63, 1.16)	0.89 (0.65, 1.22)	0.55 (0.38, 0.81)	0.55 (0.37, 0.83)
Almost every day	370 (37.0)	0.63 (0.50, 0.80)	0.69 (0.53, 0.90)	0.31 (0.22, 0.42)	0.33 (0.23, 0.48)
Missing	n=12				
Partnered sexual activity ²					
Not at all	518 (53.0)	1	1	1	1
Once or a few times	240 (24.5)	1.01 (0.80, 1.26)	0.99 (0.80, 1.23)	0.83 (0.63, 1.08)	0.85 (0.65, 1.10)
1-3 times a week	186 (19.0)	0.93 (0.72, 1.20)	0.95 (0.73, 1.24)	0.51 (0.35, 0.74)	0.55 (0.37, 0.81)
Almost every day	34 (3.5)	0.46 (0.20, 1.05)	0.43 (0.19, 0.97)	***	***
Missing	n=33				
<i>Remote sexual connection in last month</i>					
Sex over phone, video chat, or texting ³					
Not at all	888 (90.9)	1	1	1	1
Once or a few times	63 (6.4)	1.47 (1.09, 1.97)	1.13 (0.85, 1.51)	1.85 (1.36, 2.52)	1.53 (1.13, 2.06)
>1-3 times a week	26 (2.6)	1.41 (0.87, 2.24)	1.22 (0.79, 1.89)	1.11 (0.63, 1.95)	1.08 (0.63, 1.85)
Missing	n=33				
Use a hookup or dating app ³					
Not at all	947 (94.3)	1	1	1	1
Once or a few times	34 (3.4)	1.29 (0.82, 2.01)	0.89 (0.61, 1.29)	1.91 (1.36, 2.61)	1.55 (1.14, 2.10)
>1-3 times a week	23 (2.3)	2.04 (1.47, 2.83)	1.39 (0.85, 2.26)	1.48 (0.84, 2.61)	1.28 (0.75, 2.21)
Missing	n=6				
<i>Increased romantic relationship tension</i>					
No	492 (66.3)	1	1	1	1
Yes	250 (33.7)	3.06 (2.42, 3.87)	2.79 (2.19, 3.55)	3.16 (2.32, 4.31)	3.23 (2.37, 4.39)
Missing	n=4				

¹Covariates are age, gender, marital status, and income

²Partnered sexual activity defined as mutual masturbation, oral sex, or intercourse

³Few people reported sex over the phone, video chat, or texting (n=3) and using hookup or dating app (n=11) every day, so we combined this category with 1-3 times a week for the most frequent category

⁴Restricted to those who reported a romantic partner (weighted n=746).

Table 3. Adjusted log-binomial regression for the associations between frequency of social and sexual connections and outcomes of depression and loneliness, stratified by sexual relationship type (living with partner vs not living with partner)

Exposure variables	Depressive symptoms			Loneliness		
	Live with a partner aPR ¹ (95%CI)	Do not live with a partner aPR ¹ (95% CI)	p for interaction	Live with a partner aPR ¹ (95%CI)	Do not live with a partner aPR ¹ (95% CI)	p for interaction
<i>In-person social connection in last month</i>						
Hug or kiss a family member (not partner)						
<1-3 times a week	1	1		1	1	
1-3 times a week or more	0.68 (0.50, 0.91)	0.91 (0.66, 1.25)	0.2	0.63 (0.41, 0.96)	0.73 (0.50, 1.07)	0.6
In-person visit with non-household member						
<1-3 times a week	1	1		1	1	
1-3 times a week or more	0.94 (0.65, 1.37)	0.85 (0.59, 1.24)	0.7	0.91 (0.53, 1.55)	0.77 (0.51, 1.18)	0.6
<i>Remote social connection in last month</i>						
Video chat with friend/family member						
<1-3 times a week	1	1		1	1	
1-3 times a week or more	1.05 (0.81, 1.36)	0.93 (0.72, 1.21)	0.5	0.94 (0.65, 1.36)	1.20 (0.91, 1.58)	0.3
Send or receive a card or letter in the mail						
<1-3 times a week	1	1		1	1	
1-3 times a week or more	1.23 (0.79, 1.91)	1.47 (0.92, 2.34)	0.6	1.39 (0.78, 2.48)	1.00 (0.48, 2.07)	0.5
<i>In-person sexual connection in last month</i>						
Hug or kiss romantic partner						
<1-3 times a week	1	1		1	1	
1-3 times a week or more	0.68 (0.52, 0.89)	0.94 (0.64, 1.38)	0.2	0.44 (0.31, 0.64)	0.56 (0.32, 0.97)	0.5
Partnered sexual activity**						
<1-3 times a week	1	1		1	1	
1-3 times a week or more	0.81 (0.61, 1.08)	1.12 (0.69, 1.82)	0.3	0.48 (0.30, 0.76)	0.71 (0.35, 1.43)	0.4
<i>Remote sexual connection in last month</i>						
Sex over phone, video chat, or texting***						
<1-3 times a week	1	1		1	1	
1-3 times a week or more	0.89 (0.38, 2.07)	1.66 (0.97, 2.84)	0.2	0.34 (0.04, 2.70)	1.45 (0.85, 2.48)	0.2
Use a hookup or dating app***						
<1-3 times a week	1	1		1	1	
1-3 times a week or more	1.61 (0.88, 2.95)	1.67 (1.13, 2.48)	0.9	1.79 (0.71, 4.49)	2.28 (1.62, 3.22)	0.6

Increased romantic relationship tension ²						
No	1	1		1	1	
Yes	3.16 (2.39, 4.17)	2.00 (1.30, 3.08)	0.08	3.26 (2.23, 4.74)	3.27 (1.95, 5.47)	1.0

¹Covariates are age, gender, and income. Marital status excluded because of overlapping definition with stratifying variable.

²Restricted to those who reported a romantic partner (weighted n=746).

REFERENCES

1. Holmes, E.A., et al., *Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science*. Lancet Psychiatry, 2020.
2. Kawachi, I. and L.F. Berkman, *Social ties and mental health*. Journal of Urban health, 2001. **78**(3): p. 458-467.
3. Matthews, T., et al., *Social isolation, loneliness and depression in young adulthood: a behavioural genetic analysis*. Soc Psychiatry Psychiatr Epidemiol, 2016. **51**(3): p. 339-48.
4. Taylor, H.O., et al., *Social Isolation, Depression, and Psychological Distress Among Older Adults*. J Aging Health, 2018. **30**(2): p. 229-246.
5. Barari, S., et al., *Evaluating COVID-19 Public Health Messaging in Italy: Self-Reported Compliance and Growing Mental Health Concerns*. 2020.
6. Hou, Z., et al. *Assessment of public attention, risk perception, emotional and behavioural responses to the COVID-19 outbreak: social media surveillance in China*. 2020.
7. Li, S., et al., *The Impact of COVID-19 Epidemic Declaration on Psychological Consequences: A Study on Active Weibo Users*. Int J Environ Res Public Health, 2020. **17**(6).
8. Jackson, C. and M. Newall. *Americans Adapting to the Coronavirus Pause*. 2020 [cited 2020 April 28]; Available from: <https://www.ipsos.com/en-us/news-polls/axios-ipsos-coronavirus-index>.
9. Kirzinger, A., et al. *KFF Health Tracking Poll - Early April 2020: The Impact Of Coronavirus On Life In America*. 2020 [cited 2020 April 28]; Available from: <https://www.kff.org/health-reform/report/kff-health-tracking-poll-early-april-2020/>.
10. American Psychiatric Association. *New Poll: COVID-19 Impacting Mental Well-Being: Americans Feeling Anxious, Especially for Loved Ones; Older Adults are Less Anxious*. 2020 [cited 2020 April 28]; Available from: <https://www.psychiatry.org/newsroom/news-releases/new-poll-covid-19-impacting-mental-well-being-americans-feeling-anxious-especially-for-loved-ones-older-adults-are-less-anxious>.
11. GBD Disease and Injury Incidence and Prevalence Collaborators, *Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017*. Lancet, 2018. **392**(10159): p. 1789-1858.
12. Bruce, L.D., et al., *Loneliness in the United States: A 2018 National Panel Survey of Demographic, Structural, Cognitive, and Behavioral Characteristics*. Am J Health Promot, 2019. **33**(8): p. 1123-1133.
13. Yu, B., et al., *Trends in depression among Adults in the United States, NHANES 2005-2016*. J Affect Disord, 2020. **263**: p. 609-620.
14. Twenge, J.M., et al., *Age, period, and cohort trends in mood disorder indicators and suicide-related outcomes in a nationally representative dataset, 2005-2017*. J Abnorm Psychol, 2019. **128**(3): p. 185-199.
15. National Institute of Mental Health. *Major Depression*. 2019 April 28, 2020]; Available from: <https://www.nimh.nih.gov/health/statistics/major-depression.shtml>.
16. Das, A., *Loneliness does (not) have cardiometabolic effects: A longitudinal study of older adults in two countries*. Soc Sci Med, 2019. **223**: p. 104-112.
17. Cowling, B.J., et al., *Community psychological and behavioral responses through the first wave of the 2009 influenza A(H1N1) pandemic in Hong Kong*. J Infect Dis, 2010. **202**(6): p. 867-76.
18. Zhu, X., et al., *Changes in emotion of the Chinese public in regard to the SARS period*. 2008. **36**(4): p. 447-454.
19. Chong, M.Y., et al., *Psychological impact of severe acute respiratory syndrome on health workers in a tertiary hospital*. Br J Psychiatry, 2004. **185**: p. 127-33.
20. Matsuishi, K., et al., *Psychological impact of the pandemic (H1N1) 2009 on general hospital workers in Kobe*. Psychiatry Clin Neurosci, 2012. **66**(4): p. 353-60.

21. Richardson, A., et al., *The next generation of users: prevalence and longitudinal patterns of tobacco use among US young adults*. American journal of public health, 2014. **104**(8): p. 1429-1436.
22. Hall, J.P., et al., *Effect of Medicaid expansion on workforce participation for people with disabilities*. American journal of public health, 2017. **107**(2): p. 262-264.
23. Tsai, J., et al., *Homelessness among a nationally representative sample of US veterans: prevalence, service utilization, and correlates*. Social psychiatry and psychiatric epidemiology, 2016. **51**(6): p. 907-916.
24. Kohout, F.J., et al., *Two shorter forms of the CES-D depression symptoms index*. Journal of aging and health, 1993. **5**(2): p. 179-193.
25. Andresen, E.M., et al., *Screening for depression in well older adults: Evaluation of a short form of the CES-D*. American journal of preventive medicine, 1994. **10**(2): p. 77-84.
26. Zhang, W., et al., *Validating a shortened depression scale (10 item CES-D) among HIV-positive people in British Columbia, Canada*. PloS one, 2012. **7**(7).
27. Hughes, M.E., et al., *A short scale for measuring loneliness in large surveys: Results from two population-based studies*. Research on aging, 2004. **26**(6): p. 655-672.
28. Perissinotto, C.M., I.S. Cenzer, and K.E. Covinsky, *Loneliness in older persons: a predictor of functional decline and death*. Archives of internal medicine, 2012. **172**(14): p. 1078-1084.
29. Gerst-Emerson, K. and J. Jayawardhana, *Loneliness as a public health issue: the impact of loneliness on health care utilization among older adults*. American journal of public health, 2015. **105**(5): p. 1013-1019.
30. Brody, D.J., L.A. Pratt, and J.P. Hughes, *Prevalence of depression among adults aged 20 and over: United States, 2013-2016*. 2018: US Department of Health and Human Services, Centers for Disease Control and Prevention.
31. Rico-Uribe, L.A., et al., *Loneliness, social networks, and health: a cross-sectional study in three countries*. PloS one, 2016. **11**(1): p. e0145264.
32. Gordon, I., et al., *Oxytocin, cortisol, and triadic family interactions*. Physiology & behavior, 2010. **101**(5): p. 679-684.
33. Field, T., *Touch for socioemotional and physical well-being: A review*. Developmental review, 2010. **30**(4): p. 367-383.
34. Sumioka, H., et al., *Huggable communication medium decreases cortisol levels*. Scientific reports, 2013. **3**(1): p. 1-6.
35. Schoenenberg, K., A. Raake, and J. Koepe, *Why are you so slow?—Misattribution of transmission delay to attributes of the conversation partner at the far-end*. International Journal of Human-Computer Studies, 2014. **72**(5): p. 477-487.
36. Burkill, S., et al., *Using the web to collect data on sensitive behaviours: a study looking at mode effects on the British National Survey of Sexual Attitudes and Lifestyles*. PloS one, 2016. **11**(2).