TITLE: Gut microbiome diversity is an independent predictor of survival in cervical cancer patients receiving chemoradiation

Authors and Affiliations

Travis T. Sims, MD, MPH¹⁺, Molly B. El Alam, MPH²⁺, Tatiana V. Karpinets, PhD³, Stephanie Dorta-Estremera, PhD⁴, Venkatesh L. Hegde, PhD⁴, Sita Nookala, PhD⁴, Kyoko Yoshida-Court, PhD², Xiaogang Wu, PhD³, Greyson W. G. Biegert, BS², Andrea Y. Delgado Medrano, BS², Travis Solley, BS², Mustapha Ahmed-Kaddar, BS², Bhavana V. Chapman, MD², K. Jagannadha Sastry, PhD⁴, Melissa P. Mezzari, PhD⁵, Joseph F. Petrosino, PhD⁵, Lilie L. Lin, MD², Lois Ramondetta, MD¹, Anuja Jhingran, MD², Kathleen M. Schmeler, MD¹, Nadim J Ajami, PhD³, Jennifer Wargo, MD, MMSc⁶, Lauren E. Colbert, MD, MSCR^{2*}, Ann H. Klopp, MD, PhD^{2*}

¹Department of Gynecologic Oncology and Reproductive Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX, USA. ²Department of Radiation Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX, USA. ³Department of Genomic Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX, USA. ⁴Department of Thoracic Head and Neck Medical Oncology, The University of Texas MD Anderson Cancer Center and the UTHealth Graduate School of Biomedical Sciences at Houston, Houston, TX, ⁵Department of Molecular Virology and Microbiology, Alkek Center for Metagenomics and Microbiome Research, Baylor College of Medicine, Houston, TX, USA. ⁶Department of Surgical Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX, USA.

+ Authors Contributed Equally

* Shared corresponding authorship

Correspondence: L.E. Colbert or A.H. Klopp, Department of Radiation Oncology, Unit
1422, The University of Texas MD Anderson Cancer Center, 1515 Holcombe
Boulevard, Houston, TX 77030, USA. Telephone: 832-652-6033 (L.E.C.), 713-563-2444
(A.H.K); fax: 713-745-2398; e-mail: lcolbert@mdanderson.org,
aklopp@mdanderson.org.

ABSTRACT

Diversity of the gut microbiome is associated with higher response rates for cancer patients receiving immunotherapy but has not been investigated in patients receiving radiation therapy. Additionally, current studies investigating the gut microbiome and outcomes in cancer patients may not adjusted for established risk factors. Here, we sought to determine if diversity and composition of the gut microbiome was independently associated with survival in cervical cancer patients receiving chemoradiation. Our study demonstrates that the diversity of gut microbiota is associated with a favorable response to chemoradiation. Additionally, compositional variation among patients correlated with short term and long-term survival. Short term survivor fecal samples were significantly enriched in *Porphyromonas*, *Porphyromonadaceae*, and *Dialister*, whereas long term survivor samples were significantly enriched in *Escherichia Shigella*, *Enterobacteriaceae*, and *Enterobacteriales*. Moreover, analysis of immune cells from cervical tumor brush samples by flow cytometry revealed that patients with a high microbiome diversity had increased tumor infiltration of CD4+ lymphocytes as well as activated subsets of CD4 cells expressing ki67+ and CD69+ over the course of radiation therapy. The modulation of gut microbiota before chemoradiation might provide an alternative way to enhance treatment efficacy and improve treatment outcomes in cervical cancer patients.

Key words: gynecologic cancer, cervical cancer, HPV-associated cancers, microbiome, chemoradiation

MAIN

Cervical cancer continues to be one of the leading causes of cancer-associated mortality globally¹. More than 500,000 new cases of invasive cervical cancer will be diagnosed worldwide in 2020, resulting in over 300,000 deaths². Multimodality therapy consisting of definitive chemoradiation (CRT) comprising external-beam radiotherapy (EBRT) followed by intracavitary brachytherapy with concurrent systemic chemotherapy continues to be the standard of care in clinical practice for locally advanced disease³.

The fecal or gut microbiome, a diverse community of bacteria, archaea, fungi, protozoa, and viruses, is thought to influence host immunity by modulating multiple immunologic pathways, thus impacting health and disease^{4–6}. The diversity of the gut microbiome is defined as the number and relative abundance distribution of these distinct types of microorganisms colonizing within the gut⁷. Studies have suggested that dysbiosis of the gut microbiome confers a predisposition to certain malignancies and influences the body's response to a variety of cancer therapies, including chemotherapy, radiotherapy, and immunotherapy^{6,8–11}. For example, melanoma patients are more likely to have a favorable response to immune checkpoint blockade and exhibit improved systemic and antitumor immunity if they have a more diverse intestinal microbiome¹¹.

Radiotherapy promotes the activation of T cells directed against tumor antigens^{12–15}. In combination with immunotherapy, radiotherapy can maximize the antitumor immune response and promote durable disease control^{16,17}. We theorize that the gut microbiota may modulate radioresponse through immunologic mechanisms^{14,18}. Studies investigating the gut microbiome and outcomes in cancer patients often do not adjust for confounding patient and tumor characteristics. To assess this, we sought to identify independent gut microbial risk factors in cervical cancer (CC) patients receiving chemoradiation (CRT) and to evaluate their impact on

survival. We hypothesize that gut microbial differences may affect clinical outcomes in patients with cervical cancer

RESULTS

Patient Characteristics

A total of 55 patients with a mean age of 47 years (range, 29-72 years) volunteered to participate in this study. The patients received standard treatment for cervical cancer with 5 weeks of EBRT and weekly cisplatin. After completion of EBRT, patients received brachytherapy. For evaluation of treatment response, patients underwent magnetic resonance imaging (MRI) at baseline and week 5 and positron emission tomography (PET)/computed tomography (CT) 3 months after treatment completion (Fig. 1a). Most patients had stage IIB disease (51%) and squamous histology (78%). Their clinicopathologic data are summarized in Supplementary Table 1. We staged cervical cancer using the 2014 International Federation of Gynecology and Obstetrics staging system. The median cervical tumor size according to MRI was 5.4 cm (range, 1.2-11.5 cm). Thirty patients (55%) had lymph node involvement according to imaging studies. We first analyzed the bacterial 16S rDNA (16Sv4) fecal microbiota at baseline with respect to disease histology, grade, and stage. We found that the baseline α -diversity (within tumor samples) and β -diversity (between samples) of the fecal microbiome in the cervical cancer patients did not differ according to histology, grade, or stage ($P > 0.05$) (Supplementary Fig. 1a-d).

Univariate and multivariate analysis of factors affecting recurrence free survival (RFS) and overall survival (OS)

In the univariate Cox proportional hazard regression model predicting RFS, 3 covariates showed $P \leq 0.2$. As shown in Table 1, univariate analysis identified older age (Hazard Ratio (HR)

of 0.93 (95% CI = 0.87-0.98, $P = 0.0096$), Shannon diversity index (SDI) (HR of 0.51 (95% CI = 0.23-1.1, $P = 0.087$)) and body mass index (BMI) (HR of 0.92 (95% CI = 0.84-1, $P = 0.096$)) as risk factors for RFS. Multivariate survival analyses identified BMI and SDI as independent prognostic factors for RFS with a HR of 0.87 (95% CI = 0.77-0.98, $P = 0.02$) and 0.36 (95% CI = 0.15-0.84, $P = 0.018$) respectively. As shown in Table 2, univariate analysis identified SDI (HR of 0.34 (95% CI = 0.1-1.1, $P = 0.08$) and BMI (HR of 0.83 (95% CI = 0.69-1, $P = 0.055$)) as risk factors for OS. For OS, multivariate survival analyses again identified BMI and SDI as independent prognostic factors with a HR of 0.78 (95% CI = 0.623-0.97, $P = 0.025$) and 0.19 (95% CI = 0.043-0.83, $P = 0.028$) respectively.

Baseline Gut Microbiota Diversity is Associated with Favorable Responses

During the median follow-up period of 24.5 months, 7 patients died; all patients (12.7% of the total study population) died of disease (DOD). Figure 1 shows the Kaplan-Meier curves for RFS and OS. Given our univariate and multivariate analyses performed by Cox proportional hazard model identified SDI as an independent predictor for RFS and OS, we first tested the relationship between gut diversity and RFS and OS in our cohort by stratifying patients based on high and low SDI. We stratified the patients by SDI as high-diversity versus low-diversity groups based on the cutoff value of SDI (2.69) calculated by receiver operating characteristic curve (ROC). We demonstrate that patients with high fecal alpha diversity at baseline showed a trend toward prolonged RFS and OS when compared to those with low diversity ($P = 0.16$ and 0.094 , respectively) (Fig. 1a,b). Next, because our univariate and multivariate analyses performed by Cox proportional hazard model also identified BMI as an independent predictor for RFS and OS, we tested the relationship between diversity and RFS and OS in our cohort by stratifying patients based on high and low Shannon diversity metric and normal or high BMI. As shown in Figure

1d,e, when BMI and gut diversity are stratified for at baseline, patients with normal BMI and higher SDI had a longer median RFS duration ($P = 0.0027$) (Fig. 1d). OS (Fig. 1e) was longer for patients with normal BMI and higher gut diversity ($P = 0.2$).

Compositional Difference in Gut Microbiome in Response to chemoradiation

To further investigate whether the composition of gut microbiome was associated with response to CRT, we used Linear discriminant analysis (LDA) Effect Size analysis to identify bacterial genera that were differentially enriched in short term and long term cervical cancer patients ($P < 0.05$; LDA score > 3.5). In all patients, multiple taxa differed significantly at baseline between short and long term survivors. Specifically, short term survivor fecal samples were significantly enriched in *porphyromonas*, *porphyromonadaceae*, and *dialister*, whereas long term survivor samples were significantly enriched in *Escherichia Shigella*, *Enterobacteriaceae*, and *Enterobacteriales* ($P < 0.05$; LDA score > 3.5 , Fig. 2a,b). Our univariate analyses performed by Cox proportional hazard model identified *Pasteurellales*, *Haemophilus* and *Veillonella* as independent predictors for RFS and OS. We tested the relationship between these taxa and RFS and OS in our cohort by stratifying patients based on their relative abundance at baseline (Supplemental Fig. 2). We demonstrate that patients with high relative abundance of *Veillonella* at baseline showed a trend toward prolonged RFS and OS when compared to those with a low relative abundance at baseline ($P = 0.08$ and $P = 0.054$, respectively).

Association between Gut Microbiota Profile and Immune Signatures

Because the gut microbiota is thought to influence disease progression partially through modulating local immune response, we analyzed the cervical tumors in our cohort of patients via flow cytometry on tumor brushings performed before week 1, week 3 and week 5 of radiation therapy. To identify features associated with high gut diversity, Spearman correlation analysis was

conducted between immune signatures at each time point. High SDI was positively correlated with tumor infiltration of CD4 T cells at week 3, and CD4ki67+ T-cells at week 5, (Table 3 and Fig. 3a-d). These results suggest that patients with high gut diversity develop increased infiltration of activated CD4+ T-cell subsets.

DISCUSSION

The aim of this study was to identify independent gut microbial risk factors in cervical cancer patients receiving chemoradiation and to evaluate their impact on survival. We found BMI and gut diversity to be independent risk factors for RFS and OS in cervical cancer patients undergoing chemoradiation. Higher alpha gut diversity at baseline correlated with an improved RFS and OS. Our results indicate that overweight or obesity is a favorable prognostic factor independent of gut diversity. Additionally, our results demonstrate that patients with better clinical survival exhibit higher diversity as well as a distinct gut microbiome composition. Lastly, the association between gut diversity and local immune signatures highlights helper CD4+ T cells as potential mediators of antitumor immunity upon CRT treatment. Taken together, our results imply that the diversity of gut microbiota might be a shared benefit factor in those who respond well to CRT treatment.

It is now generally accepted that the gut microbiome modulates immune responses, antitumor immunity, and clinical outcomes in a variety of malignancies^{9,11,19}. The gut microbiome is thought to affect both innate and adaptive immune responses. Specifically how the gut microbiome exerts its influence continues to be explored, but this explanation may have important implications if specific taxa are found to change host response to treatment via immunomodulation⁶. In our study, T helper cell profiles at baseline correlate with gut diversity. These results confer that T cells and response to CRT are likely affected by the gut microbiota

independent of other factors such as BMI. Using multi-color flow cytometry we performed correlation analysis on individual immune signatures and microbiome diversity. The frequency of helper CD4⁺ T cells were chiefly identified. Cervical cancer is considered to be an immunogenic tumor because its origin is dependent on a persistent infection with human papillomavirus (HPV), most often HPV16 or HPV18²⁰. Previous studies have reported that the number and functional orientation of tumor-infiltrating CD4⁺ and CD8⁺ T cells and the presence of M1 type macrophages strongly correlates with survival in patients with cervical cancer after chemoradiation^{20,21}. T cells are capable of rapid antigen-specific responses and play critical roles in immune recall responses. In addition to the percentage of CD4⁺ T cell subsets, the increase in CD4 Ki67, CD4 CD69, and CD4 PD1 in patients with high microbiota diversity implies that the gut microbiome also modulates the proliferation of certain immune cell populations. Recent studies have already reported that chemoradiotherapy for cervical cancer induces unfavorable immune changes reflected by a decreased number of circulating lymphocytes, both CD4⁺ and CD8⁺ T cells, and an increased percentage in myeloid-cell populations, including myeloid-derived suppressor cells and monocytes²⁰. Our study suggest that CD4⁺ T cells infiltrating the tumor microenvironment support and encourage the activity of other immune cells by releasing T cell cytokines.

We found gut diversity to be associated with a favorable response to CRT against cervical cancer. Considering the correlation between gut diversity and local helper T cells being reshaped upon CRT treatment, we propose that patients harboring a more diverse gut microbiota at baseline may benefit from CRT to a greater extent. This might be mediated by the reprogramming of local antitumor immune responses. The significance of our study lies in that the modulation of gut microbiota before treatment might provide an alternative way to enhance the efficacy of CRT, specifically in advanced staged disease in which systemic failure of current therapies represents a

major challenge. Our results suggest that changes in the gut microenvironment contribute substantially to treatment success or failure, particularly in so-called immunogenic tumors like cervical cancer.

Our own group has previously characterized the gut microbiome of cervical cancer patients compared to healthy female controls, and have reported on differences in the relative abundance of specific taxa²². Our new findings support the hypothesis that organisms inhabiting the gut microbiome may be manipulated to improve cancer treatment response. Knowing specific gut microbial organisms that inhabit and undergo changes in patients with cervical cancer during CRT provides further insight into mechanisms that may modulate immune response and potentiate treatment outcomes in cancer patients. Researchers have already studied the treatment-enhancing utility of the gut microbiota in multiple areas of medicine^{10,23}. Additionally, there is emerging data describing the influence of the gut microbiome as it pertains to radiotherapy²⁴. Given that radiation can change the composition of the gut microbiome by altering the relative abundance of different taxa, we have to postulate whether radioresistant taxa ultimately alter the effectiveness of radiotherapy for cervical cancer^{6,25,26}. The results of our study illustrate the potential of intentionally modifying the gut microbiota to accumulate CRT-tolerant species as an interventional strategy to enhance response of cervical cancer to CRT. Furthermore, determining whether changes in the human gut microbiome during CRT affect patients' risk of treatment-related toxic effects may be an area that deserves further investigation.

The "obesity paradox", which suggest a positive association between increasing BMI as it pertains to a specific disease, was firstly reported in heart failure²⁷, but has since been described in a variety of disease processes, including other gynecologic cancers^{28–30}. Theories centered around the "obesity paradox" suggest that patients with a high BMI may be better able to withstand

cancer-induced consumption and stress compared with patients with a low BMI³¹. In uterine cancer it has been reported that the risk of recurrence differed significantly by BMI³². Specifically, a greater proportion of obese women met criteria for having a low risk of recurrence, while thin women tended to have a high-intermediate risk or recurrence. Many studies have investigated the impact of BMI on cervical cancer, but the association between weight and cervical cancer remains ambiguous³³. HPV is considered to be responsible for 99.7% of all cervical cancers³⁴, however, it has been suggested that obesity may further increase this risk^{35–37}. Other reports however do not report an association^{38,39}. For example, a review by Lane *et al.* finally refuted the relationship between cervical cancer and obesity ultimately citing a lack of evidence⁴⁰. The inconsistent conclusions among studies investigating the association between BMI and cervical cancer may be attributed to numerous factors including patient selection criteria, sample size and generalizability of the study population. Among these factors, patient selection criteria may be especially important, because tumor histology seems to be closely associated with BMI³⁷.

The strengths of this study include the use of careful clinical staging, histopathology, and reliable phylogenetic and statistical analysis to assess bacterial community compositional changes using both microbial divergence and taxon-based methods. Additionally, we followed a complete protocol for 16S analysis ranging from the sample collection method, DNA extraction, and microbiome sequencing, thus limiting artifactual variations. Although this study has yielded intriguing findings, an important limitation is the small sample size. Consequently, the sample size limited our ability to weigh statistical power. Yet despite the relatively small size of this prospective study, large statistically significant differences were still observed, and we believe the results presented herein provide solid evidence elucidating the role of the gut microbiome as it pertains to the treatment of cervical cancer. We hope the integration of these data will produce

actionable strategies geared toward targeting and manipulating the microbiome in order to ultimately improve cervical cancer therapy.

In conclusion, our study demonstrates that the diversity of gut microbiota is associated with a favorable response to chemoradiation. Additionally, compositional variation among patients correlated with short term and long-term survival. Our study demonstrates that gut diversity is a significant factor for predicting OS in CC patients undergoing CRT when BMI is accounted for, and may help explain the “obesity paradox” in cancer response. Moreover, analysis of immune cells from cervical tumor brush samples by flow cytometry revealed that an association between high microbial diversity, increased tumor infiltration of CD4⁺ lymphocytes and the activation of CD4 cells over the course of radiation therapy. The correlation between gut diversity and increased tumor infiltration of CD4⁺ lymphocytes suggest that patients harboring a more diverse gut microbiota at baseline may benefit from CRT to a greater extent. The significance of our study lies in that modulation of gut microbiota before chemoradiation might provide an alternative way to enhance treatment efficacy and improve treatment outcomes in cervical cancer patients. Additional studies exploring the relationship between gut diversity, chemoradiation, and treatment efficacy are needed to further understand the role of the gut microbiome in cervical cancer treatment.

METHODS

Participants and clinical data. Gut microbiome and cervical swab samples were collected prospectively from cervical cancer patients according to a protocol approved by The University of Texas MD Anderson Cancer Center Institutional Review Board (MDACC 2014-0543) for patients with biopsy-proven carcinoma of the cervix treated at MD Anderson and the Lyndon B. Johnson Hospital Oncology Clinic from September 22, 2015, to January 11, 2019. All patients had new

diagnoses of locally advanced, nonmetastatic carcinoma of the cervix and underwent definitive CRT with EBRT followed by brachytherapy. Patients received a minimum of 45 Gy via EBRT in 25 fractions over 5 weeks with weekly cisplatin followed by two brachytherapy sessions at approximately weeks 5 and 7 with EBRT in between for gross nodal disease or persistent disease in the parametrium. Patients with stage IB1 cancer were given CRT due to the presence of nodal disease. Clinical variables, demographics, and pathologic reports were abstracted from electronic medical records.

Sample collection and DNA extraction. Stool was collected from all patients by a clinician performing rectal exams at five time points (baseline; weeks 1, 3, and 5 of radiotherapy; and 3 months after CRT completion) using a matrix-designed quick-release Isohelix swab to characterize the diversity and composition of the microbiome over time. The swabs were stored in 20 µl of protease K and 400 µl of lysis buffer (Isohelix) and kept at -80°C within 1 h of sample collection.

16S rRNA gene sequencing and sequence data processing. 16S rRNA sequencing was performed for fecal samples obtained from all patients at four time points to characterize the diversity and composition of the microbiome over time. 16S rRNA gene sequencing was done at the Alkek Center for Metagenomics and Microbiome Research at Baylor College of Medicine. 16S rRNA was sequenced using approaches adapted from those used for the Human Microbiome Project⁴¹. The 16S rDNA V4 region was amplified via polymerase chain reaction with primers that contained sequencing adapters and single-end barcodes, allowing for pooling and direct sequencing of polymerase chain reaction products. Amplicons were sequenced on the MiSeq platform (Illumina) using the 2 x 250-bp paired-end protocol, yielding paired-end reads that

overlapped nearly completely. Sequence reads were demultiplexed, quality-filtered, and subsequently merged using the USEARCH sequence analysis tool (version 7.0.1090) (4). 16S rRNA gene sequences were bundled into operational taxonomic units at a similarity cutoff value of 97% using the UPARSE algorithm⁴². To generate taxonomies, operational taxonomic units were mapped to an enhanced version of the SILVA rRNA database containing the 16Sv4 region. A custom script was used to create an operational taxonomic unit table from the output files generated as described above for downstream analyses of α -diversity, β -diversity, and phylogenetic trends. Principal coordinates analysis was performed by institution and sample set to make certain no batch effects were present.

Flow Cytometry. Immunostaining was performed according to standard protocols⁴³. Cells were fixed using the Foxp3/Transcription Factor Staining Buffer Set (eBioscience, Waltham, MA) and stained with a 16 color panel with antibodies from Biolegend (San Diego, CA), BD Bioscience (San Jose, CA), eBioscience (Waltham, MA), and Life Technologies (Carlsbad, CA). Analysis was performed on a 5-laser, 18 color LSRFortessa X-20 Flow Cytometer (BD Biosciences, San Jose, CA). Analysis was performed using FlowJo version 10 (Flowjo LLC, Ashland, OR). We then followed a similar previously published method⁴³. Briefly, the cells were incubated with the antibodies for surface markers at 4°C in dark for 30 minutes. They were then washed twice with FACS buffer and fixed and permeabilized with FOXP3 Fix/perm Kit (ThermoFisher Scientific, Waltham, MA). Next, intracellular staining was performed by preparing the antibodies in permeabilization buffer and incubating the cells for 30 minutes at 4°C in the dark. Cells were washed with FACS buffer twice and prepared for acquisition on an LSR Fortessa X-20 analyzer at the Flow Cytometry Core at MD Anderson Cancer Center and were analyzed using FlowJo

software (FlowJo, LLC, Ashland, OR). Compensation controls were prepared using OneComp ebeads (eBioscience, Waltham, MA) and fluorescence minus one controls were used⁴³.

Statistical analyses. For microbiome analysis, rarefaction depth was set at 7066 reads. The Shannon diversity index (SDI) was used to evaluate α -diversity (within samples), and principle coordinates analysis of unweighted UniFrac distances was used to examine β -diversity (between samples). Patient and tumor characteristics were analyzed by univariate and multivariate Cox regression models for Recurrence-free survival (RFS) and Overall survival (OS) based on univariate p -value < 0.1 . Characteristics included age, body mass index (BMI), race, FIGO stage, grade, histology, nodal status, smoking status, antibiotic use and max tumor size. For each outcome of interest, a multivariate Cox regression analysis was performed to adjust for the effects of prognostic factors identified on univariate analysis as influencing survival in cervical cancer. These analyses were conducted using covariates with $p \leq 0.1$ in a stepwise fashion. We also ran a correlation analysis of alpha diversity metrics with tumor flow cytometry markers using a linear regression and Spearman's correlation. Alpha (within sample) diversity was evaluated using SDI. The relative abundance of microbial taxa, classes, and genera present in long term vs short term survivors was determined using LDA Effect Size⁴⁴, applying the one-against-all strategy with a threshold of 3.5 for the logarithmic LDA score for discriminative features and α of 0.05 for factorial Kruskal-Wallis testing among classes. Long term survivors were classified as patients who had a follow up of two years or more and were alive at time of last follow up, while short term survivors had a follow up of one year or less. LDA Effect Size analysis was restricted to bacteria present in 20% or more of the study population. Kaplan-Meier curves were generated for patients with normal BMI and overweight/obese BMI based on Cox analysis and clostridia

abundance. The significance of differences was determined using the log-rank test. Statistical significance was set at an alpha of 5% for a two-sided p-value. Analyses were conducted using Rstudio version Orange Blossom – 1.2.5033.

ACKNOWLEDGEMENTS

This research was supported in part by the Radiological Society of North America Resident/Fellow Award (to L.E.C.), the National Institutes of Health (NIH) through MD Anderson’s Cancer Center Support Grant P30CA016672, the Emerson Collective and the National Institutes of Health T32 grant #5T32 CA101642-14 (T.T.S). This study was partially funded by The University of Texas MD Anderson Cancer Center HPV-related Cancers Moonshot (L.E.C and A.K.). The human subjects who participated in this study are gratefully acknowledged.

AUTHOR CONTRIBUTION

All authors were involved in subject identification, data collection, interpretation of the statistical analysis, and review and approval of the final manuscript. The study concept was conceived by L.E.C., A.K., M.B.E.A, and T.T.S. The manuscript was written by T.T.S.

COMPETING INTERESTS

The authors report no conflicts of interest, financial or otherwise, related to the subject of this article.

ROLE OF FUNDING SOURCES

The funding sources were not involved in the research hypothesis development, study design, data analysis, or manuscript writing. Data access was limited to the authors of this manuscript.

The investigation described in this manuscript was presented in-part by Dr. Travis T. Sims during the American Society Of Clinical Oncology (ASCO) 2020 Annual Meeting, May 29-June 2, 2020.

REFERENCES

1. Islami F, Fedewa SA, Jemal A. Trends in cervical cancer incidence rates by age, race/ethnicity, histological subtype, and stage at diagnosis in the United States. *Prev Med*. 2019;123:316-323. doi:10.1016/j.ypmed.2019.04.010
2. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin*. 2019;69(1):7-34. doi:10.3322/caac.21551
3. Monk BJ, Tewari KS, Koh W-J. Multimodality Therapy for Locally Advanced Cervical Carcinoma: State of the Art and Future Directions. *J Clin Oncol*. 2007;25(20):2952-2965. doi:10.1200/JCO.2007.10.8324
4. The Human Microbiome Project Consortium. Structure, function and diversity of the healthy human microbiome. *Nature*. 2012;486(7402):207-214. doi:10.1038/nature11234
5. Cho JA, Chinnapen DJF. Targeting friend and foe: Emerging therapeutics in the age of gut microbiome and disease. *J Microbiol*. 2018;56(3):183-188. doi:10.1007/s12275-018-8037-z
6. Helmink BA, Khan MAW, Hermann A, Gopalakrishnan V, Wargo JA. The microbiome, cancer, and cancer therapy. *Nat Med*. 2019;25(3):377-388. doi:10.1038/s41591-019-0377-7
7. Jin Y, Dong H, Xia L, et al. The Diversity of Gut Microbiome is Associated With Favorable Responses to Anti-Programmed Death 1 Immunotherapy in Chinese Patients With NSCLC. *J Thorac Oncol Off Publ Int Assoc Study Lung Cancer*. 2019;14(8):1378-1389. doi:10.1016/j.jtho.2019.04.007
8. Cerf-Bensussan N, Gaboriau-Routhiau V. The immune system and the gut microbiota: friends or foes? *Nat Rev Immunol*. 2010;10(10):735-744. doi:10.1038/nri2850

9. McQuade JL, Daniel CR, Helmink BA, Wargo JA. Modulating the microbiome to improve therapeutic response in cancer. *Lancet Oncol.* 2019;20(2):e77-e91. doi:10.1016/S1470-2045(18)30952-5
10. Roy S, Trinchieri G. Microbiota: a key orchestrator of cancer therapy. *Nat Rev Cancer.* 2017;17(5):271-285. doi:10.1038/nrc.2017.13
11. Gopalakrishnan V, Spencer CN, Nezi L, et al. Gut microbiome modulates response to anti-PD-1 immunotherapy in melanoma patients. *Science.* 2018;359(6371):97-103. doi:10.1126/science.aan4236
12. Dewan MZ, Galloway AE, Kawashima N, et al. Fractionated but Not Single-Dose Radiotherapy Induces an Immune-Mediated Abscopal Effect when Combined with Anti-CTLA-4 Antibody. *Clin Cancer Res.* 2009;15(17):5379-5388. doi:10.1158/1078-0432.CCR-09-0265
13. Lugade AA, Moran JP, Gerber SA, Rose RC, Frelinger JG, Lord EM. Local Radiation Therapy of B16 Melanoma Tumors Increases the Generation of Tumor Antigen-Specific Effector Cells That Traffic to the Tumor. *J Immunol.* 2005;174(12):7516-7523. doi:10.4049/jimmunol.174.12.7516
14. Lhuillier C, Rudqvist N-P, Elemento O, Formenti SC, Demaria S. Radiation therapy and anti-tumor immunity: exposing immunogenic mutations to the immune system. *Genome Med.* 2019;11(1):40. doi:10.1186/s13073-019-0653-7
15. Lee Y, Auh SL, Wang Y, et al. Therapeutic effects of ablative radiation on local tumor require CD8⁺ T cells: changing strategies for cancer treatment. *Blood.* 2009;114(3):589-595. doi:10.1182/blood-2009-02-206870

16. Ngwa W, Irabor OC, Schoenfeld JD, Hesser J, Demaria S, Formenti SC. Using immunotherapy to boost the abscopal effect. *Nat Rev Cancer*. 2018;18(5):313-322. doi:10.1038/nrc.2018.6
17. Grassberger C, Ellsworth SG, Wilks MQ, Keane FK, Loeffler JS. Assessing the interactions between radiotherapy and antitumour immunity. *Nat Rev Clin Oncol*. 2019;16(12):729-745. doi:10.1038/s41571-019-0238-9
18. Sivan A, Corrales L, Hubert N, et al. Commensal Bifidobacterium promotes antitumor immunity and facilitates anti-PD-L1 efficacy. *Science*. 2015;350(6264):1084-1089. doi:10.1126/science.aac4255
19. Hooper LV, Littman DR, Macpherson AJ. Interactions between the microbiota and the immune system. *Science*. 2012;336(6086):1268-1273. doi:10.1126/science.1223490
20. Meir H van, Nout RA, Welters MJP, et al. Impact of (chemo)radiotherapy on immune cell composition and function in cervical cancer patients. *OncImmunology*. 2017;6(2):e1267095. doi:10.1080/2162402X.2016.1267095
21. Jordanova ES, Gorter A, Ayachi O, et al. Human leukocyte antigen class I, MHC class I chain-related molecule A, and CD8+/regulatory T-cell ratio: which variable determines survival of cervical cancer patients? *Clin Cancer Res Off J Am Assoc Cancer Res*. 2008;14(7):2028-2035. doi:10.1158/1078-0432.CCR-07-4554
22. Alexander JL, Wilson ID, Teare J, Marchesi JR, Nicholson JK, Kinross JM. Gut microbiota modulation of chemotherapy efficacy and toxicity. *Nat Rev Gastroenterol Hepatol*. 2017;14(6):356-365. doi:10.1038/nrgastro.2017.20

23. Kohl KD, Weiss RB, Cox J, Dale C, Denise Dearing M. Gut microbes of mammalian herbivores facilitate intake of plant toxins. van Dam N, ed. *Ecol Lett*. 2014;17(10):1238-1246. doi:10.1111/ele.12329
24. Muls A, Andreyev J, Lalondrelle S, Taylor A, Norton C, Hart A. Systematic Review: The Impact of Cancer Treatment on the Gut and Vaginal Microbiome in Women With a Gynecological Malignancy. *Int J Gynecol Cancer*. 2017;27(7):1550-1559. doi:10.1097/IGC.0000000000000999
25. Gerassy-Vainberg S, Blatt A, Danin-Poleg Y, et al. Radiation induces proinflammatory dysbiosis: transmission of inflammatory susceptibility by host cytokine induction. *Gut*. 2018;67(1):97-107. doi:10.1136/gutjnl-2017-313789
26. Sokol H, Adolph TE. The microbiota: an underestimated actor in radiation-induced lesions? *Gut*. 2018;67(1):1-2. doi:10.1136/gutjnl-2017-314279
27. Lavie CJ, Osman AF, Milani RV, Mehra MR. Body composition and prognosis in chronic systolic heart failure: the obesity paradox. *Am J Cardiol*. 2003;91(7):891-894. doi:10.1016/s0002-9149(03)00031-6
28. Lavie CJ, De Schutter A, Milani RV. Healthy obese versus unhealthy lean: the obesity paradox. *Nat Rev Endocrinol*. 2015;11(1):55-62. doi:10.1038/nrendo.2014.165
29. Walker JL, Piedmonte MR, Spirtos NM, et al. Laparoscopy Compared With Laparotomy for Comprehensive Surgical Staging of Uterine Cancer: Gynecologic Oncology Group Study LAP2. *J Clin Oncol*. 2009;27(32):5331-5336. doi:10.1200/JCO.2009.22.3248
30. S A-A, A S, J K. Obesity and cancer. *J Med*. 2002;33(1-4):73-86.

31. Chen H-N, Chen X-Z, Zhang W-H, et al. The Impact of Body Mass Index on the Surgical Outcomes of Patients With Gastric Cancer: A 10-Year, Single-Institution Cohort Study. *Medicine (Baltimore)*. 2015;94(42):e1769. doi:10.1097/MD.0000000000001769
32. Gunderson CC, Java J, Moore KN, Walker JL. The impact of obesity on surgical staging, complications, and survival with uterine cancer: A Gynecologic Oncology Group LAP2 ancillary data study. *Gynecol Oncol*. 2014;133(1):23-27. doi:10.1016/j.ygyno.2014.01.041
33. Poorolajal J, Jenabi E. The association between BMI and cervical cancer risk: a meta-analysis. *Eur J Cancer Prev Off J Eur Cancer Prev Organ ECP*. 2016;25(3):232-238. doi:10.1097/CEJ.0000000000000164
34. Chi DS, Berchuck A, Dizon DS, Yashar CM. *Principles and Practice of Gynecologic Oncology*. Seventh edition. Wolters Kluwer; 2017.
35. Nagaraju GP, Rajitha B, Aliya S, et al. The role of adiponectin in obesity-associated female-specific carcinogenesis. *Cytokine Growth Factor Rev*. 2016;31:37-48. doi:10.1016/j.cytogfr.2016.03.014
36. Martín-López R, Hernández-Barrera V, De Andres AL, Garrido PC, De Miguel AG, García RJ. Breast and cervical cancer screening in Spain and predictors of adherence. *Eur J Cancer Prev*. 2010;19(3):239–245. doi:10.1097/CEJ.0b013e3283372125
37. Lacey JV, Swanson CA, Brinton LA, et al. Obesity as a potential risk factor for adenocarcinomas and squamous cell carcinomas of the uterine cervix. *Cancer*. 2003;98(4):814-821. doi:10.1002/cncr.11567

38. Brinton LA, Herrero R, Reeves WC, de Britton RC, Gaitan E, Tenorio F. Risk Factors for Cervical Cancer by Histology. *Gynecol Oncol*. 1993;51(3):301-306.
doi:10.1006/gyno.1993.1294
39. Törnberg SA, Carstensen JM. Relationship between Quetelet's index and cancer of breast and female genital tract in 47,000 women followed for 25 years. *Br J Cancer*. 1994;69(2):358-361. doi:10.1038/bjc.1994.65
40. Lane G. Obesity and gynaecological cancer. *Menopause Int*. 2008;14(1):33-37.
doi:10.1258/mi.2007.007036
41. The Human Microbiome Project Consortium. A framework for human microbiome research. *Nature*. 2012;486(7402):215-221. doi:10.1038/nature11209
42. Edgar RC. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat Methods*. 2013;10(10):996-998. doi:10.1038/nmeth.2604
43. Dorta-Estremera S, Colbert LE, Nookala SS, et al. Kinetics of Intratumoral Immune Cell Activation During Chemoradiation for Cervical Cancer. *Int J Radiat Oncol Biol Phys*. 2018;102(3):593-600. doi:10.1016/j.ijrobp.2018.06.404
44. Segata N, Izard J, Waldron L, et al. Metagenomic biomarker discovery and explanation. *Genome Biol*. 2011;12(6):R60. doi:10.1186/gb-2011-12-6-r60

Figures

Fig 1. Relationship between gut diversity and BMI.

(A) Schema of the sample collection, treatment, and analyses used in the present study. Kaplan-Meier curves for (B) recurrence free survival, (C) overall survival stratified by high and low gut diversity. Kaplan-Meier curves for (D) recurrence free survival, (E) overall survival stratified by BMI and gut diversity. Cases represent patients.

Fig. 2. Compositional differences of the gut microbiome in short and long term survivors.

(A) The different abundance of bacterial genus between the two groups were identified by LefSe. It was significantly different when alpha value of the factorial Kruskal–Wallis test was <0.05 and the logarithmic LDA score was >3.0 . The left histogram showed the LDA scores of genera differentially abundant between the two groups. The taxonomy was listed, followed by its core group. Putative species (Specific OTUs) identified as significantly more enriched/depleted (Fisher/Wilcoxon test p value < 0.05) in patients with short-term vs long-term in baseline samples. (B) Cladogram representation of the significantly different taxa features from phylum (inner circle) to genus (outer circle) (C) The right heatmap showed the relative abundance of specific bacteria by phylum, class, order, family and genus between short-term and long-term survivors.

Fig 3. Correlation analysis of Shannon Diversity Index with tumor immune signatures.

(A,B,C,D) Spearman correlations between Shannon Diversity Index and CD4+, CD4+ Ki67+, CD4+ CD69+, and CD4+ PD1+. Statistical analysis was performed by Spearman correlation or Mann-Whitney tests.

Supplemental Fig 1. The fecal microbiota of individuals with cervical cancer.

The fecal microbiota of individuals with cervical cancer by demographics. Diversity (within sample diversity) was measured using the Shannon diversity metric and Beta diversity (between sample diversity) was determined by unweighted Unifrac. No differences were observed in either metric by cancer histology (A,D), grade (B,E) or cancer stage (C,F).

Supplemental Fig 2. Relationship between gut diversity and BMI.

Kaplan-Meier curves for (A) recurrence free survival, (B) overall survival stratified by relative abundance of *Pasteurellales*. Kaplan-Meier curves for (C) recurrence free survival, (D) overall survival stratified by relative abundance of *Haemophilus*. Kaplan-Meier curves for (E) recurrence free survival, (F) overall survival stratified by relative abundance of *Veillonella*. Cases represent patients.

551 **Table 1. Univariate and multivariate Cox regression analysis for recurrence-free survival**

Characteristics	Univariate model		Multivariate model	
	HR (95% CI)	P value	HR (95% CI)	P value
Age	0.93 (0.87-0.98)	0.0096	0.93* (0.88-0.99)	0.03‡
BMI (kg/m2)	0.92 (0.84-1)	0.096	0.87* (0.77-0.98)	0.02‡
Normal (18.5 to 24.9)	1 (reference)		—	—
Overweight (25 to 29.9)	0.81(0.26-2.53)	0.715	—	—
Obese (30 or more)	0.47(0.13-1.67)	0.240	—	—
Race/Ethnicity				
Asian	1 (reference)		—	—
Black	0.37(0.02-5.90)	0.479	—	—
Hispanic	0.39 (0.05-3.21)	0.382	—	—
White	0.39 (0.05-3.31)	0.390	—	—
Other	4.1309E-08(-inf - +inf)	0.998	—	—
Stage				
I	1 (reference)		—	—
II	1.50 (0.31-7.34)	0.615	—	—
III	3.99 (0.80-20.01)	0.091	—	—
IV	2.54 (0.23-28.12)	0.447	—	—
Grade				
Well	1 (reference)			
Moderate	55297546(-inf - +inf)	0.998	—	—
Poor	97336741.9(-inf - +inf)	0.998	—	—
Unknown	76285161.4(-inf - +inf)	0.998	—	—
Histology				
Squamous	1 (reference)		—	—
Adenocarcinoma/Adenosquamous	1.06(0.34-3.34)	0.918	—	—
Node Level on PET				
Common Iliac	1 (reference)		—	—
External Iliac	1.33 (0.35-4.95)	0.675	—	—
Internal Iliac	0.67 (0.07-6.89)	0.736	—	—

Characteristics	Univariate model		Multivariate model	
	HR (95% CI)	<i>P</i> value	HR (95% CI)	<i>P</i> value
Para-Aortic	1.31 (0.14-12.55)	0.818	—	—
None	0.34 (0.06-2.09)	0.245	—	—
Max Tumor Dimension on MRI	1.3 (1-1.8)	0.042	—	—
Smoking status				
Current	1 (reference)		—	—
Former	0.91 (0.10-7.84)	0.934	—	—
Never	0.89(0.11-7.17)	0.909	—	—
Antibiotic Use				
No	1 (reference)		—	—
Yes	78371200.7 (-inf - +inf)	0.998	—	—
Brachytherapy				
HDR	1 (reference)		—	—
PDR	1.41 (0.48-4.149)	0.532	—	—
Baseline Gut Alpha Diversity				
Observed OTU	0.99 (0.97-1)	0.21	—	—
Shannon	0.51 (0.23-1.1)	0.087	0.36* (0.15-0.84)	0.018‡
Simpson	0.025 (0.000036-1.7)	0.087	—	—
Inverse Simpson	0.93 (0.84-1)	0.11	—	—
Fisher	0.95 (0.88-1)	0.23	—	—
Camargo	13 (0.14-1300)	0.27	—	—
Pielou	0.02 (0.00026-1.6)	0.081	—	—

CI, Confidence interval; *HR*, hazard ratio.

*Significant hazard ratios.

‡Significant *P* value.

556 **Table 2. Univariate and multivariate Cox regression analysis for overall survival**

Characteristics	Univariate model		Multivariate model	
	HR (95% CI)	P value	HR (95% CI)	P value
Age	0.95 (0.87-1)	0.23	—	—
BMI (kg/m²)	0.83 (0.69-1)	0.055	0.78* (0.623-0.97)	0.025‡
Normal (18.5 to 24.9)	1 (reference)		—	—
Overweight (25 to 29.9)	0.23(0.08-2.32)	0.323	—	—
Obese (30 or more)	0.42 (0.06-4.56)	0.19	—	—
Race/Ethnicity				
Asian	1 (reference)		—	—
Black	4.46E-09 (-inf - +inf)	0.999	—	—
Hispanic	0.23(0.02-2.22)	0.204	—	—
White	0.17 (0.02-1.90)	0.151	—	—
Other	4.48E-09 (-inf - +inf)	0.999	—	—
Stage				
I	1 (reference)		—	—
II	1.19 (0.12-11.43)	0.881	—	—
III	1.49 (0.09-23.93)	0.776	—	—
IV	5.13 (0.32-82.34)	0.248	—	—
Grade				
Well	1 (reference)		—	—
Moderate	116103697.1 (-inf - +inf)	0.999	—	—
Poor	46065187.92(-inf - +inf)	0.999	—	—
Unknown	149251105.9(-inf - +inf)	0.999	—	—
Histology				
Squamous	1 (reference)		—	—
Adenocarcinoma/Adenosquamous	3.40 (0.69-16.90)	0.134	—	—
Node Level on PET				
Common Iliac	1 (reference)		—	—
External Iliac	1.099 (0.09-12.86)	0.306	—	—
Internal Iliac	4.83 (0.24-98.040)	0.999	—	—
Para-Aortic	5.9333E-08 (-inf - +inf)	0.354	—	—

Characteristics	Univariate model		Multivariate model	
	HR (95% CI)	P value	HR (95% CI)	P value
None	3.34(0.26-42.69)	0.940	—	—
Max Tumor Dimension on MRI	1.2 (0.77-1.8)	1.2	—	—
Smoking status				
Current	1 (reference)		—	—
Former	106318829.2 (-inf - +inf)	0.999	—	—
Never	61091037.65 (-inf - +inf)	0.999	—	—
Antibiotic Use				
No	1 (reference)		—	—
Yes	0.53 (0.06-4.56)	0.564	—	—
Brachytherapy				
HDR	1 (reference)		—	—
PDR	0.89 (-1.61-1.39)	0.884	—	—
Baseline Gut Alpha Diversity				
Observed OTU	0.98 (0.95-1)	0.14	—	—
Shannon	0.34 (0.1-1.1)	0.08	0.19* (0.043-0.83)	0.028‡
Simpson	0.0059 (1.2e-05-2.9)	0.1	—	—
Inverse Simpson	0.85 (0.7-1)	0.13	—	—
Fisher	0.91 (0.79-1)	0.15	—	—
Camargo	2200 (0.84-5800000)	0.055	—	—
Pielou	0.0036 (5e-06-2.5)	0.093	—	—

CI, Confidence interval; HR, hazard ratio.

*Significant hazard ratios.

‡Significant P value.

Table 3. Correlation of baseline gut diversity (Inverse Shannon Diversity) with phenotype of tumor infiltrating lymphocytes during chemoradiation treatment. The percent of live lymphocytes expressing each marker was correlated with baseline Shannon diversity of the gut microbiome.

	P-value*	Q value	R ₂
CD4+Ki.67+ at T4	0.004 [‡]	0.0714	
CD4+CD69+ at T3	0.004 [‡]	0.1429	
CD4+PD1+ at T3	0.0367 [‡]	0.2143	
CD4+CTLA4+ at T3	0.057	0.2857	
CD4+			0.1

*P-value correlation of immune metric with baseline gut diversity

[‡]Significant *P* value.

Supplemental Table 1. Patient and tumor characteristics (N=55)

	N	(%)
Median age, yrs (range)	48 (28-72)	—
BMI, Mean (SD), kg/m²	28.7(6.06)	—
Race/Ethnicity		
Asian	2	(36.4)
Black	4	(18.2)
Hispanic	24	(43.6)
White	24	(43.6)
Other	1	(1.8)
FIGO Stage		
IA1	1	(1.8)
IA2	0	(0)
IB1	5	(9.09)
IB2	6	(10.9)
IIA	3	(5.45)
IIB	28	(50.9)
IIIA	9	(16.3)
IIIB	0	(0)
IVA	3	(5.45)
IVB	0	(0)
Grade		
Well	4	(7.2)
Moderate	20	(36.3)
Poor	25	(45.4)
Unknown	6	(10.9)
Histology		
Squamous	43	(78.1)
Adenocarcinoma	8	(18.1)
Adenosquamous	3	(3.63)
Node Level on PET		
Common Iliac	9	(16.3)

External Iliac	23	(41.8)
Internal Iliac	5	(9.09)
Para-Aortic	3	(5.45)
None	15	(27.2)
Median cervical tumor size (cm)	5.4	—
Smoking status		
Current	4	(7.27)
Former	20	(36.3)
Never	31	(56.3)
Antibiotic Use		
No	5	(9.1)
Yes	50	(90.9)
Brachytherapy		
HDR	21	(38.2)
PDR	34	(61.8)
Concurrent Chemotherapy (cycles)		
<i>Cisplatin</i>		
(1-3)	2	(3.6)
(≥4)	51	(92.7)
<i>Carboplatin</i>		
(2)	1	(1.8)
<i>Carboplatin + Cisplatin</i>		
(2)+(2)	1	(1.8)

568 FIGO- International Federation of Gynecology and Obstetrics

569 HDR-High Dose Rate

570 PDR- Pulsed Dose Rate

571 **Supplemental Table 2. Univariate Cox regression analysis for recurrence-free survival – Alpha Diversity all**
572 **time points**

573

Characteristics	Univariate model	
	HR (95% CI)	<i>P</i> value
Observed OTU		
Baseline	0.99 (0.97-1)	0.21
Week 1	1 (0.97-1)	0.69
Week 3	0.99 (0.96-1)	0.59
Week 5	1 (0.98-1)	0.71
Week 12	1 (0.98-1)	0.77
Shannon		
Baseline	0.51 (0.23-1.1)	0.087
Week 1	0.94 (0.2-4.4)	0.94
Week 3	1.2 (0.25-5.6)	0.83
Week 5	0.83 (0.35-1.9)	0.66
Week 12	2.7 (0.13-57)	0.51
Simpson		
Baseline	0.025 (0.00036-1.7)	0.087
Week 1	13 (1.4e-05-1.2e+07)	0.13
Week 3	52 (6.6e-05-4.1e+07)	0.57
Week 5	0.31 (0.013-7.8)	0.48
Week 12	130000 (7.5e-13-2.2e+22)	0.56

Characteristics	Univariate model	
	HR (95% CI)	<i>P</i> value
Inverse Simpson		
Baseline	0.93 (0.84-1)	0.11
Week 1	0.96 (0.79-1.2)	0.69
Week 3	1 (0.95-1.1)	0.34
Week 5	1 (0.92-1.2)	0.54
Week 12	1.1 (0.81-1.4)	0.59
Fisher		
Baseline	0.95 (0.88-1)	0.23
Week 1	0.97 (0.86-1.1)	0.66
Week 3	0.96 (0.83-1.1)	0.6
Week 5	1 (0.91-1.2)	0.69
Week 12	1 (0.89-1.2)	0.81

574 *CI*, Confidence interval; *HR*, hazard ratio.

575 *Significant hazard ratios.

576 ‡Significant *P* value.

577 **Supplemental Table 3. Univariate Cox regression analysis for overall survival – Alpha Diversity all time**
 578 **points**

579

Characteristics	Univariate model	
	HR (95% CI)	<i>P</i> value
Observed OTU		
Baseline	0.98 (0.95-1)	0.14
Week 1	0.98 (0.94-1)	0.35
Week 3	0.97 (0.92-1)	0.21
Week 5	1 (0.96-1)	0.98
Week 12	NA (NA-NA)	1
Shannon		
Baseline	0.34 (0.1-1.1)	0.08
Week 1	0.48 (0.063-3.7)	0.48
Week 3	1.2 (0.25-5.6)	0.83
Week 5	0.23 (0.037-1.4)	0.11
Week 12	NA (NA-NA)	1
Simpson		
Baseline	0.0059 (1.2e-05-2.9)	0.1
Week 1	0.45 (1.1e-08-1.8e+07)	0.93
Week 3	0.009 (1.7e-07-490)	0.4
Week 5	1.4 (0.00063-3200)	0.93
Week 12	NA (NA-NA)	1

Characteristics	Univariate model	
	HR (95% CI)	<i>P</i> value
Inverse Simpson		
Baseline	0.85 (0.7-1)	0.13
Week 1	0.86 (0.62-1.2)	0.39
Week 3	0.81 (0.61-1.1)	0.15
Week 5	0.89 (0.66-1.2)	0.46
Week 12	NA (NA-NA)	1
Fisher		
Baseline	0.91 (0.79-1)	0.15
Week 1	0.91 (0.74-1.1)	0.34
Week 3	0.84 (0.64-1.1)	0.22
Week 5	0.99 (0.79-1.3)	0.94
Week 12	NA (NA-NA)	1

580 *CI*, Confidence interval; *HR*, hazard ratio.

581 *Significant hazard ratios.

582 ‡Significant *P* value.