Title: SARS-CoV-2 infection in London, England: Impact of lockdown on community point-prevalence, March-May 2020

Authors: Michael Edelstein¹ (ORCID 0000-0002-7323-0806), Chinelo Obi¹ (ORCID [0000-0002-1652-6627](https://orcid.org/0000-0002-1652-6627)), Meera Chand¹ (ORCID [0000-0001-9697-0340](https://orcid.org/0000-0001-9697-0340)), Susan Hopkins¹ (ORCID 0000-0001-5179-5702), Kevin Brown¹, Mary Ramsay¹ (ORCID [0000-0002-7156-7640](https://orcid.org/0000-0002-7156-7640))

1. National Infection Service, Public Health England, London, UK

Corresponding author: Michael Edelstein

Corresponding author email: Michael.edelstein@phe.gov.uk

Abstract

COVID-19 point prevalence PCR community testing allows disease burden estimation. In a sample of London residents, point prevalence decreased from 2.2% (95%CI 1.4;3.5) in early April (reflecting infection around lockdown implementation) to 0.2% (95%CI 0.03-1.6) in early May (reflecting infection 3-5 weeks into lockdown). Extrapolation from reports of confirmed cases suggest that 5-7.6% of total infections were confirmed by testing during this period. These data complement seroprevalence surveys improving the understanding of transmission in London.

Conflict of interest: None declared

Funding statement: This study was conducted as part of Public Health England routine COVID-19 surveillance activities. No specific funding was received.

Background

The World Health Organization declared COVID-19, a respiratory disease caused by SARS-CoV-2, a pandemic on March 11th 2020. The UK reported its first case on January 31st; and as of May 18th 2020 243,695 confirmed cases had been reported[1]. At the beginning of the study in early April, UK policy was to prioritise available tests for patients requiring hospitalisation and outbreaks in institutional settings. Evidence suggests that over 80% of COVID-19 cases are mild[2] and that the proportion who are asymptomatic when tested may be as high as 80%[3]. Therefore, using laboratory confirmation in hospitalised cases will severely underestimate disease incidence. Repeated testing among individuals in the community will help estimate trends in COVID-19 point prevalence and help to describe the characteristics of milder infection. We assessed COVID-19 point prevalence in London

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

between early April and early May 2020, which approximately reflect infection around the time of the lockdown and 3-5 weeks into lockdown.

Use of a participative cohort

Flusurvey (Flusurvey.net) is a participative surveillance cohort set up for influenza surveillance[4] and adapted for COVID19. Any member of the UK public can register and report on symptoms weekly. On March 27th 2020, all Flusurvey participants resident in London were asked for their consent and the consent of their household members, to receive a nasal swab for COVID-19 testing. Self-swabbing has previously been shown to be acceptable to Flusurvey participants[5] and uses packaging compliant with UN3733[6]. Together with the swab, Flusurvey participants and their household members were asked to report any cough, fever, shortness of breath and loss of smell in the two weeks prior to the swab using a short questionnaire. The first set of swabs and questionnaires were posted to participants on 31/03/2020 (wave 1). In the second phase of swabbing, half of the participants were randomly selected and sent a swab between 29 and 30/04/2020 (wave 2). In order to ascertain pre-lockdown point prevalence in the first wave of sampling, we divided wave 1 samples between timely (collected before 7/04/2020) and late (collected or received 7-29/04/2020) samples.

Nasal swabs were tested for viral RNA using polymerase chain reaction (PCR) at the Public Health England Respiratory Virus Unit using primers targeted to either the RdRp and/or Orf gene[7,8].

The age and sex distribution of the initial respondents were compared with the London population[9], and positivity rate by sex and age group calculated, together with 95% confidence intervals (95%CI). We described the symptoms that timely wave 1 participants reported and their association with being a case, using risk ratios together with 95%CI. We calculated the positive predictive value of each of the reported symptoms as well as for fever and cough as a cluster. We calculated the positivity rate among participants who returned their first swab late and their second swab, with 95%CI.

Characteristics of participants

1,373 individuals (582 London Flusurvey participants and 791 household members) consented to self-swabbing. By 12/05/2020, 1081 first wave samples were received (response rate: 79%) of which 17 were excluded because the participant no longer reported a London address. The participants who sent the remaining 1064 samples were distributed across each London Borough, although the sample underrepresented children (table 1). Age distribution of participants did not differ significantly by sex ($p=0.8$). Of the 698 individuals who received a second swab, 444 had returned them by 12/05/2020 (response rate 63.6%). Of the 1064 wave 1 samples received, 816 were timely and 248 were late.

Table1. Age distribution of sample compared with London population

Of the 816 timely samples from the first wave, 18 were positive in 14 households (positivity 2.2%, 95%CI 1.4-3.5). Positivity was higher among females and among 10-19 year olds (Table 2). Of the 18 positive individuals, 14 (78%) had experienced symptoms in the last 2 weeks and 4 (22%) had not. Symptom onset among symptomatic participants ranged from March 15-30th and time from symptom onset to sample collection ranged from 4-19 days. Cough was the most commonly reported symptom among cases and all symptoms were significantly associated with being COVID19 positive (table 3). The positive predictive value of these symptoms ranged 7.1-13.5% (table 3). Of the 248 late wave 1 samples, 2 (both female and symptomatic) were positive (0.8%, 95% CI 0.2-3.2) and only 1/444 samples received from wave 2 was positive (0.2%, 95% CI 0.03-1.6). The case was female, asymptomatic, not a household contact of a previously identified case, and had tested negative in early April. Of the 20 positive participants in phase 1 of swabbing, five (including three asymptomatic) were retested and all were negative. Of the 3 asymptomatic cases that were retested, none reported symptoms when asked again. Thirty-seven individuals lived in the 16 households which had a least one case. Household size ranged from 1-4 individuals. Five households had 2 cases (Table 4). No additional cases were found during wave 2 in the 5 households that were tested in both waves and had an individual who tested positive in wave 1.

Impact of the lockdown on point prevalence

This study is the first to describe changes in the point prevalence of COVID-19 infection in the community in London, and England more generally. The 2.2% point prevalence reflects infections acquired in London shortly before and after the lockdown on March 23rd 2020, and the positivity rate of 0.2% reflects infections during the second half of April, three to five weeks into the lockdown. The decrease in incidence suggests the lockdown was effective at reducing COVID-19 transmission in London. Positivity is likely to be an underestimate; apart from the usual limitations of PCR diagnostics, self-testing may be slightly less sensitive than administered swabs[10].

Around the time of our second swab, a pilot survey across England found a similar infection prevalence of 0.27% in early May[11]. In Iceland, a PCR-based screening in a random sample of the population had a 0.6% positivity[12]. In Vo, a town in the Veneto region of Italy, screening of the entire population around the time of the lockdown showed a 2.6% positivity[13]. The London point prevalence in early April was expected to be higher than other UK regions, based on the numbers of hospitalised cases per population at the time of swabbing[1].

Implications of findings

Evidence suggests that, in mild cases, the virus is detectable in the upper respiratory tract by PCR from approximately 2 days before to 10 days after the onset of symptoms in over 90% of cases[14,15]. A further 10% are positive at 14 days, with occasional reports of longer-term detection[14]. Based on these values, a community PCR point prevalence survey, which only identifies acutely infected individuals, is broadly equivalent to the number of cases acquired over a 12-day period plus an additional 10% of cases acquired in the 4 days before that. Assuming a 5 day lag between disease onset and hospital admission (and testing), and using April 3 as the most common date for the first set of swabs, our infections should correspond to those presenting to hospital and getting tested between March 23 – April 7. Over that time, 10,007 cases were reported in London[1], corresponding to a point prevalence of 0.11%. Compared with the 2.2% positivity from our survey, this suggests that, in London, only 5% of cases were being confirmed through the testing strategy at that time. During the time period around the second swabbing (April 23rd to May 8th, determined using a similar approach) there were 2254 reported cases in London, a point prevalence of 0.025% suggesting that 7.6% of infections were detected. These crude figures should be interpreted with caution because viral shedding likely declines over time, affecting test sensitivity in a more complex way. Nevertheless, this figure is consistent with a model that

estimated that between 4 and 8.4% of symptomatic cases were being confirmed through testing in hospitals[16].

Two findings were unexpected. First the positivity was higher among females than males, whereas sex-disaggregated data for COVID-19 show equal numbers of cases between men and women[17]. This could be due to a higher proportion of mild cases among females as higher severity among males is well documented[18]. Second, 45% of the cases with a date of symptom onset tested positive more than 10 days after the beginning of their symptoms; this compares published data suggesting 90% of mild cases testing negative 10 days post symptom onset[14]. These findings warrant further investigation.

The repeated questionnaire and swabbing enabled us to confirm that all retested positive cases who reported no symptoms were genuinely asymptomatic rather than pre-symptomatic.

There are limitations to our data. First, the sample is relatively small and purposive (as individuals self-select to register with the cohort) and is therefore likely to not be representative on characteristics beyond age and sex. Secondly, limited information about timing and symptoms makes reconstructing timelines among households difficult. Thirdly, we made a number of assumptions when estimating under-ascertainment, and did not fully take into account the temporal distribution of likelihood of PCR detection. Linking the swabbing results with information collected weekly in Flusurvey may enable us to reconstruct timelines for each case and to document the duration of symptoms and risk factors associated with becoming infected. This data should be interpreted in conjunction with seroprevalence studies, which are ongoing.

Ethics statement

Ethics approval for inviting participants for self-swabbing was obtained when Flusurvey was initially set up in 2008/9, and each participant was individually re-consented.

Table1. Age distribution of sample compared with London population

Age group	London		Flusurvey participants	
	n	%	n	%
<10	1215736	13.7	59	6.7%
10-19	988894	11.1	59	7%
20-44	3626354	40.7	345	43.2%
45-54	1160346	13	119	15.3%
55-64	857538	9.6	126	14.2%
65+	1059213	11.9	108	13.5%

Table2. COVID19 Positivity rate by testing period, age and sex

	Participants	Sex		Age					
		Male	Female	<10	10-19	20-44	45-54	55-64	65+
Testing early April 2020	COVID19 positive (n)	5	13	1	3	9	1	3	1
	COVID19 negative (n)	395	403	58	56	336	118	123	107
	Positivity (% 95%CI)	1.2 (0.4;2.9)	3.1 (1.7;5.3)	1.7 (0.04;9.1)	5.1 (1.1;14.1)	2.6 (1.2;4.9)	0.8 (0.02;4.6)	2.4 (0.5;6.8)	0.9 (0.02;5)
Testing mid-April 2020	COVID19 positive (n)	0	2	0	0	1	0	0	1
	COVID19 negative (n)	106	140	12	16	114	44	25	35
	Positivity (% 95%CI)	-	1.4 (0.3; 5.5)	-	-	0.9 (0.1; 6)	-	-	2.8 (0.4;18.3)
Testing Early May 2020	COVID19 positive (n)	0	1	0	0	1	0	0	0
	COVID19 negative (n)	214	229	35	40	165	74	69	60
	Positivity (% 95%CI)	-	0.4 (0.06;3.1)	-	-	0.6 (0.08; 4.2)	-	-	-

Table 3. Distribution of symptoms among Flusurvey London participants

Symptom	number of COVID positive reporting		number of COVID negative reporting		Risk Ratio	confidence interval	Positive predictive value (%)
	n	%	n	%			
Fever	7	39	45	6	6.9	3.6;13.1	13.5
Cough	8	44	92	12	3.8	2.2;6.7	8
Fever or cough	11	61	101	13	4.8	3.2; 7.3	9.8
Shortness of breath	5	28	45	6	4.9	2.2;10.9	10
Loss of smell	5	28	46	6	4.8	2.2;10.7	8

Table 4. Number of participants positive for SARS-CoV-2 infection among households with at least one participant testing positive

Household number	Number of tested household members	Number of positives in wave 1 (timely samples)	Number of positives in wave 1 (late samples)	Number of positives in wave 2
1	4	1	-	0
2	3	2		-
3	2	1	-	0
4	1	1	-	0
5	2	1	1	-
6	2	2	-	-
7	3	2	-	-
8	2	1	-	-
9	1	1	-	0
10	2	1	-	0
11	2	1	-	-
12	2	1	-	-
13	4	1	-	-
14	2	2	-	-
15	3	-	1	-
16	2	0	-	1

References

1. Public health England and NHSX. Coronavirus (COVID-19) in the UK.
<https://coronavirus.data.gov.uk/>
2. The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. Vital Surveillances: The Epidemiological Characteristics of an Outbreak of 2019 Novel Coronavirus Diseases (COVID-19) — China, 2020. China CDC Weekly. 2020, 2(8): 113-122
3. Sutton D, Fuchs K, D'Alton M, Goffman D. Universal Screening for SARS-CoV-2 in Women Admitted for Delivery. N Engl J Med. 2020 Apr 13
4. Adler AJ, Eames KT, Funk S, Edmunds WJ. Incidence and risk factors for influenza-like illness in the UK: online surveillance using Flusurvey. BMC Infect Dis. 2014 May 1;14:232
5. Wenham C, Gray ER, Keane CE, Donati M, Paolotti D, Pebody R et al. Self-Swabbing for Virological Confirmation of Influenza-Like Illness Among an Internet-Based Cohort in the UK During the 2014-2015 Flu Season: Pilot Study. J Med Internet Res. 2018 Mar 1;20(3):e71
6. Public Health England. Packaging and transport requirements for patient samples – UN3373. <https://www.gov.uk/government/publications/packaging-and-transport-requirements-for-patient-samples-un3373>
7. Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DKW et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. Euro Surveill. 2020 Jan;25(3)
8. National Centre for Viral Disease Control and Prevention. Specific primers and probes for detection 2019 novel coronavirus.
http://ivdc.chinacdc.cn/kyjz/202001/t20200121_211337.html
9. Office for national Statistics. National population projections: 2018-based.
<https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationprojections/bulletins/nationalpopulationprojections/2018based>
10. Seaman CP, Tran LTT, Cowling BJ, Sullivan SG. Self-collected compared with professional-collected swabbing in the diagnosis of influenza in symptomatic individuals: A meta-analysis and assessment of validity. Clin Virol. 2019 Sep;118:28-35
11. Office for National Statistics. Coronavirus (COVID-19) Infection Survey pilot: England, 14 May 2020.
<https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/bulletins/coronaviruscovid19infectionsurveyspilot/england14may2020>

12. Gudbjartsson DF, Helgason A, Jonsson H, Magnusson OT, Melsted P, Norddahl GL et al. Spread of SARS-CoV-2 in the Icelandic Population. *N Engl J Med*. 2020 Apr 14. doi: 10.1056/NEJMoa2006100. [Epub ahead of print]
13. Lavezzo E, Franchin E, Ciavarella C. Suppression of COVID-19 outbreak in the municipality of Vo, Italy. medRxiv preprint.
<https://www.medrxiv.org/content/10.1101/2020.04.17.20053157v1.full.pdf>
14. Liu Y, Yan LM, Wan L, Xiang TX, Le A, Liu JM et al. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis*. 2020 Mar 19. pii: S1473-3099(20)30232-2
15. Tan W, Lu Y, Zhang J, Wang J, Dan Y, Tan Z et al. Viral Kinetics and Antibody Responses in Patients with COVID-19.
<https://www.medrxiv.org/content/10.1101/2020.03.24.20042382v1>
16. Russell TW, Hellewell J, Abbott S, Golding N, Gibbs H, Jarvis CI et al. Using a delay-adjusted case fatality ratio to estimate under-reporting
https://cmmid.github.io/topics/covid19/global_cfr_estimates.html
17. Hu Y, Sun J, Dai Z, Deng H, Li X, Huang Q et al. Prevalence and severity of corona virus disease 2019 (COVID-19): A systematic review and meta-analysis. *J Clin Virol*. 2020 Apr 14;127:104371
18. Intensive care national audit and research centre. ICNARC report on COVID-19 in critical care. May 1st 2020.
<https://www.icnarc.org/DataServices/Attachments/Download/f48efee2-d38b-ea11-9125-00505601089b>