

1 **Title Page**

2

3 **Forecasting COVID-19 cases and deaths in epidemic-mitigating European**
4 **countries by Richards function-based regression analyses**

5 Cheng Long¹, Xinmiao Fu^{2,*}

6

7 ¹ Department of Orthopaedic, Sichuan University West China Hospital, Chengdu City, Sichuan Province
8 610041, China

9

10 ² Provincial University Key Laboratory of Cellular Stress Response and Metabolic Regulation, College of
11 Life Sciences, Fujian Normal University, Fuzhou City, Fujian Province 350117, China

12

13 * To whom correspondence should be addressed to Professor Xinmiao Fu (xmfu@fjnu.edu.cn)

14

15 **Running title:** Forecasting COVID-19 outbreaks in 10 European countries

16

17 **Keywords:** COVID-19, SARS-CoV-2, Richards function, coronavirus, pandemic, total
18 confirmed cases, total COVID-19 deaths

19

20 Word counts: 951

21

22

23

24

25 **Abstract:** The COVID-19 pandemic has hit many countries, and in some European
26 countries it has been mitigated since April. Here we applied Richards function to simulate
27 and forecast the course of COVID-19 epidemics in Italy, Spain, France, Germany, Turkey,
28 Belgium, Ireland, Netherlands, Portugal and Switzerland. Potential total COVID-19
29 confirmed cases in these countries were estimated to be 240400 ± 1300 , 294100 ± 4000 ,
30 178500 ± 800 , 176900 ± 700 , 155400 ± 1000 , 57900 ± 400 , 24000 ± 200 , 46200 ± 300 , 30000 ± 300
31 and 30700 ± 100 respectively. Most of these countries are predicted to approach ending
32 stage between late May and early June such that daily new cases will become minimal,
33 which may guide societal and economic restorations. In addition, total COVID-19 deaths
34 were estimated to be 33500 ± 300 , 28200 ± 200 , 27800 ± 200 , 8740 ± 80 , 4500 ± 30 , 9250 ± 70 ,
35 1530 ± 20 , 6240 ± 50 , 1380 ± 10 and 1960 ± 8 , respectively. To our best knowledge, this is the
36 first study forecasting the COVID-19 epidemic by applying the Richard function-based
37 regression analysis.
38

39

40 **Introduction**

41 The COVID-19 pandemic caused by the novel coronavirus (SARS-CoV-2) has hit many
42 countries [1], resulting in more than 4 million cases and over 300,000 deaths in the world as of
43 May 14, 2020. In European countries such as Italy and Spain, daily new cases are declining
44 since April, yet they are still as many as one thousand per day. In these epidemic-mitigating
45 countries, governments are attempting to restore normal societal and economic activity,
46 however, concerns are raised about the risk of COVID-19 resurgence. It is of significance to
47 precisely forecast when the epidemic in each country will become minimal such that societal
48 and economic restorations will not lead in the resurgence of the outbreak. Further, such
49 forecasting may also inform governments to get better preparedness for making control
50 measures and supplying medical resources. Estimations by traditional epidemiological models
51 apparently require much detailed data for analysis [2, 3]. Here we applied a simple data-driven,
52 Richards function-based approach for epidemic forecasting only based on the daily cumulative
53 confirmed COVID-19 cases. This approach has been explored for simulating 2003 SARS
54 outbreaks in several regions/countries by others [4, 5] and also for estimating COVID-19 deaths
55 in China by us recently [6].
56

57

58 **Methods**

59 The Richards function is expressed as follows [4, 5]:

$$y = a * (1 + (d - 1) * \exp((x - xc) * k))^{(1 / (1 - d))} \quad (1)$$

60 where y is the daily cumulative number of confirmed COVID-19 cases and x is day; a , d , xc and
61 k are constants. Specifically, total number of COVID-19 cases are given by parameter a in the
62 equation. Simulation was performed using Microcal Origin software with the Richard function.

63 **Results**

64 **Estimation of potential total COVID-19 confirmed cases**

65 We decided to collect the data of the cumulative confirmed cases from the website of
66 Worldometer [7] for 10 most affected and epidemic-mitigating European countries, i.e., Italy,
67 Spain, France, Germany, Turkey, Belgium, Ireland, Netherlands, Portugal and Switzerland (for
68 detail, refer to the supporting information file **original data.xlsx**). Regression analyses indicate
69 that the data of each country were well fitted with the Richards function (all R^2 values being
70 close to 0.999; **Table 1** and **Figure 1**). Potential total numbers of confirmed cases in the above
71 countries were estimated to be 240400 ± 1300 , 294100 ± 4000 , 178500 ± 800 , 176900 ± 700 ,
72 155400 ± 1000 , 57900 ± 400 , 24000 ± 200 , 46200 ± 300 , 30000 ± 300 and 30700 ± 100 respectively.
73 All parameters from regression analyses are presented in **Table S1** and can be utilized to predict
74 daily new cases in each country since May 15 according to equation 1.

75

76 In addition, we estimated the key date, on which daily new cases are lower than 0.1% of the
77 potential total confirmed cases as defined by us subjectively. It appears that most countries will
78 approach such late stage between late May and early June (refer to **Table 1**) such that daily new
79 cases will become minimal in comparison with potential total confirmed cases. In our opinion,
80 nationwide societal and economic restorations should be initiated at least not earlier than the
81 estimated key dates, although local restorations may be operated earlier according to the local
82 epidemic situation.

83 **Estimation of potential total COVID-19 deaths**

84 In several countries, crude fatality ratios are as high as over 10% such that people have raised
85 grave concerns about how many patients will die eventually. Assuming that the number of
86 deaths is proportional to the number of confirmed cases for the COVID-19 outbreak under
87 specific circumstances, we speculated that the cumulative COVID-19 deaths would also obey
88 the Richards function. Regression analyses indicate that the data of cumulative COVID-19
89 deaths in each country were well fitted with the Richards function (all R^2 values being close to
90 0.999; **Table 1** and **Figure 2**). All parameters of regression analyses are displayed in **Table S2**
91 and can be utilized to predict daily new deaths in each country since May 15 according to
92 equation 1. Potential total COVID-19 deaths in Italy, Spain, France, Germany, Turkey, Belgium,

93 Ireland, Netherlands, Portugal and Switzerland were estimated to be 33500 ± 300 , 28200 ± 200 ,
94 27800 ± 200 , 8740 ± 80 , 4500 ± 30 , 9250 ± 70 , 1530 ± 20 , 6240 ± 50 , 1380 ± 10 and 1960 ± 8 ,
95 respectively. Crude fatality ratios (total deaths among total confirmed cases) of these countries
96 could be estimated to be $13.9 \pm 0.9\%$, $9.6 \pm 0.9\%$, $15.6 \pm 0.7\%$, $4.9 \pm 0.9\%$, $2.9 \pm 0.7\%$, $16.0 \pm 0.9\%$,
97 $6.4 \pm 1.3\%$, $13.5 \pm 0.9\%$, $4.6 \pm 0.8\%$ and $6.4 \pm 0.4\%$, respectively.
98

99 **Discussion**

100 To our best knowledge, this is the first study forecasting the COVID-19 epidemic by applying
101 the Richard function-based regression analysis. It should be pointed out that all the above
102 estimates are based on the assumption that there are no significant changes regarding the current
103 interventions, including non-pharmacological interventions such as quarantine and other
104 prevention measures and also diagnosis and treatment approaches. In addition, UK and Sweden,
105 two most affected European countries with more than 240,000 and 29,000 COVID-19
106 confirmed cases as of May 15, respectively, were not subjected to analysis because their daily
107 new cases have not yet started to significantly decline (**Figure. S1**).
108

109 Collectively, our forecasting on the course of COVID-19 outbreaks in these European countries
110 may provide valuable guidance for governments and also the public to get better preparedness
111 and optimize their efforts to contain this unprecedented crisis at current critical stage. The
112 estimated key date for each country may serve as a reference for operating societal and
113 economic restorations such that the resurgence of COVID-19 can be avoided, minimized or
114 under control according to COVID-19 containment in China [8, 9]. In addition, because
115 thousands of COVID-19 patients have died in these countries, our prediction on the course of
116 COVID-19 deaths (refer to **Figure 2** and **Table S2**) may benefit the mental health service that
117 needs to be timely provided to the families of passed patients [10]. As a matter of fact, we
118 precisely estimated total COVID-19 deaths in different areas of China (including Hubei
119 Province and Wuhan city, the epicenter of the COVID-19 outbreaks in China [1]) by applying
120 both Boltzmann and Richards functions-based regression analyses [6]. Such estimation help
121 governments to evaluate the severity of the COVID-19 epidemics.
122

123 **Acknowledgments**

124 This work is support by the National Natural Science Foundation of China (No. 31972918 and
125 31770830 to XF). We thank graduate students (Zuqin Zhang, Youhui Zeng, Shuting Shi and
126 Zhijie Huang, all from Professor Fu's research group at Fujian Normal University) for data
127 collection.

128

129 **Author contribution:** X.F. designed the study; C.L. organized data and performed analysis;
130 X.F. wrote the manuscript.

131

132 **Conflict of interest:** none.

133

134

135 **References**

- 136 (1) WHO: Coronavirus disease (COVID-19) outbreak,
137 <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>.
- 138 (2) Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and international spread of
139 the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *Lancet* 2020; DOI:
140 10.1016/S0140-6736(2020)30260-30269.
- 141 (3) Yang Y, et al. Epidemiological and clinical features of the 2019 novel coronavirus outbreak in China.
142 <https://doi.org/10.1101/2020021020021675> 2020.
- 143 (4) Wang XS, Wu J, Yang Y. Richards model revisited: Validation by and application to infection dynamics. *J Theor
144 Biol* 2012; 313: 12–19.
- 145 (5) Hsieh YH. Richards Model: A Simple Procedure for Real-time Prediction of Outbreak Severity. *Modeling and
146 Dynamics of Infectious Diseases* 2009: p. 216–236 https://doi.org/10.1142/9789814261265_9789814260009.
- 147 (6) Gao Y, et al. Forecasting the cumulative number of COVID-19 deaths in China: a Boltzmann function-based
148 modeling study. *Infect Control Hosp Epidemiol* 2020: 1-3.
- 149 (7) Worldometer: COVID-19 CORONAVIRUS PANDEMIC, <https://www.worldometers.info/coronavirus/>.
- 150 (8) Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19),
151 <https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf>;
152 2020.
- 153 (9) National Health Commission of China: Update of the COVID-19 Outbreak Data (in Chinese),
154 http://www.nhc.gov.cn/xcs/yqfkdt/gzbd_index.shtml.
- 155 (10) Xiang YT, et al. Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *Lancet
156 Psychiatry* 2020; 7(3): 228-229.

157

158

159 **Table 1 Estimation of potential total COVID-19 confirmed cases and deaths in European**
160 **countries.**

| Countries | Total confirmed cases | | Key date ^a | Total deaths | | Crude 161 fatality ratio (%) ^b |
|-------------|-----------------------|----------------|-----------------------|--------------|----------------|---|
| | Number | R ² | | Number | R ² | |
| Italy | 240400±1300 | 0.999 | June 6 | 33500 ±300 | 0.999 | 13.9±0.9 |
| Spain | 294100±4000 | 0.999 | June 13 | 28200±200 | 0.999 | 9.6±0.9 |
| France | 178500±800 | 0.999 | May 18 | 27800±200 | 0.999 | 15.6±0.7 |
| Germany | 176900±700 | 0.999 | May 24 | 8740±80 | 0.999 | 4.9±0.9 |
| Turkey | 155400±1000 | 0.999 | June 8 | 4500 ±30 | 0.999 | 2.9±0.7 |
| Belgium | 57900±400 | 0.999 | June 5 | 9250±70 | 0.999 | 16.0±0.9 |
| Ireland, | 24000±200 | 0.998 | May 25 | 1530±20 | 0.997 | 6.4±1.3 |
| Netherlands | 46200±300 | 0.999 | June 4 | 6240±50 | 0.999 | 13.5±0.9 |
| Portugal | 30000±300 | 0.999 | June 9 | 1380±10 | 0.999 | 4.6±0.8 |
| Switzerland | 30700±100 | 0.999 | May 14 | 1960±8 | 0.999 | 6.4±0.4 |

162 ^a Key date is determined when daily new cases are less than 0.1% of the potential total confirmed cases.


163 ^b Crude fatality ratio was calculated as total deaths among total confirmed cases, and the standard deviation
164 was calculated according to partial differential equations.

165

166

167 Figures and legends

Figure 1



168

169 **Figure 1. Fitting the cumulative confirmed COVID-19 cases of each 170 European country to the Richards function**


171 (A, B) Plots of the cumulative number of COVID-19 confirmed cases of the indicated European countries as of May 14, 2020, with the simulation results being plotted as Magenta lines.
172 Simulation was performed using Microcal Origin software with the Richard function. All
173 parameters from regression analyses are shown in Table S1.
174

175

176

177

Figure 2



178

Figure 2. Fitting the cumulative COVID-19 deaths of each European country to the Richards function

(A, B) Plots of the cumulative number of COVID-19 confirmed cases of the indicated European countries as of May 14, 2020, with the simulation results being plotted as Magenta lines. Simulation was performed using Microcal Origin software with the Richard function. All parameters from regression analyses are shown in Table S2.

185

186