1 Impaired T cell functions along with elevated activated Tregs at the early stage of 2 asymptomatic SARS-CoV-2 infection

3

4 Jingyi Yang^{1,2,3,*}, Ejuan Zhang^{1,2,3,*}, Maohua Zhong^{1,2,3,*}, Qingyu Yang^{2,3,4}, Ke Hong^{3,4}, Ting
5 Shu^{2,3,4}, Dihan Zhou^{1,2}, Jie Xiang^{3,4}, Jianbo Xia⁶, Xi Zhou^{1,2,3,7}, Dingyu Zhang^{3,4}, Chaolin
6 Huang^{3,4,†}, You Shang^{5,3,4,†}, Huimin Yan^{1,2,3,7,8, †}

7

8 1 Joint Laboratory of Infectious Diseases and Health, Wuhan Institute of Virology & Wuhan
9 Jinyintan Hospital, Wuhan Institute of Virology, Center for Biosafety Mega-Science, Chinese
10 Academy of Sciences (CAS), Wuhan, Hubei 430023 China

11 2 State Key Laboratory of Virology, Wuhan Institute of Virology, Center for Biosafety
12 Mega-Science, CAS, Wuhan, Hubei 430071, China

13 3 Center for Translational Medicine, Jinyintan Hospital, Wuhan, Hubei 430023 China

14 4 Joint Laboratory of Infectious Diseases and Health, Wuhan Institute of Virology & Wuhan
15 Jinyintan Hospital, Wuhan Jinyintan Hospital, Wuhan, Hubei 430023 China

16 5 Department of Critical Care Medicine, Union Hospital, Tongji Medical College, Huazhong
17 University of Science and Technology, Wuhan, Hubei 430030 China

18 6 Department of Laboratory Medicine, Maternal and Child Health Hospital of Hubei Province,
19 Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei 430070
20 China

21 7 University of Chinese Academy of Sciences, Beijing 100049 China

22 8 Lead Contact

23 *Contributed equally

24 † Correspondence: hmyan@wh.iov.cn (H.Y.), you_shanghust@163.com (Y.S.),
25 88071718@qq.com (C. H)

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

1 **Summary**

2 **Background** Limited data are available on the T cell responses for the asymptomatic
3 SARS-CoV-2 infection case.

4 **Methods** An imported SARS-CoV-2 infection case in Wuhan was admitted in hospital for
5 quarantine and observation. The T cell responses were followed up by flow cytometry analysis of
6 the peripheral blood nonnuclear cells (PBMCs) at days 7, 13, 22, and 28 after admission.

7 **Findings** We found the imported SARS-CoV-2 infection in Wuhan is an asymptomatic case. His T
8 cell differentiation, proliferation and activation matched the classical kinetics of T cell responses
9 induced by viral infection, but the activation maintained at a relatively low level. Function
10 analysis indicated frequencies of IFN- γ producing CD4⁺ and CD8⁺ T cells were notably lower
11 than that of the healthy controls (HC) at day 7, and then rebound gradually. But IFN- γ ⁺CD8⁺ T
12 cells were detained at a significant lower level even at day 28, when the SARS-CoV-2 virus had
13 already become undetectable for 3 weeks. Moreover, percentage of IL-17 producing CD4⁺ T cells
14 was also detained constantly at a much lower level compared to HC. At day 7, although
15 percentage of Tregs was in normal range, the frequency of activated Treg (aTreg) was remarkably
16 as high as 4·4-fold of that in HC.

17 **Interpretation** The T cell activation in the asymptomatic SARS-CoV-2 infection experienced a
18 significant suppression and presented impairment of Th1/Th17 and CD8⁺ T cell functions. Early
19 elevation of the aTregs might play role in the activation and function of T cells in the
20 asymptomatic SARS-CoV-2 infection.

1 Introduction

2 In December 2019, an epidemic of severe acute respiratory syndrome coronavirus-2
3 (SARS-CoV-2) causing human disease named coronavirus disease (COVID-19) was firstly
4 reported in Wuhan China. Up to May 9, 2020, a total of 3,855,788 cases have been reported with
5 265,862 deaths according to WHO report. SARS-CoV-2 infection is characterized by a broad
6 range of symptoms including fever, dry cough, and general malaise in the majority of cases while
7 severe pneumonia, acute respiratory distress syndrome (ARDS), septic shock and/or multiple
8 organ failure in a minor population^{1,2}. Based on the clinical presentation, COVID-19 patients can
9 be classified into mild, moderate, severe and critical. However, a group of laboratory-confirmed
10 SARS-CoV-2 infected cases, found by assay with quantitative real-time reverse transcription PCR
11 (qRT-PCR), presented neither clinical symptom nor radiographic abnormality. This group of
12 people was designated as asymptomatic infected individuals. The suspected rate of asymptomatic
13 infections is substantially high, which might surpass 17.9%^{3,4}. Furthermore, the viral load
14 detected in the asymptomatic case was similar to that in the symptomatic patients, which suggests
15 high transmission potential of the asymptomatic individuals^{5,6}. Actually, a growing number of
16 reports have evidenced substantial asymptomatic transmissions^{4,7,8}.

17 A study of a patient with mild-to-moderate COVID-19 disease showed that the CD4⁺ T and
18 CD8⁺ T cells were activated after SARS-CoV-2 infection and before the resolution of symptoms⁹,
19 suggesting possible potent protective T cell functions against SARS-CoV-2 infection. On the other
20 hand, several studies showed that viral infection may cause significant decrease of T lymphocytes
21 and impairment of T cell function in COVID-19 patients, especially in severe COVID-19 cases^{2,10}.
22 Exhausted function of CD8⁺ T cells with the increased expression of NKG2A was also reported in
23 severe cases¹¹. But so far, how T cells behave and function following acute infection of
24 SARS-CoV-2 in asymptomatic infected individuals remains largely unknown. The suspected high
25 rate of asymptomatic infections and substantial asymptomatic transmissions assure high necessity
26 of answering the above questions^{3,4,7,8}. In this article, we reported the kinetics of CD4⁺ T and
27 CD8⁺ T cell responses in an asymptomatic SARS-CoV-2 infected case, accompanied with the
28 clinical and virological features.

1 **Methods**

2 **Study design, patients and healthy control**

3 We did a prospective study of an imported SARS-CoV-2 infection case in Wuhan, Hubei,
4 China. A man in 30s of Hubei resident, who had traveled to Thailand from Wuhan around
5 mid-January, 2020, returned by airplane and was tested positive for SARS-CoV-2 upon arrival at
6 Wuhan on March 2020. He was transferred to Wuhan Jinyintan Hospital for isolation. At day 14
7 post hospitalization, the man was discharged to a designated hotel for another 14-day-isolation.
8 We followed up the man for 28 days during the hospitalization and isolation.

9 Eleven healthy age-matched male volunteers were enrolled as healthy controls. These
10 volunteers were confirmed without ongoing or past SARS-CoV-2 infection by detection of
11 SARS-CoV-2 nucleotide acid in nasopharyngeal swab samples, plasma SARS-CoV-2 specific IgM
12 and IgG using qRT-PCR or the colloidal gold strip, respectively.

13 This study was reviewed and approved by the Medical Ethical Committee of Wuhan
14 Jinyintan hospital (approval number KY-2020-47-01). Written informed consent was obtained
15 from the patient and the healthy controls.

16 **Clinical laboratory measurements**

17 Serial nasopharyngeal swab samples with or without sputum were obtained on days 0 (2
18 hours before hospitalization), 1, 2, 5, 8, 13, 22, 28 and were tested for the expression of E gene,
19 RdRp (R) gene, and N gene of SARS-CoV-2 using qRT-PCR as described by our previous study¹².

20 Clinical laboratory investigations were performed, included series of complete blood count,
21 serum biochemical test (including liver and renal function, creatine kinase, LDH, and electrolytes),
22 coagulation profile, common pathogens , as well as interleukine 6 (IL-6).

23 **Antibody detection and T cell responses evaluation**

24 Plasma and blood cells were separated from fresh peripheral blood from the SARS-CoV-2
25 infected case and healthy volunteers. Plasma was used for SARS-CoV-2 binding IgM or IgG
26 detections. PBMCs were separated from the blood cells by Ficoll-plaque density gradient
27 centrifugation and resuspended in complete RPMI1640 medium containing 10% FBS (Gibco), 1%
28 penicillin and 1% streptomycin. PBMCs were then divided into three panels for analysis: panel 1
29 for T cell differentiation, proliferation and activation detection, panel 2 for T cell cytokine
30 production detection, and panel 3 for Treg detection. PBMCs of panel 1 and panel 3 were directly

used for staining while PBMCs of panel 2 were stimulated with 200 ng/ml PMA (Beyotime, China), 2.5 μ M ionomycin (Beyotime, China) in the presence of 1 μ M monensin (BioLegend, USA) and 2.5 μ g/ml Brefeldin A at 37°C, 5% CO₂ for 4.5 h before staining. PBMCs were stained with dead cell discrimination marker (eBioscience™ Fixable Viability Dye eFluor™ 506, FVD) and surface-staining antibodies in PBS at 4°C for 30 min. After washing with PBS, cells were fixed with fixation/permeabilization buffer (eBioscience) at 4°C overnight, and then stained with the respective panel of intracellular markers in a permeabilization buffer at 4°C for 30 min. Antibodies used for each panel were listed in supplementary table 1. A BD LSR Fortessa flow cytometer (Becton Dickinson) was used to assess the stained cells and data were analyzed using FlowJo V7.0.

Statistical analysis

Data analysis was carried out with InStat, version 5.0 (GraphPad Software, La Jolla, CA, USA) and the data of healthy controls were presented as mean $\pm$ SD with dot plots.

Results

After admission on March, 2020, the man received regular physical and clinical examinations and treatments. He was a healthy smoker taking no medications. Clinical examination revealed a temperature of 36.2 °C, a pulse rate of 85 beats per minute, a blood pressure of 156/106 mm Hg, a respiratory rate of 21 breaths per minute, and oxygen saturation 98% while breathing ambient air. SARS-CoV-2 was again detected at days 1-2 in sputum and/or nasopharyngeal swab but was undetectable since day 5 (figure 1A and Supplementary Table 2). No other respiratory pathogens and common pathogens were detected. On day 14, the man was discharged to a designated hotel for another 14-day-isolation. Within the 28-day follow-up, the man experienced no clinical symptom, no fever, no lethargy, no sore throat, no chest pain, no dyspnea, and no dry cough. Moreover, chest CT images taken during hospitalization did not show any significant abnormality. Taken together, this man was an asymptomatic SARS-CoV-2 infected individual characterized as no clinical symptom or CT abnormality but SARS-CoV-2 positive by repeated qRT-PCR tests.

Among the tested biochemical markers, only serum amyloid A (SAA) and high-sensitivity C-reactive protein (hsCRP) increased above the upper limit of normal (ULN) at day 8 and day 14. On day 1, both SAA and hsCRP were at normal level (range of SAA, 0-10mg/ml & range of

1 hsCRP, 0-5mg/ml). SAA was elevated to 11.7 mg/ml at day 8 and 15.07 mg/ml at day 14 while
2 hsCRP was increased to 7.7 mg/ml at day 8 and 13.5 mg/ml at day 14. In contrast to the
3 significant increase in moderate and severe COVID-19 patients^{2, 10, 13}, IL-6 in the subject
4 constantly kept in the normal range. SARS-CoV-2-binding IgM was presented in plasma since day
5 7 but IgG remained undetectable at all detected time points (figure 1A).

6 During the follow-up period, the cell count of total lymphocytes, CD4⁺ T cells, CD8⁺ T cells
7 as well as the ratio of CD4⁺/CD8⁺ T cells maintained at a normal level (figure S1B). By analyzing
8 the differentiation of CD4⁺ and CD8⁺ T cells, a significant and gradual increase of effector T cells
9 (CD45RO⁺CD27⁻, Teff) in both CD4⁺ and CD8⁺ T cells were observed in the follow up (figure 1B).
10 The frequencies of naïve T cells (CD45RO⁻CD27⁺, Tna), central memory T cells
11 (CD45RO⁺CD27⁺, Tcm) in either CD4⁺ or CD8⁺ T cells were not significantly changed, while the
12 effector memory subset (CD45RO⁺CD27⁻, Tem) were slightly increased (figure 1B, S2A, S3A).
13 These results matched the classical kinetics of T cell differentiation induced by viral infection. In
14 consistent, proliferation of T cells (assessed by the frequency of transcription factor Ki-67⁺), was
15 highly upregulated at day 7 in both CD4⁺ and CD8⁺ T cells. The elevated percentage of Ki-67⁺
16 cells lasted until day 22 in CD4⁺ T cells, while rapidly decreased to baseline expression at day 13
17 in CD8⁺ T cells (figure 1C). The frequencies of Ki67⁺ cells in each differentiated subsets (Teff,
18 Tna, Tcm, Tem) of CD4⁺ or CD8⁺ T cells showed the similar trend as the corresponding parent
19 populations (figure 1C, the left panel of S2 and S3).

20 The activation of CD4⁺ and CD8⁺ T cells, determined as the percentage of PD-1⁺, HLA-DR⁺
21 or CD38⁺HLA-DR⁺, were remarkably lower than the normal low limit at day 7, peaked at day 13,
22 and gradually decreased to a low level that was comparable as the beginning (figure 1D, 1E, the
23 middle and right panel of S2 & S3, S4). These results indicate that the activation of T cells in the
24 peripheral blood might be transiently suppressed after the initial infection, and then activated from
25 the reduced baseline. This might be due to the virus induced T cell suppression, or the
26 re-distribution of activated T cells towards the infected tissues, or unknown else.

27 To analyze the function of T cells, the PBMCs were stimulated by the polyclonal stimulator
28 PMA and Ionomycin for 4.5 hours *ex vivo*, and proceed by intracellular cytokine staining of IFN- γ ,
29 IL-4 and IL-17. Notable low frequencies of IFN- γ producing CD4⁺ and CD8⁺ T cells were
30 detected at day 7, compared to the corresponding cell subsets of the healthy controls (figure 2,

1 IFN- γ panel). Although the percentages of IFN- γ producing CD4⁺ and CD8⁺ T cells gradually
 2 increased from day 7 to day 28, that of CD4⁺ T cells (figure 2A) reached the low limit of healthy
 3 controls at day 13, while that of CD8⁺ T cells (figure 2B) was even persistently lower than the
 4 normal low limit throughout the whole following up period. These results imply a functional
 5 suppression of type 1 immune responses at the beginning of infection and only a partial recovery
 6 afterwards. IL-4 producing CD4⁺ T cells gradually increased from day 7 to day 22 but dropped to
 7 the baseline at day 28, indicating an activation of Th2 type immune response after viral infection
 8 (figure 2A, IL-4 panel). In contrast, frequency of IL-17 producing CD4⁺ T cells (figure 2A, IL-17
 9 panel) maintained at low level remarkably below the healthy controls, without significant increase
 10 of frequency in all the four tested time points, suggesting that viral infection persistently
 11 suppressed the Th17 type immune response. The IL-4 or IL-17 producing CD8⁺ cells were very
 12 rare and with no significant change pattern (figure 2B). These results hint that the Th1 and Th17
 13 type responses were suppressed by the viral infection at the early stage, and the function of Th1
 14 cell-mediated cellular immune responses were partially recovered after the viral clearance, while
 15 the Th17 cells were persistently inhibited.

16 Regulatory T cells (Treg) were further analyzed to address their effect on the conventional T
 17 cells. The Treg population of this case, characterized by CD3⁺CD8⁻CD4⁺CD127⁻CD25⁺, was
 18 gradually increased since day 7, reached the peak at a much higher level than the healthy controls
 19 at day 22, and then slightly decreased at day 28 (figure 3A). According to the expression of FoxP3
 20 and CD45RA, Treg cells were further divided into three subsets: resting Treg cells (rTreg,
 21 CD45RA⁺FoxP3^{lo}), activated Treg cells (aTreg, CD45RA⁺FoxP3^{hi}) and cytokine-secreting
 22 non-suppressive T cells (nonTreg, CD45RA⁺FoxP3^{lo})¹⁴. Taking the notions that the aTreg has the
 23 strongest inhibitory function, the rTreg presents much weaker inhibitory activity compared to
 24 aTreg but can be converted to aTreg¹⁴, we thus focused on the suppressive aTreg and rTreg subsets.
 25 Compared to the rTreg which maintained at a sustained level as the healthy controls, the
 26 composition of aTregs was at a high level at day 7, as much as 4-4-fold of that in healthy controls,
 27 then gradually decreased to normal range at day 22 (figure 3B). As CTLA-4 is one of the most
 28 important co-inhibitory molecules expressed by Treg, we further determined CTLA-4 expression
 29 on the total Treg and the two Treg subsets. Consistent with the previous publications, the aTregs
 30 possessed the highest CTLA-4 expression. Notably, the frequency of CTLA-4⁺ cells in aTregs was

over 4-fold of that in total Tregs or rTregs. The dynamics of CTLA-4⁺ in total Tregs and the two subsets were similar in which the CTLA-4⁺ population peaked at day 13 and then dropped but without significant change compared with healthy controls (figure 3C).

Discussion

Currently, the emerging infectious disease COVID-19 caused by SARS-CoV-2 infection is becoming a worldwide threat. The suspected high rate of asymptomatic infections in population and a significant number of asymptomatic transmissions arouse more concerns on this novel pathogen and disease in potential future spread^{3, 4, 7, 8}. Here, we report a prospective study on the T cell responses of an imported SARS-CoV-2 infection case in Wuhan, who was unexpectedly found being an asymptomatic infection case after a 28-day follow-up study during hospitalization and isolation. This longitudinal characteristics of T cell response implied a viral-induced long-term effect on the T cell activation even in the asymptomatic SARS-CoV-2 infected individuals.

T cell response is demonstrated to play vital role in clearing the viruses and infected cells in series of viral infectious diseases. On the other hand, dysfunction of CD4⁺ and CD8⁺ T cells usually happens after viral infection and results in delayed viral clearance, viral persistence or even severe infectious tissue damage, such as substantial studies reported in chronic HBV and HIV infected patients^{15, 16}. Different from the significant decrease of T lymphocytes reported in the symptomatic COVID-19 cases^{2, 17}, the T cell count did not decrease in this asymptomatic individual. The CD4⁺ and CD8⁺ T cells experienced classical kinetics of T cell differentiation induced by viral infection and the proliferation of T cells especially Teff was even very active at the early stage of infection as we detected. However, the activation of T cells was somehow kept at low level (figure 1D, E, S2 and S3). This phenomenon is quite different from that activation usually accompanied with the proliferation during acute phases of viral infection¹⁸, suggesting that the T cell activation was suppressed in this asymptomatic case though the proliferation of the T cells might be stimulated initially. Accompanied with the suppressed activation, the T cell functions might be impaired. We noted that the CD4⁺ T cell was suppressed in activation, but the accompanied function impairment only detected in the Th1 and Th17 subsets, but not in the Th2, suggesting a selective suppression on CD4 T cells (figure 2A). We observed that the CD8⁺ T cell

1 was also suppressed in activation especially at early infection stage, and the accompanied function
2 impairment was significantly presented in the IFN- γ ⁺CD8⁺ T cell subset. Furthermore, the
3 suppression of CD8⁺ T cell was even lasted until day 28, in which the IFN- γ expression was still
4 kept at much lower level compared to the healthy controls (figure 2B), suggesting a persistent
5 function impairment on CD8 T cells. As we tested the function of T cells ex-vivo under the
6 polyclonal (non-specific) stimulation, the observed functional impairment is not limited to the
7 viral-specific T cells, but to the total reservoirs of CD4⁺ and CD8⁺ cells. Thus, the function
8 impairment indicated by the decreased cytokine productions might also suggest a featured T cell
9 response bias in this asymptomatic individual. It should be noted that the persistent suppression of
10 Th17 cells, which usually was involved in the pulmonary inflammation, might be one key aspect
11 for the explanation of this asymptomatic case who did not progress into radiographically
12 detectable pulmonary inflammatory lesion. More intensive studies on the T cell response in the
13 asymptomatic case should be conducted to give more insights into the protective immune
14 responses to the SARS-CoV-2 infection.

15 In contrast to our study, several studies have reported results of T cell activation and function
16 in the different stages or severities during SARS-CoV-2 infection. Different from this
17 asymptomatic case, patients with mild-to-moderate COVID-19 disease showed remarkable
18 activation of the CD4⁺ and CD8⁺ T cells before the resolution of symptoms⁹, while the patients
19 with severe COVID-19 disease showed a significant decrease of T lymphocytes and impairment of
20 T cell function^{2, 10}. These results pointed out the complicated interactions between viral infection,
21 T cell responses and clinical outcomes, which requires more intensive studies to clarify the role of
22 T cell responses in the local immune system of the lung and in the SARS-CoV-2 infection induced
23 pneumonia.

24 For understanding the mechanism of the suppressed T cell activation and function of the
25 asymptomatic case, we focused on the regulatory T (Treg) cells, which suppress immune
26 responses to a broad range of antigens, and limit immune response by employing multiple
27 mechanisms¹⁹. According to the expression of FoxP3 and CD45RA, Treg cells are classified into
28 two suppressive subsets: a slight inhibitory CD45RA⁺FoxP3^{lo} rTreg and a strongest inhibitory
29 CD45RA-FoxP3^{hi} aTreg¹⁴. In exerting function of Tregs, CTLA-4 on aTregs can capture its
30 ligands CD80 and CD86 from the surface of antigen presenting cells (APCs), denying their

availability for co-stimulation of CD4⁺ T and CD8⁺ T cells²⁰. In this asymptomatic case, aTreg which possesses the highest inhibitory molecule CTLA-4, was markedly elevated at the early infection stage. The elevation of aTregs accompanying with the low activation and impaired function of T cells at early infection stage of this asymptomatic case suggested that aTregs might suppress T cell activation and the following function in the T cell priming stage. However, it was reported that the frequency of Tregs was significantly reduced in the severe and moderate cases^{2, 10}. The current knowledge of the kinetics of Treg activation in either the asymptomatic or the symptomatic patients is still limited. More detailed investigations on the functional regulation of Tregs in the SARS-CoV-2 infection are urgently needed.

The present study has some limitations. First, there was only one case in this longitudinal prospective study. The characteristics of T cell response should be confirmed further in larger cohorts of people with asymptomatic SARS-CoV-2 infection. Second, T cell functions were analyzed using polyclonal stimulator PMA and Ionomycin that indicated the potential function of T cell reservoir. The quantity and quality of viral-specific T cells that directly related to the viral control have to be further studied in the patients with different disease severities. Hopefully, we may have some more answers in our following cohort studies.

In conclusion, the SARS-CoV-2 asymptomatic infection induced low activation and impaired function of CD4⁺ and CD8⁺ T cells exemplified by suppressed IFN- γ production in both CD4⁺ T and CD8⁺ T cells and inhibited IL-17 production in CD4⁺ T cells. In addition, the elevation of aTreg at early infection stage suggested an aberrant Treg activation and a unique immune pathology of the SARS-CoV-2 virus. Our study shed some light on early interaction between the SARS-CoV-2 infection and host immune responses, which might give us more insights into the preventive and curative immune strategy.

24

Contributors

JY, EZ, MZ contributed to the conception, design, data acquisition, analysis, and interpretation, and drafted and critically revised the manuscript. QY, TS, DihZ, JieX, JiaX contributed to the acquisition of data. KH provided clinical care to the patient and assisted with clinical descriptions. XZ, DinZ, CH, YS made contribution to the study concept and design. HY contributed to the conception, design, data analysis, and interpretation, and drafted and critically revised the manuscript. All of the authors gave final approval and agreed to be accountable for all

1 aspects of the work.

2

3 **Declaration of interests**

4 All authors declare no competing interests.

5

References

1. Cao X. COVID-19: immunopathology and its implications for therapy. *Nature reviews Immunology*. 2020; **20**(5): 269-70.
2. Chen G, Wu D, Guo W, Cao Y, Huang D, Wang H, et al. Clinical and immunological features of severe and moderate coronavirus disease 2019. *The Journal of clinical investigation*. 2020; **130**(5): 2620-9.
3. Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Euro surveillance : bulletin Europeen sur les maladies transmissibles = European communicable disease bulletin*. 2020; **25**(10).
4. Black JRM, Bailey C, Przewrocka J, Dijkstra KK, Swanton C. COVID-19: the case for health-care worker screening to prevent hospital transmission. *Lancet*. 2020; **395**(10234): 1418-20.
5. Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected Patients. *The New England journal of medicine*. 2020; **382**(12): 1177-9.
6. Gandhi M, Yokoe DS, Havlir DV. Asymptomatic Transmission, the Achilles' Heel of Current Strategies to Control Covid-19. *The New England journal of medicine*. 2020.
7. Bai Y, Yao L, Wei T, Tian F, Jin DY, Chen L, et al. Presumed Asymptomatic Carrier Transmission of COVID-19. *Jama*. 2020.
8. Furukawa NW, Brooks JT, Sobel J. Evidence Supporting Transmission of Severe Acute Respiratory Syndrome Coronavirus 2 While Presymptomatic or Asymptomatic. *Emerging infectious diseases*. 2020; **26**(7).
9. Thevarajan I, Nguyen THO, Koutsakos M, Druce J, Caly L, van de Sandt CE, et al. Breadth of concomitant immune responses prior to patient recovery: a case report of non-severe COVID-19. *Nature medicine*. 2020; **26**(4): 453-5.
10. Qin C, Zhou L, Hu Z, Zhang S, Yang S, Tao Y, et al. Dysregulation of immune response in patients with COVID-19 in Wuhan, China. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2020.
11. Zheng M, Gao Y, Wang G, Song G, Liu S, Sun D, et al. Functional exhaustion of antiviral lymphocytes in COVID-19 patients. *Cellular & molecular immunology*. 2020; **17**(5): 533-5.

12. Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, et al. A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-19. *The New England journal of medicine*. 2020.
13. Giamarellos-Bourboulis EJ, Netea MG, Rovina N, Akinosoglou K, Antoniadou A, Antonakos N, et al. Complex Immune Dysregulation in COVID-19 Patients with Severe Respiratory Failure. *Cell host & microbe*. 2020.
14. Miyara M, Yoshioka Y, Kitoh A, Shima T, Wing K, Niwa A, et al. Functional delineation and differentiation dynamics of human CD4⁺ T cells expressing the FoxP3 transcription factor. *Immunity*. 2009; **30**(6): 899-911.
15. Yang F, Yu X, Zhou C, Mao R, Zhu M, Zhu H, et al. Hepatitis B e antigen induces the expansion of monocytic myeloid-derived suppressor cells to dampen T-cell function in chronic hepatitis B virus infection. *PLoS pathogens*. 2019; **15**(4): e1007690.
16. Morou A, Palmer BE, Kaufmann DE. Distinctive features of CD4⁺ T cell dysfunction in chronic viral infections. *Current opinion in HIV and AIDS*. 2014; **9**(5): 446-51.
17. Liu J, Li S, Liang B, Wang X, Wang H, Li W, et al. Longitudinal characteristics of lymphocyte responses and cytokine profiles in the peripheral blood of SARS-CoV-2 infected patients. *EBioMedicine*. 2020; **55**: 102763.
18. Arunkumar G, Devadiga S, McElroy AK, Prabhu S, Sheik S, Abdulmajeed J, et al. Adaptive Immune Responses in Humans During Nipah Virus Acute and Convalescent Phases of Infection. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2019; **69**(10): 1752-6.
19. Arpaia N, Green JA, Molledo B, Arvey A, Hemmers S, Yuan S, et al. A Distinct Function of Regulatory T Cells in Tissue Protection. *Cell*. 2015; **162**(5): 1078-89.
20. Maeda Y, Nishikawa H, Sugiyama D, Ha D, Hamaguchi M, Saito T, et al. Detection of self-reactive CD8(+) T cells with an anergic phenotype in healthy individuals. *Science*. 2014; **346**(6216): 1536-40.

1 **Figure legends**

2

3 **Figure 1 Differentiation, proliferation and activation of T cells in peripheral blood of the**

4 **asymptomatic patient during SARS-CoV-2 infection.** A, Timeline of SARS-CoV-2 infection.

5 SARS-CoV2 was detected by qRT-PCR in nasopharyngeal swab, sputum, and by RBD-specific

6 IgM, IgG using colloidal gold strips. B, Frequencies of naïve or effector T cells (Teff) in the CD4⁺

7 or CD8⁺ T cells. C-E, Frequencies of Ki-67⁺, PD-1⁺ or HLA-DR⁺ cells in CD4⁺ or CD8⁺ T cells of

8 the patient at indicated time points and in that of healthy donors (n=11). Gating strategies of the

9 naïve (CD45RO⁻CD27⁺), central memory (Tcm, CD45RO⁺CD27⁺), effector memory (Tem,

10 CD45RO⁺CD27⁻) and effector (Teff, CD45RO⁻CD27⁻) subsets, Ki-67⁺, PD-1⁺ or HLA-DR⁺ cells

11 in CD4⁺ T cells and CD8⁺ T cells were shown as left panels.

12

13

14 **Figure 2. Cytokine production of T cells in peripheral lymphocytes after *ex vivo* stimulation.**

15 PBMCs were isolated from the patient at days 7, 13, 22 and 28 post hospitalization and in healthy

16 donors (n=11). PBMCs were stimulated by PMA/Ionomycin for 4.5 h in the presence of BFA and

17 monensin. The production of IFN- γ , IL-4, and IL-17 by CD4⁺ (A) or CD8⁺ T cells (B) were

18 analyzed by intracellular cytokine staining and flow cytometry. IFN- γ , IL-4, and IL-17 producing

19 CD4⁺ and CD8⁺ T cells were gated based on the unstimulated cells as shown in the left panels.

20

21

22 **Figure 3. Frequencies and function of Tregs and Treg subsets in peripheral lymphocytes.**

23 PBMCs were isolated from the patient at days 7, 13, 22 and 28 post hospitalization or healthy

24 donors (n=11). A, Gating strategy and the frequency of CD25⁺CD127⁺ Tregs in CD3⁺CD8⁻CD4⁺T

25 cells. B, Gating strategies and the frequencies of resting Tregs (rTregs, CD45RA⁺Foxp3^{lo}) and

26 activated Tregs (aTregs, CD45RA⁺Foxp3^{hi}) in Tregs. C, Gating strategies and the frequencies of

27 CTLA-4⁺ cells in Tregs, rTregs and aTregs.

Figure 1

A

B

C

D

E

Figure 2

Figure 3