Case fatality risk by age from COVID-19 in a high testing setting in Latin America: Chile, March-May, 2020

Running title: COVID19 case fatality risk by age, Chile

Authors:

Eduardo A. Undurraga^{1,2*}, Gerardo Chowell³, Kenji Mizumoto^{4,5}

Affiliations:

¹ Escuela de Gobierno, Pontificia Universidad Católica de Chile, Santiago, Región Metropolitana, Chile

² Millennium Initiative for Collaborative Research in Bacterial Resistance (MICROB-R), Chile

³ Department of Population Health Sciences, School of Public Health, Georgia State University, Atlanta, Georgia, USA

⁴ Graduate School of Advanced Integrated Studies in Human Survivability, Kyoto University Yoshida-Nakaadachi-cho, Sakyo-ku, Kyoto, Japan

⁵ Hakubi Center for Advanced Research, Kyoto University, Yoshidahonmachi, Sakyo-ku, Kyoto, Japan;

Word count:

Abstract: 200/200

Main: 2173/3500

References: 50/50

Tables and Figures: 6/7 (max 4 panels per figure)

*Address for correspondence: Eduardo A. Undurraga, Escuela de Gobierno, Pontificia Universidad Católica de Chile. Vicuña Mackenna 4860, Macul CP 7820436, Región Metropolitana, Chile; email: eundurra@uc.cl; phone: +56 (2) 2354 9536. ORCID: 0000-0002-4425-1253

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Abstract

Background

Early severity estimates of COVID-19 are critically needed to better assess the potential impact of the ongoing pandemic in different socio-demographic groups. Using real-time epidemiological data from Chile, the nation in Latin America with the highest testing rate for COVID-19, we derive delay-adjusted severity estimates by age group as of May 18th, 2020.

Methods

We employed statistical methods and daily series of age-stratified COVID-19 cases and deaths reported in Chile to estimate the delay-adjusted case fatality rate across six age groups.

Results

Our most recent estimates of the time-delay adjusted case fatality rate are 0.08% (95% Credible Interval CrI:0.04-0.13%) among persons aged 0-39, 0.61% (95%CrI:0.41-0.87%) for those aged 40-49, 1.06% (95%CrI:0.76-1.40%) for those aged 50-59, 3.79% (95%CrI:3.04-4.66%) for those aged 60-69, 12.22% (95%CrI:10.40-14.38%) for those aged 70-79, and 26.27% (95%CrI:22.95-29.80%) for persons aged 80 and over. The overall time-delay adjusted case fatality rate is 1.78% (95%CrI: 1.63-1.95%) across all age groups.

Conclusions

Severity estimates from COVID-19 in Chile indicate a disproportionate impact among seniors, especially among those aged ≥ 70 years. COVID-19 is imposing a high death toll in Latin America. Case fatality rates in Chile suggest the health system is not yet overwhelmed, but the epidemic is expanding fast.

Keywords: COVID-19, Chile, death risk by age group, time-delay adjusted CFR, 2020

Background

The coronavirus disease 2019 (CoVID-19) pandemic has strained or overwhelmed health systems across the world [1, 2], with almost five million lab-confirmed cases and 320 thousand deaths as of May 18, 2020 [3, 4]. The first case in Latin America of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), the cause of CoVID-19, was reported in February 25 in São Paulo, Brazil, a travel hub in the region [5]. A few weeks later, countries in the region had imposed major epidemic control measures, such as closed borders, restricted travel, closure of schools and universities, and enforced lockdowns [6]. Despite these measures, the ongoing coronavirus pandemic has already imposed a high toll to most countries in Latin America, killing thousands in Brazil (21,408), Ecuador (3,056), and Peru (3,244) thus far [3]. In addition to already strained healthcare systems, other factors have affected the dynamic of the pandemic in the region, including migration, sociopolitical crises, struggling economies, other infectious disease outbreaks, and challenges tied to the implementation of social distancing, hygiene, and lockdown strategies, due to inadequate water and sanitation infrastructure, precarious living conditions [6-11].

Chile has seen a rapid increase in CoVID-19 cases in recent weeks, with 46,048 cases and 544 deaths [12] as of May 18, 2020. The first CoVID-19 case in Chile was reported in March 3, 2020 [12]. The Ministry of Health put in place a quick response effort, announcing restrictions on large gatherings in March 13th, and subsequently, closing all daycares, schools, and universities (March 16th), imposing border controls and telework recommendations (March 18th), closing non-essential businesses (March 19th), national night curfews (March 22nd), and implementing a strategy of intermittent lockdowns in selected municipalities since March 22nd [13-15]. Estimates for the early stage of the epidemic showed sustained transmission with an estimated reproduction number $R \sim 1.6$ (95% CI: 1.5, 1.7), and 20-day

ahead forecast suggested containment measures significantly slowed down the spread of the virus [16]. The crude case fatality rate (CFR), the number of cumulative deaths over the number of cumulative cases, in Chile (1.2%) has remained well below the global average (6.5%) [4, 12]. But in light of limited health system capacity, infectious disease experts have warned about authorities' excessive confidence over early successes [17]. In fact, models have estimated the number of ill patients could overwhelm treatment capacity late May to mid-June [18], which has not yet occurred.

SARS-CoV-2 infection can result in a wide spectrum of clinical outcomes, including asymptomatic infection, mild symptoms, hospitalization, or death [19-21]. The case fatality rate (CFR) is a commonly used estimate of the severity of an epidemic [22], as it provides a reliable benchmark for public health officials to make decisions about intensity and duration of interventions to mitigate or suppress an epidemic [23]. But obtaining CFR estimates during the course of an epidemic is challenging, as CFR is typically affected by right censoring and ascertainment bias [24-28]. Right-censoring occurs because of a time delay between onset of symptoms and death and may lead to underestimation of CFR; under ascertainment of cases occurs because mild or asymptomatic COVID-19 cases often go undetected by disease surveillance systems, which are not designed to detect all infections [25, 29-31].

Here we provide real-time estimates of adjusted age-specific CFR during the CoVID-19 epidemic in Chile, March through May 2020, to gauge the severity of the SARS-CoV-2 epidemic. In particular, Chile has tested at a higher rate (23.13 total tests per 1000 people by May 22) than any other country in Latin America [32, 33]. The health system has not yet reached its maximum capacity, with 85% and 96% of critical beds in use nationally and in the

capital Santiago [34], respectively, so our estimates are still not affected by excess deaths due to healthcare demand exceeding capacity. To our knowledge, these are the first estimates of CFR in Latin America, which could inform critical decisions by public health officials.

Methods

Data Sources

We obtained daily cumulative numbers of reported laboratory confirmed COVID-19 cases and deaths stratified by age group from March 3, 2020, through May 18, 2020, from the Ministry of Health [12]. Data on deaths by age group were missing for a few days (March 29 and 31, April 1-3, 6, 8).

Statistical analysis

For the estimation of CFR in real time, we employed the delay from hospitalization to death, h_s , which is assumed to be given by $h_s = H(s) - H(s-1)$ for $s > 0$ where $H(s)$ is a cumulative density function of the delay from hospitalization to death and follows a gamma distribution with mean 10.1 days and standard deviation (SD) 5.4 days [35]. Let π_{a,t_i} be the time-delay adjusted case fatality ratio on reported day t_i in area a , the likelihood function of the estimate π_{a,t_i} is

$$L(\pi_{a,t_i}; c_{a,t}, D_{a,t})$$

$$= \prod_{t_i} \left(\sum_{t=1}^{t_i} c_{a,t} \right) \left(\pi_{a,t_i} \frac{\sum_{t=2}^{t_i} \sum_{s=1}^{t-1} c_{a,t-s} h_s}{\sum_{t=1}^{t_i} c_{a,t}} \right)^{D_{a,t_i}} \left(1 - \pi_{a,t_i} \frac{\sum_{t=2}^{t_i} \sum_{s=1}^{t-1} c_{a,t-s} h_s}{\sum_{t=1}^{t_i} c_{a,t}} \right)^{\sum_{t=1}^{t_i} c_{a,t} - D_{a,t_i}}$$

where $c_{a,t}$ represents the number of new cases with reported day t in area a , and $D_{a,ti}$ is the cumulative number of deaths until reported day t_i in area a [36, 37]. Among the cumulative cases with reported day t in area a , $D_{a,ti}$ have died and the remainder have survived the infection. The contribution of those who have died with biased death risk is shown in the middle parenthetical term and the contribution of survivors is presented in the right parenthetical term. We assume that $D_{a,ti}$ is the result of the binomial sampling process with probability $\pi_{a,ti}$.

We estimated model parameters using a Monte Carlo Markov Chain (MCMC) method in a Bayesian framework. Convergence of MCMC chains were evaluated using the potential scale reduction statistic [38, 39]. Estimates and 95% credibility intervals (CrI) for these estimates are based on the posterior probability distribution of each parameter and based on the samples drawn from the posterior distributions. All statistical analyses were conducted in R version 3.6.1 (R Foundation for Statistical Computing, Vienna, Austria) using the ‘rstan’ package.

Results

As of May 18, a total of 46,048 COVID-19 cases and 544 deaths have been reported in Chile. Reported cases were mostly observed among persons aged 0-39 years (53.9%), followed by 40-49 year olds (17.3%), and 50-59 year olds (14.4%) (Table 1). Most reported deaths correspond to seniors, especially those aged 80 years and older (38.1%), followed 70-79 year olds (27.9%), and 60-69 year olds (16.9%) (Table 1).

Figure 1A presents the gender proportion of reported cases by age groups and cumulative morbidity ratio by gender and age group. Across age groups, the proportion of male cases is

higher than 50%, except for those aged 80 years and above (χ^2 test, p-value <0.0001).

Cumulative morbidity ratio by gender and age group is presented in Figure 1B, indicating that ratio among males is higher than among females across age groups (proportion test, p-value < 0.000) except for those aged 80 years and above (proportion test, p-value = 0.13).

Figure 2 displays the cumulative cases of CoVID-19 by age group (A through G), and the cumulative deaths by age group (H through N) over time. The figure suggests cumulative cases of CoVID-19 are growing faster than cumulative deaths. The growth curve for cumulative cases across all age groups and for persons aged 0-39 appears to increase exponentially after around day 30 (April 30th, 2020), while exponential growth in cumulative deaths for all age groups appears to occur after day 45 (May 15th, 2020).

Figure 3 shows observed and model-based posterior estimates of the crude CFR of CoVID-19 by age group (A-G) and time-delay adjusted CFR by age group (H-N). Black dots show crude case fatality ratios, and light and dark indicate 95% and 50% credible intervals for posterior estimates, respectively.

Overall, our model-based crude CFR fitted the observed data well. Crude CFR for all age groups (G) increased at an early stage of the epidemic, peaking around April 25th and was followed by a decrease in CFR for about two weeks. There is a suggestive increasing trend across most age groups starting around May 10th, 2020. Our model-based posterior estimates for the time-delay adjusted CFR are substantially higher than the crude observed CFR. The overall adjusted CFR follows a decreasing trend, except for infected patients aged 50-59 years, where the adjusted CFR increases until around April 21st.

The most recent estimates (May 18, 2020) of the time-delay adjusted CFR are 0.08% (95% CrI: 0.04-0.13%) for persons aged 0-39, 0.61% (95%CrI: 0.41-0.87%) for those aged 40-49, 1.06% (95%CrI: 0.76-1.40%) for those aged 50-59, 3.79% (95%CrI: 3.04-4.66%) for those aged 60-69, 12.22% (95%CrI: 10.40-14.38%) for those aged 70-79, and 26.27% (95%CrI: 22.95-29.80%) for persons aged 80 and over. The overall time-delay adjusted CFR is 1.78% (95%CrI: 1.63-1.95%) across all age groups (Table 2, Figure 4).

Discussion

To the best of our knowledge, this is the first study to estimate the time delay adjusted CFR by age group for COVID-19 in Latin America, a region that has yet received little attention during the ongoing coronavirus pandemic. Consistent with other recent COVID-19 research [23, 25, 40], our results show the COVID-19 epidemic in Chile disproportionately influenced seniors, especially those aged ≥ 70 years. These results suggest that an aging population could exacerbate the fatality impact of COVID-19 [41], similar to influenza and respiratory syncytial virus [42], and consistent with data available from Italy[41]. The comparatively low CFR observed in Chile during the early stages of the epidemic [17], probably reflected the age structure and socioeconomic status of initial cases. During the first weeks of the pandemic, COVID-19 cases occurred among relatively young age groups, with most transmission occurring among individuals between 20 and 60 years of age, and in high-income communities with better access to healthcare and lower prevalence of risk factors for severe COVID-19 [12, 43, 44].

Our latest estimates (as of May 18th, 2020) of the adjusted CFR among those aged 80 and over reach values as high as 26.3% (95%CrI: 23.0-29.8), an estimate that is 328-fold higher

than our estimates for those aged 0-39 (0.08%), and 2.2-fold higher than our estimates for those aged 70-79 (12.2%).

An upward trend in the crude CFR, as seen for the overall population through most of April and particularly for the 50-59 age group, suggests the transmission is spreading to more vulnerable populations. Interestingly, throughout March, most of transmission occurred among relatively young, better-off populations. In Santiago, which has more than 75% of COVID-19 cases in Chile, targeted enforced quarantines were put in place starting March 28th through mid-April in seven municipalities, six of which are among the richest in Chile [45]. Epidemic transmission thus moved during April towards lower income municipalities, where social distancing measures are more challenging to comply with, due to a higher proportion of the population participating in the informal economy, higher population density, poorer infrastructure, and lower quality healthcare [12, 46]. An upward trend in the crude CFR could also result from an increasing number of unreported cases due to saturated testing capacity. However, this is an unlikely explanation as RT-PCR testing capacity has increased with cases, maintaining an average positivity rate (positive tests / total tests) of 9.7% throughout April (SD: 2.4%). However, results show an increase in crude CFR around day 45 (May 15th, 2020), probably reflecting the exponential increase of cumulative cases around day 30 (April 30, 2020), and a substantial increase in positive test rate (average in May 1-18: 15.2% SD:4.5), which suggests there is probably an increasing proportion of under diagnosed COVID-19 cases in Chile [47].

The downward trend in the adjusted CFR at the early stage of the study period may have been influenced by reporting delays. In particular, the observed differences in our estimates of the

crude and adjusted CFR are directly due to the time-delay which we assume fixed during the course of the epidemic.

The small proportion of men (38.8%) among CoVID-19 cases in people aged 80 and over is probably attributable to the relatively small male population size for that age group; men represent only 37% of the population >80 in Chile [48]. Higher mortality among men has been reported in China and the U.S. [49, 50], our data provide the opportunity to examine the CFR by gender and age.

Our study is not exempted from limitations. Our CFR estimates are probably affected by under ascertainment, as has been estimated elsewhere [24-28], which may have pushed our estimates upwards. Infectious disease with a substantial share of asymptomatic or mild infections, such as COVID-19, may be more accurately characterized by infection-fatality risk (deaths / infected people), but those data are not yet available in Chile.

In conclusion, using real-time epidemiological data from a high COVID-19 testing setting in Latin America, we found that COVID-19 epidemic in Chile disproportionately affected seniors, especially those aged ≥ 70 years, suggesting an older population could exacerbate the death toll brought by COVID-19. COVID-19 is already imposing a high death toll in Latin America. Case fatality rates in Chile suggest the health system is not yet overwhelmed, but the epidemic is expanding fast, and healthcare demand may soon exceed capacity. These real-time estimates may help inform decisions by public health officials in the region, and underscore the need for continuous efforts to face this pandemic.

Acknowledgments

This work was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI “[grant Number 20H03940]”; the Leading Initiative for Excellent Young Researchers from the Ministry of Education, Culture, Sport, Science & Technology of Japan to [KM]; National Science Foundation NSF “[grant 1414374]” as part of the joint NSF-National Institutes of Health NIH-United States Department of Agriculture USDA Ecology and Evolution of Infectious Diseases program; UK Biotechnology and Biological Sciences Research Council “[grant BB/M008894/1]” to [GC]; and the Millennium Science Initiative of the Ministry of Economy, Development, and Tourism, Government of Chile, “[grant Millennium Initiative for Collaborative Research in Bacterial Resistance]”.

Conflict of interest

All authors report no conflicts of interest.

References

1. Walker PG, Whittaker C, Watson O, et al. The Global Impact of COVID-19 and Strategies for Mitigation and Suppression. On behalf of the imperial college covid-19 response team, Imperial College of London **2020**.
2. Flaxman S, Mishra S, Gandy A. Estimating the number of infections and the impact of non-pharmaceutical interventions on COVID-19 in 11 European countries. Imperial College COVID-19 Response Team **2020**; 30.
3. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis* **2020**; 20:P533-P4.

- 274 4. World Health Organization. Coronavirus Disease (COVID-19) Outbreak. Available from:
275 <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>. Accessed 20 May 2020.
- 276 5. Rodriguez-Morales AJ, Gallego V, Escalera-Antezana JP, et al. COVID-19 in Latin
277 America: The implications of the first confirmed case in Brazil. *Travel Med Infect Dis* **2020**;
278 epub ahead of print:101613.
- 279 6. Burki T. COVID-19 in Latin America. *Lancet Infect Dis* **2020**; 20:547-8.
- 280 7. The Lancet. The unfolding migrant crisis in Latin America. *Lancet* **2019**; 394:1966.
- 281 8. Fetzer TR, Witte M, Hensel L, et al. Global Behaviors and Perceptions at the Onset of the
282 COVID-19 Pandemic. National Bureau of Economic Research Working Paper Series **2020**;
283 No. 27082.
- 284 9. Navarro J-C, Arrivillaga-Henríquez J, Salazar-Loor J, Rodriguez-Morales AJ. COVID-19
285 and dengue, co-epidemics in Ecuador and other countries in Latin America: Pushing strained
286 health care systems over the edge. *Travel Med Infect Dis* **2020**.
- 287 10. Kirby T. South America prepares for the impact of COVID-19. *The Lancet Respiratory*
288 *Medicine* **2020**; online first.
- 289 11. World Bank. The Economy in the Time of CoVID-19. Available from:
290 <https://bit.ly/35B2r3y>. Accessed April 2020.
- 291 12. Ministry of Health. Cifras Oficiales COVID-19. Available from:
292 <https://www.gob.cl/coronavirus/cifrasoficiales/>. Accessed May 22 2020.

- 293 13. Ministry of Health. Dispone medidas sanitarias que indica por brote de COVID-19.
294 Norms 1143498, 1143591, 1746958, 1143651 Available from:
295 <https://www.leychile.cl/N?i=1143498&f=2020-04-04&p=>.
- 296 14. Ministry of Health. Dispone régimen especial de cumplimiento de jornada laboral y
297 flexibilidad horaria por brote de coronavirus (COVID-19). Norm 1143629. Available from:
298 <https://www.leychile.cl/N?i=1143629&f=2020-03-20&p=>.
- 299 15. Ministerio del Interior y Seguridad Pública. Declara estado de excepción constitucional
300 de catástrofe, por calamidad pública, en el territorio, norma 1143580 decreto108. Available
301 from: <https://www.leychile.cl/N?i=1143580&f=2020-03-26&p=>.
- 302 16. Tariq A, Undurraga EA, Laborde CC, et al. Early transmission dynamics of COVID-19 in
303 Chile: From sub-exponential ascending growth dynamics to a stationary disease wave,
304 March-April, 2020. medRxiv **2020**:2020.05.15.20103069.
- 305 17. McGowan C. Chile doctors fear complacency over Covid-19 after initial successes. The
306 Guardian. Santiago de Chile, **2020**.
- 307 18. Rainisch G, Undurraga EA, Chowell G. A dynamic modeling tool for estimating
308 healthcare demand from the COVID19 epidemic and evaluating population-wide
309 interventions. Int J Infect Dis **2020**.
- 310 19. Guan W-j, Ni Z-y, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in
311 China. N Engl J Med **2020**.
- 312 20. Cao J, Hu X, Cheng W, Yu L, Tu W-J, Liu Q. Clinical features and short-term outcomes
313 of 18 patients with corona virus disease 2019 in intensive care unit. Intensive Care Medicine
314 **2020**:1-3.

- 315 21. Wu JT, Leung K, Bushman M, et al. Estimating clinical severity of COVID-19 from the
316 transmission dynamics in Wuhan, China. *Nat Med* **2020**.
- 317 22. Reed C, Biggerstaff M, Finelli L, et al. Novel framework for assessing epidemiologic
318 effects of influenza epidemics and pandemics. *Emerg Infect Dis* **2013**; 19:85.
- 319 23. Mizumoto K, Chowell G. Estimating risk for death from 2019 novel coronavirus disease,
320 China, January-February 2020. *Emerg Infect Dis* **2020**; 26:10-3201.
- 321 24. Li R, Pei S, Chen B, et al. Substantial undocumented infection facilitates the rapid
322 dissemination of novel coronavirus (COVID-19). *Science* **2020**; 368:489-93.
- 323 25. Riou J, Hauser A, Counotte MJ, Althaus CL. Adjusted age-specific case fatality ratio
324 during the COVID-19 epidemic in Hubei, China, January and February 2020. *medRxiv* **2020**.
- 325 26. Ghani AC, Donnelly CA, Cox DR, et al. Methods for Estimating the Case Fatality Ratio
326 for a Novel, Emerging Infectious Disease. *Am J Epidemiol* **2005**; 162:479-86.
- 327 27. Jewell NP, Lei X, Ghani AC, et al. Nonn-parametric estimation of the case fatality ratio
328 with competing risks data: an application to Severe Acute Respiratory Syndrome (SARS).
329 *Statistics in medicine* **2007**; 26:1982-98.
- 330 28. Lipsitch M, Donnelly CA, Fraser C, et al. Potential biases in estimating absolute and
331 relative case-fatality risks during outbreaks. *PLoS Negl Trop Dis* **2015**; 9.
- 332 29. Undurraga EA, Halasa YA, Shepard DS. Use of expansion factors to estimate the burden
333 of dengue in Southeast Asia: A systematic analysis. *PLoS Negl Trop Dis* **2013**; 7:e2056.
- 334 30. Wang C, Liu L, Hao X, et al. Evolving Epidemiology and Impact of Non-pharmaceutical
335 Interventions on the Outbreak of Coronavirus Disease 2019 in Wuhan, China. *medRxiv* **2020**.

- 336 31. Jung S-m, Akhmetzhanov AR, Hayashi K, et al. Real-time estimation of the risk of death
337 from novel coronavirus (COVID-19) infection: inference using exported cases. Journal of
338 clinical medicine **2020**; 9:523.
- 339 32. Thomson E, Sanders P. Prognosis. Chile Charts New Path With Rolling Lockdowns,
340 Immunity Cards. Available from: [https://www.bloomberg.com/news/articles/2020-04-](https://www.bloomberg.com/news/articles/2020-04-22/with-immunity-cards-and-rolling-lockdowns-chile-forges-own-path)
341 [22/with-immunity-cards-and-rolling-lockdowns-chile-forges-own-path](https://www.bloomberg.com/news/articles/2020-04-22/with-immunity-cards-and-rolling-lockdowns-chile-forges-own-path). Accessed May 18
342 2020.
- 343 33. Hasell J, Ortiz-Ospina E, Mathieu E, Ritchie H, Roser M. To understand the global
344 pandemic, we need global testing – the Our World in Data COVID-19 Testing dataset.
345 Available from: <https://bit.ly/2ROHdts>. Accessed May 18 2020.
- 346 34. Sociedad Chilena de Medicina Intensiva SOCHIMI. Encuesta nacional sobre ocupación
347 de unidades críticas durante contingencia CoVID-19. Available from: [https://medicina-](https://medicina-intensiva.cl/site/covid/img/noticias/informe_22MY2020.pdf)
348 [intensiva.cl/site/covid/img/noticias/informe_22MY2020.pdf](https://medicina-intensiva.cl/site/covid/img/noticias/informe_22MY2020.pdf). Accessed May 22 2020.
- 349 35. Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic
350 proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess
351 cruise ship, Yokohama, Japan, 2020. Eurosurveillance **2020**; 25:2000180.
- 352 36. Nishiura H, Klinkenberg D, Roberts M, Heesterbeek JA. Early epidemiological
353 assessment of the virulence of emerging infectious diseases: a case study of an influenza
354 pandemic. PLoS One **2009**; 4.
- 355 37. Tsuzuki S, Lee H, Miura F, et al. Dynamics of the pneumonic plague epidemic in
356 Madagascar, August to October 2017. Eurosurveillance **2017**; 22.

357 38. Gamerman D, Lopes HF. Markov chain Monte Carlo: stochastic simulation for Bayesian
358 inference. CRC Press, **2006**.

359 39. Gelman A, Rubin DB. Inference from iterative simulation using multiple sequences. Stat
360 Sci **1992**; 7:457-72.

361 40. Feng Z, Li Q, Zhang Y, et al. The epidemiological characteristics of an outbreak of 2019
362 novel coronavirus diseases (COVID-19)-China, 2020. China CDC Weekly **2020**; 2:113-22.

363 41. Dowd JB, Andriano L, Brazel DM, et al. Demographic science aids in understanding the
364 spread and fatality rates of COVID-19. Proc Natl Acad Sci **2020**; 117:9696-8.

365 42. Mizumoto K, Chowell G, Simonsen L, Nishiura H. The burden of influenza and
366 respiratory syncytial viruses in Japan, 2006–2014: A region-and age-specific excess mortality
367 study. Int J Infect Dis **2019**; 79:96-7.

368 43. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients
369 with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet **2020**; 395:1054-62.

370 44. Wang B, Li R, Lu Z, Huang Y. Does comorbidity increase the risk of patients with
371 COVID-19: evidence from meta-analysis. Aging **2020**; 12:6049-57.

372 45. Sistema Nacional de Información Municipal. Datos Municipales. Available from:
373 http://datos.sinim.gov.cl/datos_municipales.php. Accessed May 19 2020.

374 46. Cociña M, Frei R, Larrañaga O. Desiguales: Orígenes, cambios y desafíos de la brecha
375 social en Chile. Programa de las Naciones Unidas para el Desarrollo (PNUD), **2017**.

47. Asahi K, Undurraga EA, Wagner R. Benchmarking the CoVID-19 pandemic across countries and states in the U.S.A. under heterogeneous testing. medRxiv **2020**:2020.05.01.20087882.
48. Instituto Nacional de Estadísticas. Estimaciones y proyecciones de la población de Chile 1992-2050. Available from: <https://www.censo2017.cl/>. Accessed April 2 2020.
49. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York City area. Jama **2020**.
50. Epidemiology Working Group for NCIP Epidemic Response. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. Zhonghua liu xing bing xue za zhi= Zhonghua liuxingbingxue zazhi **2020**; 41:145.

Tables

Table 1. Distribution of the CoVID-19 cases in Chile by sex and age groups, as of May 18, 2020

		Confirmed cases (%)	Deaths (%)	Crude fatality rate (%)
Total		46,048 (100.00)	544 (100.00)	1.18%
Sex	Male	24,523 (53.3)	NA	NA
	Female	21,525 (46.7)	NA	NA
Age-group	0-39	24,830 (53.9)	11 (2.0)	0.04
	40-49	7,953 (17.3)	31 (5.7)	0.39
	50-59	6,624 (14.4)	51 (9.4)	0.77
	60-69	3,748 (8.1)	92 (16.9)	2.45
	70-79	1,785 (3.9)	152 (27.9)	8.52
	80 and above	1,108 (2.4)	207 (38.1)	18.68

Table 2. Summary results of time-delay adjusted case fatality ratio of COVID-19 by age group in Chile, 2020 (May 18, 2020)

Age group	Latest estimate (95%CrI ^a)	Range of median estimates during the study period	Crude CFR (95% CI ^b)
0-39	0.08% (0.04-0.13%)	0.08-0.25%	0.00% (0.00-0.10%) 10/24830
40-49	0.61% (0.41-0.87%)	0.19-0.61%	0.40% (0.20-0.50%) 28/7953
50-59	1.06% (0.76-1.40%)	0.49-1.35%	0.60% (0.40-0.80%) 40/6624
60-69	3.79% (3.04-4.66%)	3.42-4.99%	2.20% (1.70-2.70%) 82/3748
70-79	12.22% (10.40-14.38%)	11.07-21.19%	7.50% (6.30-8.80%) 133/1785
80-	26.27% (22.95-29.80)	25.23-68.95%	16.70% (14.50-19.00%) 185/1108
All age groups	1.78% (1.63-1.95%)	1.72-2.56%	1.00% (0.90-1.10%) 478/46048

Notes.

^aCrI: 95% credibility intervals (CrI)

^b95%CI: 95% confidence interval

Figures

Figure 1. Gender proportion of reported COVID-19 cases by age groups (A), and cumulative morbidity ratio by gender and age group (B), March through May 18th, 2020, Chile

409 **Figure 2:** Temporal distribution of cases and deaths by age group due to COVID-19, March through May 18th, 2020, Chile.

411 **Notes**

412 Cumulative cases of CoVID-19 by the following age groups (A) 0-39, (B) 40-49, (C) 50-59, (D) 60-69, (E) 70-79, (F) 80years or more , (G) all cases and cumulative deaths
 413 by age group (H) 0-39, (I) 40-49, (J) 50-59, (K) 60-69, (L) 70-79, (M) 80 years or more, (N) all fatal cases over time. Day 1 corresponds to April 1st in 2020. As the dates of
 414 illness onset were not available, we used dates of reporting.

415 **Figure 3:** Temporal variation of risk of death caused by COVID-19, March through May 18th, 2020, Chile.

416
417

Notes.

418 Observed and posterior estimated of crude case fatality ratio of persons aged (A) 0-39, (B) 40-49, (C) 50-59, (D) 60-69, (E) 70-79, (F) 80 and more, and (G) all age groups.

419 Time-delay adjusted case fatality ratio of persons aged (H) 0-39, (I) 40-49, (J) 50-59, (K) 60-69, (L) 70-79, (M) 80 and more, and (N) all age groups. Day 1 corresponds to

420 April 1st, 2020. Black dots show crude case fatality ratio, and light and dark indicates 95% and 50% credible intervals for posterior estimates, respectively.

Figure 4. Latest estimates of time-delay adjusted risk of death caused by COVID-19 by age group, March-May 2020, Chile.

Notes

Distribution of time-delay adjusted case fatality risks derived from the latest estimates (May 8, 2020) are presented. Top to bottom: aged 0-39, aged 40-49, aged 50-59, aged 60-69, aged 70-79, aged 80- and all age groups.