Renin-angiotensin-aldosterone system inhibitors and mortality in patients with hypertension hospitalized for COVID19: systematic review & meta-analysis

Journal:	<i>Open Heart</i>
Manuscript ID	openhrt-2020-001353
Article Type:	Original research
Date Submitted by the Author:	02-Jun-2020
Complete List of Authors:	Ssentongo, Anna; Penn State College of Medicine, Trauma Surgery, Public Health Sciences Ssentongo, Paddy; Penn State College of Medicine, Public Health Sciences Heilbrunn, Emily; Penn State College of Medicine, Public Health Sciences Lekoubou, Alain; Penn State College of Medicine, Public Health Sciences Du, Ping; Penn State College of Medicine, Public Health Sciences Liao, Duanping; Penn State College of Medicine, Public Health Sciences Oh, John S; Penn State College of Medicine, Public Health Sciences Chinchilli, Vernon; Penn State College of Medicine, Public Health Sciences
Keywords:	HYPERTENSION < HYPERTENSION, ANTIHYPERTENSIVE DRUGS < HYPERTENSION, HYPERTENSIVE HEART DISEASE < HYPERTENSION
Abstract:	<p>Objective: The association between renin-angiotensin-aldosterone (RAAS) inhibitors and Coronavirus diseases 2019 (COVID-19) mortality is unclear. We aimed to explore the association of RAAS inhibitors, including angiotensin-converting inhibitors (ACEi) and angiotensin II receptor blockers (ARBs) with COVID-19 mortality in patients with hypertension.</p> <p>Methods: MEDLINE, SCOPUS, OVID, and Cochrane Library were searched for the period of January 1, 2020 to May 20, 2020. Studies reporting the association of RAAS inhibitors (ACEi and ARBs) and mortality in patients with hypertension, hospitalized for COVID-19 were extracted. Two reviewers independently extracted appropriate data of interest and assessed the risk of bias. All analyses were performed using random-effects models on log-transformed risk ratio estimates, and heterogeneity was quantified.</p> <p>Results: Data were collected on 2,065,805 individuals (mean age, 58.73 years; 53.4% male). Patients with hypertension taking RAAS inhibitors were 35% less likely to die from COVID-19 compared to patients with hypertension not taking RAAS inhibitors (pooled RR= 0.65, 95% Confidence Intervals (CI): 0.45-0.94). To explore the association of COVID-19 and specific classes of RAAS inhibitors, we conducted a subgroup analysis of ARBs and ACEi separately from studies that provided them. Pooled risk ratio estimates from ARBs and ACEi showed a lower but not significant risk of death from COVID-19 (RR=0.93, 95% CI: 0.70-1.22) and ACEi (RR=0.65, 95% CI: 0.32-1.30).</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	Conclusions: In this meta-analysis, it was discovered that taking RAAS inhibitors, significantly decreased the risk of COVID-19 mortality in patients with hypertension. This indicates a potential protective role that RAAS-inhibitors may have in COVID-19 patients with hypertension.

SCHOLARONE™
Manuscripts

medRxiv preprint doi: <https://doi.org/10.1101/2020.05.21.20107003>; this version posted June 2, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. All rights reserved. No reuse allowed without permission.

I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Renin-angiotensin-aldosterone system inhibitors and mortality in patients with hypertension hospitalized for COVID-19: a systematic review and meta-analysis

Anna E. Ssentongo, MPH^{§1,2*}, Paddy Ssentongo, MD, MPH^{1,3*}, Emily S. Heilbrunn, BS^{1*}, Alain Lekoubou, MD, Msc^{1,4}, Ping Du, MD, PhD¹, Duanping Liao, MD, PhD¹, John S. Oh, MD², Vernon M. Chinchilli, PhD¹

Word count: 1912

*Authors contributed equally

Corresponding Author

Anna Ssentongo, MPH

Penn State University

Department of Surgery

500 University Drive

Hershey, PA 17033

Pennsylvania State University

USA, Tel: 814-404-5208

Email: assentongo@pennstatehealth.psu.edu

Confidential: For Review Only

Abstract

Objective: The association between renin-angiotensin-aldosterone (RAAS) inhibitors and Coronavirus diseases 2019 (COVID-19) mortality is unclear. We aimed to explore the association of RAAS inhibitors, including angiotensin-converting inhibitors (ACEi) and angiotensin II receptor blockers (ARBs) with COVID-19 mortality in patients with hypertension.

Methods: MEDLINE, SCOPUS, OVID, and Cochrane Library were searched for the period of January 1, 2020 to May 20, 2020. Studies reporting the association of RAAS inhibitors (ACEi and ARBs) and mortality in patients with hypertension, hospitalized for COVID-19 were extracted. Two reviewers independently extracted appropriate data of interest and assessed the risk of bias. All analyses were performed using random-effects models on log-transformed risk ratio estimates, and heterogeneity was quantified.

Results: Data were collected on 2,065,805 individuals (mean age, 58.73 years; 53.4% male). Patients with hypertension taking RAAS inhibitors were 35% less likely to die from COVID-19 compared to patients with hypertension not taking RAAS inhibitors (pooled RR= 0.65, 95% Confidence Intervals (CI): 0.45-0.94). To explore the association of COVID-19 and specific classes of RAAS inhibitors, we conducted a subgroup analysis of ARBs and ACEi separately from studies that provided them. Pooled risk ratio estimates from ARBs and ACEi showed a lower but not significant risk of death from COVID-19 (RR=0.93, 95% CI: 0.70-1.22) and ACEi (RR=0.65, 95% CI: 0.32-1.30).

Conclusions: In this meta-analysis, it was discovered that taking RAAS inhibitors, significantly decreased the risk of COVID-19 mortality in patients with hypertension. This indicates a potential protective role that RAAS-inhibitors may have in COVID-19 patients with hypertension.

Key Questions:

What is already known about this subject?

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), responsible for the recent coronavirus disease 2019 (COVID-19) pandemic, interfaces with the renin-angiotensin-aldosterone system (RAAS) through angiotensin-converting enzyme 2 (ACE2). Recent studies have questioned whether RAAS inhibitors are safe in patients with COVID-19. However, observational studies involving patients hospitalized with COVID-19 that report the association of RAAS-inhibitors and COVID-19 severity or death have yielded conflicting findings.

What does this study add?

This systematic review and meta-analysis discovered that RAAS inhibitors reduced the risk of mortality by 35% in patients with hypertension hospitalized for COVID-19 (RR= 0.65, 95% CI 0.45-0.94).

How might this impact clinical practice?

Patients taking RAAS-inhibitors to manage their hypertension should continue to do as per current treatment guidelines.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), responsible for the recent coronavirus disease 2019 (COVID-19) pandemic, interfaces with the renin-angiotensin-aldosterone system (RAAS) through angiotensin-converting enzyme 2 (ACE2).¹ The current hypotheses related to the influence ACE2 may have in facilitating virus severity and mortality have been inconclusive. The increased expression of ACE2 is thought to potentially catalyze infection with COVID-19, and therefore increase the severity and risk of death.² On the contrary, it has been found that ACE2 may be protective against acute lung injury.³

ACE2 is an 805-amino-acid, homologous to the human angiotensin-converting enzyme (ACE), with 40% identity and 61% similarity.⁴ The SARS-Cov2 virus binds to the ACE2 receptor for cell entry.¹ Prior research has suggested that ACE- inhibitors (ACEi) and angiotensin-II blockers (ARBs), which are commonly used in patients with hypertension or diabetes, may raise ACE2 levels and thus could increase the risk of severe COVID-19 infection.⁵ Although ACE and ACE2 are two different enzymes with two different active sites, there are reports that ACE inhibitors affect the expression of ACE2 in the heart and kidneys.⁶ ARBs alter ACE2 expression, both at the mRNA and protein level.^{6 7} ACE2 is upregulated in both the renal vasculature tissue and cardiac tissue as a result of RAAS inhibitor exposure.⁸

Individuals with cardiovascular disease including hypertension are susceptible to SARS-CoV-2 infection,⁹ and majority depend on RAAS inhibitors for hypertension control, the potential influence of ACEi and ARB during SARS-CoV-2 infection requires urgent exploration for a clarification. Despite these theoretical uncertainties regarding whether pharmacologic regulation of ACE2 may influence the infectivity of SARS-CoV-2, there is clear potential for harm related to the withdrawal of RAAS-inhibitors in patients in otherwise stable condition.

To date, observational studies involving patients hospitalized with COVID-19 that report the association of RAAS-inhibitors and COVID-19 severity or death have yielded conflicting findings. Some studies are finding potential harmful associations of exposure to ACEi or ARBs with an increased

risk of severity in COVID-19¹⁰, and other studies failed to confirm such findings regarding a potential harmful association.^{11 12} Many individuals with hypertension take ACE inhibitors and ARBs, but the association of RAAS-inhibitors and mortality in COVID-19 patients has not been systematically reviewed using a large number of studies. The aim of this systematic review and meta-analysis is to delineate the association between use of RAAS-inhibitors and mortality in patients with COVID-19. We hypothesize that RAAS-inhibitors may increase mortality rates from the novel coronavirus that causes COVID-19.

Methods

Search Strategy and Study Selection

The authors state that all supporting data are available within the article and its online-only data supplement. We explored PubMed (MEDLINE) Scopus, the OVID databases, SCOPUS, Cochrane Library databases and medrxiv.org, using search criteria provided in the supplemental material (**Supplemental Text 1**). We invoked the Meta-analyses of Observational Studies in Epidemiology (MOOSE) during our search (**Supplemental Table 1**). We included all studies published from January 1, 2020 to May 20, 2020 that reported on the use of RAAS inhibitors (ACEi or ARBs) in patients hospitalized with COVID-19. We identified papers reporting the mortality rate in patients with and without exposure to RAAS-inhibitors. The following Medical Subject Heading (MeSH) and key words were used for the literature search of PubMed and other databases: “receptors, angiotensin” OR “angiotensin” OR “angiotensin receptors” OR “angiotensin converting enzyme inhibitors” “Renin Angiotensin Aldosterone System” OR “Angiotensin Receptor Blocker” OR “Ace Inhibitor” OR “Angiotensin Converting Enzyme Inhibitor” AND “SARS-CoV-2” OR “COVID-19” OR “Coronavirus”. Two reviewers (ESH and AES) initially screened the titles and abstracts of all papers for eligibility. We included articles that reported the rates of death in COVID-19 patients with and without taking RAAS inhibitors. No language limitation

was identified. We excluded studies that were not conducted in humans and did not report the rates of death.

Quality Assessment and Data Extraction

Two reviewers (ESH and AES) then screened full-text articles. A third reviewer (PS) was recruited in the event of a discrepancy. Data extracted included the author, year of publication, country, sample size, the number of patients in the RAAS inhibitor group that did or did not die, and the RR/OR/HR of death in the RAAS inhibitor group compared to the non-RAAS inhibitor group. Two reviewers (ESH and AES) independently assessed the quality of the included studies. The Newcastle-Ottawa Scale (NOS) was utilized for the quality assessment of included studies. NOS scale rates observational studies based on 3 parameters: selection, comparability between the exposed and unexposed groups, and exposure/outcome assessment. This scale assigns a maximum of 4 stars for selection, 2 stars for comparability, and 3 stars for exposure/outcome assessment. Studies with less than 5 stars were considered low quality, studies receiving 5 through 7 stars were considered moderate quality, and those receiving more than 7 stars were classified as high quality.

Data Analysis

The primary outcome of interest was the risk of mortality in patients with hypertension hospitalized for COVID-19. The exposure of interest was the use of RAAS inhibitors. Subgroup analysis of ACEi and ARB separately were conducted. We utilized the reported RR, HR, OR as the measures of the association between exposure to RAAS-inhibitors and the risk of mortality in COVID-19. For studies without measures of associations, we applied a generalized linear mixed model to calculate the odds ratios using the number of events and the sample size of each study group.¹³ Because the outcome of mortality was relatively rare, we combined RRs and HRs with ORs in the present meta-analysis and reported the pooled effect size as RRs as common risk estimates for all studies. We invoked a random-effects models to pool study results for the association between exposure to RAAS-inhibitors and the

risk of mortality.¹⁴ We constructed forest plots to display pooled estimates. We assessed inter-study heterogeneity using I^2 statistics, expressed as % (low (25%), moderate (50%), and high (75%)) and Cochrane's Q statistic (significance level < 0.05)^{15 16}. Assessment of potential ascertainment bias (as might be caused by publication bias) was conducted with funnel plots, by plotting the study effect size against standard errors of the effect size, and Egger's test. [12] We performed all statistical analyses with R software, version 3.4.3 (R, College Station, TX).

Results

As shown in **Figure 1**, we identified a total of 337 studies from the five databases. We identified 152 studies as duplicates and excluded, leaving 185 studies. When screening titles and abstracts, we excluded 56 studies and another 129 based on full text, which left us with 2,065,805 patients from 15 studies for qualitative analysis and 61,268 patients from 11 studies for the quantitative analysis (**Table 1**).^{10-12 17-27}

The association between taking RAAS inhibitors and COVID-19 mortality

Of the seven studies included in the meta-analysis exploring the association between COVID-19 mortality and RAAS exposure in patients with hypertension, three reported a significantly lower risk with mortality (**Figure 2**). No studies reported a significantly higher risk of mortality. The overall pooled estimates showed a 35% reduction in the risk of mortality (RR=0.65, 95% CI: 0.45-0.94). The inverse association between RAAS use and low risk of mortality from COVID-19, may indicate a potential protective role for RAAS inhibitors among COVID-19 patients with hypertension. Between-study heterogeneity was high ($I^2 = 80$, $p < 0.01$).

Subgroup analysis of the association of ARBs and ACEi with mortality from COVID-19

To explore the association of COVID-19 and specific classes of RAAS inhibitors, we conducted a subgroup analysis of ARBs and ACEi separately from studies that provided them. Three studies provided

1 risk ratios estimates (or data to calculate the RR) for ACEi (**Figure 3**) and three for ARBs (**Figure 4**).
2
3 Pooled risk ratio estimates from ARBs and ACEi showed a low risk of death from COVID-19 (RR=0.93,
4
5 95% CI: 0.70-1.22) and ACEi (RR=0.65, 95% CI: 0.32-1.30). However, the association was not
6
7 significant. Between-study heterogeneity was moderate ($I^2= 58$, $p=0.09$) from ARBs and high ($I^2= 90$,
8
9 $p<0.01$) for ACEi.
10
11
12
13

14 **Sensitivity Analyses, Publication Bias Publication and Study Quality**

15
16 We conducted two types of sensitivity analyses. First, limiting our analysis to the 7 RAAS inhibitor
17
18 studies whose sample size consisted of only individuals with hypertension, we excluded and replaced
19
20 one study at a time from the meta-analysis and calculating the pooled RR for the remaining studies. No
21
22 substantial changes from pooled RR were observed when other studies were removed in turn. The
23
24 pooled RR ranged from 0.57 to 0.72 ($p<0.0001$ for all) (**Supplementary Figure 1**). Second, by
25
26 including all studies (main analysis plus 3 studies with sample size not limited to population with
27
28 hypertension, see **table 1**), pooled RR from sensitivity analysis ranged from 0.68 to 0.82 ($p<0.0001$ for
29
30 all) (**Supplementary Figure 2**). We did not find significant funnel plot asymmetry for the Egger test for
31
32 RAAS inhibitor, ACEi and ARBs ($p > .05$). Upon visual inspection, the funnel plots appeared
33
34 symmetrical for RAAS inhibitors, ACEi and ARBs (**Supplemental Figures 3-6**). The median study
35
36 quality score was 8 out of 9 (range=7-9).
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

This systematic review and meta-analysis of cohort studies consisting of 2,065,805 patients with a global representation suggests that the treatment of hypertension with RAAS-inhibitors is associated with a lower risk of mortality in patients with COVID-19. This finding is important, for the association between RAAS-inhibitor exposure and mortality in COVID-19 patients has been inconclusive thus far. This topic has been heavily debated, and some studies even have interpolated a risk of taking RAAS-inhibitors using data from previous coronavirus outbreaks and preclinical studies. [6]

RAAS-inhibitors have been found to mitigate the risk of severe lung injury by reducing the activation of the RAAS through the inactivation of angiotensin II³ and the generation of angiotensin-(1-9)⁴ and angiotensin-(1-7)²⁸. Angiotensin- (1-7) binds to the G protein-coupled receptors Mas to mediate various physiological effects including vasorelaxation, cardio protection, anti-oxidation and inhibition of angiotensin II -induced signaling. This is one hypothesized mechanism illustrating how the treatment of chronic conditions with RAAS-inhibitors may be beneficial in COVID-19 patients.

Alternatively, it is hypothesized that the biological mechanisms of RAAS inhibitors may predispose COVID-19 patients to severe disease and mortality. This was explained by the observation that SARS-CoV-2 enters a cell by binding to the membrane-bound ACE2 receptors. Animal models suggest that ACEis and ARBs increase membrane-bound ACE2 receptors, to which then increases the availability of cells for SARS-CoV-2 to bind and cellular entry.⁶ This hypothesis has sparked a debate in populations, for many individuals taking RAAS inhibitors have grown concerned that their medications may be predisposing them to developing COVID-19, and later dying from it.²⁹ Our meta-analysis and systematic review support the notion that RAAS inhibitor exposure does *not* increase COVID-19-related mortality but rather shows beneficial effect.

Strengths and Limitations

1
2 Limitations of our study include possible selection bias in the published literature as a result of the strict
3
4 COVID-19 testing algorithm employed in the early stages of the pandemic. This may have resulted in
5
6 missed COVID-19 cases or deaths. Nevertheless, this is the largest quantitative synthesis of evidence on
7
8 the association between RAAS-inhibitor exposure and COVID-19 mortality. The regions with the
9
10 highest burden of COVID-19, including Asia, Europe, and North America, were represented thus
11
12 increasing the external validity of our findings. The sample size included in this study was also quite
13
14 large, allowing us to thoroughly cover a large population.
15
16

17 **Conclusion**

18
19
20 This systematic review and meta-analysis found that COVID-19 patients with hypertension who take
21
22 RAAS-inhibitors are protected from COVID-19 mortality compared to hypertensive patients not taking
23
24 RAAS-inhibitors. Patients taking RAAS-inhibitors to manage their chronic diseases may continue to do
25
26 as per current treatment guidelines and based on the clinical judgment of their health care providers.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Author Affiliations

¹Department of Public Health Sciences, Penn State College of Medicine, Hershey, Pennsylvania, United States of America

²Department of Surgery, Penn State College of Medicine and Milton S. Hershey Medical Center, Hershey, Pennsylvania, United States of America

³Center for Neural Engineering, Department of Engineering, Science and Mechanics, The Pennsylvania State University, Pennsylvania, United States of America

⁴Department of Neurology, Penn State College of Medicine and Milton S. Hershey Medical Center, Hershey, Pennsylvania, United States of America

Competing Interest We report no competing interest

Contributorship AS, PS, ES, and VC conceived the study. AS, EH, and PS conducted the literature search. AS and PS completed data analysis. AS, DL, PD, VC, AL JO, EH and PH interpreted the data. AS, ES, and PS wrote the manuscript. All Authors agreed to the manuscript in its final form

Acknowledgements We would like to acknowledge Melissa Butt for reviewing and proving helpful feedback.

Patient and Public Involvement: Patients or the public were not involved in the preparation or dissemination of this manuscript

Funding This study was not funded

Ethics Statement: This is a systematic review and meta-analysis and individual patient was not used. Therefore we did not need IRB or an ethics board approval.

Data Sharing Statement: All data relevant to the study are included in the article or uploaded as supplementary information

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Confidential: For Review Only

REFERENCES

1. Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020
2. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *The Lancet Respiratory Medicine* 2020
3. Imai Y, Kuba K, Rao S, et al. Angiotensin-converting enzyme 2 protects from severe acute lung failure. *Nature* 2005;436(7047):112-16.
4. Tipnis SR, Hooper NM, Hyde R, et al. A human homolog of angiotensin-converting enzyme cloning and functional expression as a captopril-insensitive carboxypeptidase. *Journal of Biological Chemistry* 2000;275(43):33238-43.
5. Ferrario CM, Ahmad S, Groban L. Mechanisms by which angiotensin-receptor blockers increase ACE2 levels. *Nature Reviews Cardiology* 2020:1-1.
6. Ferrario CM, Jessup J, Chappell MC, et al. Effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockers on cardiac angiotensin-converting enzyme 2. *Circulation* 2005;111(20):2605-10.
7. Wang X, Ye Y, Gong H, et al. The effects of different angiotensin II type 1 receptor blockers on the regulation of the ACE-AngII-AT1 and ACE2-Ang (1-7)-Mas axes in pressure overload-induced cardiac remodeling in male mice. *Journal of molecular and cellular cardiology* 2016;97:180-90.
8. Ferrario CM, Jessup J, Gallagher PE, et al. Effects of renin-angiotensin system blockade on renal angiotensin-(1-7) forming enzymes and receptors. *Kidney international* 2005;68(5):2189-96.
9. Ssentongo P, Ssentongo AE, Heilbrunn ES, et al. The association of cardiovascular disease and other pre-existing comorbidities with COVID-19 mortality: A systematic review and meta-analysis. *medRxiv* 2020
10. Zeng Z, Sha T, Zhang Y, et al. Hypertension in patients hospitalized with COVID-19 in Wuhan, China: A single-center retrospective observational study. *medRxiv* 2020
11. Mehra MR, Desai SS, Kuy S, et al. Cardiovascular Disease, Drug Therapy, and Mortality in Covid-19. *New England Journal of Medicine* 2020
12. Zhang P, Zhu L, Cai J, et al. Association of inpatient use of angiotensin converting enzyme inhibitors and angiotensin II receptor blockers with mortality among patients with hypertension hospitalized with COVID-19. *Circulation research* 2020
13. Chang B-H, Hoaglin DC. Meta-analysis of odds ratios: Current good practices. *Medical care* 2017;55(4):328.
14. Schwarzer G, Carpenter JR, Rücker G. Meta-analysis with R: Springer 2015.
15. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in medicine* 2002;21(11):1539-58.
16. Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *Bmj* 2003;327(7414):557-60.
17. Mehta N, Kalra A, Nowacki AS, et al. Association of use of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers with testing positive for coronavirus disease 2019 (COVID-19). *JAMA cardiology* 2020
18. Huang Z, Cao J, Yao Y, et al. The effect of RAS blockers on the clinical characteristics of COVID-19 patients with hypertension. *Annals of Translational Medicine* 2020;8(7)
19. Mancina G, Rea F, Ludergrani M, et al. Renin-angiotensin-aldosterone system blockers and the risk of Covid-19. *New England Journal of Medicine* 2020
20. Li J, Wang X, Chen J, et al. Association of renin-angiotensin system inhibitors with severity or risk of death in patients with hypertension hospitalized for coronavirus disease 2019 (COVID-19) infection in Wuhan, China. *JAMA cardiology* 2020
21. Meng J, Xiao G, Zhang J, et al. Renin-angiotensin system inhibitors improve the clinical outcomes of COVID-19 patients with hypertension. *Emerging microbes & infections* 2020;9(1):757-60.

1
2 22. Zhang L, Sun Y, Zeng H-L, et al. Calcium channel blocker amlodipine besylate is associated with
3 reduced case fatality rate of COVID-19 patients with hypertension. *medRxiv* 2020
4 23. Guo T, Fan Y, Chen M, et al. Cardiovascular implications of fatal outcomes of patients with
5 coronavirus disease 2019 (COVID-19). *JAMA cardiology* 2020
6 24. Bean D, Kraljevic Z, Searle T, et al. ACE-inhibitors and Angiotensin-2 Receptor Blockers are not
7 associated with severe SARS-COVID19 infection in a multi-site UK acute Hospital Trust. *medRxiv*
8 2020
9 25. Yang G, Tan Z, Zhou L, et al. Angiotensin II Receptor Blockers and Angiotensin-Converting Enzyme
10 Inhibitors Usage is Associated with Improved Inflammatory Status and Clinical Outcomes in
11 COVID-19 Patients With Hypertension. *medRxiv* 2020
12 26. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting characteristics, comorbidities, and
13 outcomes among 5700 patients hospitalized with COVID-19 in the New York City area. *Jama*
14 2020
15 27. Ip A, Parikh K, Parrillo JE, et al. Hypertension and Renin-Angiotensin-Aldosterone System Inhibitors
16 in Patients with Covid-19. *medRxiv* 2020
17 28. Donoghue M, Hsieh F, Baronas E, et al. A novel angiotensin-converting enzyme-related
18 carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9. *Circulation research*
19 2000;87(5):e1-e9.
20 29. Diaz JH. Hypothesis: angiotensin-converting enzyme inhibitors and angiotensin receptor blockers
21 may increase the risk of severe COVID-19. *Journal of Travel Medicine* 2020
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Studies of inclusion

Author	Year	Country	Continent	Sample size (n)	Study type	Study period	Mean age	Male (%)	Study Population	Covariates adjusted	Effect Estimate	Quality Score
Zhang et al 2 3 4	2020	China	Asia	3,430	Cohort	12/31/2019-2/20/2020	64	53	Hypertensive COVID-19 patients taking ACEI/ARB vs hypertensive COVID-19 not taking ACEI/ARB	Age, sex, diabetes, coronary heart disease, cerebrovascular disease, and chronic renal disease, in-hospital medications (antiviral drug and, lipid lowering drug).	HR	9
Huang et al 5 6 7	2020	China	Asia	50	Cohort	2/07/2020-3/03/2020	-	54	Hypertensive COVID-19 patients taking RAAS inhibitors vs hypertensive COVID-19 patients not taking RAAS inhibitors	-	Qualitative Review Only	8
Mehra et al 8 9 10	2020	-	Asia, Europe, & North America	8,910	Cohort	12/20/2019-3/15/2020	49	60	Hypertensive COVID-19 patients taking ACEI/ARB vs not taking ACEI/ARB	Age, sex, coronary artery disease, congestive heart failure, arrhythmia, COPD, smoking, statin	OR	8
Mancia et al 11 12 13 14	2020	Italy	Europe	37,031	Case-control	2/21/2020-3/11/202	68	64	Hypertensive COVID-19 patients taking ACEI vs hypertensive COVID-19 patients not taking ACEI, hypertensive COVID-19 patients taking ARB vs hypertensive COVID-19 patients not taking ARB	Age, Sex, comorbidities, and exposure to treatments	OR	8
Lei et al 15 16 17	2020	China	Asia	1,178	Cohort	1/15/2020-3/15/2020	56	46	Hypertensive COVID-19 patients taking ACEI/ARB vs hypertensive COVID-19 patients not taking ACEI/ARB	-	Calculated OR	8
Mehra et al 18 19 20	2020	United States	North America	18,472	Cohort	3/08/2020-4/12/2020	49	40	COVID-19 patients taking ACEI vs not taking ACEI, COVID-19 patients taking ARB vs not taking ARB, COVID-19 patients taking ACEI/ARB vs not taking ACEI/ARB	-	Calculated OR	7
Meng et al 21 22	2020	China	Asia	417	Cohort	1/11/2020-2/23/2020	57	65	Hypertensive COVID-19 patients taking ACEI/ARB vs hypertensive COVID-19 patients not taking ACEI/ARB	-	Qualitative Review Only	8
Zhang et al 23 24 25	2020	China	Asia	90	Cohort	-	-	-	COVID-19 patients taking hypertensive drugs vs not taking hypertensive drugs	Age, sex, days from symptom onset to hospital admission, and exposure to treatments	HR	9
Guo et al 26 27	2020	China	Asia	187	Cohort	1/23/2020-2/23/2020	59	49	COVID-19 patients taking ACEI vs not taking ACEI, COVID-19 patients taking ARB vs not taking ARB	-	Qualitative Review Only	7
Bean et al 28 29 30	2020	England	Europe	1,200	Cohort	3/1/2020-4/13/2020	63	42	COVID-19 patients taking ACEI/ARB vs not taking ACEI/ARB	Age, sex, hypertension, diabetes, chronic kidney disease, and ischemic heart disease/heart failure	OR	9
Yang et al 31 32	2020	China	Asia	247	Cohort	1/5/2020-3/8/2020	60	55	COVID-19 patients taking ACEI/ARB vs not taking ACEI/ARB	-	Calculated	8
Yang et al 33 34 35	2020	China	Asia	251	Cohort	1/5/2020-3/3/2020	-	-	Hypertensive COVID-19 patients taking ACEI/ARB vs hypertensive COVID-19 patients not taking ACEI/ARB vs non-hypertensive COVID-19 patients not taking ACEI/ARB	-	Calculated	8
Richardson et al 36 37	2020	United States	North America	5,700	Case series	3/1/2020-4/4/2020	63	60	Hypertensive COVID-19 patients taking ACEI/ARB vs hypertensive COVID-19 patients not taking ACEI/ARB	-	Calculated	8
Ip et al 38 39 40	2020	United States	North America	3,017	Cohort	-	-	-	Hypertensive COVID-19 patients taking ACEI/ARB vs hypertensive COVID-19 patients taking other hypertensive medications	-	Calculated	8
Puchet et al 41 42 43	2020	France	Europe	1,985,598	Cohort	2/21/2020-4/5/2020	-	-	Hypertensive COVID-19 patients taking ACEI vs hypertensive COVID-19 patients taking ARB vs hypertensive patients taking other drugs	-	Qualitative Review only	7

Figure 1: Flow Diagram

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

Author, Publication Year, Country	Risk Ratio	RR	95% CI	Weight
Mehra et al,2020,Asia, Europe, & North America		0.33	[0.23; 0.48]	33.8%
Mancia et al,2020,Italy		0.91	[0.72; 1.14]	36.2%
Li et al,2020,China		0.92	[0.52; 1.63]	30.0%
Overall (Random-Effect Model)		0.65	[0.32; 1.30]	100.0%
Heterogeneity: $I^2 = 90\%$, $\tau^2 = 0.3372$, $p < 0.01$				

Author, Publication Year, Country

Li et al,2020,China
Mancia et al,2020,Italy
Mehra et al,2020,Asia, Europe, & North America

Overall (Random-Effect Model)

Heterogeneity: $I^2 = 58\%$, $\tau^2 = 0.0346$, $p = 0.09$

Risk Ratio **RR 95% CI Weight**

0.77 [0.50; 1.18] 24.7%

0.83 [0.66; 1.04] 42.2%

1.23 [0.90; 1.69] 33.1%

0.93 [0.70; 1.22] 100.0%

Supplementary material for Ssentongo et., al 2020

Supplemental text 1: PubMed Search Terms

(((((("receptors, angiotensin"[MeSH Terms] OR "angiotensin"[All Fields])) OR "angiotensin receptors"[All Fields]) OR ("angiotensin"[All Fields] OR "(((("angiotensin converting enzyme inhibitors"[Pharmacological Action] OR "angiotensin-converting enzyme inhibitors"[MeSH Terms]) OR ("angiotensin converting enzyme inhibitors"[All Fields]) OR OR "ace inhibitor"[All Fields])) OR "angiotensin-converting enzyme inhibitors"[MeSH Terms]) OR angiotensin converting enzyme inhibitors"[All Fields])

Supplemental Table 1: Meta-Analyses and Systematic Reviews of Observational Studies (MOOSE)

	Topic	Page number
Title	Identify the study as a meta-analysis (or systematic review)	1
Abstract	Use the journal's structured format	2
Introduction	Present:	4
	The clinical problem	4
	The hypothesis	4
	A statement of objectives that includes the study population, the condition of interest, the exposure or intervention, and the outcome(s) considered	4
Sources	Describe:	
	Qualifications of searchers (eg, librarians and investigators)	5
	Search strategy, including time period included in the synthesis and keywords	5
	Effort to include all available studies, including contact with authors	5
	Databases and registries searched	5
	Search software used, name and version, including special features used (e.g. explosion)	5-6
	Use of hand searching (e.g, reference lists of obtained articles)	5
	List of citations located and those excluded, including justification	5
	Method of addressing articles published in languages other than English	5
	Method of handling abstracts and unpublished studies	5
	Description of any contact with authors	5
Study Selection	Describe	
	Types of study designs considered	5
	Relevance or appropriateness of studies gathered for assessing the hypothesis to be tested	5
	Rationale for the selection and coding of data (eg, sound clinical principles or convenience)	5
	Documentation of how data were classified and coded (eg, multiple raters, blinding, and inter-rater reliability)	5
	Assessment of confounding (e.g. comparability of cases and controls in studies where appropriate)	5-6
	Assessment of study quality, including blinding of quality assessors; stratification or regression on possible predictors of study results	5-6
	Assessment of heterogeneity	6
	Statistical methods (eg, complete description of fixed or random effects models, justification of whether the chosen models account for predictors of study results, dose-response models, or cumulative meta-analysis) in sufficient detail to be replicated	6
Results	Present	6-7
	A graph summarizing individual study estimates and the overall estimate	15
	A table giving descriptive information for each included study	Table 1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	Results of sensitivity testing (eg, subgroup analysis)	6-7
	Indication of statistical uncertainty of findings	6-7
Discussion	Discuss	8-9
	Strengths and weaknesses	8-9
	Potential biases in the review process (eg, publication bias)	8-9

*Modified from Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. JAMA 2000;283:2008–12. Copyrighted © 2000, American Medical Association. All rights reserved.

Supplementary Figure 1. Sensitivity analysis of random-effects meta-analysis of studies evaluating the association of ACEi/ARB with mortality in patients with hypertension hospitalized with COVID-19 (only study population hypertension)

Supplementary Figure 2. Sensitivity analysis of random-effects meta-analysis of studies evaluating the association of ACEi/ARB with mortality in patients with hypertension hospitalized with COVID-19 (study population with and without hypertension)

Supplementary Figure 3. Funnel plot depicting publication bias for studies evaluating the association of ACEi/ARB with mortality in patients with hypertension hospitalized with COVID-19 (only study population hypertension)

Supplementary Figure 4. Funnel plot depicting publication bias for studies evaluating the association of ACEi/ARB with mortality in patients with hypertension hospitalized with COVID-19 (study population with and without hypertension)

Supplementary Figure 5. Funnel plot depicting publication bias for studies evaluating the association of ARB with mortality in patients with hypertension hospitalized with COVID-19

Supplementary Figure 6. Funnel plot depicting publication bias for studies evaluating the association of ACEi with mortality in patients with hypertension hospitalized with COVID-19