Hand Washing Compliance and COVID-19: A Non-Participatory Observational Study among Hospital Visitors

Biniyam Sahiledengle^{1*}, Yohannes Tekalegn¹, Abulie Takele¹, Demisu Zenbaba¹, Zinash Teferu¹, Alelign Tasew¹, Tesfaye Assefa², Kebebe Bekele³, Abdi Tesemma³, Habtamu Gezahegn⁴, Tadesse Awoke⁵, Bruce John Edward Quisido²

¹Madda Walabu University Goba Referral hospital, Department of Public Health, Bale-Goba, Ethiopia

²Madda Walabu University Goba Referral hospital, Department of Nursing, Bale-Goba, Ethiopia

³Madda Walabu University Goba Referral hospital, Department of Surgery, Bale-Goba, Ethiopia

⁴Madda Walabu University Goba Referral hospital, Department of Physiology, Bale-Goba, Ethiopia

⁵University of Gondar, College of Medicine and Health Sciences, Department of Epidemiology and Biostatistics, Gondar, Ethiopia

* Corresponding Author

E-mail: biniyam.sahiledengle@gmail.com

ORCID: 0000-0002-1114-4849

Abstract

Background: Hand washing remains a key measure for intercepting the dispatch of the Coronavirus disease (COVID-19). However, hand washing must be perpetuated properly using soap and water for at least 20 seconds. In response to the current COVID-19 pandemic, various hospitals have imposed mandatory hand washing to everyone prior entering the facilities, and when leaving. This study aimed to assess the hand washing compliance among visitors of a university referral hospital.

Methods: A non-participatory observational study was conducted in the main entrance of the hospital from April 27 to May 3, 2020, to measure hand washing compliance of its visitors. The quality of hand washing was assessed via direct observation for compliance with the recommended World Health Organization (WHO) core steps. Data were collected using Open Data Kit (ODK) mobile application.

Results: A total of 1,282 hospital visitors were observed, of which 874(68.2%) were males. Full hand washing compliances were observed among 0.9% (95% CI: 0.4-1.4) of the visitors. Withal, there was no difference in the compliances between genders (0.9% vs 0.7%, $P = 0.745$).

Conclusion: Despite the fact that proper hand washing with soap and water is proven to be one of the effective ways in preventing the spread of COVID-19, a significant number of hospital visitors did not practice standard hand washing procedures. Improvements in this measure are urgently needed in the face of the current COVID-19 pandemic.

Keywords: Hand washing, Compliance, COVID-19, Ethiopia, Observation

Background

Hand washing is reckoned to be one of the paramount measures for preventing the transmissions of Severe Acute Respiratory Syndrome (SARS)-CoV-2 virus, which causes the Coronavirus disease (COVID-19). Hand washing is simple, cost-effective, and one of the first lines of defense in ceasing the spread of the current pandemic. It must be done properly using soap and water for at least 20 seconds [1-3].

Respiratory infection caused by viruses like COVID-19 spread when mucus or droplets containing the virus advances into the body through the eyes, nose, or throat. Most often, this betides through contact with contaminated hands. The risk of COVID-19 transmission is also magnified when people come in contact with an infected person, or when they abut surfaces that an infected person has contaminated [3]. Coronavirus subsist in the hands may be effaced with handwashing using soap and water, and if it's done properly. If the hands are virtuous, there will be a marginal chance that people can dispatch the virus to the eyes, nose, or mouth of other people, or when they touch their faces [4]. Currently, there is a systematic review apropos hand washing and its risk for respiratory infection, and it implied that hand cleaning can gash the risk of the infection by 16% [5]. It is therefore paramount to wash the hands often and correctly to stop the spread of COVID-19 [1, 2, 6]. The unerring way of hand washing requires five key steps: 1) wet both hands under clean running water; 2) stow a liberal amount of soap to the front and back of the hands, as well as fingertips; 3) rub in between and around the fingers; rub the back of each hand using the palm of the other hand; rub fingertips of each hand to the opposite palm of the other hand, and rub each thumb clasped to the opposite hand for at least 20 seconds; 4) rinse thoroughly under running water, and dry hands with clean paper/towel; 5) turn off the water using a paper towel [1,7,8].

Compliance of the community with the recommended hand hygiene practices has paramount importance in averting the spread of COVID-19. Be that as it may, there are ample attestations that in umpteen settings, hand washing behaviors linger as an area which requisites improvement [1, 9-12]. A recent study by Pogrebna et al found that a country's hand washing culture is a very compelling predictor of the degree of COVID-19 spread [12].

In response to COVID-19, the World Health Organization (WHO) has strongly recommended that all the state members proffer public hand hygiene stations, and ensuring obligatory usage of these when visitors or personnel enter and leave the facilities, such as hospitals and health centers[2]. Ethiopia reported its first case of COVID-19 on 13th March 2020 [13]. From that time on, the country aggressively engaged and communicated different COVID-19 prevention measures to the general public. As part of its preventive proceedings, the public was notified to regularly wash their hands with soap and water for 20 seconds using the recommended steps. Furthermore, all hospitals should ensure the availability of hand washing stations in their main entrances, and that everyone should wash their hands before entering the facilities and when leaving [14]. Still and all, the acceptance of hospital visitors toward hand washing as a routine habit – and their compliances with the prescribed hand washing techniques were uncertain. In lieu of these, concerns about the poor hand hygiene culture in Ethiopia – and their negative implications in the control of COVID-19 were the motivations behind the undertakings of this study. Accordingly, this study assessed the hand washing compliances among the visitors of Madda Walabu University Goba Referral Hospital, Southeast of Ethiopia.

Methods

Study Design and Setting

A non-participatory observation study was conducted for a period of one week (from April 27 to May 3, 2020) to determine the magnitude of hand washing compliance (HWC) of the visitors entering Madda Walabu University Goba Referral Hospital. It is the only referral hospital in Bale Zone, Southeast Ethiopia. Data were collected through direct observations and were executed in the main entrance of the referral hospital for seven consecutive days to appreciate a complete picture of the weekly HWC among its visitors. The observations were imposed in the morning and afternoon (8:30-11:00 A.M and 2:00 -4:30 P.M). These times were chosen because they were the busiest visiting hours. Due to the current COVID-19 pandemic, hand washing facilities and the necessary resources (plain soaps and clean, running waters) were available at the main entrance of the referral hospital. Accordingly, visitors were obliged to wash their hands upon arrival on the hospital premises, and when leaving. Furthermore, large 1.2 m² hand washing signs were stationed at the hospital entrance to remind visitors about washing their hands.

Study Participants and Data Collection Procedures

All visitors who washed their hands before entering and when leaving the referral hospital were observed. Anyone who needs assistance/instruction in washing his/her hands was excluded. The observation was conducted using the recommended WHO hand washing tool [7]. Trained data collectors performed all observations and recorded the data using (Open Data Kit (ODK) Collect mobile Apps. To ensure data quality, the data collectors were informed to observe one visitor at a time. They also made sure that the visitors did not recognize them in order for the observant to display natural handwashing behaviors. Each time, two data collectors were involved, and they stand beside the washing area pretending to wait for someone else, or sometimes, they let their shoes be cleaned at the nearby shoe-cleaning stands.

Prior to the beginning of the study, all data collectors were extensively trained regarding data collection protocols and hand washing steps. To establish the inter-rater reliability, a testing session was imposed to data collectors by simulating hand washing demonstrations by random individuals while they observe them. And the interrater reliability was determined to be 0.86.

As recommended by the WHO, the study recorded and analyzed the recommended hand washing steps: 1) wetting the hands with water, 2) applying enough amount of soap to cover all hand surfaces, 3) rubbing the hand's palm with the other palm - rotary movement, 4) rubbing the right palm over the left dorsum with interlaced fingers, and vice versa, 5) again palm to palm rubbing with fingers interlaced, back of fingers to opposing palms with fingers interlocked, 6) rotational rubbing of left thumb clasped in the right palm, and vice versa, 7) rotational rubbing, backward and forward with clasped fingers of the right hand in the left palm, and vice versa, 8) rinsing the hands with water, 9) drying them with a clean item/towel, and finally, 10) using a clean item when turning off the faucet [7].

Operational Definition

Full compliance: wetting of hands, covering wet hands with soap, scrubbing all surfaces, including palms, back of the hands, in between fingers, and under the fingernails for about 20 seconds, rinsing well with running water, and finally, dry of hands with a clean cloth or by waving in the air [1].

Partial compliance: washing both hands with soap specifically, wetting hands, covering wet hands with soap, washing right palm over left dorsum with interlaced fingers and vice versa, and palm to palm rubbing with fingers interlaced, rinsing well with running water, and finally, drying using a clean cloth for about 20 seconds.

Minimal compliance: washing both hands with soap for about 20 seconds, which only includes wetting the hands, covering wet hands with soap, and rinsing well with running water.

Non-compliance: rinsing both hands with water only.

Statistical Analysis

All observed data were collected using ODK and were exported to the Statistical Package for the Social Sciences (SPSS) 20.0 software program for analysis. Based on the operational definitions mentioned, the level of hand washing compliance was computed. Descriptive statistics were computed to summarize the study findings. To test whether hand washing compliance varied across the weekdays and gender, the Chi-square test (χ^2) was used. A p-value < 0.05 was considered as statistically significant. Finally, data were presented in tables and graphs, as appropriate.

Ethical approval and considerations

The study was approved by ethical review committee of Madda Walabu University Goba Referral Hospital Research and Technology Transfer Office and permission was sought from Goba Referral Hospital medical director office (*Ref.no 01/2/12594*). Data collectors were worn protective face masks and kept reasonable physical distance during data collection. The data were collected in a private condition and kept confidential.

Results

A total of 1,282 visitors were observed, of which 874 (68.2%) were males and 408 (31.8%) were females. Six hundred thirty (47.8%) visitors were observed in the morning schedules (**Table 1**).

Table 1: The Characteristics of the Study Participants and their Distributions Over the Weekdays in Goba Referral Hospital, May 2020 (n=1,282)

Variable	Category	Frequency	Percent
Sex	Male	874	68.2
	Female	408	31.8
Observation Sessions	Afternoon	669	52.2
	Morning	613	47.8
Days of the Week	Monday	226	17.6
	Tuesday	132	10.3
	Wednesday	231	18.0
	Thursday	171	13.3
	Friday	156	12.2
	Saturday	225	17.6
	Sunday	141	11.0

Hand Washing Compliance

The descriptive analysis of the hand washing steps was shown in **Table 2**. The median (Q1-Q3) duration of hand washing was 22 seconds (14-34). In total, only 0.9% (95%CI: 0.4-1.4) of the study participants were fully compliant with the recommended hand washing techniques. A significant number of hospital visitors did not wash their hands properly or they have washed their hands in less effective ways (minimal, partial, or non-compliant) (**Fig 1**). The areas of the hand most neglected during hand washing among hospital visitors were presented in **Fig 2**.

Fig 1: Bar graph including hand washing compliance of hospital visitors at the entrance of Madda Walabu University Goba Referral Hospital, Southeast Ethiopia, May 2020 (n=1,282)

Fig 2: Images of most commonly missed hand washing areas identified among hospital visitors at the entrance of Madda Walabu University Goba Referral Hospital, Southeast Ethiopia, May 2020 (n=1,282)

Table 2: Hand washing compliance in response to COVID-19 pandemic among Study participants of Madda Walabu University Goba Referral Hospital, Southeast Ethiopia (n=1,282)

Hand Washing Practice	Frequency (yes)	Percent	95%CI
Wet hands with water	1271	99.1	98.7-99.6
Apply enough soap to cover all hand surfaces	952	74.3	71.8-76.8
Rub hands palm to palm	607	47.3	44.5-50.1
Right palm over left dorsum with interlaced fingers and vice versa	176	13.7	12.0-15.7
Palm to palm with fingers interlaced	197	15.4	13.3-17.4
Backs of fingers to opposing palm with fingers interlocked	71	5.5	4.3-6.8
Rotational rubbing of left thumb clasped in the right palm and vice versa	64	5.0	3.8-6.3
Rotational rubbing, backward and forward with clasped fingers of the right hand in left the palm and vice versa	65	5.1	3.9-6.3
Rinse hands with water	1102	86.0	84.0-87.9
Use towel/ clean item to turn off the faucet	185	14.4	12.6-16.3
Duration of hand washing, Median (Q1-Q3): 22 seconds (14-34)			
< 20 seconds	596	46.5	43.9-49.0
≥ 20 seconds	686	53.5	51.0-56.1
The Overall Hand Washing Compliance Score			
Full compliance with hand washing			
Yes	11	0.9	0.4-1.4
No	1271	99.1	98.6-99.6
Partial compliance to hand washing [#]			
Yes	37	2.9	2.0-3.8
No	1245	97.1	96.2-98.0
Minimal compliance to hand washing [#]			
Yes	605	47.2	44.4-49.9
No	677	52.8	50.1-55.6
Non-compliance to hand washing [#]			
Yes	258	20.1	17.8-22.3
No	1024	78.9	77.7-82.2

[#] Visitors did not wash their hands properly or in a less effective way as described in the operational definition section

Relationship between Hand Washing Compliance and Associated Variables

As shown in **Table 3**, there is no statistically significant difference between male and female hospital visitors concerning hand washing compliance (full and partial compliance). Significant differences were noted in hand washing compliance (full compliance) in between the weekdays ($p=0.03$) (**Additional File 2**). No significant differences in hand washing were noted based on time of day ($p=0.875$) (**Additional File 2**).

Table 3: The association between hand washing compliance and sex of the respondents of Madda Walabu University Goba Referral Hospital, Southeast Ethiopia, May 2020 (n=1,282)

Hand Washing Practice	Category	Male (n=874)	Female (n=408)	p-value
Wet hands with water	Yes	866(99.1)	405(99.3)	0.745
	No	8(0.9)	3(0.7)	
Apply enough soap to cover all hand surfaces	Yes	658(75.3)	294(72.1)	0.218
	No	216(24.7)	114(27.9)	
Rub hands palm to palm	Yes	423(48.4)	184(45.1)	0.270
	No	451(51.6)	224(54.9)	
Right palm over left dorsum with interlaced fingers and vice versa	Yes	141(16.1)	35(8.6)	p-value < 0.001*
	No	733(83.9)	373(91.4)	
Palm to palm with fingers interlaced	Yes	142(16.2)	55(13.5)	0.201
	No	732(83.8)	353(86.5)	
Backs of fingers to opposing palm with fingers interlocked	Yes	58(6.6)	13(3.2)	0.012*
	No	816(93.4)	395(96.8)	
Rotational rubbing of left thumb clasped in the right palm and vice versa	Yes	48(5.5)	16(3.9)	0.229
	No	826(94.5)	392(96.1)	
Rotational rubbing, backward and forward with clasped fingers of the right hand in left palm and vice versa	Yes	51(5.8)	14(3.4)	0.068
	No	823(94.2)	394(96.6)	
Rinse hands with water	Yes	760(87.0)	342(83.8)	0.133
	No	114(13.0)	66(16.2)	
Use towel/ clean item to turn off the faucet	Yes	131(15.0)	54(13.2)	0.405
	No	743(85.0)	354(86.8)	
Overall Compliance Level				
Full compliance with hand washing	Yes	8 (0.9)	3(0.7)	0.745
	No	866(99.1)	405 (99.3)	
Partial compliance to hand washing	Yes	27 (3.1)	10 (2.5)	0.525
	No	847(96.9)	398 (97.5)	
Minimal compliance with hand washing	Yes	444(50.8)	161 (39.5)	p-value < 0.001*
	No	430 (49.2)	247 (60.5)	
Non-compliance to hand washing	Yes	162 (18.5)	96 (23.5)	0.038*
	No	712 (81.5)	312 (76.5)	

*p-value < 0.05

Discussion

Good hand washing compliance with soap and water for at least 20 seconds is effective in preventing the spread of COVID-19. In response to the current COVID-19 pandemic, the WHO recommended the establishment of hand washing stations located around public areas where people can wash their hands frequently, and somehow prevent themselves from contracting COVID-19 disease. However, public compliance towards the recommended hand washing techniques in these stations is still unknown in many low-income countries, including Ethiopia. Currently, it is now customary to see hand washing facilities in different public areas, yet the question of "Do the people comply with the correct hand washing techniques?" is a fundamental unanswered question. To the best of our knowledge, there is no study done in Sub-Saharan Africa and in Ethiopia, particularly regarding the assessment on the level of hand washing compliance among hospital visitors during this COVID-19 pandemic.

In the current study, only 0.9% of the visitors were observed for full compliance with the recommended hand washing steps. The foremost, very low compliance among hospital visitors is of great concern, and thus highlight the urgent need for interventions from the health authorities, healthcare professionals, and hospital administrators. Also, the findings underscore the need for further behavioral change strategies in increasing the hospital visitors' compliance, not merely with just the provision of hand washing stations. The level of compliance reported in this study was almost comparable from the baseline assessments in Miami 0.52% [15], and São Paulo, Brazil 1.7% [16]. Despite the similarities between the aforementioned studies, our finding is still extremely low, hence, the present study is conducted during this time of COVID-19 pandemic. Conversely, a relatively higher hand washing adherence by visitors was observed in the United Kingdom 25% [17] and 10.6% [18].

In this study, 46.5% of the study participants wash their hands for less than 20 seconds. According to the CDC, UNICEF, and other experts, a 20-second hand washing is recommended. However, the WHO recommends 40-60 seconds, and this encompasses the entire hand washing process from wetting the hands and applying soap, until the hands are fully dried, whereas, the 20-second recommendation focuses only on the process of scrubbing hands with soap [7, 8, 22, 24]. The challenges behind keeping this duration in public stations need further attention. Some suggest that if people have trouble keeping track, humming the entire "Happy Birthday" song twice before rinsing can help [23]. Failing to meet the proper hand washing techniques and accurate

duration can have serious consequences in controlling COVID-19 in counties like Ethiopia, where the healthcare system is not yet matured enough to handle such a pandemic.

Previously conducted studies came up with promising results – audible reminders [19], stylized human watching eye presentations [20], and active methods to encourage hand washing, led to marked improvements on compliance [21]. On the other hand, improving hand washing thru knowledge alone is typically insufficient to change hand washing behaviors [22]. In many cases, humans modify their behavior in a socially desirable way when being observed by others. Hence, applying this basic idea to public hand washing stations may be helpful to positively influence the behavior of individuals [20, 23].

In Ethiopia, hand washing awareness-raising campaigns by some public figures, national radio and television public service announcements, and social media graphics were intensified to raise awareness about the benefits of hand washing in relation to combating COVID-19. Unfortunately, regardless of these efforts mentioned, there are still little improvements in the hand washing habits of the community. Answering the reasons and factors behind the current low compliance is beyond the scope of this study.

There are few limitations in our study. First, we did not use the electronic hand washing observation system which offers a greater advantage over human observations. Second, despite our assumption, the presence of the data collectors may have influenced the hospital visitors' hand washing activities, and it is likely to have a 'Hawthorne effect'. Third, selectivity may be possible as the data collectors do not collect data from all people entering and leaving the health facility.

Conclusion

In conclusion, compliance with proper hand washing among the visitors of Goba Referral Hospital in that specific time period is alarmingly low. A significant number of hospital visitors did not wash their hands properly or washed them in less effective ways. In this era of COVID-19 pandemic, mandating compliance towards proper hand washing to all hospital visitors before entering the hospital and leaving it should be of paramount importance.

220 **Acknowledgments**

221 The authors would like to thank Madda Walabu University Goba referral hospital public health department staff
222 for providing their unreserved support. We would like to acknowledge Goba Referral Hospital staff for their
223 support at a time of data collection.

224

225

References

1. Centers for Disease Control and Prevention (CDC). Hand Hygiene in Healthcare Settings. 2019. Retrieved from <https://www.cdc.gov/handhygiene/index.html> (Accessed May 12, 2020).
2. World Health Organization (WHO). WHO interim recommendation on hand obligatory hand hygiene against transmission of COVID-19. 2020. Retrieved from <https://www.who.int/docs/default-source/inaugural-who-partners-forum/who-interim-recommendation-on-obligatory-hand-hygiene-against-transmission-of-covid-19.pdf> (Accessed May 12, 2020).
3. UNICEF. Everything you need to know about washing your hands to protect against coronavirus (COVID-19) Retrieved from <https://www.unicef.org/coronavirus/everything-you-need-know-about-washing-your-hands-protect-against-coronavirus-covid-19> (Accessed May 12, 2020)
4. COVID-19 and water, sanitation and hygiene message for policy makers. Retrieved from <https://www.sanitationandwaterforall.org/sites/default/files/2020-04/SWA%20messages%20on%20COVID-19.pdf> (Accessed May 12, 2020)
5. Rabie T, Curtis V. Hand washing and risk of respiratory infections: a quantitative systematic review. Tropical medicine and international health. 2006; 11(3):258-67.
6. WHO. Coronavirus disease (COVID-19) advice for the public. Retrieved from <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public> (Accessed May 12, 2020).
7. WHO. How to hand wash poster .2019. Retrieved from https://www.who.int/gpsc/5may/How_To_HandWash_Poster.pdf?ua=1 (Accessed May 12, 2020).
8. Centers for Disease Control and Prevention (CDC). Hand washing: Clean Hands Save Lives. Retrieved from <https://www.cdc.gov/handwashing/when-how-hand-washing.html> (Accessed May 12, 2020).
9. Alzyood M, Jackson D, Aveyard H, Brooke J. COVID-19 reinforces the importance of handwashing. J Clin Nurs. 2020. <https://doi.org/10.1111/jocn.15313>
10. Allegranzi B, Bagheri Nejad S, Combescure C, Graafmans W, Attar H, Donaldson L, et al. Burden of endemic health-care-associated infection in developing countries: systematic review and metaanalysis. Lancet. 2011;377(9761):228-41.
11. Loftus MJ, Guitart C, Tartari E, Stewardson AJ, Amer F, Bellissimo-Rodrigues F, et al. Hand hygiene in low- and middle-income countries. Int J Infect Dis. 2019;86:25-30.
12. Pogrebna G, Kharlamov A. The Impact of Cross-Cultural Differences in Hand washing Patterns on the COVID-19 Outbreak Magnitude (preprint).
13. Retrieved from: <https://www.afro.who.int/news/first-case-covid-19-confirmed-ethiopia> (Accessed May 23, 2020).
14. National COVID-19 management guideline handbook. FMOH. April 2020. Retrieved from <http://www.moh.gov.et/ejcc/sites/default/files/2020-04/COVID%2019%20Handbook%20for%20health%20professionals%20FMOH%202020.pdf> (Accessed May 12, 2020).
15. Birnbach DJ, Nevo I, Barnes S, Fitzpatrick M, Rosen LF, Everett-Thomas R, Sanko JS, Arheart KL. Do hospital visitors wash their hands? Assessing the use of alcohol-based hand sanitizer in a hospital lobby. American journal of infection control. 2012; 40(4):340-3.
16. Vaidotas M, Yokota PK, Marra AR, Camargo TZ, da Silva Victor E, Gysi DM, Leal F, dos Santos OF, Edmond MB. Measuring hand hygiene compliance rates at hospital entrances. American journal of infection control. 2015; 43(7):694-6.
17. Wolfe R, O'Neill E. Hand hygiene compliance by visitors to hospitals—can we do better?. American journal of infection control. 2012; 40(9):899.
18. Fakhry M, Hanna GB, Anderson O, Holmes A, Nathwani D. Effectiveness of an audible reminder on hand hygiene adherence. American journal of infection control. 2012; 40(4):320-3.
19. Fakhry M, Hanna G, Anderson O, Holmes A, Nathwani D. Effectiveness of an audible reminder on hand hygiene adherence. American Journal of Infection Control. 2012. 40:320-3
20. Pfattheicher S, Strauch C, Diefenbacher S, Schnuerch R. A field study on watching eyes and hand hygiene compliance in a public restroom. Journal of Applied Social Psychology. 2018;48(4):188-94.
21. Willison-Parry T, Haidar E, Martini L, Coates A. Hand washing adherence by visitors is poor: Is there a simple solution? American Journal of Infection Control. 2013; 41:928-9.

- 279 22. Seimetz E, Kumar S, Mosler H-J. Effects of an awareness raising campaign on intention and behavioural
280 determinants for hand washing. Health Educ Res. 2016;31(2):109-120.
- 281 23. Global Hand washing Partnership. Retrieved from [https://globalhand washing.org/about-hand washing/faqs/](https://globalhandwashing.org/about-handwashing/faqs/)
282 (Accessed May 12, 2020).
- 283 24. How Much Time You Spend Washing Your Hands Makes a Difference. Retrieved from
284 <https://www.healthline.com/health/how-long-should-you-wash-your-hands> (Accessed May 12, 2020).

285

286

287 **Supporting Information**

288 **Additional File 1:** The Association between Hand Washing Compliance during the Weekdays in Madda Walabu
 289 University Goba Referral Hospital, Southeast Ethiopia, May 2020 (n=1,282)

290 **Additional File 2:** The association between hand washing compliance and observation sessions in Madda
 291 Walabu University Goba Referral Hospital, Southeast Ethiopia, May 2020 (n=1,282)

292

Hand Washing Compliance

Full compliance to HW

0.9

Partial compliance to HW

2.9

Minimal compliance to HW

47.2

Non-compliance to HW

20.1

■ Percent

0 5 10 15 20 25 30 35 40 45 50

5.5%

backs of fingers to opposing palms with fingers interlocked

5.1%

rotational rubbing, backwards and forwards with clasped fingers of right hand in left palm and vice versa.

5.0%

rotational rubbing of left thumb clasped in right palm and vice versa