Association Between ACEIs or ARBs Use and Clinical Outcomes in COVID-19 Patients: A Systematic Review and Meta-analysis

Carlos Diaz-Arocutipa, MD^{1,2}; Jose Saucedo-Chinchay, MD²; Adrian V. Hernandez, MD, PhD^{1,3}

¹Vicerrectorado de Investigación, Universidad San Ignacio de Loyola, Lima, Peru

²Programa de Atencion Domiciliaria – EsSalud, Lima, Peru

³Health Outcomes, Policy, and Evidence Synthesis (HOPES) Group, University of Connecticut School of Pharmacy, Storrs, CT, USA

Corresponding author:

Carlos Diaz-Arocutipa, MD

Vicerrectorado de Investigación, Universidad San Ignacio de Loyola, Av. La Fontana 550, La Molina, Lima 00012, Peru

Tel: +511 994928488

Email: carlosdiaz013@gmail.com

Word count: 3274

Key points

Question: What is the association between angiotensin-converting enzyme inhibitors (ACEIs) or angiotensin receptor blockers (ARBs) use and clinical outcomes in coronavirus disease 2019 (COVID-19) patients?

Findings: In this systematic review and meta-analysis of 40 observational studies, the use of ACEIs or ARBs was not associated with higher all-cause mortality in COVID-19 patients. Additionally, ACEIs or ARBs use was independently associated with lower COVID-19 severity.

Meaning: These results support the current international guidelines to continue the use of ACEIs and ARBs in COVID-19 patients with hypertension.

Abstract

Importance: There is a controversy regarding whether or not to continue angiotensin-converting enzyme inhibitors (ACEIs) and angiotensin receptor blockers (ARBs) in patients with coronavirus disease 2019 (COVID-19).

Objective: To evaluate the association between ACEIs or ARBs use and clinical outcomes in COVID-19 patients.

Data Sources: Systematic search of the PubMed, Embase, Scopus, Web of Science, and Cochrane Central Register of Controlled Trials from database inception to May 31, 2020. We also searched the preprint servers medRxiv and SSNR for additional studies.

Study Selection: Observational studies and randomized controlled trials reporting the effect of ACEIs or ARBs use on clinical outcomes of adult patients with COVID-19.

Data Extraction and Synthesis: Risk of bias of observational studies were evaluated using the Newcastle-Ottawa Scale. Meta-analyses were performed using a random-effects models and effects expressed as Odds ratios (OR) and mean differences with their 95% confidence interval (95%CI). If available, adjusted effects were pooled.

Main Outcomes and Measures: The primary outcome was all-cause mortality and secondary outcomes were COVID-19 severity, hospital discharge, hospitalization, intensive care unit admission, mechanical ventilation, length of hospital stay, and troponin, creatinine, procalcitonin, C-reactive protein (CRP), interleukin-6 (IL-6), and D-dimer levels.

Results: 40 studies (21 cross-sectional, two case-control, and 17 cohorts) involving 50615 patients were included. ACEIs or ARBs use was not associated with all-cause mortality overall (OR 1.11, 95%CI 0.77-1.60, $p=0.56$), in subgroups by study design and using adjusted effects. ACEI or ARB use was independently associated with lower COVID-19 severity (aOR 0.56, 95%CI 0.37-0.87, $p<0.01$). No significant associations were found between ACEIs or ARBs use and hospital discharge, hospitalization, mechanical ventilation, length of hospital stay, and biomarkers.

Conclusions and Relevance: ACEIs or ARBs use was not associated with higher all-cause mortality in COVID-19. However, ACEI or ARB use was independently associated with lower COVID-19 severity. Our results support the current international guidelines to continue the use of ACEIs and ARBs in COVID-19 patients with hypertension.

Introduction

Coronavirus disease 2019 (COVID-19) is a global pandemic involving more than 185 countries.¹ This disease is caused by the severe acute respiratory coronavirus-2 (SARS-CoV-2) and was first detected in Wuhan, China in December, 2019.² The infection by SARS-CoV-2 is caused by the binding of the viral spike glycoprotein to the angiotensin-converting enzyme 2 (ACE2).³ In humans, ACE2 is ubiquitously expressed with predominance in the lungs, heart, kidneys, and vascular system.⁴ ACE2 is a major component of the renin-angiotensin system (RAS) that acts as a carboxypeptidase converting angiotensin II (Ang II) into angiotensin 1-7 (Ang 1-7).⁵ The degradation of Ang II by ACE2 regulates negatively the RAS activation and attenuates the vasoconstrictive, pro-oxidant, pro-fibrotic, and pro-inflammatory actions mediated by Ang II.⁶ The RAS is considered a complex system that requires a balanced interplay between two counter-regulatory axes (ACE2/Ang 1-7/MasR and ACE/Ang II/AT₁R).⁶ Moreover, there is evidence that ACE2 has a protective physiological role in many organs, including lungs and heart, and its imbalance can be lead to disease states.⁵

Angiotensin-converting enzyme inhibitors (ACEIs) and angiotensin receptor blockers (ARBs) are widely used in clinical practice for the treatment of hypertension, heart failure, and diabetic nephropathy.⁷ It has been hypothesized that these drugs could increase the risk of infection and severity in COVID-19 patients.⁸⁻¹⁰ However, major international cardiology societies have recommended not to discontinue the use of ACEIs and ARBs in COVID-19 patients with hypertension due to a lack of clinical evidence.¹¹

Recently, several studies that evaluated the effect of RAS inhibitors on COVID-19 have been published. Therefore, we performed a systematic review and meta-analysis to evaluate the association between ACEIs or ARBs use and clinical outcomes in COVID-19 patients.

Methods

This review was reported according to the MOOSE (Meta-analysis of Observational Studies in Epidemiology) guidelines (eTable 1)¹² and was registered in PROSPERO database (CRD42020177848).

Search strategy

We searched PubMed, Embase, Scopus, Web of Science, and Cochrane Central Register of Controlled Trials. The preprint servers medRxiv and SSNR were also searched. The search was conducted from inception to April 4, 2020, and updated on May 31, 2020. The complete search strategy is available in eTable 2. There were no restrictions on language. We conducted hand searches of reference lists of included studies and relevant reviews articles to identify further eligible studies. Additionally, clinicaltrials.gov registry was searched for finished as well as ongoing randomized controlled trials (RCTs).

Eligibility criteria

We included observational studies and RCTs that evaluated the association between ACEIs or ARBs use and at least one clinical outcome in COVID-19 patients (≥18 years) diagnosed by reverse transcription-polymerase chain reaction. Case reports, case series, systematic reviews, narrative reviews, commentaries, and abstracts were excluded.

Study Selection

Two authors (CDA and JSC) downloaded all articles from electronic search to EndNote X8 and duplicates were removed. Titles and abstracts were independently screened by two authors (CDA and JSC) to identify potentially relevant studies. Two authors (CDA and JSC) independently screened the full-text and registered reasons for the exclusion. Any disagreement was resolved by consensus.

1

2 **Outcomes**

3 The primary outcome was all-cause mortality and the secondary outcomes were COVID-
4 19 severity, hospital discharge, hospitalization, intensive care unit (ICU) admission,
5 mechanical ventilation, length of hospital stay, troponin, creatinine, procalcitonin, C-
6 reactive protein (CRP), interleukin-6 (IL-6) and D-dimer. We used author-reported
7 definitions for all outcomes.

8

9 **Data Extraction**

10 Information from each study was independently extracted by two authors (CDA and JSC)
11 using a standardized data extraction form and any disagreement was resolved by
12 consensus. If additional data was needed, we contacted the corresponding author through
13 email. We extracted the following data: author, publication year, country, study design,
14 sample size, eligibility criteria, age, sex, comorbidities, ACEIs or ARBs use, and primary
15 and secondary outcomes. If available, unadjusted and adjusted effect measures were also
16 extracted.

17

18 **Risk of bias assessment**

19 The Newcastle-Ottawa Scale (NOS) was used to assess the risk of bias in case-control
20 and cohort studies.¹³ Each study was classified in the following groups: low risk of bias (8-
21 9 points), moderate risk of bias (5-7 points), and high risk of bias (0-4 points). For cross-
22 sectional studies, we used an adapted version of NOS¹⁴ and each study was assigned in
23 the following groups: low risk of bias (8-10 points), moderate risk of bias (5-7 points), and
24 high risk of bias (0-4 points). The risk of bias was independently assessed by two authors
25 (CDA and JSC) and any disagreement was resolved by consensus.

26

Statistical analysis

We performed all meta-analyses using random-effects models. Between-study variance was estimated using the Paule-Mandel estimator.¹⁵ We pooled odds ratios (OR) and mean differences (MD) with their 95% confidence intervals (95%CI) for binary and continuous outcomes, respectively. In case of studies have only reported median and interquartile range, then mean and standard deviation were estimated using the method published by Wan et al.¹⁶ As exploratory analyses, we combined adjusted effects from studies that included a minimum set of confounding variables (age, sex, and cardiovascular comorbidities) in their multivariate models. Heterogeneity among studies was evaluated using the chi-squared test (threshold $p < 0.10$) and I^2 statistic. Heterogeneity was defined as low if $I^2 < 30\%$, moderate if $I^2 = 30-60\%$, and high if $I^2 > 60\%$. Funnel plots were used to evaluate publication bias and the Egger's test was performed to measure asymmetry of funnel plots only if 10 or more studies were included.¹⁷ Subgroup analyses were conducted according to study design (cross-sectional vs cohort). In post hoc sensitivity analyses, we adjusted all 95%CI using the Hartung-Knapp method to address possible type I error with the conventional random-effects approach.¹⁸ All meta-analyses were conducted using the *meta* package from R 3.6.3. A two-tailed $p < 0.05$ was considered as statistically significant.

Results

Study selection

Our search strategy identified initially 110 articles. After removal of duplicates, 87 articles remained. After screening of studies by title/abstract, 36 articles were excluded. After full-text revision of 51 articles, 11 articles were excluded. A total of 40 studies were selected for analysis (21 cross-sectional, two case-control, and 17 cohort studies) (Figure 1). Only one contacted author provided additional information on mortality.¹⁹

1

2 **Study Characteristics**

3 Main characteristics of the 40 included studies (n=50615) were summarized in Table 1.
 4 The exposure variable in almost all studies was the chronic use of ACEIs and ARBs (i.e.
 5 before hospital admission) as registered in medical records, although in two studies^{20,21} it
 6 was defined as in-hospital use. Also, only six studies²²⁻²⁷ reported that ACEIs and ARBs
 7 were not discontinued during hospitalization. The definition of outcomes was the same in
 8 almost all included studies. In contrast, the definition of COVID-19 severity was very
 9 heterogeneous across the studies due to different clinical guidelines used for management
 10 of COVID-19 in each country. The most common criteria for COVID-19 severity was
 11 critical/severe vs mild/moderate which was used in five studies^{24,28-31} (Table 1).

12 Our search in clinicaltrials.gov identified 16 registered RCTs (eTable 3), of which
 13 six evaluate the impact of continuation or discontinuation of ACEIs and ARBs on COVID-
 14 19 outcomes and four placebo-controlled trials assess the efficacy of ARB (losartan and
 15 valsartan) and ACEI (ramipril) in COVID-19 patients who are not previously taking a RAS
 16 inhibitor.

17

18 **Risk of bias assessment**

19 Almost all cross-sectional studies had moderate risk of bias, all case-control studies had
 20 low risk of bias, and 9 of 17 cohort studies had low risk of bias (eTables 4, 5, and 6). None
 21 of studies was scored as high risk of bias.

22

23 **All-cause mortality**

24 In 22 studies (11 cross-sectional and 11 cohorts, n=23059), the use of ACEIs or ARBs was
 25 not associated with higher odds of all-cause mortality (OR 1.11, 95%CI 0.77-1.60, p=0.56)
 26 and heterogeneity was high among studies (Figure 2). In the subgroup analysis by study

design, ACEIs or ARBs use were not associated with all-cause mortality in 11 cross-sectional studies (OR 0.86, 95%CI 0.65-1.15, $p=0.32$) and 11 cohort studies (OR 1.30, 95%CI 0.71-2.38, $p=0.40$) (Figure 2). The funnel plot did not show asymmetry and the Egger's test was not significant ($p=0.39$) (eFigure 1). ACEI use was not associated with all-cause mortality (OR 1.18, 95%CI 0.83-1.66, $p=0.35$) (eFigure 2), but ARB use was associated with increased odds of all-cause mortality (OR 1.79, 95%CI 1.07-3.00, $p=0.03$) (eFigure 3).

Six studies^{21,32-36} reported adjusted effect measures of association between ACEIs/ARBs use and all-cause mortality (eTable 7). The pooled estimate of three studies^{21,35,36} with similar adjusted variables (age, sex, and cardiovascular comorbidities) found that ACEIs or ARBs use was not associated with all-cause mortality (aHR 0.83, 95%CI 0.49-1.38, $p=0.47$) (eFigure 4). Furthermore, the pooled estimate of ACEI studies^{32,34,35} (aHR 0.97, 95%CI 0.83-1.13, $p=0.67$) and ARB studies^{32,34,35} (aHR 1.14, 95%CI 0.98-1.34, $p=0.08$) showed no association either (eFigure 5 and 6).

Secondary Outcomes

COVID-19 severity

In 18 studies (11 cross-sectional, two case-control, and five cohorts, $n=11870$), the use of ACEIs or ARBs was not associated with COVID-19 severity (OR 0.79, 95%CI 0.59-1.07, $p=0.13$) and showed high heterogeneity among studies (Figure 3). The funnel plot showed asymmetry, suggesting publication bias which was confirmed by the Egger's test ($p<0.01$) (eFigure 7). Subgroup analysis by study design showed that ACEIs or ARBs use was only associated with lower COVID-19 severity in five cohort studies (OR 0.61, 95%CI 0.39-0.95, $p=0.03$) (Figure 3). In contrast, the use of ACEI (OR 1.10, 95%CI 0.55-2.18, $p=0.79$) and ARB (OR 1.00, 95%CI 0.77-1.29, $p=0.98$) separately were not associated with COVID-19 severity (eFigure 8 and 9).

The pooled adjusted estimate of four studies^{23,24,37,38} showed that the ACEIs or ARBs use (aOR 0.56, 95%CI 0.37-0.87, $p < 0.01$) was independently associated with lower COVID-19 severity (eFigure 10). However, adjusted estimates of ACEI (aOR 0.66, 95%CI 0.37-1.18, $p = 0.15$) and ARB (aOR 0.97, 95%CI 0.79-1.20, $p = 0.81$) use separately were not associated with COVID-19 severity (eFigure 11 and 12).

Hospital discharge

In three studies^{29,31,39} (two cross-sectional and one cohort, $n = 301$), the use of ACEIs or ARBs was not associated with hospital discharge (OR 2.27, 95%CI 0.96-5.35, $p = 0.06$) (eFigure 13).

Hospitalization

In four studies^{19,32,39,40} (one cross-sectional and three cohorts, $n = 5048$), the use of ACEIs or ARBs was not associated with hospitalization (OR 1.83, 95%CI 0.95-3.52, $p = 0.07$) (eFigure 14). Likewise, the use of ACEI (OR 1.63, 95%CI 0.94-2.83, $p = 0.08$) and ARB (OR 1.48, 95%CI 0.95-2.31, $p = 0.08$) were not associated with hospitalization (eFigure 15 and 16).

ICU admission

In six studies^{19,40-44} (two cross-sectional and four cohorts, $n = 8884$), the use of ACEIs or ARBs was associated with increased odds of ICU admission (OR 1.45, 95%CI 1.17-1.80, $p < 0.01$) (eFigure 17). In contrast, the use of ACEI (OR 1.16, 95%CI 0.72-1.86, $p = 0.53$) and ARB (OR 1.26, 95%CI 0.87-1.83, $p = 0.23$) by separate were not associated with ICU admission (eFigure 18 and 19).

Mechanical Ventilation

In seven studies^{21,28,29,33,40,44,45} (three cross-sectional and four cohorts, n=6533), the use of ACEIs or ARBs was not associated with mechanical ventilation (OR 1.39, 95%CI 0.99-1.94, p=0.06) (eFigure 20).

Length of hospital stay

In five studies^{26,29,31,39,46} (four cross-sectional and one cohort, n=699), the use of ACEIs or ARBs was not associated with length of hospital stay (MD -0.96 days, 95%CI -2.50 to 0.57, p=0.22) (eFigure 21).

Troponin level

In four studies^{26-28,31} (two cross-sectional and two cohorts, n=580), the use of ACEIs or ARBs was not associated with troponin level (MD -0.01 µg/L, 95%CI -0.04 to 0.02, p=0.37) (eFigure 22).

Creatinine level

In six studies^{24,26-28,31,47} (two cross-sectional and four cohorts, n=716), the use of ACEIs or ARBs was not associated with creatinine level (MD -0.58 µmol/L, 95%CI -8.72 to 7.56, p=0.89) (eFigure 23).

Procalcitonin level

In five studies^{26-28,31,47} (two cross-sectional and three cohorts, n=651), the use of ACEIs or ARBs was not associated with procalcitonin level (MD -0.02 ng/mL, 95%CI -0.05 to 0.01, p=0.21) (eFigure 24).

CRP level

In five studies^{26,28,29,31,47} (two cross-sectional and three cohorts, n=709), the use of ACEIs or ARBs was not associated with CRP level (MD -6.39 mg/L, 95%CI -16.19 to 3.41, p=0.20) (eFigure 25).

IL-6 level

In four studies^{26,27,31,47} (two cross-sectional and two cohorts, n=601), the use of ACEIs or ARBs was not associated with IL-6 level (MD -4.41 pg/mL, 95%CI -13.24 to 4.42, p=0.33) (eFigure 26).

D-dimer level

In six studies^{26-29,31,47} (two cross-sectional and one cohort, n=751), the use of ACEIs or ARBs was not associated with D-dimer level (MD -0.91 nmol/L, 95%CI -2.77 to 0.94, p=0.33) (eFigure 27).

Sensitivity analyses

The results of the sensitivity analyses are reported in the eTable 8. Overall, the results showed that ACEIs or ARBs use was independently with lower COVID-19 severity, ARB use was independently associated with higher mortality, and ACEIs or ARBs use was associated with higher ICU admission.

Discussion

We found that the use of ACEIs or ARBs was not significantly associated with all-cause mortality in COVID-19 patients, and when analyzed by study design or when using adjusted effects. In contrast, ACEIs or ARBs use was independently associated with lower COVID-19 severity. Although ACEIs or ARBs use was associated with an increased odds of ICU admission, this effect disappeared when ACEIs and ARBs were analyzed

individually. No significant associations were found between ACEIs or ARBs use and other clinical outcomes or biomarkers. Risk of bias was low or moderate across studies.

It has been proposed that RAS play a crucial role in the pathogenesis of infection by SARS-CoV-2, since it uses the ACE2 receptor to enter into cells, with the subsequent downregulation of this surface protein.⁵ The reduction of ACE2 expression in infected cells can lead to a tissue and systemic RAS imbalance with a predominance of the dangerous ACE/Ang II/AT₁R axis.⁵ This phenomenon can be particularly harmful in the elderly population since they have already a lower level of ACE2 expression compared to young people.⁴⁸ This could partly explain the higher mortality observed in older patients with COVID-19.⁴⁹ Recent evidence from a Chinese cohort of 12 COVID-19 patients showed that circulating Ang II levels were markedly elevated compared to healthy controls and linearly associated with viral load and lung injury.⁵⁰ Moreover, in an animal experiment of acute lung injury induced by acid, the SARS-CoV spike protein enhances the pulmonary Ang II levels and lung injury severity.⁵¹ Altogether, these data suggest that SARS-CoV-2 can mediate the damage to lungs and possibly to other organs through the absence of degradation of Ang II. Therefore, RAS modulators such as ACEIs and ARBs can be used as potential therapeutic agents in COVID-19 patients. This is currently under investigation in several ongoing clinical trials.

In general, we found that the use of ACEIs or ARBs had a neutral effect on all-cause mortality and other clinical outcomes in COVID-19 patients. Two studies^{21,52} with larger samples and adjustment for confounders reported a significant reduction of all-cause mortality and severity in these patients. Nowadays, there is controversy regarding the use of ACEIs and ARBs in patients with COVID-19 and hypertension. Initially it was suggested that the use of these drugs could increase ACE2 expression; however, there is conflicting evidence about the effect of ACEIs and ARBs on ACE2 tissue expression in animal models.⁵³ Besides, studies in humans showed no effect of ACEIs and ARBs

administration on ACE2 protein levels in urine and plasma.^{54,55} Furthermore, a recent Mendelian randomization study revealed a lack of association between genetically proxy ACE inhibition and lung ACE2 expression or circulating ACE2 levels.⁵⁶ Likewise, in a study on human myocardial samples, there was no significant difference in ACE2 expression in tissue samples with and without exposure to ACEI.⁵⁷ Overall, these findings suggest that ACEIs and ARBs are unlikely to raise ACE2 in humans. Thus, it seems reasonable that ACEIs and ARBs could exert its effect on COVID-19 mainly through inhibition of the ACE/Ang II/AT₁R axis.

The lung is the target organ in COVID-19; however, other organs may potentially be involved. A recent study reported that acute cardiac injury, manifested as elevated troponin levels, was present in 20% of COVID-19 patients and was independently associated with worse outcomes.⁵⁸ Although the pathophysiological basis of this finding is not fully understood, it has been proposed that SARS-CoV-2 can cause cardiac injury through several mechanisms: direct viral damage, systemic inflammatory response, microangiopathy, and myocardial infarction.⁵⁹ Similarly, acute kidney injury was observed in up to 27% of COVID-19 patients,⁶⁰ this is probably related to alterations in renal microvasculature, kidney cell viral infection, and systemic inflammation.⁶¹ Dysregulation of the immune system is key in the pathogenesis of COVID-19 leading, in some cases, to an overproduction of pro-inflammatory cytokines (interleukin-6, interleukin-1 β , and tumor necrosis factor-alpha) resulting in what has been called a cytokine storm.⁶² Likewise, the procoagulant-anticoagulant balance has been found to be impaired in COVID-19, leading to formation of microthrombi and marked elevation of D-dimer.⁶³

A recent meta-analysis found that elevation of troponin, creatinine, D-dimer, and procalcitonin were significantly associated with a higher risk of critical disease or mortality in COVID-19 patients.⁶⁴ The RAS imbalance and the loss of ACE2 expression observed in COVID-19, with the subsequent reduction of Ang 1-7 and elevation of Ang II levels, can

contribute to the tissue and systemic damage caused by SARS-CoV-2. Thus, considering that RAS inhibitors are capable of regulating both tissue and systemic RAS, it has been suggested that could have a beneficial effect in COVID-19. However, our study did not find a significant association of ACEIs or ARBs use on troponin, inflammatory markers (procalcitonin, CRP, and IL-6), creatinine, and D-dimer levels in COVID-19 patients. Further research is needed to clarify the potential therapeutic role of RAS inhibitors on multiorgan dysfunction associated with COVID-19.

We excluded two large observational studies by Mehra et al.⁶⁵ (n=8910) and by Mehra et al.⁶⁶ (n=96032) because of several concerns of the quality of their registry data in open letters by researchers worldwide⁶⁷ and acknowledged by the two journals in expressions of concern.^{68,69}

There are three previous systematic reviews examining the effects of ACEI/ARB use on COVID-19 patients. Zhang et al.⁷⁰ found that ACEI/ARB exposure was not associated with a higher risk of severe infection or mortality. However, only 12 studies and unadjusted estimates were combined. Guo et al.⁷¹ showed that ACEI/ARB use was associated with lower mortality in COVID-19 patients although only included six studies were included. Mackey et al.⁷² only conducted a narrative synthesis of 14 studies, concluding that there is no evidence of association between ACEI/ARB use with more severe COVID-19 disease. Compared to these reviews, our study included 40 studies and evaluated 13 outcomes. Additionally, to our knowledge, our review is the first that pooled adjusted effect estimates for mortality and COVID-19 severity.

Our study has some limitations. First, given most of the studies did not use adjusted effects, there is an increased risk of bias in their pooled effect measures. Thus, these results should be considered with caution. However, we also reported meta-analyses of adjusted estimates of a few available studies. Second, the majority of the included studies were of cross-sectional design, thus causality cannot be concluded due to

the methodological limitations of this design. Third, heterogeneity was high in most of the evaluated outcomes. Possible reasons for heterogeneity include sample size, differences in outcome definitions, heterogeneous population, among others. Fourth, given that discontinuation of ACEIs or ARBs during hospitalization was not reported consistently across studies, this could influence the significance of pooled estimates. Finally, we could not adequately evaluate the effects of ACEIs and ARBs by separate, since were mainly reported as aggregate due to scarcity of studies.

Conclusions

In conclusion, the use of ACEIs or ARBs was not associated with higher all-cause mortality in COVID-19 patients, based on the meta-analysis of cross-sectional and cohort studies and also using adjusted effects. ACEI or ARB use was independently associated with lower COVID-19 severity. Also, there was no evidence of association between ACEIs or ARBs use and nearly all secondary clinical outcomes and biomarkers. Although these results are not conclusive, our review supports current international guidelines to continue the use of RAS inhibitors in COVID-19 patients with hypertension. More studies are needed to determine the potential beneficial effect of ACEIs and ARBs in patients with COVID-19.

Author Contributions: Diaz-Arocutipa had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Diaz-Arocutipa. *Acquisition, analysis, or interpretation of data:*

Diaz-Arocutipa, Saucedo-Chinchay, Hernandez. *Drafting of the manuscript:* Diaz-

Arocutipa. *Critical revision of the manuscript for important intellectual content:* Diaz-

Arocutipa, Saucedo-Chinchay, Hernandez. *Statistical analysis:* Diaz-Arocutipa,

Hernandez. *Administrative, technical, or material support:* Diaz-Arocutipa. *Study supervision:* Diaz-Arocutipa, Hernandez.

Funding: None.

Conflict of Interest Disclosures: None.

REFERENCES

1. COVID-19 Dashboard by the Center for Systems Science and Engineering at Johns Hopkins University. Available at <https://coronavirus.jhu.edu/map.html> (accessed May 30, 2020).
2. Zhou P, Yang X-L, Wang X-G, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020;579(7798):270-273. doi:10.1038/s41586-020-2012-7
3. Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. *Science*. 2020;367(6485):1444-1448. doi:10.1126/science.abb2762
4. South AM, Diz DI, Chappell MC. COVID-19, ACE2, and the cardiovascular consequences. *Am J Physiol Heart Circ Physiol*. 2020;318(5):H1084-h1090. doi:10.1152/ajpheart.00217.2020
5. Gheblawi M, Wang K, Viveiros A, et al. Angiotensin converting enzyme 2: SARS-CoV-2 receptor and regulator of the renin-angiotensin system. *Circ Res*. 2020. doi:10.1161/circresaha.120.317015
6. Santos RAS, Oudit GY, Verano-Braga T, Canta G, Steckelings UM, Bader M. The renin-angiotensin system: going beyond the classical paradigms. *Am J Physiol Heart Circ Physiol*. 2019;316(5):H958-h970. doi:10.1152/ajpheart.00723.2018

- 1 7. Barrios V, Coca A, Escobar C, Enrique R, Rincón LM. Use of angiotensin-converting
2 enzyme inhibitors and angiotensin receptor blockers in clinical practice. *Expert Rev*
3 *Cardiovasc Ther.* 2012;10(2):159-166. doi:10.1586/erc.11.185
- 4 8. Diaz JH. Hypothesis: angiotensin-converting enzyme inhibitors and angiotensin receptor
5 blockers may increase the risk of severe COVID-19. *J Travel Med.* 2020.
6 doi:10.1093/jtm/taaa041
- 7 9. Esler M, Esler D. Can angiotensin receptor-blocking drugs perhaps be harmful in the
8 COVID-19 pandemic? *J Hypertens.* 2020;38(5):781-782.
9 doi:10.1097/hjh.0000000000002450
- 10 10. Patel AB, Verma A. COVID-19 and Angiotensin-Converting Enzyme Inhibitors and
11 Angiotensin Receptor Blockers: What Is the Evidence? *JAMA.* 2020.
12 doi:10.1001/jama.2020.4812
- 13 11. Bavishi C, Maddox TM, Messerli FH. Coronavirus Disease 2019 (COVID-19) infection
14 and renin angiotensin system blockers. *JAMA Cardiol.* 2020.
15 doi:10.1001/jamacardio.2020.1282
- 16 12. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in
17 epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in
18 Epidemiology (MOOSE) group. *JAMA.* 2000;283(15):2008-2012.
19 doi:10.1001/jama.283.15.2008
- 20 13. Wells G, Shea B, O'Connell D, et al. The Newcastle-Ottawa scale (NOS) for assessing
21 the quality of nonrandomised studies in meta-analysis. Available at
22 http://www.ohri.ca/programs/clinical_epidemiology/oxfordasp (accessed April 4, 2020).
- 23 14. Herzog R, Alvarez-Pasquin MJ, Diaz C, Del Barrio JL, Estrada JM, Gil A. Are
24 healthcare workers' intentions to vaccinate related to their knowledge, beliefs and
25 attitudes? A systematic review. *BMC Public Health.* 2013;13:154. doi:10.1186/1471-2458-
26 13-154

15. Veroniki AA, Jackson D, Viechtbauer W, et al. Methods to estimate the between-study variance and its uncertainty in meta-analysis. *Res Synth Methods*. 2016;7(1):55-79. doi:10.1002/jrsm.1164
16. Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol*. 2014;14:135. doi:10.1186/1471-2288-14-135
17. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(7109):629-634. doi:10.1136/bmj.315.7109.629
18. Hartung J, Knapp G. A refined method for the meta-analysis of controlled clinical trials with binary outcome. *Stat Med*. 2001;20(24):3875-3889. doi:10.1002/sim.1009
19. Rentsch CT, Kidwai-Khan F, Tate JP, et al. Covid-19 testing, hospital admission, and intensive care among 2,026,227 United States veterans aged 54-75 years. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.04.09.20059964
20. Chen M, Fan Y, Wu X, et al. Clinical characteristics and risk factors for fatal outcome in patients with 2019-coronavirus infected disease (COVID-19) in Wuhan, China (2/27/2020). Available at SSRN: <https://ssrn.com/abstract=3546069>. doi:10.2139/ssrn.3546069
21. Zhang P, Zhu L, Cai J, et al. Association of inpatient use of angiotensin converting enzyme inhibitors and angiotensin II receptor blockers with mortality among patients with hypertension hospitalized with COVID-19. *Circ Res*. 2020. doi:10.1161/CIRCRESAHA.120.317134
22. Ashraf MA, Shokouhi N, Shirali E, et al. COVID-19 in Iran, a comprehensive investigation from exposure to treatment outcomes. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.04.20.20072421
23. De Spiegeleer A, Bronselaer A, Teo JT, et al. The effects of ARBs, ACEIs and statins on clinical outcomes of COVID-19 infection among nursing home residents. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.05.11.20096347

24. Feng Z, Li J, Yao S, et al. The use of adjuvant therapy in preventing progression to severe pneumonia in patients with Coronavirus Disease 2019: a multicenter data analysis. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.04.08.20057539
25. Guo T, Fan Y, Chen M, et al. Cardiovascular implications of fatal outcomes of patients with Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol.* 2020. doi:10.1001/jamacardio.2020.1017
26. Li J, Wang X, Chen J, Zhang H, Deng A. Association of renin-angiotensin system inhibitors with severity or risk of death in patients with hypertension hospitalized for Coronavirus Disease 2019 (COVID-19) infection in Wuhan, China. *JAMA Cardiol.* 2020. doi:10.1001/jamacardio.2020.1624
27. Meng J, Xiao G, Zhang J, et al. Renin-angiotensin system inhibitors improve the clinical outcomes of COVID-19 patients with hypertension. *Emerg Microbes Infect.* 2020;9(1):757-760. doi:10.1080/22221751.2020.1746200
28. Huang Z, Cao J, Yao Y, et al. The effect of RAS blockers on the clinical characteristics of COVID-19 patients with hypertension. *Ann Transl Med.* 2020;8(7):430. doi:10.21037/atm.2020.03.229
29. Tan N-D, Qiu Y, Xing X-B, Ghosh S, Chen M-H, Mao R. Associations between angiotensin converting enzyme inhibitors and angiotensin II receptor blocker use, gastrointestinal symptoms, and mortality among patients with COVID-19. *Gastroenterology.* 2020. doi:10.1053/j.gastro.2020.05.034
30. Yan H, Valdes AM, Vijay A, et al. Role of drugs affecting the renin-angiotensin-aldosterone system on susceptibility and severity of COVID-19: a large case-control study from Zhejiang province, China. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.04.24.20077875
31. Yang G, Tan Z, Zhou L, et al. Angiotensin II receptor blockers and angiotensin-converting enzyme inhibitors usage is associated with improved inflammatory status and

- 1 clinical outcomes in COVID-19 patients with hypertension. *medRxiv [Preprint]*. 2020.
2 doi:10.1101/2020.03.31.20038935
- 3 32. Giorgi Rossi P, Marino M, Formisano D, Venturelli F, Vicentini M, Grilli R.
4 Characteristics and outcomes of a cohort of SARS-CoV-2 patients in the province of
5 Reggio Emilia, Italy. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.04.13.20063545
- 6 33. Jung S-Y, Choi JC, You S-H, Kim W-Y. Association of renin-angiotensin-aldosterone
7 system inhibitors with COVID-19-related outcomes in Korea: a nationwide population-
8 based cohort study. *Clin Infect Dis*. 2020. doi:10.1093/cid/ciaa624
- 9 34. Khera R, Clark C, Lu Y, et al. Association of angiotensin-converting enzyme inhibitors
10 and angiotensin receptor blockers with the risk of hospitalization and death in hypertensive
11 patients with Coronavirus Disease-19. *medRxiv [Preprint]*. 2020.
12 doi:10.1101/2020.05.17.20104943
- 13 35. Lee H-Y, Ahn J, Kang CK, et al. Association of angiotensin II receptor blockers and
14 angiotensin-converting enzyme inhibitors on COVID-19-related outcome (4/1/2020).
15 Available at SSRN: <https://ssrn.com/abstract=3569837>. doi:10.2139/ssrn.3569837
- 16 36. Tedeschi S, Giannella M, Bartoletti M, et al. Clinical impact of renin-angiotensin system
17 inhibitors on in-hospital mortality of patients with hypertension hospitalized for COVID-19.
18 *Clin Infect Dis*. 2020. doi:10.1093/cid/ciaa492
- 19 37. Bravi F, Flacco ME, Carradori T, et al. Predictors of severe or lethal COVID-19,
20 including angiotensin converting enzyme inhibitors and angiotensin II receptor blockers, in
21 a sample of infected Italian citizens. *medRxiv [Preprint]*. 2020.
22 doi:10.1101/2020.05.21.20109082
- 23 38. Rhee SY, Lee J, Nam H, Kyoung D-S, Kim DJ. Effects of a DPP-4 inhibitor and RAS
24 blockade on clinical outcomes of patients with diabetes and COVID-19. *medRxiv*
25 *[Preprint]*. 2020. doi:10.1101/2020.05.20.20108555

39. Zeng Z, Sha T, Zhang Y, et al. Hypertension in patients hospitalized with COVID-19 in Wuhan, China: a single-center retrospective observational study. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.04.06.20054825
40. Mehta N, Kalra A, Nowacki AS, et al. Association of use of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers with testing positive for Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol.* 2020. doi:10.1001/jamacardio.2020.1855
41. Argenziano MG, Bruce SL, Slater CL, et al. Characterization and clinical course of 1000 patients with COVID-19 in New York: retrospective case series. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.04.20.20072116
42. Benelli G, Buscarini E, Canetta C, et al. SARS-COV-2 comorbidity network and outcome in hospitalized patients in Crema, Italy. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.04.14.20053090
43. Kim L, Garg S, O'Halloran A, et al. Interim analysis of risk factors for severe outcomes among a cohort of hospitalized adults identified through the U.S. Coronavirus Disease 2019 (COVID-19)-associated Hospitalization Surveillance Network (COVID-NET). *medRxiv [Preprint]*. 2020. doi:10.1101/2020.05.18.20103390
44. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting characteristics, comorbidities, and outcomes Among 5700 patients hospitalized with COVID-19 in the New York City area. *JAMA.* 2020. doi:10.1001/jama.2020.6775
45. Regina J, Papadimitriou-Olivgeris M, Burger R, et al. Epidemiology, risk factors and clinical course of SARS-CoV-2 infected patients in a Swiss university hospital: an observational retrospective study. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.05.11.20097741
46. Zhou X, Zhu J, Xu T. Clinical characteristics of coronavirus disease 2019 (COVID-19) patients with hypertension on renin-angiotensin system inhibitors. *Clin Exp Hypertens.* 2020:1-5. doi:10.1080/10641963.2020.1764018

47. Chen Y, Yang D, Cheng B, et al. Clinical characteristics and outcomes of patients with diabetes and COVID-19 in association with glucose-lowering medication. *Diabetes Care*. 2020;dc200660. doi:10.2337/dc20-0660
48. Chen J, Jiang Q, Xia X, et al. Individual Variation of the SARS-CoV2 Receptor ACE2 Gene Expression and Regulation. Preprints 2020, 2020030191.
49. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395(10229):1054-1062. doi:[https://doi.org/10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3)
50. Liu Y, Yang Y, Zhang C, et al. Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury. *Sci China Life Sci*. 2020;63(3):364-374. doi:10.1007/s11427-020-1643-8
51. Kuba K, Imai Y, Rao S, et al. A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury. *Nature Medicine*. 2005;11(8):875-879. doi:10.1038/nm1267
52. Bean D, Kraljevic Z, Searle T, et al. Treatment with ACE-inhibitors is associated with less severe disease with SARS-Covid-19 infection in a multi-site UK acute Hospital Trust. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.04.07.20056788
53. Sriram K, Insel PA. Dangers of ACE inhibitor and ARB usage in COVID-19: evaluating the evidence. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.03.25.20043927
54. Epelman S, Tang WH, Chen SY, Van Lente F, Francis GS, Sen S. Detection of soluble angiotensin-converting enzyme 2 in heart failure: insights into the endogenous counter-regulatory pathway of the renin-angiotensin-aldosterone system. *J Am Coll Cardiol*. 2008;52(9):750-754. doi:10.1016/j.jacc.2008.02.088
55. Mariana CP, Ramona PA, Ioana BC, et al. Urinary angiotensin converting enzyme 2 is strongly related to urinary nephrin in type 2 diabetes patients. *Int Urol Nephrol*. 2016;48(9):1491-1497. doi:10.1007/s11255-016-1334-8

56. Gill D, Arvanitis M, Carter P, et al. ACE inhibition and cardiometabolic risk factors, lung ACE2 and TMPRSS2 gene expression, and plasma ACE2 levels: a Mendelian randomization study. *medRxiv [Preprint]*. 2020:2020.2004.2010.20059121. doi:10.1101/2020.04.10.20059121
57. Tucker NR, Chaffin M, Bedi KC, et al. Myocyte specific upregulation of ACE2 in cardiovascular disease: implications for SARS-CoV-2 mediated myocarditis. *medRxiv [Preprint]*. 2020:2020.2004.2009.20059204. doi:10.1101/2020.04.09.20059204
58. Shi S, Qin M, Shen B, et al. Association of cardiac injury with mortality in hospitalized patients with COVID-19 in Wuhan, China. *JAMA Cardiol.* 2020. doi:10.1001/jamacardio.2020.0950
59. Tersalvi G, Vicenzi M, Calabretta D, Biasco L, Pedrazzini G, Winterton D. Elevated troponin in patients with Coronavirus Disease 2019: possible mechanisms. *J Card Fail.* 2020. doi:10.1016/j.cardfail.2020.04.009
60. Ali H, Daoud A, Mohamed MM, et al. Survival rate in acute kidney injury superimposed COVID-19 patients: a systematic review and meta-analysis. *Renal Failure.* 2020;42(1):393-397. doi:10.1080/0886022X.2020.1756323
61. Su H, Yang M, Wan C, et al. Renal histopathological analysis of 26 postmortem findings of patients with COVID-19 in China. *Kidney Int.* 2020. doi:10.1016/j.kint.2020.04.003
62. Felsenstein S, Herbert JA, McNamara PS, Hedrich CM. COVID-19: immunology and treatment options. *Clin Immunol.* 2020;215:108448. doi:10.1016/j.clim.2020.108448
63. Connors JM, Levy JH. COVID-19 and its implications for thrombosis and anticoagulation. *Blood.* 2020. doi:10.1182/blood.2020006000
64. Zheng Z, Peng F, Xu B, et al. Risk factors of critical & mortal COVID-19 cases: a systematic literature review and meta-analysis. *J Infect.* 2020. doi:10.1016/j.jinf.2020.04.021

- 1 65. Mehra MR, Desai SS, Kuy S, Henry TD, Patel AN. Cardiovascular Disease, Drug
2 Therapy, and Mortality in Covid-19. *NEJM*. 2020. doi:10.1056/NEJMoa2007621
- 3 66. Mehra MR, Desai SS, Ruschitzka F, Patel AN. Hydroxychloroquine or chloroquine with
4 or without a macrolide for treatment of COVID-19: a multinational registry analysis. *Lancet*.
5 2020. doi:10.1016/S0140-6736(20)31180-6
- 6 67. A mysterious company's coronavirus papers in top medical journals may be
7 unraveling. Available at [https://www.sciencemag.org/news/2020/06/mysterious-company-](https://www.sciencemag.org/news/2020/06/mysterious-company-s-coronavirus-papers-top-medical-journals-may-be-unraveling)
8 [s-coronavirus-papers-top-medical-journals-may-be-unraveling](https://www.sciencemag.org/news/2020/06/mysterious-company-s-coronavirus-papers-top-medical-journals-may-be-unraveling) (accessed June 02, 2020).
- 9 68. Rubin EJ. Expression of Concern: Mehra MR et al. Cardiovascular Disease, Drug
10 Therapy, and Mortality in Covid-19. *NEJM*. 2020. doi:10.1056/NEJMe2020822
- 11 69. Expression of concern: Hydroxychloroquine or chloroquine with or without a macrolide
12 for treatment of COVID-19: a multinational registry analysis. *Lancet*. 2020.
13 doi:10.1016/S0140-6736(20)31290-3
- 14 70. Zhang X, Yu J, Pan L-y, Jiang H-y. ACEI/ARB use and risk of infection or severity or
15 mortality of COVID-19: A systematic review and meta-analysis. *Pharmacol Res*.
16 2020;158:104927. doi:10.1016/j.phrs.2020.104927
- 17 71. Guo X, Zhu Y, Hong Y. Decreased mortality of COVID-19 with renin-angiotensin-
18 aldosterone system inhibitors therapy in patients with hypertension: a meta-analysis.
19 *Hypertension*. 2020. doi:10.1161/HYPERTENSIONAHA.120.15572
- 20 72. Mackey K, King VJ, Gurley S, et al. Risks and impact of angiotensin-converting
21 enzyme inhibitors or angiotensin-receptor blockers on SARS-CoV-2 infection in adults.
22 *Ann Intern Med*. 2020. doi:10.7326/m20-1515
- 23 73. Peng YD, Meng K, Guan HQ, et al. [Clinical characteristics and outcomes of 112
24 cardiovascular disease patients infected by 2019-nCoV]. *Zhonghua Xin Xue Guan Bing Za*
25 *Zhi*. 2020;48(0):E004. doi:10.3760/cma.j.cn112148-20200220-00105

- 1 74. Liu Y, Huang F, Xu J, et al. Anti-hypertensive angiotensin II receptor blockers
2 associated to mitigation of disease severity in elderly COVID-19 patients. *medRxiv*
3 *[Preprint]*. 2020. doi:10.1101/2020.03.20.20039586
- 4 75. Feng Y, Ling Y, Bai T, et al. COVID-19 with different severity: a multi-center study of
5 clinical features. *Am J Respir Crit Care Med*. 2020. doi:10.1164/rccm.202002-0445OC
- 6 76. Ip A, Parikh K, Parrillo JE, et al. Hypertension and renin-angiotensin-aldosterone
7 system inhibitors in patients with Covid-19. *medRxiv [Preprint]*. 2020.
8 doi:10.1101/2020.04.24.20077388
- 9 77. Wang Y, Lu X, Chen H, et al. Clinical course and outcomes of 344 intensive care
10 patients with COVID-19. *Am J Respir Crit Care Med*. 2020. doi:10.1164/rccm.202003-
11 0736LE
- 12 78. Reynolds HR, Adhikari S, Pulgarin C, et al. Renin–angiotensin–aldosterone system
13 inhibitors and risk of Covid-19. *NEJM*. 2020. doi:10.1056/NEJMoa2008975
- 14 79. Mancia G, Rea F, Ludergrani M, Apolone G, Corrao G. Renin–angiotensin–
15 aldosterone system blockers and the risk of Covid-19. *NEJM*. 2020.
16 doi:10.1056/NEJMoa2006923
- 17 80. Jurado A, Martin MC, Abad-Molina C, et al. COVID-19 in Spain: age, Interleukin-6, C
18 Reactive Protein and lymphocytes as key clues from a multicentre retrospective study.
19 *medRxiv [Preprint]*. 2020. doi:10.1101/2020.05.13.20101345
- 20 81. Conversano A, Melillo F, Napolano A, et al. RAAs inhibitors and outcome in patients
21 with SARS-CoV-2 pneumonia: a case series study. *Hypertension*. 2020.
22 doi:10.1161/hypertensionaha.120.15312
- 23 82. Baker KF, Hanrath AT, Schim van der Loeff I, et al. COVID-19 management in a UK
24 NHS Foundation Trust with a high consequence infectious diseases centre: a detailed
25 descriptive analysis. *medRxiv [Preprint]*. 2020. doi:10.1101/2020.05.14.20100834

Figure legends

Figure 1. Flow diagram of study selection

Figure 2. Forest plot showing the association between ACEIs or ARBs use and all-cause mortality in COVID-19 patients

Figure 3. Forest plot showing the association between ACEIs or ARBs use and COVID-19 severity

Figure 1

Figure 2

Figure 3

Table 1. Characteristics of included studies

Study	Country	Study design	Sample size	Eligibility criteria	Age, years	Male	Comorbidities	ACEIs or ARBs use	Outcomes	Definition of mortality	Definition of COVID-19 severity
Guo et al, ²⁵ 2020	China	Cross-sectional	187	Patients with COVID-19 and who were treated and discharged or died during hospitalization	58.5 ± 14.6 ^b	49%	Hypertension (33%), diabetes (15%), CAD (11.2%)	10%	All-cause mortality	In-hospital death	NR
Peng et al, ⁷³ 2020	China	Cross-sectional	112	Patients with COVID-19 and cardiovascular diseases	62 (55 - 67) ^c	47%	Hypertension (82%), diabetes (20%), CAD (55%), HF (36%)	20%	All-cause mortality, COVID-19 severity	In-hospital death	Critical vs mild/severe
Meng et al, ²⁷ 2020	China	Cross-sectional	42	Patients with COVID-19 and hypertension	64.5 (55.8-69) ^c	57%	Hypertension (100%), diabetes (14%), CAD (19%)	40%	All-cause mortality, COVID-19 severity, troponin, creatinine, procalcitonin, IL-6, D-dimer	In-hospital death	Severe vs moderate
Liu et al, ⁷⁴ 2020	China	Cross-sectional	78	Adult patients (≥18 years) with COVID-19 and hypertension	65.2 ± 10.7 ^b	55%	Hypertension (100%)	28%	COVID-19 severity	NR	Severe vs mild
Zeng et al, ³⁹ 2020	China	Cross-sectional	75 ^a	Patients with COVID-19	67 ± 11 ^b	47%	Hypertension (100%), diabetes (31%), CVD (21%)	37%	All-cause mortality, COVID-19 severity, hospitalization, hospital discharge, length of hospital stay	In-hospital death	Severe vs non-severe

Yun et al, ⁷⁵ 2020	China	Cross-sectional	113 ^a	Patients with COVID-19	53 (40-64) ^c	57%	Hypertension (100%)	29%	COVID-19 severity	NR	Critical/severe vs moderate
Li et al, ²⁶ 2020	China	Cross-sectional	362	Patients with COVID-19 and hypertension	66 (59-73) ^c	52%	Hypertension (100%), diabetes (35%), CAD (17%), HF (3%)	32%	All-cause mortality, COVID-19 severity, length of hospital stay, troponin, creatinine, procalcitonin, CRP, IL-6, D-dimer	In-hospital death	Severe vs non-severe
Argenziano et al, ⁴¹ 2020	USA	Cross-sectional	1000	Consecutive patients with COVID-19 who received emergency or inpatient care	61.7 ± 17.5 ^b	60%	Hypertension (60%), diabetes (37%), CAD (13%), HF (10%)	28%	ICU admission	NR	NR
Chen et al, ²⁰ 2020	China	Cross-sectional	123	Patients with COVID-19	57.7 ± 12.7 ^b	43%	Hypertension (33%), diabetes (11%), CAD (12%)	9%	All-cause mortality	In-hospital death	NR
Ashraf et al, ²² 2020	Iran	Cross-sectional	100	Hospitalized patients with COVID-19	58 (48-68) ^c	64%	Hypertension (26%), diabetes (26%), CAD (19%)	19%	COVID-19 severity	NR	Critical vs non-critical
Richardson et al, ⁴⁴ 2020	USA	Cross-sectional	1366 ^a	Hospitalized patients with COVID-19	63 (52-75) ^c	60%	Hypertension (100%)	30%	All-cause mortality, ICU admission, mechanical ventilation	In-hospital death	NR

Ip et al, ⁷⁶ 2020	USA	Cross-sectional	1129 ^a	Hospitalized patients with COVID-19 and hypertension	NR	NR	Hypertension (100%)	41%	All-cause mortality	In-hospital death	NR
Tedeschi et al, ³⁶ 2020	Italy	Cross-sectional	311 ^a	Hospitalized patients with COVID-19 and hypertension	76 (67-83) ^c	72%	Hypertension (100%), diabetes (24%), CVD (42%)	56%	All-cause mortality	In-hospital death	NR
Wang et al, ⁷⁷ 2020	China	Cross-sectional	344	Patients admitted to ICU with COVID-19	64 (52-72) ^c	52%	Hypertension (41%), diabetes (19%), CVD (12%)	18%	All-cause mortality	In-hospital death	NR
Reynolds et al, ⁷⁸ 2020	USA	Cross-sectional	2573 ^a	Patients with COVID-19	64 (54-75) ^c	52%	Hypertension (100%)	50%	COVID-19 severity	NR	ICU admission, mechanical ventilation, or death vs none
Mancia et al, ⁷⁹ 2020	Italy	Cross-sectional	6272 ^a	Patients with COVID-19	68 ± 13 ^b	63%	Hypertension (58%), CAD (8%), HF (5%)	46%	COVID-19 severity	NR	Critical/fatal vs mild/mode rate
Jurado et al, ⁸⁰ 2020	Spain	Cross-sectional	290 ^a	Hospitalized patients with COVID-19	NR	NR	Hypertension (100%)	67%	COVID-19 severity	NR	Severe vs mild/mode rate
Regina et al, ⁴⁵ 2020	Switzerland	Cross-sectional	200	Hospitalized adult patients (≥18 years) with COVID-19	70 (55-81) ^c	60%	Hypertension (44%), diabetes (22%), CAD (18%), CKD (14%)	26%	Mechanical ventilation	NR	NR
Tan et al, ²⁹ 2020	China	Cross-sectional	100	Hospitalized patients with COVID-19 and hypertension	NR	51%	Hypertension (100%), diabetes (28%), CAD (18%), CKD (9%)	31%	All-cause mortality, COVID-19 severity, hospital discharge,	In-hospital death	Critical/severe vs mild/mode rate

									mechanical ventilation, length of hospital stay, CRP, D-dimer		
Zhou et al, ⁴⁶ 2020	China	Cross-sectional	36 ^a	Hospitalized patients with COVID-19 and hypertension	64.8 ± 10.1 ^b	53%	Hypertension (100%)	42%	All-cause mortality, length of hospital stay	In-hospital death	NR
Conversano et al, ⁸¹ 2020	Italy	Cross-sectional	191	Hospitalized adult patients with COVID-19 pneumonia	63.4 ± 14.9 ^b	69%	Hypertension (50%), diabetes (15%), CAD (15%), HF (5%), CKD (26%)	36%	All-cause mortality	In-hospital death	NR
Yan et al, ³⁰ 2020	China	Case-control	610	Consecutive adult patients with COVID-19	48.7 ± 14.2 ^b	51%	Hypertension (22%), diabetes (10%), CVD (3%)	42%	COVID-19 severity	NR	Critical/severe vs mild/moderate
Bravi et al, ³⁷ 2020	Italy	Case-control	543 ^a	Patients with COVID-19 and hypertension	NR	NR	Hypertension (100%)	83%	COVID-19 severity	NR	Severe or very severe/lethal vs mild
Mehta et al, ⁴⁰ 2020	USA	Cohort	1735	Patients with COVID-19	NR	57%	Hypertension (93%), diabetes (46%), CAD (22%), HF (17%)	12%	Hospitalization, ICU admission, mechanical ventilation	NR	NR
Yuchen et al, ⁴⁷ 2020	China	Cohort	71 ^a	Patients with COVID-19, hypertension, and diabetes	67 (61-76) ^c	NR	Hypertension (100%), diabetes (100%)	45%	All-cause mortality, creatinine, procalcitonin, CRP,	In-hospital death	NR

									IL-6, D-dimer		
Rhee et al, ³⁸ 2020	Korea	Cohort	832	Patients with COVID-19 and diabetes	NR	53%	Hypertension (68%), diabetes (100%), CVD (27%), CKD (19%)	39%	COVID-19 severity	NR	Intensive care or death vs mild
Kim et al, ⁴³ 2020	USA	Cohort	2491	Hospitalized patients with COVID-19	62 (50-75) ^c	53%	Hypertension (57%), diabetes (33%), CAD (14%), HF (11%), CKD (16%)	30%	All-cause mortality, ICU admission	In-hospital death	NR
Khera et al, ³⁴ 2020	USA	Cohort	10196	Adult patients (≥18 years) with COVID-19 and hypertension	NR	54%	Hypertension (100%), diabetes (48%), CAD (5%), HF (27%), CKD (27%)	59%	All-cause mortality, hospitalization	In-hospital death	NR
Jung et al, ³³ 2020	Korea	Cohort	5179	Adult patients (≥18 years) with COVID-19	44.6 ± 18 ^b	44%	Hypertension (22%), diabetes (17%), CAD (1%), HF (4%), CKD (5%)	15%	All-cause mortality, mechanical ventilation	In-hospital death	NR
Huang et al, ²⁸ 2020	China	Cohort	50	Hospitalized patients with COVID-19 and hypertension	61.7 ± 12.9 ^b	54%	Hypertension (100%), diabetes (8%), CAD (2%)	40%	All-cause mortality, COVID-19 severity, mechanical ventilation, troponin, creatinine, procalcitonin	In-hospital death	Critical/severe vs mild/moderate

									in, CRP, D-dimer		
De Spiegele er et al, ²³ 2020	Belgiu m	Cohort	154	Residents at two elderly care homes with COVID-19	86 ± 7 ^b	33%	Hypertension (25%), diabetes (18%)	20%	COVID-19 severity	NR	Long-stay hospital admission or death vs none
Baker et al, ⁸² 2020	UK	Cohort	311	Hospitalized patients with COVID-19	75 (60- 83) ^c	55%	Hypertension (42%), diabetes (27%), CAD (21%), HF (14%), CKD (24%)	25%	All-cause mortality	28-day death	NR
Yang et al, ³¹ 2020	China	Cohort	126 ^a	Patients with COVID- 19 and hypertension	66 (61- 73) ^c	49%	Hypertension (100%), diabetes (30%), CVD (18%)	34%	All-cause mortality, COVID-19 severity, hospital discharge, length of hospital stay, troponin, creatinine, procalciton in, CRP, IL-6, D- dimer	In- hospital death	Critical/se vere vs mild/mode rate
Rentsch et al, ¹⁹ 2020	USA	Cohort	585	Patients with laboratory results consistent with SARS-CoV-2 or COVID-19	66.1 (60.4- 71) ^c	95%	Hypertension (72%), diabetes (44%)	44%	All-cause mortality, hospitaliza tion, ICU admission	In- hospital death	NR
Bean et al, ⁵² 2020	UK	Cohort	205	Patients with symptoms that required hospitalization with COVID-19	63 ± 20 ^b	52%	Hypertension (51%), diabetes (30%), CAD (15%)	23%	COVID-19 severity	NR	In-hospital death or required critical care

											support vs none
Feng et al, ²⁴ 2020	China	Cohort	65 ^a	Consecutive adult patients with COVID-19	47 (36-58) ^c	50%	Hypertension (100%), diabetes (31%), CVD (8%)	25%	COVID-19 severity, creatinine	NR	Critical/severe vs mild/moderate
Zhang et al, ²¹ 2020	China	Cohort	1128	Patients (18-74 years) with COVID-19 and hypertension	64 (56-69) ^c	53%	Hypertension (100%), diabetes (21%), CAD (12%)	17%	All-cause mortality, mechanical ventilation	28-day all-cause mortality	NR
Giorgi et al, ³² 2020	Italy	Cohort	2653	Symptomatic patients with COVID-19	63.2 ^b	50%	Hypertension (18%), diabetes (12%), CAD (7%), HF (6%)	31%	All-cause mortality, hospitalization	Death	NR
Benelli et al, ⁴² 2020	Italy	Cohort	411	Consecutive patients with COVID-19	66.8 ± 16.4 ^b	87%	Hypertension (47%), diabetes (16%), CVD (23%)	27%	All-cause mortality, ICU admission	In-hospital death	NR
Lee et al, ³⁵ 2020	Korea	Cohort	8266	Hospitalized patients with COVID-19	44.4 ± 19.1 ^b	38%	Hypertension (19%), diabetes (17%), CAD (6%), HF (1%)	12%	All-cause mortality	60-day death	NR

Abbreviations: ACEIs, angiotensin-converting enzyme inhibitors; ARBs, angiotensin receptor blockers; COVID-19, coronavirus disease 2019; CAD, coronary artery disease; HF, heart failure; CVD, cardiovascular disease; CKD, chronic kidney disease; NR, not reported; CRP, c-reactive protein; IL-6, interleukin-6; ICU, intensive care unit; USA, United States of America; UK, United Kingdom.

^aSubgroup from original population with data on ACEIs/ARBs use

^bMean ± SD

^cMedian (IQR)