

# Resource requirements for reintroducing elective surgery in England during the COVID-19 pandemic: a modelling study

A. J. Fowler<sup>1</sup>, T. D. Dobbs<sup>2,3</sup>, Y. I. Wan<sup>1</sup>, R. Laloo<sup>4</sup>, S. Hui<sup>1</sup>, D. Nepogodiev<sup>5</sup>, A. Bhangu<sup>5</sup>, I. S. Whitaker<sup>2,3</sup>, R. M. Pearse<sup>1\*</sup>, T. E. F. Abbott<sup>1\*</sup>

\* Joint senior authors

1. William Harvey Research Institute, Queen Mary University of London, UK.
2. Reconstructive Surgery & Regenerative Medicine Research Group, Institute of Life Sciences, Swansea University Medical School, Swansea, UK
3. Welsh Centre for Burns and Plastics, Morriston Hospital, Swansea, UK
4. Leeds Vascular Institute, Leeds General Infirmary, UK.
5. Academic Department of Surgery, University of Birmingham, UK

Correspondence to:

Tom Abbott, PhD

William Harvey Research Institute,  
Queen Mary University of London,  
Critical Care Research Office,  
The Royal London Hospital,  
London E1 1BB  
United Kingdom.

e-mail: [t.abbott@qmul.ac.uk](mailto:t.abbott@qmul.ac.uk)

Tel: +44 20 3594 0351

Abstract word count: 300/300

Manuscript word count: 3292/3500

Keywords: Surgery; Anaesthesia; COVID-19; Public Policy

## Abstract

*Background:* The response to COVID-19 has required cancellation of all but the most urgent surgeries, including many cancer operations. We estimated the number of cancelled surgical procedures in the National Health Service (NHS) in England due to COVID-19 and how this deficit would change over time once elective surgery was reintroduced.

*Methods:* Modelling study using Hospital Episode Statistics (HES) data from 2014 to 2019. Using NHS England definitions, surgical procedures were grouped into four classes of urgency. We calculated time-weighted average numbers of surgical procedures from 1<sup>st</sup> March 2020 and extrapolated to 28<sup>th</sup> February 2021 informed by activity in previous years. We estimated the procedure deficit using multiple conservative assumptions and then modelled the reintroduction of elective surgery between 1<sup>st</sup> June 2020 and 28<sup>th</sup> February 2021, estimating the resources required to achieve this. Costs of surgery were calculated using NHS reference costs. Estimates are reported with 95% confidence intervals.

*Findings:* 4,547,534 (3,318,195 - 6,250,771) patients with pooled mean age of 53.5 years were expected to undergo surgery in the NHS in England between 1<sup>st</sup> March 2020 and 28<sup>th</sup> February 2021. Due to COVID-19, 749,248 (513,565 - 1,077,448) surgical procedures were cancelled by 31<sup>st</sup> May 2020. As current guidelines require a gradual reintroduction of elective surgery, this deficit will increase further and 2,270,178 (1,453,057 - 3,363,472) patients will be awaiting surgery by 28th February 2021. The cost of these delayed procedures is £4,688,318,443 (£2,726,364,240 - £7,070,166,056). However, the safe delivery of surgery during the pandemic will require substantial extra resources including personal protective equipment and universal preoperative screening, leading to additional costs of £606,252,901 (£521,159,931 - £730,720,808).

*Interpretation:* Reintroduction of elective surgery during the pandemic response in NHS England will be associated with substantial treatment delays for many patients, and a large increase in treatment costs.

*Funding:* NIHR (DRF-2018-11-ST2-062) to AJF.

## Introduction

The outbreak of novel coronavirus SARS-CoV-2 and the disease COVID-19, were first reported in November 2019 in Wuhan, China,<sup>1</sup> and the World Health Organisation declared a global pandemic in March 2020. The healthcare response to the COVID-19 pandemic has required rapid changes to the provision of secondary healthcare services, including the creation of new hospitals.<sup>2,3</sup> Critical care capacity has increased substantially in high income countries, requiring re-deployment of staff and equipment from other departments.<sup>4</sup> Surgical services represent a large portion of healthcare activity, accounting for 5 million hospital admissions to the United Kingdom (UK) National Health Service (NHS) every year.<sup>5,6</sup> In March, National Health Service (NHS) England issued guidance to postpone non-urgent surgery from 15<sup>th</sup> April.<sup>7</sup> Only very urgent and emergency surgical treatments continued, with the cancellation of almost all planned cancer and non-cancer surgery.<sup>8,9</sup> However, the number of surgical procedures cancelled during the pandemic response, and the excess of untreated surgical disease, remain unknown.<sup>2,8</sup>

As the NHS response to the pandemic develops, there is a planned reintroduction of elective surgery with a particular focus on cancer surgery.<sup>10</sup> However, the risk of SARS-CoV-2 transmission within hospitals is a concern and further complicated by uncertain rates of asymptomatic infection in the general population (17% to 56%).<sup>11,12</sup> Asymptomatic infection rates may be as high as 24% among healthcare staff.<sup>13,14</sup> New procedures introduced to protect patients and staff from virus transmission have reduced capacity for surgery.<sup>2</sup> NHS plans to prevent in-hospital infection include preoperative patient screening, establishing 'clean' hospitals, routine staff testing and/or universal personal protective equipment (PPE).<sup>8,15</sup> However, the optimum strategy, resource requirements and time-scale for reintroducing elective surgery remain uncertain.<sup>15</sup> The number of patients requiring hospital treatment for COVID-19 infection is decreasing but critical care utilisation remains in excess of pre-pandemic levels.<sup>16</sup> Capacity for surgical care remains limited in operating rooms, post-anaesthetic recovery units and critical care units whilst the incidence of serious postoperative complications amongst patients with COVID-19 is very high.<sup>4,17</sup> A clear understanding of the number and type of overdue surgical procedures is urgently needed to inform national healthcare policies for reintroducing surgical services.

We used Hospital Episode Statistics (HES) data over five years from 2014 to 2019, to estimate the number of surgical procedures postponed in the NHS in England due to COVID-19. We then modelled multiple scenarios for the reintroduction of routine surgical activity over a nine-month period from June 2020. We provide estimates of the cumulative national deficit of surgical procedures and the resource implications for 'catching up' with postponed procedures, in terms of hospital beds, critical care beds, PPE for staff, preoperative screening of patients for COVID-19, and the associated financial cost.

## Methods

An unabridged description of the methods is provided in the appendix.

### *Data source*

We used aggregated HES data for Admitted Patient Care (APC) between 1<sup>st</sup> April 2014 and 31<sup>st</sup> March 2019 (<https://digital.nhs.uk/data-and-information>). HES provide detailed data describing hospital care in England, including the type of surgical procedure categorised by an Office for Population Censuses Surveys Classification of Interventions and Procedures (OPCS) codes. We used freely available anonymised data and research ethics and information governance approvals were therefore not required.

### *Outcomes*

The primary outcome measure was the deficit of surgical activity, defined as the number of cancelled surgical procedures, attributable to the COVID-19 pandemic response. The secondary outcomes were the associated resource requirements, defined as: hospital bed days, number of critical care admissions, number of investigations for preoperative COVID-19 screening, the amount of PPE required for operating theatre staff and the additional financial cost.

### *Modelling analysis*

We prospectively designed and published a statistical analysis plan before data analysis took place.<sup>18</sup> We used R version 3.6.1 (R Core Team, Vienna, Austria) to analyse the data. We stratified surgical procedures according to a classification published by NHS England on 17<sup>th</sup> March 2020, which described operations that should be stopped or continued during the initial pandemic response.<sup>7</sup> Surgical procedures were divided into four classes of decreasing urgency: Class 1 – Emergency operations needed within 72 hours; Class 2 – Urgent surgery that can be deferred for up to 4 weeks; Class 3 – Semi-urgent surgery that can be delayed for up to 3 months; and Class 4 – Elective surgery that can be delayed for more than 3 months. We used a pre-specified data-driven approach to identify emergency, urgent and elective surgical procedures based on waiting times and expert review of procedure coding (supplementary tables 1 and 2). We estimated the monthly volume of surgical activity that would have been expected had there not been a pandemic until 28<sup>th</sup> February 2021, by

calculating the annual change in activity during the five-year period from 1<sup>st</sup> April 2014 to 31<sup>st</sup> March 2019, and extrapolated using a linear growth assumption. We adjusted for monthly variation in admissions and estimated the average age for each class of surgery.

#### *Surgical procedure volume*

We estimated the number of postponed or cancelled surgical procedures from the 1<sup>st</sup> March 2020 up to 1<sup>st</sup> June 2020 according to several assumptions. First, that class 1 (emergency) surgery would continue at the pre-pandemic rate. Second, that class 2 surgery would continue at a reduced rate. We used four scenarios, where 20%, 40%, 60% and 80% of class 2 surgical procedures were assumed to have continued. Third, that 50% of class 3 and 4 procedures continued in March and then stopped completely in April and May, reflecting between-hospital heterogeneity in the timing of stoppage of surgery. The results were presented as the deficit of surgical procedures between 1<sup>st</sup> March and 31<sup>st</sup> May 2020 with a 95% confidence interval. Fourth, that widespread reintroduction of surgical activity would start from 1<sup>st</sup> June 2020 and continue to increase to pre-pandemic levels. For each of the four scenarios of class 2 procedures, we assumed a linear increase in activity over the three months from 1<sup>st</sup> June to 31<sup>st</sup> August 2020, with class 3 and 4 surgical procedures remaining cancelled until 31<sup>st</sup> August. In the final model we assumed that 80% of class 2 procedures continued. We repeated the analysis in an iterative fashion, by adding class 3 procedures on 1<sup>st</sup> September and class 4 procedures on 1<sup>st</sup> December. Fifth, that pre-pandemic levels of surgical activity would be reached by 28<sup>th</sup> February 2021 (supplementary figure 1). We presented the estimated number of surgical procedures carried out each month between 1<sup>st</sup> March 2020 and 28<sup>th</sup> February 2021 and a rolling deficit of surgical activity compared to the expected volume of surgery according to the previous five-year average. We did not include assumptions about a second peak of COVID-19, nor did we attempt to model the impact of reduced operating theatre utilisation due to enhanced infection control procedures.

#### *Hospital admissions*

We calculated the total number of bed-days, weighted by procedure frequency, by multiplying the median length of stay by the number of inpatient admissions. We estimated

the proportion of patients that would require postoperative critical care using a conservative assumption of 1% of patients undergoing inpatient surgery.<sup>19-21</sup>

#### *Preoperative screening tests*

We estimated preoperative COVID-19 screening requirements according to three scenarios. First, all patients would have two outpatient preoperative COVID-19 Polymerase Chain Reaction (PCR) tests. Second, all patients would have one outpatient and one inpatient preoperative SARS-CoV-2 PCR tests, requiring an additional day in hospital for isolation. Third, all patients would have one outpatient and one inpatient preoperative SARS-CoV-2 PCR tests, and patients undergoing class 1 and 2 procedures would also have a computed tomography scan of the chest, in line with guidance from the Royal College of Radiologists.<sup>22, 23</sup>

#### *Personal protective equipment*

We estimated the amount of PPE required in operating theatres according to two scenarios.<sup>24</sup> First, that eight staff members would be present for every procedure (two surgeons, two anaesthetists, one operating department practitioner and three scrub staff). Second, that four staff members would be present for every procedure (one surgeon, one anaesthetist, one operating department practitioner and one scrub staff), each requiring an FFP3 mask, a fluid repellent gown, two pairs of gloves and eye protection.

#### *Estimated financial cost*

We estimated the cost of reintroducing surgical activity from 1<sup>st</sup> June 2020 and the total deficit of surgical procedures on 28<sup>th</sup> February 2021 by combining the cost of each surgical procedure, the cost of associated preoperative COVID-19 screening and the cost of PPE. The total costs of surgical procedures was calculated by matching the OPCS v4.7 code with the Health Resource Group coding and summing the associated procedure cost according to the national schedule of NHS costs.<sup>5, 25</sup> The costs of screening tests were calculated according to £19 per SARS-CoV-2 PCR and £69 per CT scan and a range of between £222 and £346 per additional bed day.<sup>26-28</sup> The costs of PPE was calculated as £2.90 per FFP3 mask, £14.90 per 100 gloves, £3 per fluid resistant gown and £2.90 per piece of eye protection.<sup>28, 29</sup>

*Role of the funding source*

AJF holds a National Institute for Health Research Doctoral Research fellowship (DRF-2018-11-ST2-062). The funder had no role in the study design, data collection, analysis, interpretation, or writing the report. AJF and TA had full access to all the data in the study and all authors had final responsibility for the decision to submit for publication.

## Results

We identified 1073 OPCS codes for Class 1 – 4 surgical procedures, representing a total of 22,513,872 surgical admissions between 1<sup>st</sup> April 2014 and 31<sup>st</sup> March 2019 (Figure 1). The monthly median number of procedures was 382,768 (Interquartile range [IQR]: 22,890) (supplementary table 3). If growth in the number of surgical procedures had continued according to a pre-pandemic trajectory, we estimate that 4,547,534 (95%CI: 3,318,195 to 6,250,771) would have been performed between 1<sup>st</sup> March 2020 and 28<sup>th</sup> February 2021 (supplementary table 4). Patients aged over 60 years accounted for 32.7% of class 1, 43.9% of class 2, 53.7% of class 3 and 45.6% of class 4 surgical activity (supplementary table 5).

### *Surgical procedure volume*

We estimate that 749,248 (95%CI: 513,565 to 1,077,448) surgical procedures will have been cancelled between 1<sup>st</sup> March and 31<sup>st</sup> May 2020 in comparison with observed pre-pandemic levels of activity (supplementary table 6). We modelled the widespread reintroduction of class 2 surgical activity from 31<sup>st</sup> May 2020, for four scenarios of incrementally increasing activity (20% - 80%) that would reach pre-pandemic levels by 31<sup>st</sup> August 2020 (supplementary table 7). Even if class 2 - 4 surgical procedures are reintroduced in a stepwise fashion between 1<sup>st</sup> June 2020 and 28<sup>th</sup> February 2021 to reach the predicted pre-pandemic level of activity, the number of cancelled surgical procedures would be 2,270,178 (95%CI: 1,453,057 to 3,363,472) procedures (table 1, figures 2-3).

### *Hospital admissions*

The total bed days associated with the deficit of surgical activity on 31<sup>st</sup> May 2020 is 973,495 (95%CI: 624,030 to 1,423,713) days (supplementary table 8). If widespread reintroduction of surgical activity occurs from 1<sup>st</sup> June 2020, the total number of bed days associated with the cumulative deficit of surgical activity up to 28<sup>th</sup> February 2021 3,238,850 (95%CI: 1,945,237 to 4,847,579) days (table 2). The total number of critical care admissions associated with the deficit of surgical activity on 31<sup>st</sup> May 2020 is 2,756 (95%CI: 2,020 to 3,916) (supplementary table 8). If widespread reintroduction of surgical activity occurs from 1<sup>st</sup> June 2020, the estimated total number of critical care admissions associated with the cumulative deficit of surgery up to 28th February 2021 will be 8,378 admissions (95%CI: 5,890 to 12,152).

### *Preoperative screening tests*

The estimated resource requirement for preoperative COVID-19 screening associated with the deficit of surgical activity on 31<sup>st</sup> May 2020 include: 1,390,106 (95%CI: 939,958 to 2,018,500) SARS-CoV-2 PCR tests, 20,606 (95%CI: 16,736 to 25,602) CT scans and 275,604 (95%CI: 202,024 to 391,603) bed days for preoperative isolation. The screening resource requirements associated with the reintroduction of surgical activity from 1<sup>st</sup> June 2020 is provided in table 2.

### *Personal protective equipment*

The estimated amount of PPE associated with the deficit of surgical activity on 31<sup>st</sup> May 2020 is 11,120,848 (95% CI: 7,519,664 to 16,148,000) items, assuming four persons per theatre, and 22,241,696 (95% CI: 15,039,328 to 32,296,000) items assuming eight persons pre theatre. The estimated total amount of PPE associated with the cumulative deficit of surgical activity by 28th February 2021 is between 36,322,848 (95%CI: 23,248,912 to 53,815,552) items, assuming four persons per theatre and 72,645,696 items (95%CI: 46,497,824 to 107,631,104) assuming eight persons per theatre (supplementary table 8).

### *Financial cost*

The procedure cost for the deficit of surgical activity on 31<sup>st</sup> May 2020 is £1,359,552,496 (95%CI: £840,846,557 to £2,009,926,307) (table 2 and supplementary table 10). If elective surgery recommences between 1<sup>st</sup> June 2020 and 28<sup>th</sup> February, the total costs associated with performing reintroduced surgery, including the costs of the procedure, PPE and screening, are between £5,068,816,991 (95%CI: £4,170,359,336 to £6,302,032,861) and £5,347,860,447 (95%CI: £4,408,986,578 to £6,641,524,634) (supplementary table 9). The total cost of cancelled operations by 28<sup>th</sup> February 2021, including the costs of the procedure, PPE and screening tests, are between £5,045,354,856 (95%CI: £2,966,949,647 to £7,592,843,343) and £5,231,861,474 (95%CI: £3,092,857,610 to 7,865,929,880 (supplementary table 10).

An unabridged description of the results is provided in the appendix.

## Discussion

The principal finding of this study is that, without a substantial increase in capacity, accumulated delays in surgical care will lead to a backlog of more than two million overdue or cancelled surgical procedures in the NHS in England by 28<sup>th</sup> February 2021. This is equivalent to 45% of the total number of surgeries performed in England each year before the COVID-19 pandemic, and is larger than the existing waiting list to start surgical treatment in March 2020, which already stood at over 1.8 million patients.<sup>30</sup> The cost of clearing this post-pandemic waiting list is in excess of £5 billion, or 4.2% of the total NHS England budget.<sup>31</sup> Additional costs of delivering surgical services under strict infection control procedures exceed £600 million, and include personal protective equipment, preoperative screening and extra bed-days in hospital. We have not attempted to model the reduced operating theatre utilisation efficiency, suggesting the true excess cost may be far greater. Nor does our estimate account for additional costs of contracts with non-NHS providers, or the costs of postoperative complications among surgical patients with COVID-19.<sup>4, 17</sup> In order to reduce the size of any future backlog of overdue surgery, elective surgical services would need to be reintroduced at the earliest opportunity and at the maximum capacity possible. Whether this can be achieved safely remains unclear.

Postoperative mortality is much higher in patients with COVID-19 infection, compared to the expected pre-pandemic rate.<sup>17</sup> An international cohort study of more than 1000 patients with COVID-19 found the incidence of postoperative mortality was more than one in four patients, compared to an expected mortality rate of 1 in 30 patients undergoing emergency surgery.<sup>17</sup> This will inevitably lead to higher resource use, longer in-patient lengths of stay and greater financial costs. Furthermore, emerging data from the UK suggests that as many as one in twelve healthcare workers had asymptomatic infection during the first peak of the pandemic, and were at risk of transmitting the virus to other staff and patients.<sup>14</sup> Consequently, many surgeons may choose to avoid surgery in patients with COVID-19 infection, to reduce the active reservoir inside the hospital and to mitigate the higher-risk of postoperative mortality. When surgery does go ahead, the benefits must be weighed against the potential harm of bringing vulnerable patients into contact with SARS-CoV-2 and the potential risks to staff members. It is likely that preoperative screening for the virus and PPE for healthcare workers will become universal. We estimate that by February 2021, over

165,000 SARS-CoV-2 PCR tests will be required every week for preoperative screening alone. Permanent infrastructure for testing will need to be included in strategic plans for pathology services to support elective surgery. There are no national data regarding the use of protective equipment at individual institutions. However, many societies have exercised caution and have advocated for the deployment of full PPE for all high risk procedures and examinations, including all aerosol generating procedures.<sup>24</sup>

There are no published data on the rate of cancellation of surgery within the NHS in England during the COVID-19 pandemic response, of which we are aware. However, the results of our analysis are consistent with data from an international survey, which reported that the majority of both urgent cancer and non-cancer surgery were postponed during the peak of the pandemic.<sup>9</sup> We provide estimates of the likely volume of cancelled operations and the resources that may be required to reintroduce surgery and clear the backlog of cases. Delays to the re-introduction of surgical procedures may result in increased secondary morbidity and mortality, not directly attributable to COVID-19 infection, but due to increased volume of undiagnosed surgical disease and delays to curative therapy.<sup>2</sup> If delays in surgical care result in higher incidence of more advanced disease, it is likely that postoperative morbidity and mortality will also increase, as well as critical care bed utilisation and hospital length of stay. Strategies for undertaking safe surgery after the pandemic, during a prolonged pandemic response, or if COVID-19 becomes endemic, include reorganisation of referral pathways, restructuring of the surgical workforce, dedicated surgical critical care resources, isolated ‘cold’ hospitals and prioritisation of certain patient groups.<sup>2</sup> If second or subsequent peaks of COVID-19 occur, it is likely surgical services will again be curtailed, with further increases in the number of cancelled procedures. NHS leaders should make plans for the future continuity of surgical care.

Our analysis has several strengths. We analysed robust national data that describe the whole healthcare setting of a single country according to a prospectively published statistical analysis plan. Our results purposely provide conservative estimates of procedure volume and associated resource requirements. We report procedures at the admission level, rather than the episode level, which will likely underestimate the volume of surgery. We used multiple sources to calculate postoperative critical care admission, which did not

include preoperative admission, nor increased critical care utilisation for patients with COVID-19. Our estimates of PPE only include usage in operating theatre, but not on postoperative wards or in clinics. Our cost estimates do not take account of potential future reorganisation of care pathways, the use of 'hot' and 'cold' sites or use of private sector hospitals. Our analysis also has limitations. We made a series of assumptions based on NHS England guidance, which we applied to a series of plausible scenarios regarding the continuation of surgical activity during the pandemic.<sup>7, 10</sup> However, the reality of the volume, type and timing of continuation and reintroduction of surgery may differ from our model. We will only be able to test this in retrospect once actual numbers of procedures carried out are published. We did not take into account a potential second peak of COVID-19.<sup>32</sup> It is likely that the volume of emergency surgery that has continued during the pandemic is lower than pre-pandemic levels, perhaps due to patients avoiding hospital or clinicians using alternative management strategies.<sup>33</sup> The age distribution of the surgical population is skewed.<sup>6</sup> However, due to limitations of the data source, we were not able to adjust our analysis for age. It is likely that, unfortunately, some patients waiting for surgery will have died while waiting to have their surgery. The numbers of these cases are unknown and we were unable to account for these in the analysis. We will update our assumptions and analyses as new data become available.

In conclusion, accumulated delays in surgical care due to COVID-19 will lead to a backlog of more than two million overdue or cancelled surgical procedures in the NHS in England by 28<sup>th</sup> February 2021. These delays in care are likely to be associated with increased mortality. The procedure cost of clearing this waiting list is in excess of £4.6 billion, with additional costs of at least £600 million for personal protective equipment and preoperative screening during the pandemic.

## Contributions

AF, TD and TA were responsible for study design. RL, SH, TD, YW and AF were responsible for data collection. AF and TA were responsible for data analysis. AF, TD, RP and TA were responsible for data interpretation. TA wrote the first draft of the manuscript. All authors revised the manuscript for important intellectual content and approved the final version.

## Declarations of interest

AJF reports a grant from the National Institute for Health Research (DRF-2018-11-ST2-062). TDD reports funding from the Welsh Clinical Academic Training (WCAT) Fellowship. IW reports active grants from the American Association of Plastic Surgeons and the European Association of Plastic Surgeons; is an editor for Frontiers of Surgery, associate editor for the Annals of Plastic Surgery, editorial board of BMC Medicine and numerous other editorial board roles. RP reports grants from NIHR, during the conduct of the study; grants and personal fees from Edwards Life Sciences, grants and non-financial support from Intersurgical UK, outside the submitted work; and has given lectures and/or performed consultancy work for Nestle Health Sciences, BBraun, Intersurgical, GlaxoSmithKline and Edwards Lifesciences, and holds editorial roles with the British Journal of Anaesthesia, the British Journal of Surgery and BMJ Quality and Safety. TA is a member of the associate editorial board of the British Journal of Anaesthesia. All other authors report no relationships or activities that could appear to have influenced the submitted work.

## Data sharing

The data source is freely available online.

## Acknowledgements


AF holds a National Institute for Health Research Doctoral Research fellowship.

## References


1. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020; **395**(10223): 507-13.
2. Soreide K, Hallet J, Matthews JB, et al. Immediate and long-term impact of the COVID-19 pandemic on delivery of surgical services. *The British journal of surgery* 2020.
3. Abu Hilal M, Besselink MG, Lemmers DHL, Taylor MA, Triboldi A. Early look at the future of healthcare during the COVID-19 pandemic. *The British journal of surgery* 2020.
4. CovidSurg Collaborative. Global guidance for surgical care during the COVID-19 pandemic. *The British journal of surgery* 2020.
5. Abbott TEF, Fowler AJ, Dobbs TD, Harrison EM, Gillies MA, Pearse RM. Frequency of surgical treatment and related hospital procedures in the UK: a national ecological study using hospital episode statistics. *British journal of anaesthesia* 2017; **119**(2): 249-57.
6. Fowler AJ, Abbott TEF, Prowle J, Pearse RM. Age of patients undergoing surgery. *The British journal of surgery* 2019; **106**(8): 1012-8.
7. NHS England. Clinical guide to surgical prioritisation during the coronavirus pandemic, 2020.
8. Spinelli A, Pellino G. COVID-19 pandemic: perspectives on an unfolding crisis. *The British journal of surgery* 2020.
9. CovidSurg Collaborative, Nepogodiev D, Bhangu A. Elective surgery cancellations due to the COVID-19 pandemic: global predictive modelling to inform surgical recovery plans. *The British journal of surgery* 2020.
10. NHS England. Important - for action - second phase of NHS response to COVID19. In: trusts CeoNTaf, editor. NHS England website: NHS England; 2020.
11. Arons MM, Hatfield KM, Reddy SC, et al. Presymptomatic SARS-CoV-2 Infections and Transmission in a Skilled Nursing Facility. *The New England journal of medicine* 2020.
12. Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Euro surveillance : bulletin European sur les maladies transmissibles = European communicable disease bulletin* 2020; **25**(10).
13. Shields A et al. SARS-CoV-2 seroconversion in health care workers. *MedRxiv* 2020.
14. Treibel T et al. COVID-19: PCR screening of asymptomatic health-care workers at London hospital. *Lancet* 2020.
15. Mayol J, Fernandez Perez C. Elective surgery after the pandemic: waves beyond the horizon. *The British journal of surgery* 2020.
16. Intensive Care National Audit and Research Centre. ICNARC report on COVID-19 in critical care 08 May 2020. 2020.
17. CovidSurg Collaborative. Mortality and pulmonary complications in patients undergoing surgery with perioperative SARS-CoV-2 infection: an international cohort study. *Lancet* 2020 (in press).
18. Fowler AJ, Dobbs TD, Abbott TEF. Estimated surgical requirements in England after COVID-19: a modelling study using hospital episode statistics. 2020. <https://http://www.qmul.ac.uk/ccpmg/sops--saps/statistical-analysis-plans-saps/09/05/2020>.
19. International Surgical Outcomes Study Group. Global patient outcomes after elective surgery: prospective cohort study in 27 low-, middle- and high-income countries. *British journal of anaesthesia* 2016; **117**(5): 601-9.
20. Pearse RM, Moreno RP, Bauer P, et al. Mortality after surgery in Europe: a 7 day cohort study. *Lancet* 2012; **380**(9847): 1059-65.
21. Intensive Care National Audit and Research Centre. Summary Statistics. 2019. <https://http://www.icnarc.org/Our-Audit/Audits/Cmp/Reports/Summary-Statistics> (accessed 20/05/2020 2020).

22. Lima DS, Ribeiro MAF, Jr., Gallo G, Di Saverio S. Role of chest CT in patients with acute abdomen during the COVID-19 era. *The British journal of surgery* 2020.
23. The Royal College of Radiologists. Statement on use of CT chest to screen for COVID-19 in pre-operative patients. 14/05/2020 2020. <https://http://www.rcr.ac.uk/college/coronavirus-covid-19-what-rkr-doing/clinical-information/role-ct-chest/role-ct-screening-0> (accessed 17/05/2020).
24. Jessop ZM, Dobbs TD, Ali SR, et al. Personal Protective Equipment (PPE) for Surgeons during COVID-19 Pandemic: A Systematic Review of Availability, Usage, and Rationing. *The British journal of surgery* 2020.
25. NHS Improvement. National Cost Collection for the NHS. 2020. <https://improvement.nhs.uk/resources/national-cost-collection/> (accessed 11/05/2020).
26. National Institute for Health and Care Excellence. Costing statement: Implementing the NICE guideline on Transition between inpatient hospital settings and community or care home settings for adults with social care needs (NG27), 2015.
27. NHS Improvement. Reference costs 2017/18: highlights, analysis and introduction to the data. 2018.
28. NHS Improvement. National Tariff Payment System: National prices and prices for emergency care services. In: NHS Improvement, editor.; 2020.
29. Medical supplies company. 2020. <http://www.medistock.co.uk> (accessed 17/05/2020 2020).
30. NHS England. NHS referral to treatment (RTT) waiting times data March 2020, 2020.
31. The Kings Fund. The NHS budget and how it has changed. 2020. <https://http://www.kingsfund.org.uk/projects/nhs-in-a-nutshell/nhs-budget> (accessed 23/05/2020 2020).
32. Leung K, Wu JT, Liu D, Leung GM. First-wave COVID-19 transmissibility and severity in China outside Hubei after control measures, and second-wave scenario planning: a modelling impact assessment. *Lancet* 2020; **395**(10233): 1382-93.
33. Cano-Valderrama O, Morales X, Ferrigni CJ, et al. Reduction in emergency surgery activity during COVID-19 pandemic in three Spanish hospitals. *The British journal of surgery* 2020.


## Figures



**Figure 1.** Number of hospital admissions for surgery in England between 1<sup>st</sup> April 2015 to 31<sup>st</sup> March 2019, categorised by class of surgery according to NHS England definition (class 1, emergency surgery; class 2, urgent surgery with a waiting time < 4 weeks; class 3, semi-urgent surgery with a waiting time < 3 months; class 4 surgery, elective surgery with a waiting time > 3 months). 56 procedure codes had no mean time recorded so were unclassified, these accounted for a total of 73 admissions and so were excluded from analysis. OPCS codes used to define surgery are provided in supplementary table 1.



**Figure 2.** Top panel: estimated number of surgical procedures performed per month between 1<sup>st</sup> February 2020 and 31<sup>st</sup> March 2021 (blue) and expected number of surgical procedures performed per month based on time-weighted five-year average between 1<sup>st</sup> April 2014 and 31<sup>st</sup> March 2019 (red). Solid line indicates mean number of surgical procedures per months and dotted lines with shaded area indicate 95% confidence intervals. Bottom panel: predicted cumulative number of cancellations between 1<sup>st</sup> February 2020 and 31<sup>st</sup> March 2021. Class1, emergency surgery; class 2, urgent surgery with a waiting time < 4 weeks; class 3, semi-urgent surgery with a waiting time < 3 months; class 4, elective surgery with a waiting time > 3 months. This assumes a stepwise reactivation of surgical activity from 1<sup>st</sup> June 2020 onwards; reaching predicted pre-pandemic levels of surgical activity by 28<sup>th</sup> February 2021.



**Figure 3.** Estimated number of surgical procedures performed per month between 1<sup>st</sup> February 2020 and 31<sup>st</sup> March 2021, stratified by class of surgery. Class 1, emergency surgery; class 2, urgent surgery with a waiting time < 4 weeks; class 3, semi-urgent surgery with a waiting time < 3 months; class 4, elective surgery with a waiting time > 3 months. This assumes a stepwise reactivation of surgical activity from 1<sup>st</sup> June 2020 onwards; reaching predicted pre-pandemic levels of surgical activity by 28<sup>th</sup> February 2021. Solid line indicates mean number of procedures performed per month and dotted lines with shaded area indicate 95% confidence intervals.

## Tables

Class of Surgery	Mar	April	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.	Jan.	Feb.
<b>Procedures carried out each month</b>												
<b>Class 1</b>	72999 (69230 to 76769)	67776 (63324 to 72227)	70844 (66191 to 75497)	69027 (64493 to 73561)	70502 (65871 to 75133)	67738 (63289 to 72187)	68575 (64071 to 73079)	72048 (67316 to 76781)	71308 (66624 to 75991)	72665 (67892 to 77438)	72593 (67825 to 77361)	66472 (62106 to 70838)
<b>Class 2</b>	35290 (29202 to 42773)	26566 (21365 to 34249)	27629 (22220 to 34767)	35770 (28767 to 45010)	35854 (28835 to 45117)	33825 (27203 to 42563)	34965 (28120 to 43998)	36731 (29540 to 46220)	36542 (29388 to 45982)	31840 (25607 to 40066)	35094 (28223 to 44160)	33840 (27215 to 42582)
<b>Class 3</b>	84128 (69109 to 110982)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	37557 (30940 to 52842)	78907 (65004 to 111020)	117752 (97005 to 165674)	136801 (112698 to 192476)	150781 (124216 to 212146)	145392 (119776 to 204563)
<b>Class 4</b>	58018 (30779 to 86570)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	28346 (13346 to 44249)	62486 (29419 to 97542)	90380 (42551 to 141083)	
<b>Total</b>	250435 (198320 to 317094)	94342 (84689 to 105656)	98473 (88411 to 110264)	104797 (93260 to 118571)	106356 (94706 to 120250)	101563 (90492 to 114750)	141097 (123131 to 169919)	187686 (161860 to 234021)	225602 (193017 to 287647)	269652 (219543 to 354229)	320954 (249683 to 431209)	336084 (251648 to 459066)
<b>Cumulative deficit of cancelled surgical procedures</b>												
<b>Class 1</b>	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)	0 (0 to 0)
<b>Class 2</b>	7058 (5840 to 8554)	13699 (11181 to 16911)	20606 (16736 to 25602)	25972 (21051 to 32354)	29557 (23935 to 36866)	31249 (25295 to 38994)	31249 (25295 to 38994)	31249 (25295 to 38994)	31249 (25295 to 38994)	31249 (25295 to 38994)	31249 (25295 to 38994)	31249 (25295 to 38994)
<b>Class 3</b>	84128 (69109 to 110982)	226801 (186645 to 311721)	375187 (308887 to 520497)	528871 (435494 to 736727)	682918 (562400 to 953468)	828247 (682124 to 1157943)	940918 (774944 to 1316469)	1019825 (839949 to 1427490)	1059076 (872284 to 1482715)	1059076 (872284 to 1482715)	1059076 (872284 to 1482715)	1059076 (872284 to 1482715)
<b>Class 4</b>	58019 (30779 to 86570)	176272 (86453 to 271165)	299260 (144356 to 463151)	426639 (204326 to 661991)	554319 (264438 to 861301)	674773 (321148 to 1049331)	799287 (379770 to 1243699)	930089 (441352 to 1447882)	1060218 (502617 to 1651015)	1087239 (511875 to 1697192)	1149726 (541294 to 1794735)	1179853 (555478 to 1841763)
<b>Total</b>	149205 (105728 to 206106)	416772 (284279 to 599797)	695053 (469979 to 1009250)	981482 (660871 to 1431072)	1266794 (850773 to 1851635)	1534269 (1028567 to 2246268)	1771454 (1180009 to 2599162)	1981163 (1306596 to 2914366)	2150543 (1400196 to 3172724)	2177564 (1409454 to 3218901)	2240051 (1438873 to 3316444)	2270178 (1453057 to 3363472)

**Table 1.** Model for reintroduction of surgical activity between 1<sup>st</sup> March 2020 and 28<sup>th</sup> February 2021, assuming: continued class 1 (emergency) activity at pre-pandemic levels; continued class 2 (urgent) activity at 80% of pre-pandemic levels and increasing on 1<sup>st</sup> June to reach pre-pandemic levels by 31<sup>st</sup> August; reintroduction of class 3 activity on 1<sup>st</sup> September and reaching pre-pandemic levels by 31<sup>st</sup> November; and reintroducing class 4 activity on 30<sup>th</sup> November and reaching pre-pandemic levels by 28<sup>th</sup> February 2021. The top panel shows the number of procedures carried out in each month between 1<sup>st</sup> March 2020 and 28<sup>th</sup> February 2021. The bottom panel shows the cumulative deficit of surgical activity on the last day of each month. Numbers are presented as predicted time-weighted average with 95% confidence intervals.

Item	Mar	April	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.	Jan.	Feb.	Total
Admissions for surgery	250435 (198320 to 317094)	94342 (84689 to 105656)	98473 (88411 to 110264)	104797 (93260 to 118571)	106356 (94706 to 120250)	101563 (90492 to 114750)	141097 (123131 to 169919)	187686 (161860 to 234021)	225602 (193017 to 287647)	269652 (219543 to 354229)	320954 (249683 to 431209)	336084 (251648 to 459066)	2237041 (1848760 to 2822676)
Bed days	698462 (595236 to 820803)	454083 (412773 to 500911)	474174 (431088 to 523006)	491418 (443677 to 546593)	499658 (451340 to 555433)	477998 (431982 to 531046)	525021 (471145 to 595103)	592262 (528495 to 682443)	627685 (557245 to 733731)	688111 (590814 to 826141)	772068 (638855 to 951898)	777615 (622802 to 979032)	7078555 (6175452 to 8246140)
Day cases	93608 (64146 to 130298)	5579 (4487 to 7020)	5802 (4666 to 7301)	7512 (6041 to 9452)	7529 (6055 to 9474)	7103 (5713 to 8938)	26192 (21433 to 35760)	47316 (38828 to 65426)	66772 (54857 to 92806)	96829 (72054 to 138554)	130405 (90568 to 189678)	148579 (98080 to 218542)	643226 (466928 to 913249)
Crit care admissions	3758 (3418 to 4172)	2921 (2702 to 3153)	3052 (2824 to 3295)	3044 (2807 to 3298)	3103 (2863 to 3361)	2977 (2747 to 3223)	3206 (2939 to 3534)	3565 (3250 to 3989)	3728 (3380 to 4228)	3909 (3511 to 4481)	4083 (3626 to 4736)	3869 (3399 to 4531)	41214 (37464 to 46001)
SARS-CoV 2 PCR test	500870 (396640 to 634188)	188684 (169378 to 211312)	196946 (176822 to 220528)	209594 (186520 to 237142)	212712 (189412 to 240500)	203126 (180984 to 229500)	282194 (246262 to 339838)	375372 (323720 to 575294)	451204 (439034 to 708458)	539304 (499366 to 862418)	641908 (503296 to 918132)	672168 (3697520 to 5645352)	4474082
CT scans	108289 (98432 to 119542)	94342 (84689 to 105656)	98473 (88411 to 110264)	104797 (93260 to 118571)	106356 (94706 to 120250)	101563 (90492 to 114750)	103540 (92191 to 117077)	108779 (96856 to 123001)	107850 (96012 to 121973)	104505 (93499 to 117504)	107687 (96048 to 121521)	100312 (89321 to 113420)	1246493 (1113917 to 1403529)
Excess bed days for screening	156827 (134174 to 186796)	88763 (80202 to 98636)	92671 (83745 to 102963)	97285 (87219 to 109119)	98827 (88651 to 110776)	94460 (84779 to 105812)	114905 (101698 to 134159)	140370 (123032 to 168595)	158830 (138160 to 194841)	172823 (147489 to 215675)	190549 (159115 to 241531)	187505 (153568 to 240524)	1593815 (1381832 to 1909427)
Excess Cost (low estimate) [£]	60917896 (51331857 to 73306367)	33233274 (29948569 to 37047417)	34693202 (31268820 to 38668761)	36624321 (32735356 to 41226551)	37190194 (32735356 to 41835160)	35533442 (33260605 to 39944496)	43149658 (38098096 to 50502225)	52630228 (46037257 to 63324449)	59484894 (51655269 to 633069475)	65637503 (55526232 to 82339430)	73608691 (60525260 to 9408331)	73549599 (58975843 to 95373146)	606252901 (521159931 to 730720808)
Excess cost (high estimate) [£]	89478274 (75186695 to 108008756)	47673180 (42975619 to 53123314)	49768036 (44870653 to 55448901)	52501434 (46944430 to 59072343)	53315249 (47699870 to 59947522)	50942562 (45602548 to 57241166)	62532680 (55189632 to 73321633)	76866377 (67183634 to 92746721)	87389922 (75811383 to 107697808)	96880730 (81804477 to 121974232)	108916925 (89341983 to 139725733)	109030988 (87176249 to 141904452)	885296357 (759787173 to 1070212581)

**Table 2.** Resource requirements each month for reintroduction of surgical activity from 1<sup>st</sup> March 2020 until 28<sup>th</sup> February 2021. Total number of procedures is presented as a time-weighted average with 95% confidence intervals derived from a linear growth model at procedure group level. Costs are presented in pounds sterling and include only excess costs associated with specific measures required for surgical time. Low estimate: four members of

staff in personal protective equipment, bed day cost of £222. High estimate: eight members of staff in personal protective equipment, bed day cost of £346. A complete breakdown of costs is in supplementary table 9. Costs are provided in Pounds Sterling (£).