

## **Title**

Burnout among healthcare professionals during COVID-19 pandemic: a cross-sectional study

## **Running title**

Staff burnout and COVID-19

## **Authors**

Mohammad Jalili, MD

Professor, Emergency Medicine Department, Tehran University of Medical Sciences, Tehran, Iran

Mahtab Niroomand, MD (Corresponding Author)

Assistant Professor, Department of Internal Medicine, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Address: 7th Floor, Bldg. No.2 SBUMS, Arabi Ave, Daneshjoo Blvd, Velenjak, Tehran, Iran.

Postal Code: 19839-63113

Phone No.: +989126191851

Email: [m.niroomand@sbmu.ac.ir](mailto:m.niroomand@sbmu.ac.ir)

Fahimeh Hadavand, MD

Assistant Professor, Infectious Diseases and Tropical Medicine Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Kataun Zeinali, MSc

Educational Supervisor, Shohadaye Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Akbar Fotouhi, MD, PhD

Professor, Department of Epidemiology and Biostatistics, Tehran University of Medical Sciences, Tehran, Iran

## **Financial Support:**

None

**Word Count:** 3324

## **Abstract**

*Background:* The unpredictable nature of the new COVID-19 pandemic and the already alarming incidence of healthcare workers being affected can have a significant impact on the psychological well-being of the staff.

*Objective:* To describe the prevalence of burnout among healthcare professionals and the associated factors.

*Design:* Cross-sectional survey.

*Setting:* Eight university affiliated hospitals in the capital city of Tehran, Iran.

*Participants:* All healthcare workers at the study sites who had been taking care of COVID-19 patients.

*Measurements:* Age, gender, marital status, having children, hospital, job category, experience, and work load, as well as the level of burnout in each subscale.

*Results:* 326 persons (53.0%) experienced high levels of burnout. The average score in emotional exhaustion, depersonalization and lack of personal accomplishment was 26.6, 10.2, and 27.3, respectively. The level of burnout in the three subscales varied based on the personal as well as work related factors and gender was the only variable that was associated with high levels of all three domains.

*Limitations:* There was no control group and thus we cannot claim a causal relationship between COVID-19 and the observed level of burnout. Not all confounding factors might have been accounted for.

*Conclusions:* Burnout is prevalent among healthcare workers caring for COVID-19 patients. Age, gender, job category, and site of practice contribute to the level of burnout that the staff experience.

*Funding source:* None

## **Introduction**

Since December 2019, when the first cases of human infection with the novel coronavirus occurred (1), the emerging infectious disease (now called COVID-19) has rapidly spread worldwide, affecting people in 210 countries and territories with the current tally exceeding 7.5 million infected people and more than 423,000 deaths (2). In addition to the lives it has claimed globally, the pandemic has led to high levels of panic and anxiety throughout the world (3-6).

Although varying in different countries, healthcare workers comprise a notable proportion of the people who contracted the illness. According to some reports, 10% of confirmed cases of COVID-19 involved healthcare providers (7). A recent report by health authorities of Iran mentioned that around 10,000 healthcare professionals have contracted COVID-19 and at least 43 are known to have died of this disease (8). This high rate of infection and mortality has a tremendous impact on the healthcare system.

Burnout is a psychological syndrome first described by Maslach et al (9) and defined as a state of psychological, emotional, and physical stress in response to prolonged exposure to occupational stress. It includes feelings of emotional exhaustion (depletion of emotional resources), depersonalization (developing cynical attitudes about patients), and reduced professional accomplishment (a sense of negative evaluation of oneself).

The deadly and uncontrollable nature of COVID-19, for which there is no known effective treatment, together with relatively high rate of infection and mortality among healthcare providers can provoke the feelings of anxiety and stress in medical staff. Issues such as social stigmatization, shortage of personal protection equipment supplies, and heavy workload on the staff can aggravate this situation. Therefore, this pandemic is expected to have substantial psychological impact on healthcare providers (10). Burnout can have serious consequences for both patients and the healthcare professionals. It not only results in poor physical and

mental health outcomes, lack of motivation, absenteeism, and low morale in the staff, but also leads to deterioration of the quality of care provided by the affected staff with resulting poor outcomes for patients. Several systematic reviews of the literature have found that high levels of burnout in health care professionals are associated with less-safe patient care (11,12).

These consequences impose immense costs on the society (13,14).

Health authorities need more information on the magnitude of this problem and its associated factors in order to better prepare for future infectious disease outbreaks, and also to adapt sound interventions and strategies to prevent further deepening of this dreadful situation.

Identification of risk factors which make certain professionals more susceptible to burnout would be an important step. These may include outbreak exposure and work related factors as well as personal characteristics.

This study was conducted on healthcare workers providing care for COVID-19 patients and aimed to explore the level of burnout among this population and also to examine factors associated with the development of this psychological sequel.

## **Methods**

### *Study Design and Participants*

This cross-sectional study was conducted during the current COVID-19 pandemic to evaluate the level of burnout among healthcare providers who were taking care of covid-19 patients at 8 university-affiliated hospitals. Altogether, these hospitals cover about one-third of Tehran province with a catchment area of about 4000000. The study was conducted about 2 months after the onset of the outbreak and when the disease had not yet been controlled.

Study participants included all healthcare providers (i.e. physicians, residents, interns, and nurses) who had taken care of COVID-19 patients at anytime during the first 2 months of the outbreak. Exclusion criteria were those with no patient contact during the outbreak or those unwilling to participate.

### *IRB Approval*

The Institutional Review Board granted approval and the requirement for written informed consent was waived based on the recognition that answering the survey instrument implied consent. Participation was voluntary and anonymity was assured. All personal information was kept confidential. Furthermore, researchers analyzed only de-identified data.

### *Data Collection*

The data collection instrument comprised of two parts: The first part of the tool asked questions pertaining socio-demographic and work-related characteristics. Participants were requested to indicate their age, gender, marital status, number of children, job title, place of work, and years of experience. This section also asked whether the respondent had been involved in the care of corona patients and if yes for how many shifts a month and how many hours per shift. The second part of the study tool was a translated version of Maslach Burnout Inventory (MBI) for Human Services Survey. In order to limit the study to burnout related to COVID-19, the phrase “due to COVID-19” was added to each item. MBI is an internationally

recognized, validated, self-report questionnaire for measuring the severity of workplace burnout (15), using the three dimensions of emotional exhaustion, depersonalization, and personal accomplishment. The questionnaire has 22 items and each item is answered on a five-point Likert scale. This tool has been extensively used in many studies in different parts of the world and the Persian translation has also been validated previously (16).

The study used a convenience sampling method for recruitment. Invitation to participate in the study was made through professional as well as informal networks. The message included an invitation explaining the purpose of the study, the name and contact details of the principal investigator, and a live link to the host survey platform (E Poll). The online self-administered questionnaire instructed participants to respond to the statements in the tool in relation to the COVID-19 outbreak. Verbal reminders through face-to-face contact and phone calls were performed by a site coordinator at each hospital.

#### *Statistical Analysis*

Burnout is expressed by scores of each of the three MBI subscales, with a high score meaning a high level of burnout. Each subscale score is calculated by adding up all scores of all items in that subscale, with the notion that the items on personal accomplishment domain are reversely scored (9,15). Scores range from 0 to 36 for emotional exhaustion, from 0 to 20 for depersonalization, and from 0 to 32 for personal accomplishment subscale. The standard cut-off values were used to define low, moderate, and high levels in each dimension (9,15).

We calculated the “average daily workload” by multiplying the number of shifts per month by the number of work hours per shift divided by 30 (number of days of a month). We also defined “high burnout” as obtaining a moderate or high score in either emotional exhaustion or depersonalization axes.

Upon completion of the study period, the data was downloaded from the online survey tool entered into an excel spreadsheet and anonymized. This data was, then, imported into the statistical software for analysis.

Descriptive analyses were conducted to examine all baseline characteristics of the participants and the outcome variables. Means and standard deviations (SDs) were calculated for continuous variables, while frequencies and percentages were produced for categorical variables. All reported p values are 2-tailed, and  $p < 0.05$  was considered statistically significant. In order to explore the risk factors associated with the development of burnout, multivariable logistic regression models were used. Odds ratios and corresponding 95% confidence intervals (CI) were reported.

#### *Role of the Funding Source*

No financial support or funding was provided for this study.



## **Results**

We invited a total of 1,002 healthcare workers to participate in the study, from which 645 people completed the survey (response rate: 64.4%). Thirty respondents declared that they had not taken care of COVID-19 patients, leaving a total of 615 completed questionnaires for analysis.

Table 1 illustrates the baseline characteristics as well as the mean MBI scores in each subscale in the studied population.

Table 1. Baseline characteristics of participants (n=615)

Variable	Value
Age in years, Mean $\pm$ SD, (range)	34.4 $\pm$ 7.8, (22-67)
Gender, n (%)	
Female	403 (65.5)
Male	208 (33.8)
Prefer not to say	4 (0.7)
Marital status, n (%)	
Married	363 (59.0)
Single	242 (39.4)
Prefer not to say	10 (1.6)
Home living children, n (%)	
Yes	229 (37.2)
No	386 (62.8)
Job category, n (%)	
Specialist	68 (11.1)
Resident	166 (27.0)
Intern	19 (3.1)
Nurse	300 (48.8)
Other	62 (10.1)
Years in practice, Mean $\pm$ SD	8.8 $\pm$ 6.7
days working over past month, Mean $\pm$ SD	18.1 $\pm$ 9.1
hours working per shift, Mean $\pm$ SD	13.3 $\pm$ 6.5
Maslach Burnout Inventory Score, Mean $\pm$ SD	
Total score	64.2 $\pm$ 7.7

---

Emotional Exhaustion subscale score	26.6 ±7.4
Depersonalization subscale score	10.2 ±2.2
Personal Accomplishment subscale score	27.3 ±3.9

---

n: number; SD: standard deviation; COVID-19: corona virus disease 2019

Our findings showed that 55 (8.9%), 252 (41.0%), and 308 (50.1%) people experienced low, moderate, and high levels in the emotional exhaustion axis, respectively. In the depersonalization axis, the values for low, moderate, and high levels of burnout were 5 (0.8%), 529 (86.0%), and 81(13.2%), respectively. In addition, only one (0.8%) participant scored low in personal accomplishment axis, while 88 (14.3%) and 526 (85.5%) people were shown to experience moderate or high levels, respectively.

Table 2 illustrates how the level of burnout in each subscale varies between different groups of participants in terms of sociodemographic variables and job category. Among the studied characteristics, gender was the only variable that was associated with high levels of all three domains of burnout. High emotional exhaustion was significantly more prevalent in females and those without children, as well as people from different job categories (most common in residents and then nurses). High levels of depersonalization were significantly more common in males and younger ( $\leq 36$  years) participants.

Table 2. Participants' level of burnout in each dimension by sociodemographic characteristics and job category, number (%)

	Emotional Exhaustion				Depersonalization				Personal Accomplishment			
	Low	Moderate	High	P	Low	Moderate	High	P	Low	Moderate	High	P
<b>Age</b>												
≤36 year	36(8.9)	154(38)	215(53.1)	0.105	3(0.7)	338(83.5)	64(15.8)	0.011	1(0.2)	59(14.6)	345(85.2)	0.723
>36 year	19(9.0)	98(46.7)	93(44.3)		2(1.0)	191(91.0)	17(8.1)		0 (0)	29(13.8)	181(86.2)	
<b>Gender</b>												
Female	30(7.4)	148(36.7)	225(55.8)	<0.001	3(0.7)	359(89.1)	41(10.2)	0.003	0(0.0)	45(11.2)	358(88.8)	0.003
Male	25(12.0)	102(49.0)	81(38.9)		2(1.0)	166(79.8)	40(19.2)		1(0.5)	42(20.2)	165(79.3)	
<b>Marital status</b>												
Married	40(11.0)	142(39.1)	181(49.9)	0.436	4(1.1)	318(87.6)	41(11.3)	0.077	1(0.3)	57(15.7)	305(84.0)	0.327
Single	15(6.2)	105(43.4)	122(50.4)		1(0.4)	202(83.5)	39(16.1)		0(0.0)	30(12.4)	212(87.6)	
Other	0(0.0)	5(50.0)	5(50.0)		0(0.0)	9(90.0)	1(10.0)		0(0.0)	1(10.0)	9(90.0)	
<b>Having children</b>												
Yes	24(10.5)	104(45.4)	101(44.1)	0.027	2(0.9)	206(90.0)	21(9.2)	0.033	1(0.4)	43(18.8)	185(80.8)	0.010
No	31(8.0)	148(38.3)	207(53.6)		3(0.8)	323(83.7)	60(15.5)		0(0)	45(11.7)	341(88.3)	

Job category												
Specialist	7(10.3)	31(45.6)	30(44.1)	0.008	0(0.0)	65(95.6)	3(4.4)	0.657	0(0.0)	8(11.8)	60(88.2)	<0.001
Resident	7(4.2)	59(35.5)	100(60.2)		2(1.2)	134(80.7)	30(18.1)		0(0.0)	7(4.2)	159(95.8)	
Intern	1(5.3)	14(73.7)	4(21.1)		0(0.0)	18(94.7)	1(5.3)		0(0.0)	3(16.8)	16(84.2)	
Nurse	26(8.7)	115(38.3)	159(53.0)		3(1.0)	257(85.7)	40(13.3)		1(0.3)	47(15.7)	252(84.0)	
Other	14(22.6)	33(53.2)	15(24.2)		0(0.0)	55(88.7)	7(11.3)		0(0.0)	23(37.1)	39(62.9)	

p: p value

We noted that of all 615 participants in the study, who had completed the MBI questionnaire, 326 persons (53.0%) experienced high levels of burnout (defined as high levels in either domains of emotional exhaustion and depersonalization). Table 3 shows the distribution of high burnout in different groups of participants based on their sociodemographic characteristics and work-related factors.

Table 3. Frequency of high burnout in different groups of participants (n=615)

Variable	High burn out level, n (%)		P value
	No	Yes	
<b>Age</b>			
≤36 years	177 (43.7)	228 (56.3)	0.041
>36 years	110 (52.4)	100 (47.6)	
<b>Gender</b>			
Female	169 (41.9)	234 (58.1)	0.002
Male	116 (55.8)	92 (44.2)	
Other	2 (50)	2 (50)	
<b>Marital status</b>			
Married	172 (47.4)	191 (52.6)	0.877
Single	110 (45.5)	132 (54.5)	
Other	5 (50.0)	5 (50.0)	
<b>Home living children</b>			
Yes	120 (52.4)	109 (47.6)	0.030
No	167 (43.3)	219 (56.7)	
<b>Job category</b>			
Intern	14 (73.7)	5 (26.3)	<0.001
Resident	58 (34.9)	108 (65.1)	
specialist	37 (54.4)	31 (45.6)	
Nurse	134 (44.7)	166 (55.3)	
Other	44 (71.0)	18 (29.0)	
<b>Hospital</b>			
Shohada	105 (57.1)	79 (42.9)	0.002



---

Imam Hussein	107 (41.8)	149 (58.2)	
Masih Daneshvari	42 (50)	42 (50)	
Other	33 (36.3)	58 (63.7)	
Years in practice			
≤ 5 years	112 (41.9)	155 (58.1)	0.042
> 5 years	175 (50.3)	173 (49.7)	
Average daily workload*			
≤ 8 hours	202 (47.4)	224 (52.6)	0.561
> 8 hours	83 (44.9)	102 (55.1)	

---

\*There were 4 missing values.

The results of multiple logistic regression analysis of different variables to detect their impact on the development of high burnout levels have been presented in Table 4. As can be seen in the table, of different variables included in analysis, age, gender, job category and hospital were significantly associated with burnout status; while marital status, having children, number of years in practice, and the average daily workload were not associated with this outcome.

Table 4. Bivariate associations between high level of burnout and sociodemographic and work-related factors (n=611)

Variable	OR	95% CI for OR		P value
		Lower limit	Upper limit	
Age*	0.97	0.95	0.99	0.010
Gender				
• Female	1			0.015
• Male	0.65	0.45	0.92	
Job category				
• Intern	1			<0.001
• Resident	6.64	2.16	20.41	0.001
• specialist	4.13	1.18	4.40	0.026
• Nurse	4.56	1.54	13.48	0.006
• Other	1.77	0.53	5.90	0.350
Hospital				
• Shohada	1			0.011
• Imam Hussein	1.86	1.25	2.78	0.002
• Masih Daneshvari	1.29	0.74	2.23	0.364
• Other	1.96	1.08	3.53	0.026

n: number; yr: year; SD: standard deviation; CI: confidence interval; OR: odds ratio

\*Per 1 year increase

## Discussion

The results of this cross-sectional survey demonstrated high levels of burnout in healthcare workers looking after COVID-19 patients. We also found that younger age and female gender were predisposing factors for burnout and that the level of burnout varied significantly by site of practice and job category (with residents being the most vulnerable group).

The results of this study showed that high levels of burnout are present among healthcare professionals who take care of COVID-19 patients. Other studies have also investigated the psychological impact of COVID-19 and other infectious disease outbreaks on hospital staff.

A recent rapid review of the existing literature retrieved 59 papers that described the emotional reactions of healthcare providers dealing with patients during an outbreak of viral infectious disease. The investigators concluded that staff in contact with these patients had greater levels of acute or posttraumatic stress (odds ratio 1.71, 95% confidence interval 1.28 to 2.29) and psychological distress (1.74, 1.50 to 2.03) as compared with lower risk controls (17). Most of these studies used instruments other than MBI, and only eight were on COVID-19. Poon et al, using State-Trait Anxiety Inventory, conducted a survey to identify anxiety levels among 1926 front-line health care workers in Hong Kong during SARS outbreak. They also used the emotional exhaustion dimension of the MBI. Mean anxiety levels (51.1 versus 47.1;  $P < 0.001$ ) and the number of burnout symptoms (7.3 versus 5.1;  $P < 0.001$ ) among front-line health care workers were higher than those among controls; anxiety scores correlated with burnout scores (18). In another study by Fiksenbaum and colleagues, an on-line survey of 333 nurses (315 women, 18 men) working in the province of Ontario, Canada was performed to investigate the role of perceived SARS threat and organizational support in predicting emotional exhaustion and stated anger (19). In their study, mean emotional exhaustion score was 4 ( $\pm 1.85$ ). In a cross-sectional survey conducted in China, using General Health Questionnaire, Dai et al evaluated mental health status of a convenient sample

of 4357 healthcare workers (20). They found that 1,704 workers (39.1%) had psychological distress (score $\geq$ 3). In another survey, Lai and colleagues assessed the magnitude of mental health outcomes of 1257 healthcare workers treating patients exposed to COVID-19 in 34 hospitals in China. They found that 634 people (50.4%) reported symptoms of depression, 560 (44.6%) anxiety, 427 (34.0%) insomnia, and 899 (71.5%) distress (21). Moreover, Liu and colleagues, using the Zung Self-rating Anxiety Scale, estimated the prevalence of anxiety among medical staff to be 12.5% (22). Xing et al in a cross-sectional survey using Symptom Checklist-90 (SCL-90) investigated the mental health status of 548 medical personnel dealing with COVID-19 at 12 hospitals in eight provinces and cities of China (23). According to the findings of their study, overall average score of SCL-90 and mean values of factors (somatization, obsessive-compulsive, anxiety, phobic anxiety, and psychoticism) among medical personnel were significantly higher than that of the norm group. Zhu et al evaluated stress, depression, and anxiety among 5062 health workers using Impact of Event Scale-Revised, Patient Health Questionnaire-9, and Generalized Anxiety Disorder 7-item. In their study 29.8 percent of participants reported stress, 13.5% depression, and 24.1% anxiety (24). In a study by Maunder on 587 workers at Toronto Hospital, 30.4% had high emotional exhaustion (score $\geq$ 27) (25). To the best of our knowledge, no prior study used MBI to assess the psychological sequel of COVID-19; therefore, a comparison of the results of this study with that of previous studies is not feasible. Burnout is a very well known consequence of healthcare for the staff and is strongly linked to significant outcomes such as patient safety (11,12); Moreover, MBI is a validated instrument to assess the level of burnout. Therefore, we believe that MBI is a practical and credible method for assessment of the psychological impact of COVID-19 pandemic on staff.

Our study lacked a control group but a comparison of our findings with other studies on burnout in our country before the COVID-19 pandemic shows a much higher level of

burnout. A systematic review of the studies which examined the prevalence of burnout among nurses in Iran, published in 2018, found that the overall prevalence of burnout among Iranian nurses was 36% (95% CI, 20-53%) (26). In our study, more than 55% of the nurses experienced high burnout during the COVID-19 pandemic. In a cross-sectional study conducted on 208 primary health care providers in Iran, the average score of the participants in emotional exhaustion dimension was 17.19, while in our study the average score was around 27 in this subscale (27). A large study to assess the level of job burnout among 1807 healthcare providers in Iran found that mean scores ( $\pm$ SD) in emotional exhaustion, depersonalization, and personal accomplishment subscales were 8.9 ( $\pm$  9.0), 23 ( $\pm$  2.9), and 34 ( $\pm$  8.6), respectively (28); while in our study these values were much higher: 26.6 ( $\pm$ 7.4), 10.2 ( $\pm$ 2.2), and 27.3 ( $\pm$ 3.9), respectively.

Several outbreaks of infectious diseases have occurred over the past two decades and they represent a serious threat to both health services and staff. Healthcare providers, who need to take care of a large number of potentially infectious victims, are under great physical and psychological pressure. Therefore, it is not surprising to expect a high rate of infection as well as mental health problems among the staff. Chen et al interviewed 13 medical staff at a hospital in China during the COVID-19 outbreak and noted that “getting infected was not an immediate worry to staff” but that “they were afraid of bringing the virus to their home”.

They also stated that issues such as ambiguity regarding “how to deal with patients when they were unwilling to be quarantined at the hospital or did not cooperate with medical measures because of panic or a lack of knowledge about the disease” and “shortage of protective equipment” were also contributing to poor mental health of medical staff (29). While our study was not meant to discover the reasons for this phenomenon, several issues can be speculated to have resulted in this state of high burnout. Heavy workload, fear of contracting the disease or transmitting it to the family, as well as lack of staff support systems are often

cited complaints. Whether these are linked to the development of high burnout or not and the contribution of each factor can be the subject of future studies. This will help administrators adapt the best approach to maintain staff mental health.

Our study found age and gender to be important predisposing factors for high burnout. Several studies have shown that staff who are women (3,21,24,30-32) or younger (33-37) are more vulnerable to psychological distress. Our findings are in accordance with these studies. However, in two studies on personnel taking care of patients with covid-19, older age was a risk factor for psychological symptoms (23,36).

While some studies have found out that having children (38,39) and having an increased contact with affected patients (18-21,25,30-32,34-36,38-45,47,48) are predisposing factors, our study failed to find such a relationship. Moreover, several studies have shown that being less experienced (3,30,42,47) puts the staff at higher risk. While we found a significant difference between the experience of those with and without high burnout, this variable did not remain in the final regression model, pointing to potential co-linearity between age and experience.

Within the job categories, residents were found in this study to be at higher risk of developing high burnout. Nurses were the second at-risk group. In previous studies, nurses were generally at higher risk than doctors (18,21,34,35,38,39,48,49). Lai et al, for example, reported more severe degrees of all measurements of mental health symptoms in nurses exposed to COVID-19 (21). However, two studies reported the opposite (50,51). A survey of 99 residents during AH1N1 influenza virus outbreak in a third level hospital in Mexico City conducted in 2009 documented high levels of burnout among this population (33). The reasons why residents were so vulnerable to burnout needs further study, but issues such as heavy workload, changes in duty rosters to accommodate the new needs, and cancellation of

vacations, as well as having less access to personal protective equipment may have caused this unfavorable state.

The level of burnout in our study was also affected by the location of work. In a study by Maunder on 769 healthcare workers during SARS outbreak, the investigators found a significantly higher levels of burnout in the staff of one of the two studied hospitals (median score 19, interquartile (IQR) range 10–29, in one hospital versus median score 16, IQR 9– 23, in the other hospital) (25). Dai et al found that working in primary hospitals were poor prognostic factors (20). In our study, working in the hospitals which were not designated as COVID-19 centers was accompanied by higher levels of burnout. Several reasons might account for this difference: working conditions affects perceived threat; designated hospitals are better prepared and equipped for healthcare delivery to COVID-19 patients; working in non-designated hospitals is not rewarded by recognition and incentives found in COVID-19 designated hospitals.

#### *Strengths and limitations*

Our study has several strengths. Burnout is a well recognized psychological consequence of working in the healthcare section and its impact on staff physical and emotional health, patient safety, and quality of care is documented. The instrument used in this study is well validated across geographical areas. We used a relatively large sample with an acceptable response rate.

This study also faces some limitations. Although we tried to consider several personal (e.g. marital status and having dependent children) and work-related parameters (e.g. workload, job category), our study might have failed to deal with all confounding factors. We specifically referred to COVID-19 in our questions; however, a comparison with other clinicians or the general population has not been performed and we cannot claim that this level of burnout is solely attributable to COVID-19 pandemic.



### *Suggestions for future research*

Other studies, using qualitative methods, are necessary to delve into the reasons for this observed phenomenon. In-depth interviews can bring important issues into consideration, which can be the subject of further studies. The necessity of implementation and evaluation of various measures and strategies to overcome this high prevalence of burnout among staff, through interventional studies, cannot be overemphasized. Last but not least, further assessment of the long term sequel of COVID-19 pandemic through follow up studies deserves particular attention.

### *Conclusion*

Our study highlighted the high prevalence of burnout among healthcare workers who are caring for patients during the new pandemic. Several personal and work related factors contribute to the level of burnout that the staff experience. These include younger age, female gender, working as a resident or nurse, and working in non-COVID-designated hospitals.

## References

1. Jin Y-H, Cai L, Cheng Z-S, Cheng H, Deng T, Fan Y-P, et al. A rapid advice guideline for the diagnosis and treatment of 2019 novel coronavirus (2019-nCoV) infected pneumonia (standard version). *Military Medical Research*. 2020;7(1):4.
2. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>.
3. Chong MY, Wang W-C, Hsieh WC, et al. Psychological impact of severe acute respiratory syndrome on health workers in a tertiary hospital. *Br J Psychiatry*. 2004;185:127–133.
4. Sim K, Chua HC. The psychological impact of SARS: a matter of heart and mind. *CMAJ*. 2004;170(5):811– 812.
5. World Health Organization. Summary table of SARS cases by country, 1 November 2002—2007 August 2003. Geneva (CH): WHO; 2003.
6. SARS Commission. Spring of fear, volumes 1, 2 and 3. Toronto (ON): SARS Commission (Canada); 2006
7. In Memoriam: Healthcare Workers Who Have Died of COVID-19. Available through:[https://www.medscape.com/viewarticle/927976#vp\\_1](https://www.medscape.com/viewarticle/927976#vp_1). Accessed 16 Apr 2020.
8. E. B. Ing1, Q. (A.) Xu2, A. Salimi3 and N. Torun. Physician deaths from corona viru(COVID-19) disease. *Occupational Medicine*, 16 May2020;
9. Maslach C, Jackson S, Leiter M. Maslach burnout inventory manual. Mountain View, CA: CPP. Inc., and Davies-Black, 1996.
10. Jianming Guo , Lianming Liao , Baoguo Wang , Xiaoqiang Li, Lianrui Guo , Zhu Tong, Qinghua Guan, Mingyue Zhou, Yingfeng Wu , Jian Zhang , Yongquan Gu. Psychological effects of COVID-19 on hospital staff: a national cross-sectional survey of China mainland, *The Lancet Psychiatry*,2020

11. Dewa CS, Loong D, Bonato S, Trojanowski L. The relationship between physician burnout and quality of healthcare in terms of safety and acceptability: a systematic review. *BMJ Open*. 2017;7(6):e015141.doi:10.1136/bmjopen-2016-015141
12. Hall LH, Johnson J, Watt I, Tsipa A, O'Connor DB. Healthcare staff wellbeing, burnout, and patient safety: a systematic review. *PLoS One*. 2016;11(7):e0159015. doi:10.1371/journal.pone.0159015
13. Dewa CS, Jacobs P, Thanh NX, Loong D. An estimate of the cost of burnout on early retirement and reduction in clinical hours of practicing physicians in Canada. *BMC Health Serv Res*. 2014;14:254.
14. Shanafelt TD, Dyrbye LN, West CP, Sinsky CA. Potential impact of burnout on the US physician workforce. *Mayo Clin Proc*. 2016;91:1667-1668.
15. Christina Maslach, Susan E. Jackson, The measurement of experienced burnout. *Journal of Occupational Behaviour*.1981; 2: 99-113
16. Saba Moalemi, Zahra Kavosi, Najimeh Beygi, Azizallah Deghan, Aliasghar Karimi, Mohammad Mahdi Parvizi. Evaluation of the Persian Version of Maslach Burnout Inventory- Human Services Survey among Iranian Nurses: Validity and Reliability. *GMJ*.2018;7:e995
17. Steve Kisely, Nicola Warren, Laura McMahon, Christine Dalais, Irene Henry, Dan Siskind. Occurrence, prevention, and management of the psychological effects of emerging virus outbreaks on healthcare workers: rapid review and meta-analysis. *BMJ*. 2020; 369: m1642. Published online 2020 May 5. doi: 10.1136/bmj.m1642
18. Poon E, Liu KS, Cheong DL, Lee CK, Yam LYC, Tang WN. Impact of severe acute respiratory syndrome on anxiety levels of frontline health care workers. *Hong Kong Med J* 2004;10:325-30

19. Fiksenbaum L, Marjanovic Z, Greenglass E, Coffey S. Emotional exhaustion and state anger in nurses who worked during the SARS outbreak: The role of perceived threat and organizational support. *Canadian Journal of Community Mental Health*. 2006;25(2):89-103
20. Dai Y, Hu G, Xiong H. Psychological impact of the coronavirus disease 2019 (COVID-19) outbreak on healthcare workers in China. medRxiv 2020.03.03.20030874 [Preprint]. 2020.
21. Lai J, Ma S, Wang Y, et al. Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw Open*. 2020;3(3):e203976. Published 2020 Mar 2. doi:10.1001/jamanetworkopen.2020.3976
22. Liu CY, Yang YZ, Zhang XM, et al. The prevalence and influencing factors in anxiety in medical workers fighting COVID-19 in China: a cross-sectional survey. *Epidemiol Infect*. 2020;148:e98. Published 2020 May 20. doi:10.1017/S0950268820001107
23. Xing J, Sun N, Xu J, Geng S, Li Y. Study of the mental health status of medical personnel dealing with new coronavirus pneumonia. *PLoS One*. 2020;15(5):e0233145. Published 2020 May 19. doi:10.1371/journal.pone.0233145
24. Zhu Z, Xu S, Wang H. COVID-19 in Wuhan: Immediate Psychological Impact on 5062 Health Workers. medRxiv 2020 2020.02.20.20025338 [Preprint].
25. Maunder RG, Lancee WJ, Balderson KE, et al. Long-term psychological and occupational effects of providing hospital healthcare during SARS outbreak. *Emerg Infect Dis*. 2006;12(12):1924-1932. doi:10.3201/eid1212.060584
26. Rezaei S, Karami Matin B, Hajizadeh M, Soroush A, Nouri B. Prevalence of burnout among nurses in Iran: a systematic review and meta-analysis. *Int Nurs Rev*. 2018;65(3):361-369. doi:10.1111/inr.12426

27. Khosravi A, Ranjbar M, Erfan A, Sadeghi Z, khojasteh L, Amiri M. "Burnout in Primary Health Care Providers in Mazandaran Province." *International Journal of Health Studies* [Online], 3.4 (2017): n. pag. Web. 31 May. 2020
28. Bazmi E, Alipour A, Yasamy MT, et al. Job Burnout and Related Factors among Health Sector Employees. *Iran J Psychiatry*. 2019;14(4):309-316
29. Chen Q, Liang M, Li Y, Guo J, Fei D, Wang L, He L, Sheng C, Cai Y, Li X, Wang J, Zhang Z. Mental health care for medical staff in China during the COVID-19 outbreak. *Lancet Psychiatry*. 2020 Apr;7(4):e15-e16. doi: 10.1016/S2215-0366(20)30078-X. Epub 2020 Feb 19.
30. Tang L, Pan L, Yuan L, Zha L. Prevalence and related factors of posttraumatic stress disorder among medical staff members exposed to H7N9 patients. *Int J Nurs Sci* 2016;4:63-7. doi:10.1016/j.ijnss.2016.12.002
31. Bukhari EE, Temsah MH, Aleyadhy AA, et al. Middle East respiratory syndrome coronavirus (MERS-CoV) outbreak perceptions of risk and stress evaluation in nurses. *J Infect Dev Ctries* 2016;10:845-50. doi:10.3855/jidc.6925
32. Huang L, Xu F, Liu H. Emotional responses and coping strategies of nurses and nursing college students during COVID-19 outbreak. medRxiv 2020.03.05.20031898 [Preprint].2020.
33. Austria-Corrales F, Cruz-Valdés B, Herrera-Kiengelher L, Vázquez-García JC, Salas-Hernández J. [Burnout syndrome among medical residents during the influenza A H1N1 sanitary contingency in Mexico.] *Gac Med Mex* 2011;147:97-103.
34. Nickell LA, Crighton EJ, Tracy CS, et al. Psychosocial effects of SARS on hospital staff: survey of a large tertiary care institution. *CMAJ* 2004;170:793-8. doi:10.1503/cmaj.1031077

35. Tam CWC, Pang EPF, Lam LCW, Chiu HF. Severe acute respiratory syndrome (SARS) in Hong Kong in 2003: stress and psychological impact among frontline healthcare workers. *Psychol Med* 2004;34:1197-204. doi:10.1017/S0033291704002247
36. Wu P, Fang Y, Guan Z, et al. The psychological impact of the SARS epidemic on hospital employees in China: exposure, risk perception, and altruistic acceptance of risk. *Can J Psychiatry* 2009;54:302-11. doi:10.1177/070674370905400504
37. Sim K, Chong PN, Chan YH, Soon WS. Severe acute respiratory syndrome-related psychiatric and posttraumatic morbidities and coping responses in medical staff within a primary health care setting in Singapore. *J Clin Psychiatry* 2004;65:1120-7. doi:10.4088/JCP.v65n0815
38. Koh D, Lim MK, Chia SE, et al. Risk perception and impact of Severe Acute Respiratory Syndrome (SARS) on work and personal lives of healthcare workers in Singapore: what can we learn? *Med Care* 2005;43:676-82. doi:10.1097/01.mlr.0000167181.36730.cc
39. Maunder RG, Lancee WJ, Rourke S, et al. Factors associated with the psychological impact of severe acute respiratory syndrome on nurses and other hospital workers in Toronto. *Psychosom Med* 2004;66:938-42. doi:10.1097/01.psy.0000145673.84698.18.
40. Lee SM, Kang WS, Cho A-R, Kim T, Park JK. Psychological impact of the 2015 MERS outbreak on hospital workers and quarantined hemodialysis patients. *Compr Psychiatry* 2018;87:123-7. doi:10.1016/j.comppsy.2018.10.003
41. Kim JS, Choi JS. Factors influencing emergency nurses' burnout during an outbreak of Middle East Respiratory Syndrome Coronavirus in Korea[Korean Soc Nurs Sci]. *Asian Nurs Res (Korean Soc Nurs Sci)* 2016;10:295-9. doi:10.1016/j.anr.2016.10.002
42. Oh N, Hong N, Ryu DH, Bae SG, Kam S, Kim KY. Exploring Nursing Intention, Stress, and Professionalism in Response to Infectious Disease Emergencies: The Experience of Local Public Hospital Nurses During the 2015 MERS Outbreak in South Korea[Korean Soc

Nurs Sci]. *Asian Nurs Res (Korean Soc Nurs Sci)* 2017;11:230-6.

doi:10.1016/j.anr.2017.08.005

43. Matsuishi K, Kawazoe A, Imai H, et al. Psychological impact of the pandemic (H1N1)

2009 on general hospital workers in Kobe. *Psychiatry Clin Neurosci* 2012;66:353-60.

doi:10.1111/j.1440-1819.2012.02336.x

44. Styra R, Hawryluck L, Robinson S, Kasapinovic S, Fones C, Gold WL. Impact on health

care workers employed in high-risk areas during the Toronto SARS outbreak. *J Psychosom*

*Res* 2008;64:177-83. doi:10.1016/j.jpsychores.2007.07.015

45. Wu P, Fang Y, Guan Z, et al. The psychological impact of the SARS epidemic on

hospital employees in China: exposure, risk perception, and altruistic acceptance of risk. *Can*

*J Psychiatry* 2009;54:302-11. doi:10.1177/070674370905400504

46. Liu X, Kakade M, Fuller CJ, et al. Depression after exposure to stressful events: lessons

learned from the severe acute respiratory syndrome epidemic. *Compr Psychiatry* 2012;53:15-

23. doi:10.1016/j.comppsy.2011.02.003

47. Lancee WJ, Maunder RG, Goldbloom DS, Coauthors for the Impact of SARS Study.

Prevalence of psychiatric disorders among Toronto hospital workers one to two years after

the SARS outbreak. *Psychiatr Serv* 2008;59:91-5. doi:10.1176/ps.2008.59.1.91

48. Goulia P, Mantas C, Dimitroula D, Mantis D, Hyphantis T. General hospital staff worries,

perceived sufficiency of information and associated psychological distress during the

A/H1N1 influenza pandemic. *BMC Infect Dis* 2010;10:322. doi:10.1186/1471-2334-10-322.

49. Wong TW, Yau JK, Chan CL, et al. The psychological impact of severe acute respiratory

syndrome outbreak on healthcare workers in emergency departments and how they cope. *Eur*

*J Emerg Med* 2005;12:13-8. doi:10.1097/00063110-200502000-00005

50. Chan AOM, Huak CY. Psychological impact of the 2003 severe acute respiratory syndrome outbreak on health care workers in a medium size regional general hospital in Singapore. *Occup Med(Lond)* 2004;54:190-6. doi:10.1093/occmed/kqh027
51. Lung F-W, Lu Y-C, Chang Y-Y, Shu BC. Mental symptoms in different health professionals during the SARS attack: A follow-up study. *Psychiatr Q* 2009;80:107-16. doi:10.1007/s11126-009-9095-5