1 Research article

2 Willingness and acceptability of anxiety and depression screening 3 among women living with HIV in Ethiopia

4
5 Abdilahi Yousuf¹, Ramli Musa², Muhammad Lokman Md. Isa³, Siti Roshaidai Mohd Arifin*⁴

6
7
8 ¹ College of medicine and health sciences, Jijiga University, Jijiga, Ethiopia.

9 ² Department of psychiatric, Kuliyah of medicine, IIUM, Kuantan, Malaysia.

10 ³ Department of Basic Medical Sciences, Kuliyah of Nursing, IIUM, Kuantan, Malaysia.

11 ⁴ Academic and industrial linkages, Kuliyah of Nursing, IIUM, Kuantan, Malaysia.

12 *Corresponding author: roshaidai@iium.edu.my

13

14 Abstract

15 **Introduction:** Integration of related health services, such as screening of depression in HIV
16 care is crucial for improving the quality of care and enhancing the use of scarce resources in
17 developing countries. However, commonly these interrelated health services are commonly
18 provided separately and there are many missed opportunities. Little is known about the client-
19 related non-implementation issues. This study aims at examining the acceptability of anxiety
20 and depression screening among women living with HIV.

21 **Material and methods:** This was a facility based cross-sectional study which included
22 women living with HIV attending two hospitals in Jijiga town, Ethiopia. The study
23 participants were identified using systematic random sampling method. An exit interview was
24 conducted with the use of a pretested questionnaire. The gathered data was analysed using
25 SPSS version 23 and multivariate logistic regression model was used to determine factors
26 associated with the factors associated with the acceptance of anxiety and depression
27 screening.

28 **Results:** A total of 409 women participated in this study. Though, only 115 (28.1%) were
29 aware about the existence of anxiety and depression screening, 357 (87.3%) accepted to
30 receive the screening for anxiety and depression. Requisite of partner approval was the most
31 common reason for not accepting the screening of anxiety and depression 21 (40.4%). When
32 used multivariate logistic regression model; holding college level education, divorced and
33 were living without partner, being self employed by occupation, aware of the existing
34 screening service, source of information from health care providers and history of previous
35 screening were significantly associated with acceptance of depression screening.

36 **Conclusion:** This research has shown that women participant living with HIV were willing to
37 undergo the screening for depression, hence future interventions should focus on the
38 integration of mental health screening in HIV clinical setup.

39 *Key words: Depression, HIV, AIDS, acceptability, screening.*

40 **Introduction**

41

42 Depression, a common mental disorder which affects more than 300 million individuals of all
43 age groups in this world wide. This disorder which is more common in women than men, is
44 the primary cause of disability and one of the main contributors of global disease burden (1)
45 including suicide (2,3).

46 Depression affects about 20% of women during their lifetime, with reproductive health
47 problems being a period of high vulnerability (4). It is also predicted that depression which is
48 estimated to be the second leading cause of disease and disability in the world in 2020, will
49 be twice common in women than men (5–9).

50 Anxiety and depression is common among people living with HIV infection (5,10–18). High
51 and low-income countries have presented higher rates of depression in people living with
52 HIV when compared with HIV negative individuals. The level of depression mostly appears
53 to be related to the severity of symptoms of HIV infection (19).

54 In developing countries, depression is also one of the main challenges that should be
55 addressed when dealing with HIV prevention strategies, as the two illnesses are intensely
56 associated and aggravate each other. In recent evidences, depression and depressive
57 symptoms are found to be the key predictors of poor adherence to HIV treatment (20) and
58 negatively impact on clinical outcomes (21).

59 So far, depression is highly prevalent in Ethiopia, and it is considered as one of the common
60 psychiatric disorders affecting women as its associates are gender-specific (5–10,14,22,23).
61 One of the findings conducted in Ethiopia showed that there is high prevalence of depression
62 on people living with HIV/AIDS and stated that 38.94% of the studied participants were
63 depressed (7). Another systematic review conducted in Ethiopia has presented that the pooled
64 prevalence of depression was higher among People Living with HIV than the general
65 population (24).

66 Regardless of its prevalence among people living with HIV, depression and anxiety are
67 commonly under diagnosed by health care providers. More than half of those eligible
68 patients, miss the opportunity to be screened for anxiety and depression disorders by doctors
69 (25).

70 Furthermore, people with chronic illnesses are usually unwilling to seek health professional
71 support and counselling. A mong people who manifest depressive symptoms, less number is
72 reported to seek treatment in their course of disorder (26,27).

73 Even though the significance of depression screening for the intervention and care of HIV
74 patients in health facilities has been long recognized, yet little is known about the
75 acceptability and predictive factors related with the acceptance of anxiety and depression
76 screening among women attending HIV care services. Therefore, this study will examine the
77 acceptability and the predictive factors associated with screening of anxiety and depression
78 among women Living with HIV. The findings of this study can be a baseline for the
79 introduction of anxiety and depression screening in HIV care services of health facilities in
80 Ethiopia.

81

82 **Methods and materials**

83 **Study design:**

84 A facility-based cross-sectional study was carried out in Antiretroviral Therapy clinics of two
85 hospitals. This study was the first phase of extensive mixed-method research conducted
86 among women with HIV.

87

88 **Study population:**

89 The study population was women aged 18 years or older who are attending HIV treatment
90 service in two hospitals.

91

92 **Sample size determinations and sampling procedure:**

93 A single population proportion formula, was used to estimate the sample size of women
94 participants to be studied with a 95% CI and 5% margin of error. (P) the prevalence of
95 depression from a baseline study in Addis Ababa, Ethiopia (i.e. 41.2% of studied respondents
96 were depressive). Expecting a 10% of non-response rate the total sample size was 409
97 patients.

98 A systematic sampling procedure was used for the selection of the study participants. To
99 determine the sampling interval, the total study population who had an appointment during
100 the data collection was divided by the total sample size, then the initial point was randomly
101 selected.

102 **Data collection**

103 A trained nurses conducted an interview with women attending the antiretroviral treatment
104 centres. A pretested structured questionnaire was used. Participants were asked about their
105 socio-demographic status, their clinical and laboratory investigation results were also
106 retrieved and questions regarding about their awareness on depression screening, their
107 previous test and their acceptance of the screening were asked.

108 **Screening**

109 For the initial screening of the patients, a Hospital Anxiety and Depression Scale (HADS)
110 was used. It is a tool developed by Zigmond and Snaith to be practiced for general
111 population. Unlike other depression screening tools, this questionnaire is easy to deal
112 separately any comorbidity or impairment secondary to another disease (28). This assessment
113 tool has seven items for anxiety and seven items for depression and it takes mostly to finish
114 around 2 to 5 minutes. As recommended, both anxiety and depression separately were scored.
115 For the scales of anxiety and depression; scores which are less than 7 shows that the
116 participants are not depressed, for the score 8-10 are mild cases and 11-14 is reported as
117 moderate and 15-21 scores shows severe cases. This questionnaire is validated and published
118 in different languages including Ethiopian local language.

119 **Data analysis**

120 The gathered data was analysed using IBM SPSS statistics version 22. A univariate analysis
121 was conducted using chi-square statistics to identify the predicting factors associated with the
122 outcome variable, acceptance of anxiety and depression screening. Multivariate logistic
123 regression was performed to identify independent variables associated with the outcome
124 variable. Variables with p-value 0.005 or less were included in the model. Odds Ratio (OR)
125 and Adjusted Odds Ratio (AOR) with their Confidence Interval (95%CI) were calculated to
126 determine the strength of the association.

127 **Ethical issues**

128 Ethical approval of the study was obtained from International Islamic University of Malaysia
129 Research Ethics Committee (IREC). Permission letter was also gained from Jijiga University
130 board of review. A written informed consent was attained from each study participant after

clarifying the purpose of the study & all other relevant information about the study. Privacy and confidentiality of women participants was strictly maintained.

Results

Characteristics of the respondents

A total of 409 women participated in this study. The mean age of the study participants was 34.5 years with standard deviation of 8.75 years. Majority of the respondents were urban dwellers 377 (92.2%) and Amhara by ethnicity 144 (35.2%). Among the total participants, 129 (31.5%) were married, 195 (47.7%) were followers of Orthodox Christian. Regarding the educational background, most of them 190 (46.5%) did not attend any formal education, almost half of the respondents 207 (50.6%) were unemployed, and almost half of them 206 (50.4%) reported their family income was less than 1400 Ethiopian birr which is equal to 37.5 US dollars. Among the interviewed participants, almost half of them were nulliparous women 208 (50.9%) as shown in (Table 1).

Table 1 Socio-demographic characteristics of participants (n= 409).

Variable	Categories	Frequency	Percent (%)
Residence	Urban	377	92.2
	Rural	32	7.8
Age	15-24	57	13.9
	25-49	339	82.9
	>49	13	3.2
Marital Status	Married	129	31.5
	Single	120	29.3
	Widowed	82	20.0
	Divorced	78	19.1
No of living children	Nulliparous/No children	208	50.9
	1-3	160	39.1
	4-6	41	10.0
Education	No formal education	190	46.5
	Primary education	97	23.7
	Secondary level	83	20.3
	College/University	39	9.5
Ethnicity	Amhara	144	35.2
	Somali	137	33.5
	Oromo	86	21.0
	Southern Nations	42	10.3
Religion	Orthodox	195	47.7
	Islam	165	40.3
	Catholic	12	2.9
	Protestant	37	9
Occupation	Un employed	206	50.4
	Self employed	117	28.6
	Student	44	10.8

	Govt employee	42	10.3
Income in ETB	<1400	206	50.4
	1400-3800	129	31.5
	>3800	74	18.1

Acceptance of anxiety and depression screening

Among the 409 interviewed participants, only 120 (29.3%) were aware about the existence of anxiety and depression screening. Majority of them 50 (41.7%) mentioned that the HCP have introduced them screening. The other sources of information included health education sessions 32 (26.7%), followed by information obtained from other peer members 38 (31.7%). When asked about their experience of previous screening, most of the women 366 (89.5%) claimed that they had never been tested previously. However, majority of the participants, 357 (87.3%) accepted to receive the screening. The remaining respondents presented various reasons for their withdrawal of the screening which included requirement of partner approval (40.4%), longer waiting time (38.5%) and fear of stigma (21.2%).

Factors associated with the acceptance of anxiety and depression screening

To identify the predicting factors of acceptance of depression and anxiety screening, we analysed the association between socio-demographic characteristics, awareness of the existing screening, history of previous screening and acceptance of women participants.

Majority of the participants who accepted the screening were holding college level education (OR: 0.32; 95 % C.I: 0.09-1.11), divorced (OR: 1.33; 95 % C.I: 0.65-2.73), were self employed by occupation (OR: 0.93; 95 % C.I: 0.48-1.79), aware of the existing screening service (OR: 0.22; 95 % C.I: 0.09-0.58), and were offered the screening by health care providers (OR: 1.04; 95 % C.I: 0.81-2.02). Moreover, a larger proportion of the participants who accepted the screening had a history of previous screening of depression and anxiety (OR: 2.34; 95 % C.I: 1.08-5.09) as shown in (Table: 2).

To adjust for potential cofounders, a multivariable logistic regression was used. Participants who had a college level of education were more likely to accept the screening offer as indicated in (Table: 2). Being aware about the screening service was also associated with accepting the screening and respondents that were invited to the screening by HCP were two times more likely to accept the test than those who heard from other source of information. In addition to this, women who had a previous history of screening practice were three times more likely to accept the screening.

186

187

188 Table 2: Association between socio-demographic and clinical characteristics with acceptance of
189 depression and anxiety screening among participants.

190

	Anxiety & Depression screening		OR [95% CI]	AOR (95% CI)
	Accepted	Not accepted		
Marital Status				
Married	108	21	1	1
Single	117	3	0.13 [0.04-0.46]	0.09 [0.02-0.38]
Widowed	70	12	0.88 [0.41-1.91]	0.71 [0.29-1.73]
Divorced	62	16	1.33 [0.65-2.73]	1.09 [0.48-2.47]
Education				
No formal education	152	38	1	1
Primary	89	7	0.31 [0.14-0.73]	0.27 [0.11-0.71]
Secondary	79	4	0.20 [0.07-0.59]	0.29 [0.90-0.96]
College/University	37	3	0.32 [0.09-1.11]	0.35 [0.06-1.93]
Occupation				
Un employed	176	30	1	1
Student	43	1	0.14 [0.02-1.03]	1.45 [0.13-6.93]
Self employed	101	16	0.93 [0.48-1.79]	1.46 [0.64-3.34]
Govt employee	37	5	0.79 [0.29-2.18]	1.12 [0.31-4.08]
Aware of depression and anxiety screening (n= 409)				
No	242	47	1	1
Yes	115	5	0.22 [0.09 -0.58]	0.10 [0.01-0.66]
Source of information (n=115)				
Health Care Provider	31	11	1.04 [0.81-2.02]	2.62 [0.26-6.03]
Health education session	27	10	0.37 [0.03-4.21]	0.20 [0.01-0.66]
Heard from peer group	23	13	1	1
Ever screened (n=409)				
No	324	42	1	1
Yes	33	10	2.34 [1.08- 5.09]	3.25 [3.98-8.48]

191

192

193

194 **Discussion and conclusion**

195

196 In this study we examined the acceptance of anxiety and depression screening among women
197 living with HIV who attend the antiretroviral treatment care. The findings of this study has
198 shown that the acceptance rate for screening of anxiety and depression among the study
199 participants was 87.3% and this indicates relatively lower proportion when compared to a
200 study conducted on depression and anxiety among pregnant women in Kilimanjaro, Tanzania
201 in which 96.2% of the respondents accepted the screening. (11) and another similar study on
202 prevalence of depressive symptoms on women in Sabah, Malaysia were 92.2% in which
203 women accepted to undergo screening for the study (29).

204 Probably, the reason for the variation of the findings might be the difference in the study
205 population, since the later studied on pregnant women who possibly had more frequent visit
206 and follow up in their health facilities. One unanticipated finding shows regardless of the
207 acceptability level of the participants, the awareness of the respondents about the availability
208 of anxiety and depression screening was low (28.1%), when compared to other studies
209 conducted earlier in different settings. The reason could be explained non-existence of proper
210 mental health education sessions in the facility where the patients were expected to receive
211 information.

212 The results of this study shows that the majority of the participants who accepted the
213 screening were holding college level education. Hereafter, education denote the formal
214 education obtained in academics and not merely the health related knowledge of the women
215 participants. This finding is consistent with other studies which presented that positive
216 association between higher educational level and women's attitude towards anxiety and
217 depression interventions (30,27). It is common in Ethiopia, that low service utilization is
218 typically associated with the poor literacy among women. In line with this, it is found that
219 HIV patients with no formal education was more likely to develop distress (31). As a result,
220 education and awareness pay a major role in reducing the unmet needs in health service
221 utilization including depression screening.

222 The respondents who were divorced, and those who were self employed by occupation were
223 more likely to accept the screening. Contrary to expectations, and not common in low income
224 countries where women empowerment is low, this finding may determine the autonomy of
225 women for taking decisions in regard to health seeking when they are economically
226 independent (32). However, among the married women, their claim of partner approval was
227 the main reason of opting out the screening offer. Thus, in marital union, male involvement in
228 decision making for the screening uptake was acknowledged.

229 Most of the study participants has mentioned health care provider's as their main source of
230 information to take part the screening. Although, this study did not initially identify the
231 reasons for the high acceptance rate, the health provider's initiated counselling could be the
232 motive for the high acceptance rate. One of the HCP's drive could be ascribed to their
233 intention to practice the protocol of Ethiopia's national guideline for comprehensive HIV
234 prevention, care and treatment which suggests the screen for depression among the HIV
235 patients in regard to. (33). Despite, the HCPs recognize that the screening was prominently
236 required service, however it was not practiced in the studied health set up. This is common in
237 major health facilities where depression is often unnoticed and not practiced (34). This could

be attributed to the reality on the ground which is related with the practicality, availability of validated tools, less number of trained professionals, and overall the guiding principle of the hospital implementation plan. Depending on the regional difference, these limited service provision was recognized to increase a large proportion of patients, remain un-detected in the ART.

Moreover, a larger proportion of the participants who accepted the screening had a history of previous screening of depression and anxiety. This similar to other studies which concluded that women who had a previous engagement in screening were more receptive and willing to accept the screening (35–37). Yet, the type of screening tool and their decision process determines their choice to accept the screening. As several studies has published that women understandings and decision manners commonly regulates their response and acceptance of the screening (38–40). Thus, in order to reduce the frequent challenges of integrating mental health with the comprehensive HIV care and minimize the unmet need of HIV women to be screened for depression, recent findings suggest that ART attendees should be invited for depression screening and eligible clients should be linked for further examination and treatment (41).

Conclusion

As the screening of symptomatic depression in HIV patients has recorded an apparent health implication for the adherence of treatment and care, health care providers should consider a routine check-up for the HIV patient's neurocognitive functioning and wellbeing. This research has shown that women participant living with HIV were willing to undergo the screening for depression and anxiety. Therefore, all concerned bodies like health organizations should enhance efforts that promote mental health interventions in order to address vulnerable community member like people living with comorbid diseases; HIV and depression.

Abbreviations

AIDS	Acquired Immune Deficiency Syndrome
ART	Antiretroviral Therapy
AOR	Adjusted Odds Ratio
HADS	Hospital Anxiety and Depression Scale
HCP	Health Care Provider
HIV	Human Immunodeficiency Virus
OR	Odds Ratio

Conflict of interest

There is no conflict of interest among authors in this article.

Ethics approval and consent to participate

All the study procedures were according the ethical standard of the institution's human ethical procedure. The ethical approval of this study was obtained from the IIUM Research Ethics Committee (IREC) (Reference: IIUM/305/14/11/2/IREC/2018-042. Dated: 04/01/2019). A permission letter was also gained from the Jijiga University board of review.

The women participants in this study were provided a written document approved by IREC and informed consent then confirmed by their signatures for the use and dissemination of the study findings.

Consent for publication

Not applicable

Data availability

The qualitative data included in this study is available from the authors upon request.

Funding

This study has not received any funds or grants from any institution or public and private sectors.

Authors contribution

AY and SR: conception and design, data collection, statistical analysis, interpretation of data and drafting the manuscript. RM and ML: critical revision of the manuscript. All authors have contributed and approved this manuscript.

Acknowledgements

The authors would like to acknowledge the WLHIV who participated this study and the medical staffs working in the ART for their assistance through the data collection process.

References

1. World Health Organization. Depression and other common mental disorders. Global health estimates. 2017. 8-12 p.
2. Rukundo GZ, Mishara BL, Kinyanda E. Burden of Suicidal Ideation and Attempt among Persons Living with HIV and AIDS in Semiurban Uganda. AIDS Res Treat. 2016;2016.
3. Zarei N, Joulaei H. The Impact of Perceived Stigma , Quality of Life , and Spiritual Beliefs on Suicidal Ideations among HIV-Positive Patients. AIDS Res Treat. 2018;2018.
4. Ajinkya S, Jadhav P, Srivastava N. Depression during pregnancy: Prevalence and obstetric risk factors among pregnant women attending a tertiary care hospital in Navi Mumbai. Ind Psychiatry J [Internet]. 2013;22(1):37. Available from: <http://www.industrialpsychiatry.org/text.asp?2013/22/1/37/123615>
5. Mohammed M, Mengistie B, Dessie Y, Godana W. Prevalence of Depression and Associated Factors among HIV Patients Seeking Treatments in ART Clinics at Harar Town , Eastern Ethiopia. J AIDS Clin Res. 2015;6(6):1–6.
6. Tebikew Yeneabat AB, Amare T. Factors associated with depressive symptoms in people living with HIV attending antiretroviral clinic at Fitcha Zonal Hospital , Central Ethiopia: cross-sectional study conducted in 2012. Dovepress. 2017;2125–31.
7. Eshetu DA, Meseret SW, Kebede MA, Techane GN, Gizachew KD, Tegegne MT, et al. Prevalence of Depression and Associated Factors among HIV / AIDS Patients

306 Attending ART Clinic at Debrebirhan Referral Hospital , North Showa , Amhara
307 Region , Ethiopia. Clin Psychiatry. 2015;1(1):3.

308 8. Tesfaw G, Ayano G, Awoke T, Assefa D, Birhanu Z, Miheretie G. Prevalence and
309 correlates of depression and anxiety among patients with HIV on- follow up at Alert
310 Hospital , Addis Ababa ,. BMC Psychiatry. 2016;1–8.

311 9. Duko B, Geja E, Zewude M, Mekonen S. Prevalence and associated factors of
312 depression among patients with HIV / AIDS in Hawassa , Ethiopia , cross sectional
313 study. Ann Gen Psychiatry. 2018;1–7.

314 10. Abebe H, Shumet S, Nassir Z, Agidew M, Abebaw D. Prevalence of Depressive
315 Symptoms and Associated Factors among HIV-Positive Youth Attending ART
316 Follow-Up in Addis Ababa , Ethiopia. AIDS Res Treat. 2019;

317 11. Samwel J, Id N, Watt MH, Minja L, Id BAK, Id BTM, et al. Depression and anxiety
318 among pregnant women living with HIV in Kilimanjaro region , Tanzania. PLoS One
319 [Internet]. 2019;1–16. Available from: <http://dx.doi.org/10.1371/journal.pone.0224515>

320 12. S. M. Meffert, T. C. Neylan, C. E. McCulloch, L. Maganga, Y. Adamu, F. Kiweewa, J.
321 Maswai, J. Owuoth, C. S. Polyak JAA and VGV. East African HIV care: depression
322 and HIV outcomes. Glob Ment Heal. 2019;(April).

323 13. Onu C, Ongeru L, Bukusi E, Cohen CR, Neylan TC, Oyaro P, et al. Interpersonal
324 psychotherapy for depression and posttraumatic stress disorder among HIV-positive
325 women in Kisumu , Kenya: study protocol for a randomized controlled trial. BMC
326 Trials [Internet]. 2016;1–8. Available from: [http://dx.doi.org/10.1186/s13063-016-](http://dx.doi.org/10.1186/s13063-016-1187-6)
327 1187-6

328 14. Gebrezgiabher BB, Abraha TH, Hailu E, Siyum H, Mebrahtu G, Gidey B, et al.
329 Depression among Adult HIV / AIDS Patients Attending ART Clinics at Aksum Town
330 , Aksum , Ethiopia: A Cross-Sectional Study. Depress Res Treat. 2019;

331 15. Prasithsirikul W, Chongthawonsatid S, Ohata PJ, Keadpudsa S, Klinbuayaem V,
332 Rerksirikul P, et al. Depression and anxiety were low amongst virally suppressed ,
333 long-term treated HIV- infected individuals enrolled in a public sector antiretroviral
334 program in Thailand. AIDS Care. 2016;(October).

335 16. Zhu Q, Huang D, Lv J, Guan P. Prevalence of perinatal depression among HIV-
336 positive women: a systematic review and meta-analysis. BMC Psychiatry.
337 2019;19(330):1–12.

338 17. Nyirenda M, Chatterji S, Rochat T, Mutevedzi P, Newell M. Prevalence and correlates
339 of depression among HIV-infected and -affected older people in rural South Africa. J
340 Affect Disord [Internet]. 2013;151(1):31–8. Available from:
341 <http://dx.doi.org/10.1016/j.jad.2013.05.005>

342 18. Logie C, James L, Tharao W, Loutfy M. Associations between HIV-related stigma,
343 racial discrimination, gender discrimination, and depression among HIV-positive
344 African, Caribbean, and Black women in Ontario, Canada. AIDS Patient Care STDS
345 [Internet]. 2013;27(2):114–22. Available from:
346 <http://www.ncbi.nlm.nih.gov/pubmed/23373665>

347 19. WHO. HIV / AIDS and mental health Report by the Secretariat. Geneva; 2008.

348 20. Reda AA, Biadgilign S. Determinants of Adherence to Antiretroviral Therapy among

- 349 HIV-Infected Patients in Africa. *AIDS Res Treat.* 2012;2012.
- 350 21. Bouhnik AD, Preau M, Vincent E, Carrieri MP, Gallais H, Lepeu G, et al. Depression
351 and clinical progression in HIV-infected drug users treated with highly active
352 antiretroviral therapy. *Antivir Ther.* 2005;10(1):53–61.
- 353 22. Amare T, Getinet W, Shumet S, Asrat B. Prevalence and Associated Factors of
354 Depression among PLHIV in Ethiopia□: Systematic Review and Meta-Analysis ,
355 2017. *AIDS Res Treat.* 2018;
- 356 23. Berhe H, Sciences H. PREVALENCE OF DEPRESSION AND ASSOCIATED
357 FACTORS AMONG PEOPLE LIVING WITH HIV/AIDS IN TIGRAY, NORTH
358 ETHIOPIA: A CROSS SECTIONAL HOSPITAL BASED STUDY. *Int J Pharm Sci*
359 *Res.* 2013;4(2):765–75.
- 360 24. Amare T, Getinet W, Shumet S, Asrat B. Prevalence and Associated Factors of
361 Depression among PLHIV in Ethiopia□: Systematic Review and Meta-Analysis ,
362 2017. *AIDS Res Treat.* 2018;2018.
- 363 25. Organizations WH. WHO recommendations on adolescent sexual and reproductive
364 health and rights. World Health Organization; 2018.
- 365 26. Sikkema KJ, Dennis AC, Watt MH, Choi KW, Yemeke TT, Joska JA. Improving
366 mental health among people living with HIV: a review of intervention trials in low-
367 and middle-income countries. *Glob Ment Heal [Internet].* 2015;2:e19. Available from:
368 http://www.journals.cambridge.org/abstract_S2054425115000175
- 369 27. Sanchez K, Eghaneyan BH, Trivedi MH. Depression Screening and Education□:
370 Options to Reduce Barriers to Treatment (DESEO): protocol for an educational
371 intervention study. *BMC Health Serv Res.* 2016;16(322):1–10.
- 372 28. Stern AF. The Hospital Anxiety and Depression Scale. *Occup Med (Chic Ill).*
373 2014;64(5):393–4.
- 374 29. Mohamad Yusuff AS, Tang L, Binns CW, Lee AH. Prevalence of antenatal depressive
375 symptoms among women in Sabah, Malaysia. *J Matern Neonatal Med [Internet].*
376 2016;29(7):1170–4. Available from:
377 <http://www.tandfonline.com/doi/full/10.3109/14767058.2015.1039506>
- 378 30. Hossein Shahnazi, Sahar Sabooteh AH. The impact of education intervention on health
379 belief model constructs regarding anxiety of nulliparous pregnant women. *J Educ*
380 *Health Promot.* 2015;
- 381 31. Basha EA, Derseh BT, Gebre Y, Haile E, Tafere G. Factors Affecting Psychological
382 Distress among People Living with HIV / AIDS at Selected Hospitals of North Shewa
383 Zone , Amhara Region , Ethiopia. *AIDS Res Treat.* 2019;2019.
- 384 32. Haque A, Dash SK, Abdul M, Chowdhury B. Maternal health care seeking behavior□:
385 the case of Haor (wetland) in Bangladesh. *BMC Public Health.* 2016;1–10.
- 386 33. Federal Ministry Of Health. NATIONAL GUIDELINES FOR COMPREHENSIVE
387 HIV PREVENTION , CARE AND TREATMENT. Addis Ababa: FMOH; 2017. 78-
388 79 p.
- 389 34. Bess KD, Adams J, Watt MH, O'Donnell JK, Gaynes BN, Thielman NM, et al.
390 Providers' attitudes towards treating depression and self-reported depression treatment

practices in HIV outpatient care. *AIDS Patient Care STDS* [Internet]. 2013;27(3):171–80. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3595951&tool=pmcentrez&rendertype=abstract>

35. Horowitz JA, Murphy CA, Gregory KE, Wojcik J. A Community-Based Screening Initiative to Identify Mothers at Risk for Postpartum Depression. *J Obstet Gynecol Neonatal Nurs*. 2010;52–61.

36. Venkatesh KK, Nadel H, Blewett D. Implementation of universal screening for depression during pregnancy: Feasibility and impact on obstetric care. *Am J Obstet Gynecol* [Internet]. 2016; Available from: <http://dx.doi.org/10.1016/j.ajog.2016.05.024>

37. Waerden JEB Van Der, Hoefnagels C, Jansen MWJ, Hosman CMH. Exploring recruitment, willingness to participate, and retention of low-SES women in stress and depression prevention. *BMC Public Health*. 2010;

38. Robinson J, Khan N, Fusco L, Malpass A, Lewis G DC. Why are there discrepancies between depressed patients' global rating of change and scores on the patient health questionnaire depression module? A qualitative study of primary care in England. *BMJ Open*. 2017;7(4).

39. Malpass A, Dowrick C, Gilbody S, Robinson J, Wiles N, Duffy L et al. Usefulness of PHQ-9 in primary care to determine meaningful symptoms of low mood: a qualitative study. *Br J Gen Pr*. 2016;643(66):78–84.

40. Velloza J, Njoroge J, Ngure K, Thuo N, Kiptinness C, Momanyi R, et al. Cognitive testing of the PHQ-9 for depression screening among pregnant and postpartum women in Kenya. *BMC Psychiatry*. 2020;20(31):1–14.

41. Ng'oma M, Meltzer-Brody S, Chirwa E SR. “ Passing through difficult times ”: Perceptions of perinatal depression and treatment needs in Malawi - A qualitative study to inform the development of a culturally sensitive intervention. *PLoS One*. 2019;1–20.