Vitamin D insufficiency and deficiency and mortality from respiratory diseases in a cohort of older adults: potential for limiting the death toll during and beyond the COVID-19 pandemic

Hermann Brenner, MD^{1,2,3,4} Bernd Holleczek, PhD^{1,5} Ben Schöttker, PhD^{1,4}

¹Division of Clinical Epidemiology and Aging Research, Germany Cancer Research Center (DKFZ), Heidelberg, Germany

²Division of Preventive Oncology, Germany Cancer Research Center (DKFZ) and National Center for Tumor Diseases (NCT), Heidelberg, Germany

³German Cancer Consortium (DKTK), Germany Cancer Research Center (DKFZ), Heidelberg, Germany

⁴Network Aging Research, University of Heidelberg, Heidelberg, Germany

⁵Saarland Cancer Registry, Saarbrücken, Germany

Corresponding Author

Ben Schöttker, PhD, MPH, MSc

Division of Clinical Epidemiology and Ageing Research

German Cancer Research Center

Im Neuenheimer Feld 581

69120 Heidelberg

Germany

Phone: +49-6221-421351

Fax : +49-6221-421302

Email: b.schoettker@dkfz.de

Abstract

Background. The COVID-19 pandemic goes along with increased mortality from acute respiratory disease, and measures to limit the spread of the infection go along with increased risk of vitamin D deficiency, especially among high risk groups. It has been suggested that vitamin D₃ supplementation might help to reduce respiratory disease mortality.

Methods. We assessed the prevalence of vitamin D insufficiency and deficiency, defined by 25(OH)D blood levels of 30-50 and <30 nmol/L, respectively, and their association with mortality from respiratory diseases during 15 years of follow-up in a cohort of 9,548 adults aged 50-75 years from Saarland, Germany.

Results. Vitamin D insufficiency and deficiency were common (44% and 15%, respectively). Compared to sufficient vitamin D status, respiratory disease mortality was 2.1 (95%-CI 1.3-3.2)- and 3.0 (95%-CI 1.8-5.2)-fold increased, respectively. Although significant increases were seen in both women and men, they were much stronger among women, with 8.5 (95% CI 2.4-30.1) and 2.3 (95% CI 1.1-4.4)-fold increase of respiratory disease mortality in case of vitamin D deficiency among women and men, respectively (p-value for interaction =0.041). Overall, 41% (95% CI 20%-58%) of respiratory disease mortality was statistically attributable to vitamin D insufficiency or deficiency.

Conclusion. Vitamin D insufficiency and deficiency are common and account for a large proportion of respiratory disease mortality in older adults, supporting suggestions that vitamin D₃ supplementation might make a major contribution to limit the burden of the COVID-19 pandemic, particularly among women.

Introduction

The Corona Virus Disease 2019 (COVID-19) pandemic goes along with strongly increased respiratory disease mortality. It has been suggested that vitamin D₃ supplementation could be a potentially promising and safe approach to reduce risk of COVID-19 infections and deaths (1). Meta-analyses of randomized clinical trials (RCTs) have shown that vitamin D₃ supplementation reduces the risk of acute respiratory tract infections (2). Risk reduction with regular (daily or weekly) supplementation of physiological doses of vitamin D was especially strong (by 70%) among people with vitamin D deficiency, but significant risk reduction (by 25%) was also found among people with higher vitamin D levels.

People with pre-existing major diseases, such as diabetes or cancer are at increased risk of dying from severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infections. At the same time, prevention of and care for these diseases have been and keep being strongly compromised by current measures to limit the COVID-19 pandemic. Meta-analyses of clinical trials have demonstrated that vitamin D₃ supplementation has the potential to also reduce cancer mortality by approximately 13% (3).

We previously assessed the prevalence of vitamin D insufficiency and deficiency and their association with all-cause mortality and mortality from cardiovascular, cancer and respiratory diseases in a cohort of 9,548 adults aged 50-75 years from Saarland, Germany (4-8). We aim to present considerably updated and sex-specific follow-up data of 15 years here and to calculate the proportion of respiratory disease mortality that is attributable to vitamin D insufficiency and deficiency. Furthermore, we discuss potential implications for prevention in the context of the ongoing COVID-19 pandemic.

Methods

Study design

This investigation is based on the ESTHER study, an ongoing statewide cohort study from Saarland, Germany, details of which have been reported elsewhere (4-9). Briefly, 9,940 men and women, aged 50-75 years at baseline, were recruited by their general practitioners during a routine health check-up between 2000 and 2002. Blood samples were taken at baseline at the general practitioners' offices. Information on socio-demographic and lifestyle characteristics and medical history were obtained by questionnaires from participants and their general practitioners, and the distribution of those characteristics was similar to the distribution in the respective age categories in the German National Health Survey conducted in a representative sample of the German population in 1998 (9). The ESTHER study was approved by the ethics committees of the University of Heidelberg and the state medical board of Saarland, Germany. Written informed consent was issued by all participants.

Variable assessment

Information on socio-demographic characteristics, lifestyle and diet were obtained by a comprehensive self-administered questionnaire from the study participants at baseline. Height and weight were assessed and documented on a standardized form by the general practitioners during the health check-up. Furthermore, blood and urine samples were taken during the health check-up, centrifuged, sent to the study center and stored at -80°C until analysis.

The most abundant and stable vitamin D metabolite in blood samples, 25-hydroxyvitamin D (25(OH)D) levels was measured from stored serum samples taken at recruitment. The laboratory methods used are described in detail elsewhere (4). In brief, 25(OH)D levels in women were measured with the Diasorin-Liaison analyzer (Diasorin Inc., Stillwater, USA). Analyses in men were conducted in the context of a separate research project several years

later (when the Diasorin measurements were no longer offered) with IDS-iSYS (Immunodiagnostic Systems GmbH, Frankfurt Main, Germany). Both immunoassays were standardized retrospectively to the gold standard method liquid chromatography tandem-mass-spectrometry.

Deaths until end of 2016 were identified by inquiry at the residents' registration offices and death certificates of deceased study participants were provided by local health authorities. The leading cause of death with an ICD-10 code was available for 98.9% of deceased study participants and was coded with ICD-10 codes R98-R99 "unknown cause of death" for 4.4% of deceased participants. These individuals were not excluded and censored at the time of death for cause specific mortality outcomes.

Statistical methods

Participants of the ESTHER baseline examination (n = 9,940) were excluded from this investigation if no blood sample was available (n = 368) or if they could not be followed-up for mortality (n = 24), which resulted in a total sample size of n = 9,548 subjects for this analysis.

We used the US-American Institute of Medicine's cut-offs to define adequate vitamin D status (> 50 nmol/L), vitamin D insufficiency (30-50 nmol/L) and vitamin D deficiency (< 30 nmol/L) (10). We assessed prevalences of vitamin D insufficiency and deficiency in the total study population and according to age, sex, life-style factors and major diseases and tested for differences between groups by χ^2 -test.

We compared all-cause, cardiovascular disease, cancer, and respiratory disease mortality between subjects with vitamin D insufficiency or deficiency and subjects with adequate 25(OH)D levels and estimated hazard ratios (HR) with 95% confidence intervals (95%CI) by multivariable Cox proportional hazards models. We used an age, sex and season adjusted model and a full model that was adjusted for potential confounders, which are listed in **Table 1**. Additional adjustment for potential intermediates (cardiovascular disease, history of

cancer, diabetes mellitus, hypertension, asthma, total serum cholesterol and serum c-reactive protein levels) did not lead to substantially different results (data not shown). Age and BMI were modelled as continuous variables and all other variables were modelled with the categories shown in **Table 1**. Furthermore, sex-specific analyses were performed and statistical tests on interaction were carried out. In addition, we estimated the population attributable fraction (PAF) of respiratory disease mortality from the prevalences of vitamin D insufficiency and deficiency and their associations with respiratory disease mortality, as derived from our study. The PAF of mortality is the share of mortality in a population that is statistically attributable to a risk factor and that could be avoided by entirely eliminating that risk factor (here: vitamin D insufficiency or deficiency) (11).

Missing values for covariates ranged from 0% to 5.8% (for fish consumption). Missing covariate values were imputed with multiple imputation using the Markov Chain Monte Carlo (MCMC) method with 200 burn-in iterations. Twenty data sets were generated. The imputation model consisted of all variables of the full model (modelled as used in the full model) but not the outcome data and the imputation was carried out stratified by sex.

All statistical tests were two-sided and the alpha level of significance was set to 0.05. All statistical analyses were conducted with the software package SAS, version 9.4 (Cary, North Carolina, USA).

Results

The study population included 43.8% men, mean age was 62.1 years (**Table 1**). Among the 9,548 participants included in the study, 4186 (43.8%) had vitamin D insufficiency (25(OH)D levels of 30 - <50 nmol/L) and 1438 (15.1%) had vitamin D deficiency (25(OH)D levels <30 nmol/L) (**Figure 1A**). **Table 1** also provides a description of further characteristics of the study population at baseline as well as prevalences of vitamin D insufficiency and deficiency according to those characteristics. Both vitamin D insufficiency and deficiency were statistically significantly more frequent among females, with higher age, BMI, in subjects with low physical activity and those who consumed fish less than once per week. A seasonal variation with higher prevalences in winter than in summer months was also observed. Moreover, the prevalence of vitamin D deficiency was significantly higher among current smokers.

Overall, 2,363 (24.7%) study participants died during a median of 15.3 years of follow-up and both vitamin D insufficiency and deficiency were associated with significantly increased all-cause mortality compared to sufficient vitamin D status (full model HRs [95%CI]: 1.2 [1.1-1.3] and 1.7 [1.5-1.9], respectively) (**Table 2**). Vitamin D deficiency was also associated with significant increases in CVD and cancer mortality by 52% and 38%, respectively (full model results). However, vitamin D insufficiency and deficiency were particularly strongly associated with respiratory disease mortality with full model HRs of 2.1 (95%CI: 1.3-3.2) and 3.0 (95%CI: 1.8-5.2), respectively. Overall, 41% (95%CI: 20%-58%) of all deaths from respiratory diseases were statistically attributable to 25(OH)D levels < 50 nmol/L (**Figure 1B**).

Table 3 shows the results of the sex-specific analyses. For all-cause, cardiovascular disease and cancer mortality, only modest, statistically non-significant differences were seen between women and men. Although significant increases were seen for respiratory disease mortality in both women and men, they were much stronger among women, with 8.5 (95% CI 2.4-

149 30.1) and 2.3 (95% CI 1.1-4.4)-fold increase of respiratory disease mortality in case of
150 vitamin D deficiency among women and men, respectively (p-value for interaction =0.024).
151

Discussion

In this large population-based cohort study from Saarland, Germany, the majority of participants aged 50-75 years at baseline had vitamin D insufficiency or deficiency, and these conditions were associated with increased mortality. In particular, mortality from respiratory diseases was 2.1- and 3.0-fold increased in subjects with vitamin D insufficiency or deficiency, respectively, compared to participants with sufficient vitamin D status. Significant associations with respiratory disease mortality were seen among both women and men, but they were particularly strong for women. Overall, 41% of deaths from respiratory diseases were statistically attributable to vitamin D insufficiency or deficiency, and could possibly be avoided by overcoming these conditions assuming causality of the association.

The assumption of causality of vitamin D₃ effects on mortality obviously requires most careful discussion. Although we made the best attempts to adjust potential confounding factors in our analyses, the major limitation of our study is that we cannot exclude the possibility of residual confounding by imperfect measurement of confounding variables, such as smoking or physical activity, or omission of unknown confounders in this observational study. As addressed in detail elsewhere (6), interpretation of the evidence is further complicated by the fact that vitamin D deficiency could be considered both a consequence of poor health as well as a risk factor for increased vulnerability to acute disease and poor outcomes of chronic diseases among people with poor health. It is therefore paramount to critically evaluate our findings in the light of additional criteria and evidence, such as biological mechanisms and plausibility, and, in particular, in the light of data from RCTs providing vitamin D₃ supplementation.

Deaths from respiratory disease are mostly deaths from exacerbations of COPD which are caused by acute infections in the majority of cases or from acute respiratory infections causing pneumonia. Vitamin D₃ is thought to protect from occurrence and poor outcomes of respiratory infections by several mechanisms, including enhanced physical barriers (maintenance of junction integrity), cellular innate immunity, and adaptive immunity (1).

179 Innate and adaptive immunity are being influenced by sex hormones (12), which may explain
180 the observed interaction of sex and 25(OH)D levels with respiratory disease mortality.
181 According to data from the US-American National Health and Nutrition Survey women have a
182 higher inflammation burden than men (the age-range was 40-90+ years (13)). Especially
183 postmenopausal women, like those included in the ESTHER study, have a high inflammatory
184 burden because a decline in estrogen level during menopause is associated with an
185 increased expression of pro-inflammatory cytokines, incl. interleukin 6 and tumor necrosis
186 factor (TNF) α (12,14,15). A cytokine storm as an adverse immune response to a SARS-
187 CoV-2 infection is currently a major hypothesis for the underlying cause of a large proportion
188 of COVID-19 deaths (16). Sufficient 25(OH)D levels are suggested to contribute to
189 prevention of the cytokine storm (17, 18). This may be especially important in
190 postmenopausal women who tend to have both lower 25(OH)D levels and a higher
191 inflammatory burden than men.

192 Although specific data for the role of vitamin D₃ in protecting from SARS-CoV-2 infections
193 and their consequences are yet to be published it appears plausible to assume that these
194 mechanisms would be relevant for this infection in a similar manner as for other severe viral
195 respiratory diseases, such as influenza. In a meta-analysis of individual participant data of 25
196 RCTs that included 11,321 participants, aged 0–95 years, vitamin D₃ supplementation was
197 shown to reduce the risk of acute respiratory tract infection (OR 0.88, 95% CI 0.81–0.96) (2).
198 The best effects were shown for daily or weekly vitamin D₃ supplementation without
199 additional bolus doses (OR 0.81, 95% CI 0.72 to 0.91). The protective effects were
200 particularly strong in those with baseline 25-hydroxyvitamin D levels <25 nmol/L (OR 0.30,
201 95% CI 0.17– 0.53), suggesting an approximately 3-fold risk among people with vitamin D
202 deficiency not receiving vitamin D₃ supplementation compared to people with vitamin D
203 deficiency receiving supplementation. In a recent meta-analysis of RCTs on vitamin D₃
204 supplementation for patients with COPD, the risk of acute exacerbations was estimated to be
205 reduced by 61% (95% CI 36%-77%) (19). In remarkable consistency with our results, these

206 meta-analyses of RCTs provide strong evidence for the preventive potential of vitamin D₃
207 supplementation against acute respiratory infections and COPD exacerbations in particular.

208 A first study posted on a pre-print server on May 13, 2020 suggests that the protective
209 effects of vitamin D₃ on other acute respiratory tract infections may be translated to COVID-
210 19 infections (20). Vitamin D deficiency and vitamin D₃ treatment data were available for 499
211 COVID-19 patients from Chicago for the year prior COVID-19 testing. Being likely vitamin D
212 deficient (defined as being vitamin D deficient at last available time point without increase of
213 vitamin D treatment) at the time of COVID-19 testing was associated with a 1.8-fold
214 increased risk of being tested positive for COVID-19 ($p < 0.02$) as compared to likely vitamin D
215 sufficient.

216 It is worth noting that beneficial effects of vitamin D₃ supplementation against manifestation
217 or exacerbation of acute respiratory infection during an epidemic would be expected to go
218 beyond individual protection of those using supplementation, as limiting such manifestation
219 and exacerbation would also be expected to reduce the potential of spread of the disease to
220 other persons and relieve the overload of the medical system by the epidemic.

221 To our knowledge, no previous vitamin D₃ supplementation RCTs have addressed mortality
222 from respiratory disease as primary endpoint, and no meta-analysis of results for this specific
223 endpoint have been reported, which most likely reflects the relatively small share of deaths
224 from this endpoint among all deaths. In our cohort of older adults these deaths accounted for
225 5.2% of all deaths. Even though this proportion is expected to be higher during the COVID-
226 19 pandemic, the majority of deaths still occur from other diseases, and the summary effect,
227 benefit-harm ratio and cost-effectiveness with respect to all relevant outcomes therefore
228 deserve most careful attention for any general prevention efforts.

229 In that respect, vitamin D₃ supplementation appears to be a particularly promising approach,
230 especially for population groups with high prevalence of vitamin D insufficiency or deficiency,
231 such as the elderly and those with severe comorbidities (which essentially coincide with
232 population groups at highest risk of severe course and death from SARS-CoV-2 infection

(21)): The personal, health care and societal costs of the vitamin D₃ intervention are negligibly low compared to the very high costs of currently employed “general population measures”, such as extensive testing for the infection and the lockdown of large proportions of economic and social life, including the delay or omission of much of routine medical care for other relevant diseases. In fact, some of these measures are expected to severely aggravate vitamin D insufficiency or deficiency, especially in high risk groups, such as restrictions of spending time outdoors for the total population (as practiced, for example, in France and Spain) or certain high risk groups, such as nursing home residents (as practiced in many countries including Germany). Such restrictions dramatically reduce opportunities to maintain adequate vitamin D levels through endogenous synthesis by relevant sun exposure.

Vitamin D₃ supplementation has been demonstrated to be safe in numerous large scale studies, and the risk of harm seems to be negligible compared to the risk of harmful side effects of the aforementioned and other general population measures, such as delayed diagnosis and treatment of cancer, myocardial infarction or stroke, withheld or deferred delivery of surgical or other medical services, or health risks related to unemployment and loneliness (22-25). On the contrary, one expected “side effect” would be reducing total cancer mortality by 13%, as suggested by a recent meta-analysis of RCTs (3). For Germany, with currently approximately 230,000 deaths from cancer per year (26), this would translate in prevention of approximately 30,000 cancer deaths each year, suggesting substantial additional benefit besides lowering the COVID-19 burden during the COVID-19 pandemic and beyond.

In conclusion, our results, along with evidence from meta-analyses from RCTs regarding results of vitamin D₃ supplementation on various outcomes, suggest that vitamin D₃ supplementation could make a major contribution to lowering mortality from respiratory and other diseases during and beyond the COVID-19 pandemic, in particular among women. The Endocrine Society recommends 1,500-2,000 IU vitamin D₃/day for adults of any age at high risk for vitamin D deficiency (27). The costs for such supplementation are in the order of 30 €

per person per year, or even half that amount when sufficient vitamin D supply is ensured by carefully dosed sun exposure during the summer months. Along with expected savings from prevented respiratory and other diseases, this would make vitamin D₃ supplementation a particularly cost-effective and most likely cost-saving measure, whose currently still widely neglected potential should receive increased attention and should more widely be utilized in the fight against the COVID-19 pandemic and beyond.

Acknowledgements

The ESTHER study was funded by grants from the Saarland state Ministry for Social Affairs, Health, Women and Family Affairs (Saarbrücken, Germany), the Baden-Württemberg state Ministry of Science, Research and Arts (Stuttgart, Germany), the Federal Ministry of Education and Research (Berlin, Germany) and the Federal Ministry of Family Affairs, Senior Citizens, Women and Youth (Berlin, Germany). The authors have no conflicts of interest to disclose. The author contributions were as follows: HB designed the research and drafted the manuscript, BS conducted statistical analyses, and BH critically read and commented the manuscript and added aspects to the discussion. All authors contributed to the data collection for this project.

References

1. Grant WB, Lahore H, McDonnell SL, Baggerly CA, French CB, Aliano JL, et al. Evidence that vitamin D supplementation could reduce risk of influenza and COVID-19 infections and deaths. *Nutrients* 2020;12(4).
2. Martineau AR, Jolliffe DA, Hooper RL, Greenberg L, Aloia JF, Bergman P, et al. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *BMJ* 2017;356:i6583
3. Keum N, Lee DH, Greenwood DC, Manson JE, Giovannucci E. Vitamin D Supplements and Total Cancer Incidence and Mortality: a Meta-analysis of randomized controlled trials. *Ann Oncol* 2019;30(5):733-43.
4. Schöttker B, Haug U, Schomburg L, Köhrle J, Perna L, Müller H, et al. Strong associations of 25-hydroxyvitamin D levels with all-cause, cardiovascular, cancer and respiratory disease mortality in a large cohort study. *Am J Clin Nutr* 2013;97:782-93.
5. Schöttker B, Jorde R, Peasey A, Thorand B, Jansen EH, Groot Ld, et al; Consortium on Health and Ageing: Network of Cohorts in Europe and the United States. Vitamin D and mortality: meta-analysis of individual participant data from a large consortium of cohort studies from Europe and the United States. *BMJ* 2014;348:g3656
6. Schöttker B, Brenner H. Vitamin D as a resilience factor, helpful for survival of potentially fatal conditions: a hypothesis emerging from recent findings of the ESTHER cohort study and the CHANCES consortium. *Nutrients* 2015;7(5):3264-78.
7. Brenner H, Jansen L, Saum KU, Holleczer B, Schöttker B. Vitamin D supplementation trials aimed at reducing mortality have much higher power when focusing on people with low serum 25-hydroxyvitamin D concentrations. *J Nutr.* 2017;147(7):1325-1333.
8. Schöttker B, Hagen L, Zhang Y, Gao X, Holleczer B, Gao X, et al. Serum 25-hydroxyvitamin D levels as an aging marker: strong associations with age and all-cause mortality independent from telomere length, epigenetic age acceleration, and 8-isoprostane levels. *J Gerontol A Biol Sci Med Sci.* 2019;74(1):121-128.
9. Löw M, Stegmaier C, Ziegler H, Rothenbacher D, Brenner H. [Epidemiological investigations of the chances of preventing, recognizing early and optimally treating chronic diseases in an elderly population (ESTHER study)]. *Dtsch Med Wochenschr* 2004;129:2643-7.
10. Slomski A. IOM endorses vitamin D, calcium only for bone health, dispels deficiency claims. *JAMA* 2011;305:453-4, 456.
11. World Health Organization health statistics and information systems. Metrics: Population attributable fraction (PAF). https://www.who.int/healthinfo/global_burden_disease/metrics_paf/en/. Accessed May 2, 2020.
12. Gubbels Bupp MR, Potluri T, Fink AL, et al. The Confluence of Sex Hormones and Aging on Immunity. *Front Immunol* 2018; 9: 1269.
13. Yang Y, Kozloski M. Sex differences in age trajectories of physiological dysregulation: inflammation, metabolic syndrome, and allostatic load. *J Gerontol A Biol Sci Med Sci* 2011;66:493–500.
14. Straub RH. The complex role of estrogens in inflammation. *Endocr Rev* 2007; 28(5): 521–74
15. Wang Y, Mishra A, Brinton RD. Transitions in metabolic and immune systems from pre-menopause to post-menopause: implications for age-associated neurodegenerative diseases. *F1000Res.* 2020;9:F1000 Faculty Rev-68. Published 2020 Jan 30. doi:10.12688/f1000research.21599.1

- 326 16. Meftahi GH, Jangravi Z, Sahraei H, Bahari Z. The possible pathophysiology mechanism
327 of cytokine storm in elderly adults with COVID-19 infection: the contribution of "inflame-
328 aging". *Inflamm Res*. 2020 Jun 11:1–15. doi: 10.1007/s00011-020-01372-8. Epub ahead
329 of print.
- 330 17. Daneshkhah A, Agrawal V, Eshein A, Subramanian H, Kumar Roy H, Backman V. The
331 Possible Role of Vitamin D in Suppressing Cytokine Storm and Associated Mortality in
332 COVID-19 Patients. *medRxiv* 2020.04.08.20058578; doi:
333 <https://doi.org/10.1101/2020.04.08.20058578>.
- 334 18. Quesada-Gomez JM, Castillo ME, Bouillon R. Vitamin D Receptor stimulation to reduce
335 Acute Respiratory Distress Syndrome (ARDS) in patients with Coronavirus SARS-CoV-2
336 infections. *J Steroid Biochem Mol Biol* 2020 Jun 11:105719. doi:
337 [10.1016/j.jsbmb.2020.105719](https://doi.org/10.1016/j.jsbmb.2020.105719). Epub ahead of print.
- 338 19. Li X, He J, Yu M, Sun J. The efficacy of vitamin D therapy for patients with COPD: a
339 meta-analysis of randomized controlled trials. *Ann Palliat Med* 2020;9(2):286-97.
- 340 20. Meltzer DO, Best TJ, Zhang H, Vokes T, Arora V, Solway J. Association of Vitamin D
341 Deficiency and Treatment with COVID-19 Incidence *medRxiv* 2020.05.08.20095893; doi:
342 <https://doi.org/10.1101/2020.05.08.20095893>.
- 343 21. Zheng Z, Peng F, Xu B, et al. Risk factors of critical & mortal COVID-19 cases: A
344 systematic literature review and meta-analysis. *J Infect* 2020; [Epub ahead of print].
345 doi:10.1016/j.jinf.2020.04.021
- 346 22. Metzler B, Siostrzonek P, Binder RK, Bauer A, Reinstadler SJ. Decline of acute coronary
347 syndrome admissions in Austria since the outbreak of COVID-19: the pandemic response
348 causes cardiac collateral damage. *Eur Heart J* 2020 Apr 16. [Epub ahead of print].
- 349 23. Tapper EB, Asrani SK. COVID-19 pandemic will have a long-lasting impact on the quality
350 of cirrhosis care. *J Hepatol* 2020 Apr 13. [Epub ahead of print].
- 351 24. Baddour K, Kudrick LD, Neopane A, Sabik LM, Peddada SD, Nilsen ML, et al. Potential
352 impact of the COVID-19 pandemic on financial toxicity in cancer survivors. *Head Neck*.
353 2020 Apr 24. [Epub ahead of print].
- 354 25. Søreide K, Hallet J, Matthews JB, Schnitzbauer AA, Line PD, Lai PBS, et al. Immediate
355 and long-term impact of the COVID-19 pandemic on delivery of surgical services. *Br J*
356 *Surg* 2020 Apr 30. [Epub ahead of print].
- 357 26. Zentrum für Krebsregisterdaten (ZfKD) im Robert Koch-Institut, Gesellschaft der
358 epidemiologischen Krebsregister e.V. (GEKID) (Eds.). *Krebs in Deutschland für*
359 *2015/2016*. Berlin, Robert Koch-Institut, 2019.
- 360 27. Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Evaluation, treatment, and prevention
361 of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol*
362 *Metab* 2011;96(7):1911-30.

Table 1 – Distribution of characteristics of the study population, and prevalence of vitamin D deficiency and insufficiency by those characteristics

Characteristic	Proportion of characteristic in total population [%]	Prevalence of vitamin D insufficiency among subjects with characteristic [%]	Prevalence of vitamin D deficiency among subjects with characteristic [%]	p
Total cohort	100.0	43.8	15.1	
Sex				<0.0001
Female	56.2	52.6	15.7	
Male	43.8	32.6	14.2	
Age (years)				<0.0001
50-64	61.3	42.4	14.2	
65-69	22.9	45.2	15.5	
70-75	15.8	47.3	17.7	
Month of recruitment (Season)				<0.0001
Jan-Feb	21.7	45.7	24.4	
Mar-Apr	13.9	51.6	21.2	
Mai-Jun	14.2	47.8	12.4	
Jul-Aug	17.0	35.6	6.9	
Sep-Oct	17.0	37.0	8.1	
Nov-Dec	16.2	47.0	15.6	
School education				<0.0001
≤ 9 years	75.0	44.7	15.3	
9 – 11 years	14.1	44.3	13.5	
≥ 12 years	11.0	36.1	15.4	
Smoking				<0.0001
Never	50.5	47.9	14.5	
Former	32.6	37.6	12.2	
Current	16.9	43.2	21.4	
BMI (kg/m ²)				<0.0001
< 30	74.5	42.1	14.1	
≥ 30	25.5	48.9	17.9	
Physical activity ^a				<0.0001
Low	67.1	45.9	16.7	
Moderate or high	32.9	39.6	11.6	
Fish consumption at least once per week				0.027
No	33.5	44.6	15.6	
Yes	66.5	43.2	14.2	

Abbreviations: BMI, body mass index

^a Defined by ≤ 1 h/week of vigorous physical activity that causes sweating

Table 2. Mortality from major causes of death among people with vitamin D deficiency and insufficiency compared to people with sufficient vitamin D status.

Cause of death	25(OH)D [nmol/L]	Participants	Number of deaths	Mortality ^a	Age, sex and season adjusted model		Full model	
					Hazard ratio (95% CI)	p-value	Hazard ratio (95% CI) ^b	p-value
Any cause	> 50	3924	873	15.6	Ref		Ref	
	30-50	4186	1010	17.1	1.28 (1.16-1.41)	<0.0001	1.20 (1.09-1.32)	0.0002
	< 30	1438	480	24.9	1.91 (1.70-2.15)	<0.0001	1.67 (1.48-1.89)	<0.0001
Cardiovascular disease	> 50	3924	313	5.6	Ref		Ref	
	30-50	4186	344	5.8	1.18 (1.00-1.39)	0.0452	1.10 (0.94-1.30)	0.2329
	< 30	1438	158	8.2	1.73 (1.41-2.12)	<0.0001	1.52 (1.23-1.86)	<0.0001
Cancer	> 50	3924	328	5.9	Ref		Ref	
	30-50	4186	344	5.8	1.17 (1.00-1.38)	0.0491	1.10 (0.94-1.29)	0.2320
	< 30	1438	153	7.9	1.58 (1.28-1.92)	<0.0001	1.38 (1.13-1.70)	0.0020
Respiratory disease ^c	> 50	3924	34	0.6	Ref		Ref	
	30-50	4186	58	1.0	2.24 (1.45-3.52)	0.0004	2.06 (1.32-3.21)	0.0015
	< 30	1438	31	1.6	3.69 (2.18-6.21)	<0.0001	3.04 (1.79-5.17)	<0.0001

Abbreviations: 25(OH)D, 25-hydroxyvitamin D; CI, confidence interval; Ref, reference

^a Mortality rate per 1000 person-years.

^b Estimate from Cox proportional hazards regression model adjusted for sex, age, season of blood draw, school education, smoking, body mass index, physical activity, and fish consumption.

^c Allocated to the following ICD-10-code groups: n=11, J18 - pneumonia; n=2, J41-J42 - chronic bronchitis; n=3, J43 - emphysema; n=75, J44 - COPD; n=2, J45 - asthma; n=12, J60-J70 - Lung diseases due to external agents; n=14, J80-J84 - Other respiratory diseases principally affecting the interstitium; n=1, J90-J94 Other diseases of pleura; n=2, J95-J99 - Other diseases of the respiratory system.

Table 3. Sex specific analysis on the associations of vitamin D deficiency and insufficiency with mortality from major causes of death.

Cause of death	25(OH)D [nmol/L]	Women					Men					p-value for interaction
		N _{total}	N _{deaths}	Mortality ^a	Hazard ratio (95% CI) ^b	p-value	N _{total}	N _{deaths}	Mortality ^a	Hazard ratio (95% CI) ^b	p-value	
Any cause	> 50	1701	235	9.4	Ref		2223	638	20.8	Ref		
	30-50	2821	558	13.7	1.24 (1.06-1.45)	0.0070	1365	452	24.7	1.19 (1.05-1.35)	0.0071	0.713
	< 30	843	246	21.0	1.76 (1.45-2.12)	<0.0001	595	234	30.8	1.66 (1.42-1.96)	<0.0001	0.703
Cardiovascular disease	> 50	1701	81	3.2	Ref		2223	232	7.5	Ref		
	30-50	2821	190	4.7	1.17 (0.90-1.53)	0.245	1365	154	8.4	1.11 (0.90-1.38)	0.3159	0.770
	< 30	843	90	7.7	1.78 (1.30-2.44)	0.0003	595	68	9.0	1.37 (1.02-1.83)	0.0335	0.198
Cancer	> 50	1701	95	3.8	Ref		2223	233	7.6	Ref		
	30-50	2821	188	4.6	1.12 (0.87-1.44)	0.3922	1365	156	8.5	1.09 (0.88-1.34)	0.4262	0.770
	< 30	843	64	5.5	1.22 (0.87-1.70)	0.2450	595	89	11.7	1.58 (1.22-2.07)	0.0007	0.198
Respiratory disease	> 50	1701	3	0.1	Ref		2223	31	1.0	Ref		
	30-50	2821	23	0.6	4.28 (1.27-1.42)	0.0189	1365	35	1.9	1.89 (1.14-3.15)	0.0136	0.201
	< 30	843	16	1.4	8.47 (2.38-30.12)	0.0010	595	15	2.0	2.25 (1.14-4.42)	0.0190	0.041

Abbreviations: 25(OH)D, 25-hydroxyvitamin D; CI, confidence interval; Ref, reference

^a Mortality rate per 1000 person-years.

^b Estimate from Cox proportional hazards regression model adjusted for sex, age, season of blood draw, school education, smoking, body mass index, physical activity, and fish consumption.

Figure 1. Prevalence of vitamin D deficiency and insufficiency (A) and proportion of deaths from respiratory diseases statistically attributable to these conditions (B)