Depressive symptoms and single nucleotide polymorphisms

predict clinical recurrence of inflammatory bowel disease

Short title: Depression and IBD course

Authors

Sebastian Bruno Ulrich Jordi ^{1,2}, Brian Matthew Lang ³, Bianca Auschra ⁴, Roland von Känel ⁴, Luc Biedermann ², Thomas Greuter ², Philipp Schreiner ², Gerhard Rogler ², Niklas Krupka ¹, Michael Christian Sulz ⁵, Benjamin Misselwitz ^{1,2*}, Stefan Begré ^{6,7*} on behalf of the Swiss IBD cohort study group[#]

¹ Clinic for Visceral Surgery and Medicine, Inselspital Bern and Bern University, Bern, Switzerland

² Department of Gastroenterology and Hepatology, University Hospital Zurich and Zurich University, Zurich, Switzerland

³ Clinic for Transplantation Immunology and Nephrology (Swiss Transplant Cohort Study), University Hospital of Basel, Basel, Switzerland

⁴ Department of Consultation-Liaison Psychiatry and Psychosomatic Medicine, University Hospital Zurich and University of Zurich, Zurich, Switzerland

⁵ Department of Gastroenterology and Hepatology, Kantonsspital St. Gallen, St. Gallen, Switzerland

⁶ Neurology, Department of Biomedical Research, Bern University Hospital, University of Bern, Bern, Switzerland

⁷ ISFOM - Institute of Stress Diseases and Stressmanagement, Zurich, Switzerland

* Both authors contributed equally and share last authorship

Members of the SIBDCS study group: Claudia Anderegg; Peter Bauerfeind; Christoph Beglinger; Stefan Begré; Dominique Belli; José M. Bengoa; Luc

27 Biedermann; Beat Bigler; Janek Binek; Mirjam Blattmann; Stephan Boehm; Jan
 28 Borovicka; Christian P. Braegger; Nora Brunner; Patrick Bühr; Bernard Burnand;
 29 Emanuel Burri; Sophie Buyse; Matthias Cremer; Dominique H. Criblez; Philippe de
 30 Saussure; Lukas Degen; Joakim Delarive; Christopher Doerig; Barbara Dora; Gian
 31 Dorta; Mara Egger; Tobias Ehmann; Ali El-Wafa; Matthias Engelmann; Jessica Ezri;
 32 Christian Felley; Markus Fliegner; Nicolas Fournier; Montserrat Fraga; Pascal Frei;
 33 Remus Frei; Michael Fried; Florian Froehlich; Christian Funk; Raoul Ivano Furlano;
 34 Suzanne Gallot-Lavallée; Martin Geyer; Marc Girardin; Delphine Golay; Tanja
 35 Grandinetti; Beat Gysi; Horst Haack; Johannes Haarer; Beat Helbling; Peter
 36 Hengstler; Denise Herzog; Cyrill Hess; Klaas Heyland; Thomas Hinterleitner;
 37 Philippe Hiroz; Claudia Hirschi; Petr Hruz; Rika Iwata; Res Jost; Pascal Juillerat;
 38 Vera Kessler Brondolo; Christina Knellwolf; Christoph Knoblauch; Henrik Köhler;
 39 Rebekka Koller; Claudia Krieger-Grübel; Gerd Kullak-Ublick; Patrizia Künzler;
 40 Markus Landolt; Rupprecht Lange; Frank Serge Lehmann; Andrew Macpherson;
 41 Philippe Maerten; Michel H. Maillard; Christine Manser; Michael Manz; Urs Marbet;
 42 George Marx; Christoph Matter; Valérie McLin; Rémy Meier; Martina Mendanova;
 43 Christa Meyenberger; Pierre Michetti; Benjamin Misselwitz; Darius Moradpour;
 44 Bernhard Morell; Patrick Mosler; Christian Mottet; Christoph Müller; Pascal Müller;
 45 Beat Müllhaupt; Claudia Mürger-Beyeler; Leilla Musso; Andreas Nagy; Michaela
 46 Neagu; Cristina Nichita; Jan Niess; Natacha Noël; Andreas Nydegger; Nicole Obialo;
 47 Carl Oneta; Cassandra Oropesa; Ueli Peter; Daniel Peternac; Laetitia Marie Petit;
 48 Franziska Piccoli-Gfeller; Julia Beatrice Pilz; Valérie Pittet; Nadia Raschle; Ronald
 49 Rentsch; Sophie Restellini; Jean-Pierre Richterich; Sylvia Rihs; Marc Alain Ritz;
 50 Jocelyn Roduit; Daniela Rogler; Gerhard Rogler; Jean-Benoît Rossel; Markus
 51 Sagmeister; Gaby Saner; Bernhard Sauter; Mikael Sawatzki; Michela Schächli;

52 Michael Scharl; Martin Schelling; Susanne Schibli; Hugo Schlauri; Sybille Schmid
 53 Uebelhart; Jean-François Schnegg; Alain Schoepfer; Frank Seibold; Mariam Seirafi;
 54 Gian-Marco Semadeni; David Semela; Arne Senning; Marc Sidler; Christiane
 55 Sokollik; Johannes Spalinger; Holger Spangenberger; Philippe Stadler; Michael
 56 Steuerwald; Alex Straumann; Bigna Straumann-Funk; Michael Sulz; Joël Thorens;
 57 Sarah Tiedemann; Radu Tutuian; Stephan Vavricksa; Francesco Viani; Jürg Vögtlin;
 58 Roland vVon Känel; Alain Vonlaufen; Dominique Vouillamoz; Rachel Vulliamy; Jürg
 59 Wermuth; Helene Werner; Paul Wiesel; Reiner Wiest; Tina Wylie; Jonas Zeitz;
 60 Dorothee Zimmermann.

61

62 **Correspondence**

63 Stefan Begré, MD

64 Address: ISFOM - Institute of Stress Diseases and Stressmanagement,
 65 Weinbergstrasse 139, CH-8006 Zurich, Switzerland

66 E-mail address: stefan.begre@dbmr.unibe.ch

67

68

69 **Authors' contributions**

70 The study was conceived and designed by S.B.U.J., B.M. and S.B.
 71 Statistical analyses were performed by S.B.U.J. and B.M.L. The manuscript was
 72 written by S.B.U.J., B.M. and S.B. The following authors have contributed to data
 73 interpretation and revised the manuscript for important intellectual content: B.M.L.,
 74 R.v.K., B.A., L.B., T.G., P.S., G.R., N.K. and M.C.S. All authors approved the final
 75 version of the manuscript.

76

77 **Abbreviations**

78 **aHR:** adjusted hazard ratio

79 **AIC:** Akaike information criterion

80 **BMI:** body mass index

81 **CD:** Crohn's disease

82 **CDAI:** Crohn's Disease Activity Index

83 **CHRNA5:** cholinergic receptor nicotinic alpha 5 subunit

84 **CI:** confidence interval

85 **EIM:** extraintestinal manifestations

86 **ENS:** enteric nervous system

87 **HADS:** Hospital Anxiety and Depression Scale

88 **HR:** hazard ratio

89 **IBD:** inflammatory bowel disease

90 **IC:** indeterminate colitis

91 **LD:** linkage disequilibrium

92 **MTWAI:** Modified Truelove and Witts Severity Index

93 **NACHR:** nicotinic acetylcholine receptors

94 **OR:** odds ratio

95 **PCLO:** piccolo presynaptic cytomatrix protein

96 **PSC:** primary sclerosing cholangitis

97 **R²:** square of the correlation coefficient

- 98 **SIBDC**: Swiss IBD cohort
- 99 **SNP**: single nucleotide polymorphism
- 100 **t_n**: follow-up time *n* of recording
- 101 **TNF**: tumor necrosis factor
- 102 **UC**: ulcerative colitis
- 103

Abstract

Background and Aims

Inflammatory bowel disease (IBD) patients are at high risk for depression. We examined interrelations between genetic risk factors for depression, depressive symptoms and IBD flares.

Methods

In 1973 patients (1137 Crohn's disease, 836 ulcerative colitis) of the Swiss IBD cohort study (SIBDC), 62 single nucleotide polymorphisms (SNPs) preselected for associations with depression, stress, pain and smoking were screened for cross-sectional associations with depression (hospital anxiety and depression subscale for depression, HADS-D $\geq$ 11). Logistic regression and Cox proportional hazards models were built to test for effects of depressive symptoms on disease course and genetic risk factors on depression and disease course. As endpoints we used active disease (CDAI $\geq$ 150 or MTWAI $\geq$ 10) and two published composite flare definitions: FNCE: physician reported flare, non-response to therapy, new complication or extraintestinal manifestation and AFFSST: active disease, physician reported flare, fistula, stenosis and new systemic therapy.

Results

Depressive symptoms were a strong risk factor for disease related endpoints including active disease (adjusted hazard ratio, aHR: 3.25, p<0.001), AFFSST (aHR: 1.62, p<0.001) and FNCE (aHR: 1.35, p=0.019). Rs588765's TC alleles and rs2522833's C allele were associated with depressive symptoms at baseline (odds

ratio, OR: 0.43, $q=0.050$ and OR: 1.73, $q=0.059$, respectively). Rs588765-TC remained protective regarding presence of depression (aHR: 0.67, $p=0.035$) and was associated with fewer active disease states (aHR: 0.72, $p=0.045$) during follow-up.

Conclusions

In IBD, genetics, depressive symptoms and inflammatory activity are intimately related: Depressive symptoms were a predictor of clinical deterioration and rs588765-TC was protective for depression and high IBD activity.

Funding

This work was supported by the Swiss National Science Foundation (SNSF).

Introduction

Inflammatory bowel diseases (IBD) are frequent conditions affecting approximately 2.2 million people in Europe and 1.5 million Americans with an increasing incidence worldwide.¹ IBD, comprising the subtypes Crohn's disease (CD), ulcerative colitis (UC) and indeterminate colitis (IC) are immunologically mediated diseases which can severely impair affected individuals' lives. The disease course of IBD is variable with periods of quiescent disease which are interrupted by IBD flares, i.e. episodes of more active inflammation.

The exact cause of IBD remains elusive, but likely involves an interplay of genetic, microbial, immunological and environmental factors.² A significant part of IBD risk is heritable and more than 230 single nucleotide polymorphisms (SNP) associated with IBD have been identified.³ These SNPs partially explain an individual's vulnerability to develop IBD but have not been useful to predict the clinical disease course in IBD patients.⁴ Many environmental factors have been implicated as trigger factors for IBD flares including diet, medication and smoking.⁵

Psychiatric conditions such as stress, depression and anxiety have been associated with IBD and the risk to experience depression is higher for IBD patients compared to healthy individuals (incidence rate ratio: 1.6).⁶ Furthermore, stress and depression overlap with pain⁷, pain sensitivity⁸ and smoking.⁹ Stress, depression and anxiety have been suggested as triggers of flares and clinical deterioration in IBD.^{10,11}

Depressive disorders are heritable conditions with heritability rates of approximately 40%¹² and many individual SNPs have been associated with the risk for depression.¹³ An example for a possible genetic origin of the observed overlap of depression, smoking and pain sensitivity are neuronal nicotinic acetylcholine

receptors (nAChRs). Recent findings suggest that nAChRs are involved in psychiatric disorders such as depression although mechanistic details remain unclear.¹⁴ Moreover, nAChRs had been associated with nicotine and alcohol dependence,^{15,16} also among SIBDC patients.¹⁷ Currently, nAChRs are investigated as pharmacological targets for depression, anxiety, pain and nicotine addiction.¹⁸ For these reasons, SNPs within nAChR genes might influence the occurrence of flares in IBD via their association with depression and/ or their link to smoking.

However, the role of genetic risk factors for depressive symptoms has not yet been clarified for individuals at high risk for depression such as IBD patients and also the interplay between genetic and environmental risk factors of depression remains unclear.

We are using data from the Swiss IBD cohort study (SIBDC)¹⁹ to explore the relationship between depression, individual SNPs and the risk for IBD deterioration. SIBDC was started in 2006 as a prospective cohort study that recruits nationwide in Switzerland.^{19,20} The impact of depressive symptoms, anxiety and social support on IBD flares has been studied in SIBDC before;^{10,21} we were aiming to validate, generalize and extend these findings, and add a genetic context.

Methods

Design of the SIBDC

The SIBDC was started in 2006 as a prospective cohort study that recruits nationwide in Switzerland.^{19,20} New patients are continuously recruited and patients are followed yearly. IBD-related clinical data is collected by physicians at enrolment and yearly follow-ups. In addition, clinical, sociodemographic and psychosocial data are acquired using patient questionnaires at enrolment and yearly follow-ups. All SIBDC patients consenting to genetic analysis with appropriate biomaterial available were genotyped and only genotyped patients were included in our study.

SIBDCS is funded by the Swiss National Science Foundation and has been approved by the local ethics committee of each participating center (institutional review board approval No. EK-1316, approved on February 5, 2007 and KEK Zurich, March 9, 2020; BASEC 2018-02068). SIBDC goals and methodology are described elsewhere.^{19,20} All patients provided written informed consent prior to inclusion into the SIBDCS. Analysis of patient data for the current study was approved by the scientific board of SIBDCS.

Patient characteristics

We exported clinical and sociodemographic variables acquired at enrolment and follow-ups: Disease activity was measured using the Crohn's disease activity index CDAI or the modified Truelove and Witts activity index (MTWAI) for CD and UC/IC patients, respectively. For intercomparable measures, CDAI or MTWAI score values were Z-score normalized to a disease activity score (disease activity score = $(s_i - \bar{s}) / \sigma_s$) or cutoff values for active disease were used (see below). The following

binary variables were created by distinguishing between patients that had any of the respective included criteria and patients that did not: *Complications* included colorectal cancer, colon dysplasia, intestinal lymphoma, osteopenia/osteoporosis, anemia (not as an adverse event of medical therapies), deep vein thrombosis, pulmonary embolism, nephrolithiasis, gallstones,²² malabsorption syndrome, massive hemorrhage, perforation or peritonitis. *Extraintestinal manifestations* (EIM) comprised peripheral arthritis/arthritis, uveitis/iritis, pyoderma gangrenosum, erythema nodosum, aphthous oral ulcers/stomatitis, ankylosing spondylitis/sacroiliitis and primary sclerosing cholangitis (PSC). Any stenosis localized in the esophagus, duodenum/jejunum, ileum, large bowel, rectum or anus was summarized in the *stenosis* variable. The *fistula* variable comprised IBD-related fistula and abscesses and anal fissure, whereas the variable *surgery* included any abdominal or fistula and abscess-related surgery. *TNF inhibitors* referred to treatments with infliximab, certolizumab pegol or adalimumab (no patient with golimumab treatment was reported at the time of data export). The variable *number of current therapies* was created by counting presently administered therapies which could be any of the following: any steroids, any antibiotics, 5-aminosalicylic acid, sulfasalazine, azathioprine, 6-mercaptopurine, methotrexate, cyclosporine, tacrolimus, infliximab, adalimumab, certolizumab pegol, cholestyramine, *E. coli* Nissle (Mutaflor®), ursodeoxycholic acid, bisphosphonates, the probiotic VSL#3®, or any other medication that was noted by physicians. A patient was considered a smoker when he or she declared to have a current habit of smoking independent of the quantity.

230

231 Outcome measures

232 Depressive symptoms were measured with the Hospital Anxiety and Depression
233 Scale's (HADS) subscale for depression (HADS-D).²³ Following well-established
234 criteria, the presence of depression was defined as HADS-D ≥ 11 , indicating probable
235 moderate or severe depression.²⁴⁻²⁶ This was done for both a constant (at baseline)
236 and a time-varying depression variable. In this study, the term *depression* is used
237 with reference to this definition unless stated otherwise.

238 Active disease was defined by CDAI ≥ 150 ²⁷ or MTWAI ≥ 10 ²⁸, respectively. An IBD
239 flare was measured as a composite endpoint of clinical variables. IBD is a complex
240 disease and we considered two different flare definitions, following the examples of
241 two prior SIBDC studies.^{10,21} The first composite clinical endpoint *active disease,*
242 *physician reported flare, fistula, stenosis, surgery or new systemic therapy*
243 (AFFSST) was reached upon occurrence/persistence of active disease, defined by
244 CDAI ≥ 150 ²⁷ or MTWAI ≥ 10 ²⁸, a physician reported flare, new fistula, new stenosis,
245 new surgery, intake of systemic steroids and/or start of therapy with a new TNF
246 inhibitor (also changes between TNF inhibitors) at follow-up.¹⁰

247 The second composite clinical endpoint *physician reported flare, non-response to*
248 *any administered therapy with consequent transition to a more aggressive*
249 *medication, new complication or new EIM* (FNCE) was defined as the occurrence of
250 any one of these events at a follow-up examination.²¹

251 For both composite endpoints we counted the first occurrence of each defining
252 feature after enrolment as event. Both composite endpoints were reached upon the
253 first occurrence of any of these events.

For data analyses with survival techniques, all endpoints (the individual flare features, the FNCE and AFFSST composite endpoints and depression) were coded with right censoring. Thereby, a time interval t_{n-1} - t_n was defined where t_n corresponds to the n^{th} follow-up visit when the event was documented first. This way, we describe a time interval during which an event had occurred $[t_{n-1}, t_n]$ and then assumed an event's occurrence at $t_{n-1}+(t_n-t_{n-1})/2$ on average.

Selection of candidate SNPs

Allelic state of a number of SNPs were determined in SIBDC patients. We selected all 95 SNPs available without a link to inflammation or IBD diagnosis. These *non-inflammatory SNPs* had been selected with three objectives: 48 SNPs were selected for an association with stress and/or depressive symptoms, based on a thorough literature search in PubMed and SNPedia. Moreover, 30 SNPs with a suspected association with pain were included. Furthermore, 16 SNPs associated with smoking in all 3 genome wide association studies available in 2015 were included.^{17,29-31} Both, pain⁷ and smoking^{11,37} have a strong reciprocal associations with depression and anxiety.

For calculation of linkage disequilibrium (LD) we used the square of the correlation coefficient (r^2) between two SNPs. Groups were formed whenever two or more SNPs shared an r^2 value of 0.5 or higher, representing moderate LD.³² The SNP with the strongest correlation to group members was chosen as the representing tag-SNP for the respective group. If there were only two SNPs in a group, the SNP with the lowest number of missing data points in the respective group was selected as tag SNP.

Screening for depression-associated SNPs

For the screening of SNPs associated with depression, we used methods of the *SNPassoc* R package³³ version 1.9-2, developed and established for whole genome association studies. We calculated the association between each of our SNP and depression while simultaneously selecting the genetic model (codominant, dominant, recessive, overdominant, log-additive) for each SNP which best describes this relationship via likelihood ratio test comparison. We filtered the list of SNPs to those with a Bonferroni corrected p-value (q-value) <0.1. Associations were confirmed by multivariable analyses with depressive symptoms at enrolment as endpoint. For multivariable analyses with depressive symptoms as dependent variable in this study, a set of control variables was defined (set 1: *sex, diagnosis, age, time since diagnosis, BMI, smoking status, alcohol consumption, standardized CDAI/MTWAI score, abdominal surgery prior to enrolment, systemic steroids, TNF inhibitors, number of current therapies*). These variables were tested for association with depressive symptoms at enrolment using univariable logistic regression analysis.

Models included all control variables and were calculated for every SNP individually. To minimize the risk of obtaining significance due to overfitting we also created a simpler secondary model by automated variable elimination. In this process the optimal model regarding the AIC was calculated. For this we used the *glmulti* R package³⁴ version 1.0.7.1.

Time-to-event analyses

To confirm associations with SNPs at enrolment (cross-sectional data), we constructed time-to-event models using the *Survival* R package³⁵ version 3.1-12. We implemented univariable and multivariable Cox proportional hazards models with the endpoint of interest being presence of depressive symptoms (HADS ≥ 11) at a follow-up examination, independent of a patient's HADS score value at enrolment.

Clinical disease course was described by the absence or presence of either active disease or flares according to composite endpoint definitions (see above). To assess the impact of depressive symptoms and depression-associated SNPs on the clinical course of IBD,¹⁰ we coded a time-varying control variable for depression (HADS-D score ≥ 11) that reflected a patient's depression status over time.³⁶ To ensure an unambiguous time sequence of time-varying depression status and clinical endpoint measurements (e.g. flares), a conservative coding approach was chosen: only if a recording of depressive symptoms preceded a flare, an association was made. Depression status was defined forwardly, while flares were defined backwardly. For example, measures for depression at time point t_n defined the time-varying depression variable for the interval t_n until t_{n+1} , while recordings of clinical variables at time point t_n defining a clinical endpoint variable (e.g. flare) were assumed to have happened at the time point in the middle between the last and the current follow-up (calculated as $t_{n-1} + (t_n - t_{n-1})/2$). As a result, concurrent depression and flares will not be detected as an association.

For these time-to-event analyses with clinical endpoints we adapted the set of control variables by removing variables that could be proxy-variables for clinical endpoints (set 2: *sex, diagnosis, time since diagnosis, age, BMI, disease related surgery prior to enrolment, smoking status, daily alcohol consumption*). As before,

325 univariable and multivariable Cox proportional hazards models were implemented.

326 All calculations were performed using R^{37} version 3.6.1.

Results

Study participants

For our analysis, data from 1973 genotyped SIBDC patients (1137 with CD, 836 with UC/IC) were available with regular follow-ups up to 4250 days. Baseline characteristics of our cohort sample are presented in **Table 1**, confirming a mixed study population with mild, moderate and severe disease. Depression (HADS-D ≥ 11) at enrolment tended to be more prevalent in CD compared to UC (9.5% in CD vs. 6.5% in UC/IC, $p=0.052$), consistent with earlier findings.³⁸ Smoking,³⁹ fistulae, stenosis, EIMs, complications, previous abdominal surgery and therapy with TNF inhibitors was more common in CD patients and CD patients had significantly longer disease courses and lower BMIs. On the other hand, UC/IC patients had a higher number of different therapies at enrolment and were more often treated with systemic steroids.

Depressive symptoms at enrolment were significantly associated with several control variables including BMI ($p=0.01$), daily alcohol consumption ($p=0.017$), disease activity ($p<0.001$) and number of current therapies ($p=0.004$, **Table 2**).

Screening of SNPs for an association with depressive symptoms

SNPs were tested for linkage disequilibrium (**Supplementary figure 1**) and SNPs with a pair-wise correlation coefficient $r^2 \geq 0.5$ aggregated in 16 groups. From each group only one representative SNP was used henceforward, reducing the number of SNPs from 95 to 62 (**Supplementary table 1**).

All SNPs were screened for an association with depression (HADS-D ≥ 11) at enrolment in three different models: univariable, complete multivariable and reduced

multivariable model after automated parameter elimination (**Supplementary table 2**). Of 62 SNPs tested, two SNPs, rs588765 and rs2522833, were significantly associated with depressive symptoms at enrolment with $q < 0.1$ (Bonferroni corrected). In the univariable model, for rs2522833 the association was strongest in a log-additive inheritance model with the C allele as an associated risk allele (odds ratio, OR: 1.71, CI: 1.25-2.34, $q = 0.043$). For rs588765, an overdominant genetic pattern yielded the strongest significance identifying TC as an associated protective allele combination (OR: 0.45, CI: 0.28-0.74, $q = 0.066$, **Table 3** and **Supplementary figure 2**). For both SNPs, the associations detected in the univariable models stayed significant ($q < 0.1$) in multivariable models correcting for possible confounders (rs2522833: $q = 0.045$; rs588765: $q = 0.061$, **Supplementary figure 3**), also after parameter elimination (rs2522833: $q = 0.059$; rs588765: $q = 0.050$, **Supplementary figure 4**). This indicates a robust association between both SNPs and depressive symptoms

SNPs effects on depression-free survival

To assess the effects of both SNPs on the hazard for continuance or new onset of depressive symptoms (HADS-D ≥ 11) over time, we implemented Cox proportional hazards models. In this time-to-event analyses, the TC allele combination of rs588765 had significant protective effects regarding depressive symptoms in the univariable model (hazard ratio (HR)): 0.68, confidence interval (CI): 0.46-0.97, $p = 0.042$). The association with rs588765-TC remained robust in the multivariable analysis (adjusted hazard ratio (aHR): 0.67, CI: 0.46-0.97, $p = 0.035$; **Table 4** and **Figure 1; panel A**). Protective effects of rs588765-TC were independent of disease

type and similar for CD and UC patients, although non-significant in each subgroup (CD: aHR: 0.70, $p=0.155$; UC: aHR: 0.58, $p=0.100$).

For the C-allele of rs2522833, no significant association was detected (aHR: 1.10, CI: 0.86-1.41, $p=0.455$, **Table 4** and **Supplementary figure 5; panel A**).

Depressive symptoms as a hazard for IBD disease course

Patients with depressive symptoms (HADS-D ≥ 11) had a significantly higher hazard of experiencing active disease (CDAI ≥ 150 /MTWAI ≥ 10) during follow-up (HR: 3.18, $p<0.001$, **Table 5** and **Supplementary figure 6; panel A**); this association remained significant after correction for confounders (aHR: 3.25, $p<0.001$, **Table 5** and **Figure 2; panel A**). We also observed a significant negative impact of depressive symptoms on the endpoints *new surgery* (aHR: 1.92, $p>0.001$), *systemic steroid therapy* (aHR: 1.55, $p=0.033$) and *primary non-response to therapy* (aHR: 1.79, $p=0.044$, **Figure 2; panel E-G**, univariable results: **Supplementary figure 7**). In other words, patients were more likely to experience the respective endpoints when depressive symptoms were present before, even when corrected for possible confounders.

Overall, for all clinical endpoints, patients with depressive symptoms had a higher hazard for clinical deterioration, even though this effect did not reach significance in all cases (**Figure 2**). Furthermore, depressive symptoms also tended to increase the hazard for individual EIMs and significant effects for new PSC (aHR: 3.88, $p=0.032$) and new uveitis/iritis (aHR: 2.01, $p=0.046$) in multivariable models were observed (**Figure 3; panel E and G**, univariable results: **Supplementary figure 8**).

To further investigate the impact of depressive symptoms on IBD disease course,¹⁰ we assessed two previously used composite endpoints indicating clinical

deterioration: AFFSST¹⁰ (active disease, physician reported flare, new fistula, new stenosis, surgery or new systemic therapy, see methods) and FNCE²¹ (physician reported flare, non-response to therapy, new complication or EIM).

Patients with depressive symptoms were also more likely to reach these composite endpoints: in the univariable analysis, the AFFSST endpoint (HR: 1.57, CI: 1.23-1.99, $p < 0.001$, **Table 5**) as well as the FNCE endpoint (HR: 1.37, CI: 1.07-1.74, $p = 0.012$, **Table 5**) were significantly associated with preceding depressive symptoms. Correction for control variables barely changed results and significance was maintained (**Figure 4**).

Effects of SNPs on IBD activity over time

We tested whether rs588765-TC or rs2522833's C allele could be directly implicated as protective or risk factors for clinical deterioration (**Table 6**). In univariable models, the TC allele combination of rs588765 showed significant protective effects against active disease (CDAI ≥ 150 / MTWAI ≥ 10 , HR: 0.78, CI: 0.61-0.99, $p = 0.042$). These significant protective effects of rs588765's TC alleles remained significant after controlling for confounders (**Figure 1; panel B**).

In contrast, the C allele of rs2522833 was not a significant risk factor for active disease (univariable: HR: 1.08, CI: 0.92-1.28, $p = 0.344$, multivariable: **Supplementary figure 5; panel B**).

Genetic inheritance patterns of rs588765 and rs2522833

In a sensitivity analysis we examined the impact of different allele variants of rs588765 on outcomes of interest. Similarly as for initial screening, the

overdominance model (TC vs. TT or CC) seemed to provide the best fit for Cox proportional models with depressive symptoms over time and active disease as endpoints (**Supplementary tables 3 and 4, Supplementary figure 8; panel A and C**). Similarly, for rs2522833, the log-additive inheritance model which uses the number of C alleles (0, 1 or 2) seemed to be the best fit regarding active disease (**Supplementary table 4, Supplementary figure 8; panel D**), but not regarding depressive symptoms over time (**Supplementary table 3, Supplementary figure 8; panel B**). No formal statistical testing comparing the fits of genetic models was done.

Discussion

In our cross-sectional analysis of depressive symptoms in IBD patients, we identified variants of two SNPs (rs588765-TC, rs2522833's C allele) that were significantly ($q < 0.10$) associated with depression in IBD. The presence of depressive symptoms was a strong risk factor for subsequent clinical deterioration in time-to-event analyses, confirming and extending earlier findings by showing robust negative effects on numerous clinical endpoints as well as two previously published composite endpoints for clinical deterioration.^{10,21} Finally, we could demonstrate that the TC allele of rs588765 was protective regarding depression and IBD activity in a time-to-event analysis.

IBD patients are a high-risk group for developing depression.⁶ In line with these findings, we observed an elevated point prevalence of depressive symptoms in our cohort (77/814, 9.5% among CD patients and 40/618, 6.5% among UC/IC patients) compared to the general population. For instance, point prevalence in a non-clinical Swiss sample with comparable assessment was estimated to be only 3.1%,⁴⁰ similar to findings in the USA⁴¹ and the UK.²⁶

We found a significant negative association between depressive symptoms and patients with a specific variant in one of the preselected SNPs (rs588765-TC, ORs of 0.43, $q = 0.05$) in a cross-sectional analysis with enrolment data. This is an observation that, to the best of our knowledge, had not been described before. Moreover, the time-to-event analysis indicated a protective effect of the same allele combination regarding new or persistent depressive symptoms over time (aHR: 0.67, $p = 0.035$), thus independently confirming protective properties of rs588765-TC. Finally, protective effects of rs588765-TC regarding depression translated to

protection regarding IBD flares (CDAI ≥ 150 /MTWAI ≥ 10 , aHR: 0.72, $p=0.045$). All analyses were corrected for smoking status and alcohol consumption indicating that protection regarding depression and flares is independent of these features.

Rs588765 is part of the chr15q25.1 region and represents a well-established tag-SNP on the CHRNA5-CHRNA3-CHRNA4 gene cluster^{42,43} that encodes the nAChR subunits $\alpha 5$, $\alpha 3$, and $\beta 4$. Rs588765 is located in the neuronal acetylcholine receptor subunit $\alpha 5$ gene (CHRNA5).⁴⁴ CHRNA5's gene product serves as a subunit of pentameric nAChRs.⁴⁵ Rs588765 impacts the expression of CHRNA5 in human brain tissue⁴⁶ with subjects homozygous with two minor alleles (TT) showing the highest CHRNA5 mRNA expression.⁴⁶ NACHRs are ligand-gated ion channels that mediate signal transmission at synapses and modulate the release of different neurotransmitters.⁴⁷ Addition of an $\alpha 5$ subunit to nAChRs $\alpha 4\beta 2$, the most frequent receptor subtype in the brain, increases the rate of receptor desensitization and calcium permeability.⁴⁷ Similarly to the brain, CHRNA5 seems to mediate excitatory nicotinic cholinergic transmission in the enteric nervous system (ENS).⁴⁸ Rs588765 had been associated with nicotine and alcohol dependence,^{15,16} also among SIBDC patients.¹⁷ Moreover, the associated CHRNA5-CHRNA3-CHRNA4 region has also been suggested to contain risk genes for affective disorders such as major depressive disorder⁴⁹⁻⁵¹ and recent findings suggest that nAChRs in general are involved in psychiatric disorders such as depression.¹⁴

The genetic model of overdominance we stated best fitting for rs588765 refers to advantageous effects of the heterozygous variant compared to homozygous variants.⁵² Examples include protective effects of hemoglobin mutation against malaria of in human sickle cell anaemia.⁵³ For rs588765, functional interactions of genetic variants associated with the T and C allele, respectively, seem possible as a

variant of epistasis. Another plausible explanation for overdominance is the concept of an inverted U-shaped response curve.⁵⁴

The second SNP (rs2522833) associated with depressive symptoms is part of the piccolo presynaptic cytomatrix protein (PCLO) gene. The presynaptic cytomatrix describes a complex electron-dense meshwork of proteins involved in transmitter release.^{55,56} Moreover, PCLO has been implicated to potentially contributing to short-term neuronal plasticity.⁴⁸ The PCLO gene has been associated with depressive conditions such as major depressive disorder⁵⁷ and the C allele of rs2522833 specifically might increase an individual's vulnerability regarding depression.⁵⁸ Our current results align well with these earlier findings and the C allele was a significant ($q < 0.1$) log-additive risk factor for depressive symptoms at baseline (ORs: 1.73, $q = 0.059$). However, our study did not detect significance for rs2522833 in the time-to-event analyses.

In our analysis, depressive symptoms remained a strong and highly significant risk factor for future clinical deterioration according to several clinical measures including the well-established disease scores CDAI and MTWAI and two pre-defined, partially overlapping composite flare definitions (AFFSST¹⁰ and FNCE²¹, respectively). Depressive symptoms also increased the likelihood for future occurrence of nearly all other IBD-related endpoints tested over time (Figures 2 and 3), even though without uniform statistical significance. This suggests that the specific definition for a flare is largely irrelevant for the association with depression.

A growing body of evidence indicates that stress in general^{11,21} and depressive symptoms specifically¹⁰ are associated with clinical flares in IBD patients. Whether the psychological state of an IBD patient can indeed directly increase gut inflammation remains an important question in IBD research.⁵⁹ Our results support a

direct impact of depressive symptoms on gut inflammation: i) we show strong associations between rs588765, depressive symptoms and IBD flares (Tables 3-6); ii) depressive symptoms remain a strong risk factor for flares according to various definitions (CDAI/MTWAI: aHR: 3.25, $p < 0.001$, AFFSST: aHR: 1.62, $p < 0.001$, FNCE: aHR: 1.35, $p = 0.019$) and iii) the TC allele of rs588765 with protective effects regarding depressive symptoms (aHR: 0.67, $p = 0.035$) was also protective against IBD flares (CDAI/MTWAI: aHR: 0.72, $p = 0.045$). The combination of clinical and genetic data strengthens this point; however, indirect effects cannot be entirely excluded. For instance, depression might be a marker for subclinical inflammation or might affect reporting of symptoms by patients. Another interesting possibility could be an effect of rs588765-TC on CHRNA5 expression in the gut and the brain simultaneously, thus independently explaining depressive symptoms and flares. Further mechanistic studies would be necessary to better understand our observation.

Our study has several strengths and limitations. Strengths include the excellent clinical characterization regarding IBD-related outcomes and the sequential measurements of the key variable *depressive symptoms* and the clinical endpoint variables over a long time enabling the combination of cross-sectional and time-to-event analyses. Limitations are the relatively small sample size for genetic analyses, yielding in lower statistical power for some analyses. Moreover, missing data points, e.g. for therapy or surgery data, further reduced statistical power for some analyses. Furthermore, even though serial measurements for HADS-D were available and a HADS-D score of ≥ 11 represents a well-established cut off strongly indicating depression,²³⁻²⁶ a diagnosis of depression normally requires direct clinical assessment by a psychiatrist. Furthermore, for validation of our data, independent

confirmation in another IBD cohort would have been desirable. Finally, in 2019, a large meta-analysis for the risk of depression including 807,553 individuals was published. This study identified 102 independent variants (not rs588765, rs2522833) of which 87 were replicated in an independent sample.¹³ Therefore, future studies with a larger number of SNPs will now be possible.

In conclusion, our study confirms an intimate relationship between depressive symptoms and IBD disease course. The prevalence of depression among IBD patients is high and active disease further increases the hazard for depression.⁶ Vice versa, depressive symptoms also independently increased the hazard for IBD flares over time. Moreover, two SNPs (rs588765-TC, rs2522833's C allele) were associated with depressive symptoms among IBD patients and one of those SNPs (rs588765-TC) was also associated with high IBD activity. These findings are in accordance with a multidirectional relationship between genetics, depression and IBD disease course. Therefore, a comprehensive therapeutic concept for IBD patients, targeting inflammation and depression might further improve patient care.⁶⁰

546 **Acknowledgments**

547 The authors would like to thank SIBDC patients and SIBDC staff for their
548 commitment and are grateful to the SIBDCS Scientific Committee for granting
549 approval for this study.

550

References

1. Jairath V, Feagan BG. Global burden of inflammatory bowel disease. *Lancet Gastroenterol Hepatol* 2020;**5**:2-3.
2. Zhang YZ, Li YY. Inflammatory bowel disease: Pathogenesis. *World J Gastroenterol* 2014;**20**:91-9.
3. Turpin W, Goethel A, Bedrani L, Croitoru M, K. Determinants of ibd heritability: Genes, bugs, and more. *Inflamm Bowel Dis* 2018;**24**:1133-48.
4. Cleyneen I, Boucher G, Jostins L, et al. Inherited determinants of crohn's disease and ulcerative colitis phenotypes: A genetic association study. *Lancet* 2016;**387**:156-67.
5. Ananthakrishnan AN, Bernstein CN, Iliopoulos D, et al. Environmental triggers in ibd: A review of progress and evidence. *Nat Rev Gastroenterol Hepatol* 2018;**15**:39-49.
6. Bernstein CN, Hitchon CA, Walld R, et al. Increased burden of psychiatric disorders in inflammatory bowel disease. *Inflamm Bowel Dis* 2019;**25**:360-8.
7. Goesling J, Clauw DJ, Hassett AL. Pain and depression: An integrative review of neurobiological and psychological factors. *Curr Psychiatry Rep* 2013;**15**:421.
8. Thompson T, Correll CU, Gallop K, Vancampfort D, Stubbs B. Is pain perception altered in people with depression? A systematic review and meta-analysis of experimental pain research. *J Pain* 2016;**17**:1257-72.
9. Fluharty M, Taylor AE, Grabski M, Munafo MR. The association of cigarette smoking with depression and anxiety: A systematic review. *Nicotine Tob Res* 2017;**19**:3-13.
10. Mikocka-Walus A, Pittet V, Rossel JB, von Kanel R, Swiss IBDCSG. Symptoms of depression and anxiety are independently associated with clinical recurrence of inflammatory bowel disease. *Clin Gastroenterol Hepatol* 2016;**14**:829-35 e1.
11. Sun Y, Li L, Xie R, et al. Stress triggers flare of inflammatory bowel disease in children and adults. *Front Pediatr* 2019;**7**:432.
12. Sullivan PF, Neale MC, Kendler KS. Genetic epidemiology of major depression: Review and meta-analysis. *Am J Psychiatry* 2000;**157**:1552-62.
13. Howard DM, Adams MJ, Clarke TK, et al. Genome-wide meta-analysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. *Nat Neurosci* 2019;**22**:343-52.
14. Rahman S, Engleman EA, Bell RL. Chapter six - recent advances in nicotinic receptor signaling in alcohol abuse and alcoholism. In: Rahman S, editor. *Progress in molecular biology and translational science*: Academic Press; 2016: 183-201.
15. Hallfors J, Loukola A, Pitkaniemi J, et al. Scrutiny of the chrna5-chrna3-chrnb4 smoking behavior locus reveals a novel association with alcohol use in a finnish population based study. *Int J Mol Epidemiol Genet* 2013;**4**:109-19.
16. Criado JR, Gizer IR, Edenberg HJ, Ehlers CL. Chrna5 and chrna3 variants and level of neuroticism in young adult mexican american men and women. *Twin Res Hum Genet* 2014;**17**:80-8.
17. Lang BM, Biedermann L, van Haaften WT, et al. Genetic polymorphisms associated with smoking behaviour predict the risk of surgery in patients with crohn's disease. *Aliment Pharmacol Ther* 2018;**47**:55-66.
18. Romanelli MN, Gratteri P, Guandalini L, et al. Central nicotinic receptors: Structure, function, ligands, and therapeutic potential. *ChemMedChem* 2007;**2**:746-67.
19. Pittet V, Michetti P, Mueller C, et al. Cohort profile update: The swiss inflammatory bowel disease cohort study (sibdcs). *Int J Epidemiol* 2019;**48**:385-6f.

- 597 20. Pittet V, Juillerat P, Mottet C, *et al.* Cohort profile: The swiss inflammatory bowel
598 disease cohort study (sibdcs). *Int J Epidemiol* 2009;**38**:922-31.
- 599 21. Camara RJ, Lukas PS, Begre S, *et al.* Effects of social support on the clinical course of
600 crohn's disease. *Inflamm Bowel Dis* 2011;**17**:1277-86.
- 601 22. Fagagnini S, Heinrich H, Rossel JB, *et al.* Risk factors for gallstones and kidney stones
602 in a cohort of patients with inflammatory bowel diseases. *PLoS One*
603 2017;**12**:e0185193.
- 604 23. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr*
605 *Scand* 1983;**67**:361-70.
- 606 24. Snaith RP. The hospital anxiety and depression scale. *Health Qual Life Outcomes*
607 2003;**1**:29.
- 608 25. Stern AF. The hospital anxiety and depression scale. *Occup Med (Lond)* 2014;**64**:393-
609 4.
- 610 26. Crawford JR, Henry JD, Crombie C, Taylor EP. Normative data for the hads from a
611 large non-clinical sample. *Br J Clin Psychol* 2001;**40**:429-34.
- 612 27. Lashner B, Brzezinski A. Crohn's disease. *GI/Liver Secrets* 2010:297-303.
- 613 28. Lichtiger S, Present DH, Kornbluth A, *et al.* Cyclosporine in severe ulcerative colitis
614 refractory to steroid therapy. *N Engl J Med* 1994;**330**:1841-5.
- 615 29. Tobacco, Genetics C. Genome-wide meta-analyses identify multiple loci associated
616 with smoking behavior. *Nat Genet* 2010;**42**:441-7.
- 617 30. Liu JZ, Tozzi F, Waterworth DM, *et al.* Meta-analysis and imputation refines the
618 association of 15q25 with smoking quantity. *Nat Genet* 2010;**42**:436-40.
- 619 31. Thorgeirsson TE, Gudbjartsson DF, Surakka I, *et al.* Sequence variants at chrnb3-
620 chrna6 and cyp2a6 affect smoking behavior. *Nat Genet* 2010;**42**:448-53.
- 621 32. Carlson CS, Eberle MA, Rieder MJ, *et al.* Selecting a maximally informative set of
622 single-nucleotide polymorphisms for association analyses using linkage
623 disequilibrium. *Am J Hum Genet* 2004;**74**:106-20.
- 624 33. Gonzalez JR, Armengol L, Guino E, Sole X, Moreno V. Snpassoc: Snps-based whole
625 genome association studies. Version 1.9-2 edn., 2014.
- 626 34. Calcagno V. Glmulti: Model selection and multimodel inference made easy. Version
627 1.0.7.1 edn., 2019.
- 628 35. Therneau TM. A package for survival analysis in r. Version 3.1-12 edn., 2020.
- 629 36. Zhang Z, Reinikainen J, Adeleke KA, Pieterse ME, Groothuis-Oudshoorn CGM. Time-
630 varying covariates and coefficients in cox regression models. *Ann Transl Med*
631 2018;**6**:121.
- 632 37. R Core Team. R: A language and environment for statistical computing. Vienna,
633 Austria: R Foundation for Statistical Computing, 2019.
- 634 38. Geiss T, Schaefert RM, Berens S, Hoffmann P, Gauss A. Risk of depression in patients
635 with inflammatory bowel disease. *J Dig Dis* 2018;**19**:456-67.
- 636 39. Lunney PC, Kariyawasam VC, Wang RR, *et al.* Smoking prevalence and its influence
637 on disease course and surgery in crohn's disease and ulcerative colitis. *Aliment*
638 *Pharmacol Ther* 2015;**42**:61-70.
- 639 40. Baer N. *Depressionen in der schweizer bevölkerung : Daten zur epidemiologie,*
640 *behandlung und sozial-beruflichen integration*: Schweizerisches
641 Gesundheitsobservatorium; 2013.
- 642 41. Kessler RC, Wang PS. Epidemiology of depression. *Handbook of depression, 2nd ed*
643 New York, NY, US: The Guilford Press; 2009: 5-22.

- 644 42. Saccone NL, Culverhouse RC, Schwantes-An TH, *et al.* Multiple independent loci at
645 chromosome 15q25.1 affect smoking quantity: A meta-analysis and comparison with
646 lung cancer and copd. *PLoS Genet* 2010;**6**.
- 647 43. Ji X, Gui J, Han Y, *et al.* The role of haplotype in 15q25.1 locus in lung cancer risk:
648 Results of scanning chromosome 15. *Carcinogenesis* 2015;**36**:1275-83.
- 649 44. Bergen AW, Javitz HS, Krasnow R, *et al.* Nicotinic acetylcholine receptor variation and
650 response to smoking cessation therapies. *Pharmacogenet Genomics* 2013;**23**:94-103.
- 651 45. Paterson D, Nordberg A. Neuronal nicotinic receptors in the human brain. *Prog*
652 *Neurobiol* 2000;**61**:75-111.
- 653 46. Wang JC, Grucza R, Cruchaga C, *et al.* Genetic variation in the chrna5 gene affects
654 mrna levels and is associated with risk for alcohol dependence. *Mol Psychiatry*
655 2009;**14**:501-10.
- 656 47. Wang JC, Cruchaga C, Saccone NL, *et al.* Risk for nicotine dependence and lung
657 cancer is conferred by mrna expression levels and amino acid change in chrna5. *Hum*
658 *Mol Genet* 2009;**18**:3125-35.
- 659 48. Garza A, Huang LZ, Son JH, Winzer-Serhan UH. Expression of nicotinic acetylcholine
660 receptors and subunit messenger rnas in the enteric nervous system of the neonatal
661 rat. *Neuroscience* 2009;**158**:1521-9.
- 662 49. Bhalala OG, Nath AP, Consortium UKBE, Inouye M, Sibley CR. Identification of
663 expression quantitative trait loci associated with schizophrenia and affective
664 disorders in normal brain tissue. *PLoS Genet* 2018;**14**:e1007607.
- 665 50. Jackson KJ, Fanous AH, Chen J, Kendler KS, Chen X. Variants in the 15q25 gene
666 cluster are associated with risk for schizophrenia and bipolar disorder. *Psychiatr*
667 *Genet* 2013;**23**:20-8.
- 668 51. Broms U, Wedenoja J, Largeau MR, *et al.* Analysis of detailed phenotype profiles
669 reveals chrna5-chrna3-chrnb4 gene cluster association with several nicotine
670 dependence traits. *Nicotine Tob Res* 2012;**14**:720-33.
- 671 52. Lippman ZB, Zamir D. Heterosis: Revisiting the magic. *Trends Genet* 2007;**23**:60-6.
- 672 53. Ferreira A, Marguti I, Bechmann I, *et al.* Sickle hemoglobin confers tolerance to
673 plasmodium infection. *Cell* 2011;**145**:398-409.
- 674 54. Comings DE, MacMurray JP. Molecular heterosis: A review. *Mol Genet Metab*
675 2000;**71**:19-31.
- 676 55. Gundelfinger ED, Fejtova A. Molecular organization and plasticity of the cytomatrix
677 at the active zone. *Curr Opin Neurobiol* 2012;**22**:423-30.
- 678 56. Parthier D, Kuner T, Korber C. The presynaptic scaffolding protein piccolo organizes
679 the readily releasable pool at the calyx of held. *J Physiol* 2018;**596**:1485-99.
- 680 57. Mbarek H, Milaneschi Y, Hottenga J-J, *et al.* Genome-wide significance for pclo as a
681 gene for major depressive disorder. *Twin Research and Human Genetics*
682 2017;**20**:267-70.
- 683 58. Woudstra S, van Tol MJ, Bochdanovits Z, *et al.* Modulatory effects of the piccolo
684 genotype on emotional memory in health and depression. *PLoS One* 2013;**8**:e61494.
- 685 59. Mikocka-Walus AA, Turnbull DA, Moulding NT, *et al.* Controversies surrounding the
686 comorbidity of depression and anxiety in inflammatory bowel disease patients: A
687 literature review. *Inflamm Bowel Dis* 2007;**13**:225-34.
- 688 60. Tarricone I, Regazzi MG, Bonucci G, *et al.* Prevalence and effectiveness of psychiatric
689 treatments for patients with ibd: A systematic literature review. *J Psychosom Res*
690 2017;**101**:68-95.

Tables

Table 1

Variables at enrolment Median (1 st quartile-3 rd quartile); min-max		CD	UC/IC	P-value
Diagnosis N=1973		1137 (57.6%)	836 (42.4%)	
Sex N=1973	Male	567/1137 (49.9%)	452/836 (54.1%)	0.072
	Female	570/1137 (50.1%)	384/836 (45.9%)	
Age (years) N=1973		37.2 (24.6-50.0); 6.5-85.5	38.5 (26.8-49.0); 4.6-81.8	0.458
Time since diagnosis (years) N=1970		6.7 (1.7-15.4); 0- 52.5	4.9 (1.4-11.7); 0- 49.5	< 0.001
BMI N=1942		22.5 (19.9-25.7); 12.3-48.1	23 (20.7-25.9); 13.1-48.8	0.003
Smoking status N=1643	Smoker	301/929 (32.4%)	73/714 (10.2%)	< 0.001
	Non-smoker	628/929 (67.6%)	641/714 (89.8%)	
Daily alcohol consumption N=1434	Yes	703/820 (85.7%)	512/614 (83.4%)	0.251
	No	117/820 (14.3%)	102/614 (16.6%)	
Disease activity N=1973	CDAI N= 1137	34.0 (11.0-80.0); 0.0-450.0		-
	MTWAI N=836		2.0 (-0.0-3.4); 0.0-19.0	-
	Standardized CDAI/MTWAI #	-0.3 (-0.7-0.6); -0.9- 7.3	-0.3 (-0.9-0.6); - 0.9-4.6	0.009
Active disease (CDAI ≥150 or MTWAI ≥10) N=1973	Yes	83/1137 (7.3%)	71/836 (8.5%)	0.373
	No	1054/1137 (92.7%)	765/836 (91.5%)	
Systemic steroids: yes N=1956	Yes	199/1124 (17.7%)	219/832 (26.3%)	< 0.001
	No	925/1124 (82.3%)	613/832 (73.7%)	
TNF inhibitors: yes N=1956	Yes	344/1124 (30.6%)	95/832 (11.4%)	< 0.001
	No	780/1124 (69.4%)	737/832 (88.6%)	
Number current therapies N=1956		1 (1-2); 0-6	2 (1-3); 0-7	< 0.001
Disease related surgery prior	Yes	550/1137 (48.4%)	93/836 (11.1%)	< 0.001

to enrolment N=1973	No	587/1137 (51.6%)	743/836 (88.9%)	
Fistula at or prior to enrolment N=1973	Yes	468/1137 (41.2%)	45/836 (5.4%)	< 0.001
	No	669/1137 (58.8%)	791/836 (94.6%)	
Stenosis at or prior to enrolment N=1973	Yes	366/1137 (32.2%)	18/836 (2.1%)	< 0.001
	No	771/1137 (67.8%)	818/836 (97.6%)	
EIM at or prior to enrolment N=1973	Yes	498/1137 (43.8%)	265/836 (31.7%)	< 0.001
	No	639/1137 (56.2%)	571/836 (68.3%)	
Complications at or prior to enrolment N=1973	Yes	562/1137 (49.4%)	356/836 (42.6%)	0.003
	No	575/1137 (50.6%)	480/836 (57.4%)	
rs588765 N=1973	CC	448/1137 (39.4%)	278/836 (33.2%)	0.018
	TC	509/1137 (44.8%)	406/836 (48.6%)	
	TT	180/1137 (15.8%)	152/836 (18.2%)	
rs2522833 N=1973	AA	419/1137 (36.9%)	300/836 (35.9%)	0.883
	AC	524/1137 (46.1%)	388/836 (46.4%)	
	CC	194/1137 (17.1%)	148/836 (17.7%)	
HADS score (A and D) N=1432		10 (5-16); 0-36	9 (5-15); 0-39	0.117
Depression (HADS-D ≥11) N=1432	Yes	77/814 (9.5%)	40/618 (6.5%)	0.052
	No	737/814 (90.5%)	578/618 (93.5%)	
Anxiety (HADS-A ≥11) N=1432	Yes	156/814 (19.2%)	93/619 (15.0%)	0.048
	No	658/814 (80.8%)	526/619 (85.0%)	
# Standardization of score values was obtained by norming values in reference to the respective score mean and standard deviation (disease activity score = $(s_i - \bar{s}) / \sigma_s$). This results in comparable values between the different scores (CDAI/ MTWAI) that indicate a patients score value in reference to the mean score value of the respective score and taking each score's internal variance into account.				

Table 1: Study participants. Characteristics of study participants at enrolment stratified for diagnosis. Abbreviations: CD: Crohn's disease, UC: ulcerative colitis, IC: indeterminate colitis, BMI: body mass index, CDAI: Crohn's Disease Activity Index, MTWAI: Modified Truelove and Witts Severity Index, TNF: tumor necrosis factor, T: thymine, C: cytosine, A: adenine, HADS: Hospital Anxiety and Depression Scale, HADS-D: subscale for depressive symptoms, HADS-A: subscale for anxiety symptoms.

Table 2

Univariable logistic regression analyses - odds ratios for depression in IBD patients

Dependent variable: depression (HADS depression sub score ≥ 11)

Sex: male	0.91 (CI: 0.58-1.43)
Diagnosis: UC/IC (vs. CD)	0.79 (CI: 0.50-1.26)
Age (years)	1.01 (CI: 1.00-1.03)
Time since diagnosis (years)	1.00 (CI: 0.97-1.02)
BMI	1.07 (CI: 1.02-1.11) ***
Smoking status: smoker	1.42 (CI: 0.90-2.26)
Daily alcohol consumption: yes	3.12 (CI: 1.24-7.83) **
Disease activity index (standardized CDAI/MTWAI)	1.80 (CI: 1.53-2.13) ***
Disease related surgery prior to enrolment	1.00 (CI: 0.64-1.62)
Systemic steroids: yes	1.55 (CI: 0.95-2.54) *
TNF inhibitors: yes	1.30 (CI: 0.78-1.59)
Number current therapies	1.01 (CI: 1.00-1.03) ***

Table 2: Variables associated with depression (HADS-D ≥ 11) at baseline.

Univariable logistic regression models showing the association of possible confounders for multivariable models with depression. Results are presented as odds ratios; p-values are indicated as follows: *: $p < 0.1$, **: $p < 0.05$, ***: $p < 0.01$. 1043 patients were analyzed. Abbreviations: CI: 95% confidence interval, IBD: inflammatory bowel disease, HADS: Hospital Anxiety and Depression Scale, UC: ulcerative colitis, IC: indeterminate colitis, CD: Crohn's disease, BMI: body mass index, CDAI: Crohn's Disease Activity Index, MTWAI: Modified Truelove and Witts Severity Index, TNF: tumor necrosis factor.

711 **Table 3**

Association analysis - odds ratios for depression in IBD patients

Dependent variable: depression at enrolment (HADS depression sub score ≥ 11)			
	Univariable	Multivariable without variable elimination	Multivariable with variable elimination
rs588765: TC (vs. CC/TT)	0.45 (CI: 0.28-0.74) *	0.43 (CI: 0.26-0.72) *	0.43 (CI: 0.25-0.72) **
rs2522833: number of C alleles (0-2)	1.71 (CI: 1.25-2.34) **	1.75 (CI: 1.26-2.44) **	1.73 (CI: 1.25-2.39) *

712 **Table 3: SNPs associated with depression (HADS-D ≥ 11).** Table showing the
713 analysis results of the two SNPs significantly ($q < 0.1$) associated with depression at
714 enrolment in IBD patients. Results are presented as odds ratios; q-values are
715 indicated as follows: *: $q < 0.1$, **: $q < 0.05$, ***: $q < 0.01$. Analyses: logistic regression
716 analysis. Abbreviations: CI: 95% confidence interval, IBD: inflammatory bowel
717 disease, HADS: Hospital Anxiety and Depression Scale, T: thymine, C: cytosine.

718

719 **Table 4**

Cox proportional hazards model - hazard ratios for new or persistent depression among IBD patients over time

Endpoint variable: depression (HADS depression sub score ≥11)			
	Univariable	Multivariable (rs588765)	Multivariable (rs2522833)
rs588765: TC (vs. CC/TT)	0.68 (CI: 0.46-0.99) **	0.67 (CI: 0.46-0.97) **	
rs2522833: number of C alleles (0-2)	1.12 (CI: 0.87-1.44)		1.10 (CI: 0.86-1.41)
Sex: male		0.80 (CI: 0.54-1.19)	0.79 (CI: 0.54-1.18)
Diagnosis: UC/IC (vs. CD)		0.82 (CI: 0.51-1.30)	0.80 (CI: 0.50-1.27)
Time since diagnosis (years)		0.99 (CI: 0.97-1.01)	0.99 (CI: 0.97-1.01)
Age (years)		1.03 (CI: 1.01-1.04) ***	1.03 (CI: 1.01-1.04) ***
BMI		1.00 (CI: 0.95-1.04)	1.00 (CI: 0.96-1.04)
Smoking status: smoker		1.63 (CI: 1.09-2.42) **	1.65 (CI: 1.11-2.44) **
Daily alcohol consumption: yes		1.21 (CI: 0.71 -2.07)	1.24 (CI: 0.73 -2.12)
Disease activity (standardized CDAI/MTWAI)		1.53 (CI: 1.31-1.79) ***	1.52 (CI: 1.30-1.78) ***
Disease related surgery prior to enrolment		1.15 (CI: 0.74-1.78)	1.14 (CI: 0.73-1.78)
Systemic steroids: yes		0.99 (CI: 0.60-1.63)	0.99 (CI: 0.60-1.64)
TNF inhibitors: yes		1.26 (CI: 0.80-1.99)	1.25 (CI: 0.79-1.96)
Number current therapies		1.07 (CI: 0.87-1.31)	1.06 (CI: 0.86-1.30)

Table 4: Predictors for new or persistent depression (HADS-D ≥ 11) over time.

Results are presented as hazard ratios for new or persistent depression among IBD patients over time; p-values are indicated as follows: *: $p < 0.1$, **: $p < 0.05$, ***: $p < 0.01$. 919 patients were analyzed and 114 times the endpoint was reached. Analysis: Cox proportional hazards model. Abbreviations: CI: 95% confidence interval, IBD: inflammatory bowel disease, HADS: Hospital Anxiety and Depression Scale, T: thymine, C: cytosine, UC: ulcerative colitis, IC: indeterminate colitis, CD: Crohn's disease, BMI: body mass index, CDAI: Crohn's Disease Activity Index, MTWAI: Modified Truelove and Witts Severity Index, TNF: tumor necrosis factor.

730 **Table 5**

Cox proportional hazards model - hazard ratios for active disease and clinical deterioration (flares) in IBD patients over time

	Endpoint variable: active disease (CDAI/MTWAI)		Endpoint variable: combined clinical endpoint (AFFSST)		Endpoint variable: combined clinical endpoint (FNCE)	
	Univariable	Multivariable	Univariable	Multivariable	Univariable	Multivariable
Preceding depression over time (yes)	3.18 (CI: 2.19-4.60) ***	3.25 (CI: 2.10-5.03) ***	1.57 (CI: 1.23-1.99) ***	1.62 (CI: 1.27 -2.08) ***	1.37 (CI: 1.07-1.74) **	1.35 (CI: 1.05 -1.72) **
Sex: male		0.86 (CI: 0.62-1.20)		0.95 (CI: 0.83-1.10)		0.89 (CI: 0.77-1.03)
Diagnosis: UC/IC (vs. CD)		1.73 (CI: 1.21-2.47) ***		0.89 (CI: 0.76-1.05)		1.09 (CI: 0.93-1.28)
Time since diagnosis (years)		1.01 (CI: 0.99-1.03)		1.00 (CI: 0.99-1.01)		1.00 (CI: 0.99-1.01)
Age (years) #		1.00 (CI: 0.99-1.01) ***		0.99 (CI: 0.98-1.00) ***		1.00 (CI: 0.99-1.00)
BMI		0.98 (CI: 0.95-1.02)		1.00 (CI: 0.98-1.01)		1.00 (CI: 0.98-1.02)
Disease related surgery prior to enrolment		1.60 (CI: 1.10 -2.32) **		1.02 (CI: 0.87 -1.20)		0.95 (CI: 0.80 -1.12)
Smoking		1.43 (CI: 1.04 -1.97) **		1.04 (CI: 0.87 -1.24)		1.14 (CI: 0.98 -1.32)

status: smoker		1.00-2.03) **		0.88-1.22)		0.97-1.34)
Daily alcohol consumption: yes		0.79 (CI: 0.51 -1.22)		1.17 (CI: 0.96 -1.43)		1.11 (CI: 0.90 -1.36)

Values of 1 with simultaneous significance were obtained by rounding and are not equal to 1

Table 5: Predictors for active disease and clinical deterioration (flare) in IBD

patients over time. Results are presented as hazard ratio for active disease or clinical deterioration (flare) respectively, in IBD patients over time. Active disease was defined as CDAI ≥ 150 or MTWAI ≥ 10 in CD and UC patients, respectively. A flare was defined by the AFFSST or FNCE composite endpoint; p-values are indicated as follows: *: $p < 0.1$, **: $p < 0.05$, ***: $p < 0.01$. Analysis: Cox proportional hazards model. Abbreviations: CI: 95% confidence interval, IBD: inflammatory bowel disease, HADS: Hospital Anxiety and Depression Scale, UC: ulcerative colitis, IC: indeterminate colitis, CD: Crohn's disease, BMI: body mass index, CDAI: Crohn's Disease Activity Index, MTWAI: Modified Truelove and Witts Severity Index, AFFSST: active disease, physician reported flare, new fistula, new stenosis, surgery or new systemic therapy, FNCE: physician reported flare, non-response to therapy, new complication or EIM.

746 **Table 6**

Cox proportional hazards model - hazard ratios for active disease in IBD patients over time

Endpoint variable: active disease (CDAI/MTWAI)				
	rs588765		rs2522833	
	Univariable (rs588765)	Multivariable (rs588765)	Univariable (rs2522833)	Multivariable (rs2522833)
rs588765: TC (vs. CC/TT)	0.78 (CI: 0.61-0.99) **	0.72 (CI: 0.52 - 0.99) **		
rs2522833: number of C alleles (0-2)			1.08 (CI: 0.92-1.28)	1.12 (CI: 0.90 - 1.39)
Sex: male		0.85 (CI: 0.61-1.18)		0.84 (CI: 0.61-1.17)
Diagnosis: UC/IC (vs. CD)		1.73 (CI: 1.19-2.50) ***		1.70 (CI: 1.17-2.46) ***
Time since diagnosis (years)		1.01 (CI: 0.99-1.03)		1.01 (CI: 0.99-1.03)
Age (years)		1.00 (CI: 0.98-1.01)		1.00 (CI: 0.98-1.01)
BMI		0.99 (CI: 0.95-1.03)		0.99 (CI: 0.95-1.03)
Disease related surgery prior to enrolment		1.66 (CI: 1.14 - 2.42) ***		1.63 (CI: 1.12 - 2.38) **
Smoking status: smoker		1.63 (CI: 1.15-2.31) ***		1.64 (CI: 1.16-2.33) ***

Daily alcohol consumption:		0.90 (CI: 0.59 -		0.88 (CI: 0.58 -
yes		1.38)		1.35)

Table 6: Predictors for active disease in IBD patients over time. Results are presented as hazard ratio for active disease in IBD patients over time. Active disease was defined as CDAI ≥ 150 or MTWAI ≥ 10 in CD and UC patients, respectively; p-values are indicated as follows: *: $p < 0.1$, **: $p < 0.05$, ***: $p < 0.01$. Analysis: Cox proportional hazards models. Abbreviations: CI: 95% confidence interval, IBD: inflammatory bowel disease, CDAI: Crohn's Disease Activity Index, MTWAI: Modified Truelove and Witts Severity Index, C: cytosine, T: thymine, UC: ulcerative colitis, IC: indeterminate colitis, CD: Crohn's disease, BMI: body mass index.

Figures

Figure 1

Figure 1: Kaplan Meier curves showing how rs588765's genotype impacts the hazards for depression and IBD activity over time. A: Percentage of IBD patients without depression stratified for rs588765 genotype (overdominant). **B:** Percentage of IBD patients without active disease defined as CDAI ≥ 150 or MTWAI ≥ 10 in CD and UC patients, respectively and stratified for rs588765 genotype (overdominant). Analysis: multivariable Cox proportional hazards models. Abbreviations: HR: hazard ratio, CI: confidence interval, T: thymine, C: cytosine, IBD: inflammatory bowel disease, CDAI: Crohn's Disease Activity Index, MTWAI: Modified Truelove and Witts Severity Index.

Figure 2

Figure 2: Kaplan Meier curves showing how depression increases the hazards for future adverse outcomes over time. Percentage of IBD patients without experiencing the respective outcome (A-H) stratified for depression. Analysis: multivariable Cox proportional hazards models. Abbreviations: HR: hazard ratio, CI: confidence interval, TNF: tumor necrosis factor, CDAI: Crohn's Disease Activity Index, MTWAI: Modified Truelove and Witts Severity Index.

Figure 3

Figure 3: Kaplan Meier curves showing how depression increases the hazards for future EIMs over time. Percentage of IBD patients without experiencing the respective outcome (A-H) stratified for depression. Analysis: multivariable Cox proportional hazards models. Abbreviations: HR: hazard ratio, CI: confidence interval, PSC: primary sclerosing cholangitis.

Figure 4

Figure 4: Kaplan Meier curves showing how depression increases the hazard for future clinical deterioration (flare, composite endpoints) over time. A: Percentage of IBD patients without a flare according to the AFFSST-definition, stratified according to the presence of depression. **B:** Percentage of IBD patients without a flare according to the FNCE-definition, stratified according to the presence of depression. Analysis: multivariable Cox proportional hazards models. Abbreviations: HR: hazard ratio, CI: confidence interval, IBD: inflammatory bowel disease, AFFSST: active disease, physician reported flare, new fistula, new stenosis, surgery or new systemic therapy, FNCE: physician reported flare, non-response to therapy, new complication or EIM.