Using past and current data to estimate potential crisis service use in mental healthcare after the COVID-19 lockdown: South London and Maudsley data

Robert Stewart^{1,2}, Matthew Broadbent²

¹King's College London (Institute of Psychiatry, Psychology and Neuroscience), London, UK

²South London and Maudsley NHS Foundation Trust, London, UK

Abstract

The lockdown policy response to the COVID-19 pandemic in the UK has a potentially important impact on provision of mental healthcare with uncertain consequences over the 12 months ahead. Past activity may provide a means to predict future demand. Taking advantage of the Clinical Record Interactive Search (CRIS) data resource at the South London and Maudsley NHS Trust (SLaM; a large mental health service provider for 1.2m residents in south London), we carried out a range of descriptive analyses to inform the Trust on patient groups who might be most likely to require inpatient and home treatment team (HTT) crisis care. We considered the 12 months following UK COVID-19 lockdown policy on 16th March, drawing on comparable findings from previous years, and quantified levels of change in service delivery to those most likely to receive crisis care. For 12-month crisis days from 16th March in 2015-19, we found that most (over 80%) were accounted for by inpatient care (rather than HTT), most (around 75%) were used by patients who were current or recent Trust patients at the commencement of follow-up, and highest numbers were used by patients with a previously recorded schizophreniform disorder diagnosis. For current/recent patients on 16th March there had been substantial reductions in use of inpatient care in the following 31 days in 2020, more than previous years; changes in total non-inpatient contact numbers did not differ in 2020 compared to previous years, although there had been a marked switch from face-to-face to virtual contacts.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Background

The COVID-19 pandemic may have a profound impact on health services through the direct effects of the virus itself, through social distancing policies and their psychological impact, and through the disrupting impact of social distancing policies on delivery of mental and physical healthcare. Mental healthcare services have had to be radically reconfigured to cope with suspected or confirmed COVID-19 infections in inpatient and outpatient settings, staff sickness or self-isolation, the need to minimise face-to-face contacts, and the need to accommodate increasing pressures on acute medical care from cases of viral pneumonia.

We have previously begun reporting on the mental healthcare impact of the UK COVID-19 pandemic, taking advantage of the Clinical Record Interactive Search (CRIS) data platform that receives 24-hourly updates from its source electronic records. Analysing activity in key services from February to mid-May 2020, and comparing periods before and after 16th March 2020, Community Mental Health Teams showed relatively stable caseloads and total contact numbers, but a substantial shift from face-to-face to virtual contacts, while Home Treatment Teams showed the same changeover but reductions in caseloads and total contacts (1).

Methods

The Biomedical Research Centre (BRC) Case Register at the South London and Maudsley NHS Foundation Trust (SLaM) has been described previously (2;3). SLaM serves a geographic catchment of four south London boroughs (Croydon, Lambeth, Lewisham, Southwark) with a population of around 1.2 million residents, and has used a fully electronic health record (EHR) across all its services since 2006. SLaM's BRC Case Register was set up in 2008, providing researcher access to de-identified data from SLaM's EHR via the Clinical Record Interactive Search (CRIS) platform and within a robust security model and governance framework (4). CRIS has been extensively developed over the last 10 years with a range of external data linkages and natural language processing resources (3). Of relevance to the work presented here, CRIS is updated from SLaM's EHR every 24 hours and thus provides relatively 'real-time' data. SLaM's EHR is itself immediately updated every time an entry is made, which include date-stamped fields indicating patient contacts ('events') and those indicating acceptance of a referral or a discharge from a given service (or SLaM care

more generally). CRIS data have been substantially enhanced by a suite of natural language processing (NLP) algorithms to extract structured data on named entities from free text fields; a catalogue describing these is available on the CRIS site (5). CRIS has supported over 200 peer reviewed publications to date, and has received approval as a data source for secondary analyses (Oxford Research Ethics Committee C, reference 18/SC/0372).

In order to investigate the composition of service user profiles associated with crisis service use over the 12 months following 16th March 2020 based on previous years' data, for each recent year (2015-2019), we took the 16th March as an index date and characterised patients who received any crisis care (inpatient or HTT) in the 12 months following each date (i.e. up to and including 15th March in the following year). To begin with, we divided these patients into three cohorts based on their status at the start of follow-up: i) Recent – those active to SLaM at any point in the 3 months up to 16th March that year; ii) Past – those previously active to SLaM but not within the 3 months up to 16th March that year; iii) New – those not previously known to SLaM before the 16th March that year. 'Active' here was defined on the basis of an open and accepted referral (i.e. regardless of whether a team had been assigned or a contact made). For each follow-up period, we investigated the proportion of 'crisis care' accounted for by inpatient vs. HTT-only. We compared total crisis person-days, inpatient person-days, and HTT-only person-days between the three cohorts for each year.

Further exploratory descriptive analyses of group (i) above divided the 12-month follow-up period into quarters to investigate whether crisis service use was higher in earlier compared to later periods after each index date. Finally, contributions to crisis days over each 12-month period for group (i) were described specifically for seven diagnostic groups, defined on the basis of any previous primary or secondary diagnosis received prior to each index data (thus overlapping rather than mutually exclusive): a) serious mental illness (ICD-10 codes F2x, F30x, F31x, F32.3, F33.3); b) schizophrenia-like disorders (F2x); c) bipolar disorder (F31x); d) any affective disorder (F3x); e) any organic disorder (F0x); f) any alcohol or substance use disorder (F1x); g) any personality disorder (F60x, F61x, or ascertained via an NLP algorithm extracting diagnostic statements in documentation).

In order to quantify and contextualise the recent changes in service contact experienced by current/recent service users in the early stages of lockdown, for each previous year we ascertained patients on March 16th who had been 'active' to SLaM in the preceding 3

calendar months (i.e. Group (i) above). Taking these cohorts, we ascertained total inpatient days and the following clinical contacts by non-inpatient teams: i) face-to-face contacts attended; ii) virtual contacts attended (by email, fax, mail, phone, online, or video link); iii) total contacts, as the sum of these two. These were quantified and compared for the 31 days up to and including 15th March, and for the 31 days from 16th March onward, for the years 2015 to 2020. Further analyses were then carried out restricting the comparisons to those with diagnoses recorded prior to 16th March in a given year, applying the same seven diagnostic groups as described above.

Results

Total crisis days for given 12-month periods are displayed and compared in Table 1, and were around the 350,000 to 360,000 range, tending to be lower in more recent years. Around 80-85% of crisis days were accounted for by inpatient days, reducing gradually from 2015-19. Around 75% of crisis days were used by current/recent patients at the start of each follow-up period (reducing from 77% to 73% from 2015-19); this proportion was lower for HTT days, with around 50-55% being used by recent patients (reducing over the years) and the remainder split fairly evenly between past and new patients.

In further analyses focusing on current/recent patients at each index date, dividing the 12-month follow-up period into quarters, crisis person-days were observed to be relatively stable over that period for current/recent patients (Figures 1a-c). Descriptions of crisis day usage by diagnostic groups of interest are displayed in Figures 2a and 2b. In summary, highest numbers of crisis days were accounted for by people with previously recorded schizophreniform diagnoses; use of crisis services had been decreasing in SMI, schizophreniform, bipolar and organic disorder diagnostic groups, had been relatively stable in patients with previously recorded affective disorder or alcohol/substance use disorder, and had been increasing in those with a previous personality disorder diagnosis. The proportion of crisis days accounted for by inpatient care (as opposed to HTT) was fairly similar between diagnostic groups, although marginally lower for those with previous affective disorder diagnoses; also, no substantial or consistent changes over the 5 years were observed.

Service contact changes for current/recent patients for the 31 day periods before and after 16th March are compared by year in Table 2. In previous years, a consistent 3-6% reduction in

inpatient person days was observed from periods before to after the index date, compared to the substantially larger 26% decrease observed in 2020. Face-to-face contacts showed a more heterogeneous 8-24% decrease over the same comparison period, compared to the 65% decrease in 2020, while virtual contacts had decreased by 3-22% in previous years compared to a 117% increase in 2020. Total contacts decreased by 21% in 2020, which was within the 8-23% range for previous years.

Service contact changes are further displayed by previous diagnosis in Figures 3a-d. For inpatient days, most diagnostic groups showed substantial differences in changes between 2020 and previous years, apart from bipolar disorder where these were not marked. Patterns across years for changes in face-to-face, virtual, and total contacts did not differ substantially between diagnostic groups. Reductions in total contacts in 2020 were least marked for bipolar and schizophreniform disorders and most marked for organic disorders, but all of these changes were within the range for the previous 5 years.

Discussion

We report a series of descriptive analyses carried out at SLaM, carried out to inform the Trust on patients who might be most likely to require inpatient and other (HTT) crisis care in the 12 months following UK COVID-19 lockdown policy on 16th March, based on findings from previous years, as well as to quantify levels of change in service delivery around that date for the group of current/recent service users most likely to be future users of crisis care. Key findings were as follows: i) 80-85% of crisis days were spent in inpatient care rather than under HTT; ii) around 75% of crisis days over a given 12 months were used by patients who were currently or recently under the care of the Trust at the commencement of that follow-up period; iii) highest numbers of crisis days were used by patients with a previously recorded schizophreniform disorder (ICD-10 F2x); iv) in recent/current patients, there had been substantial reductions in use of inpatient care in the 31 days after 16th March 2020, substantially more than in previous years; v) changes in total contacts from non-inpatient teams did not differ substantially in 2020 compared to previous years, although there had been a marked change from face-to-face to virtual contacts – this held for all diagnostic groups investigated.

Considering limitations, it is important to bear in mind that the data are derived from a single site. Because complete data are being provided for that site with no hypothetical source population intended, calculation of confidence intervals was not felt to be appropriate for the descriptive data provided in this report; applicability to other mental healthcare providers cannot therefore be inferred and would need specific investigation, although we recommend the broad approach as a methodology worth considering. Profiles of services and catchment morbidity are also likely to vary between sites; for example, the very high predominance of people with psychotic disorders in those requiring inpatient and/or HTT input may well be related to SLaM's particular inner urban catchment. Finally, the core task of predicting levels of future service activity, and the characteristics of those requiring highest-cost services, is clearly complex. On the one hand, there has been a marked consistency in the users of crisis services in past years; on the other hand, the COVID-19 pandemic has placed unique challenges on mental health services, those using them and those not yet using them but in need of them. Therefore it will be important to set up an adequate monitoring system to assess service use as close to real-time as possible and to identify any unexpected divergences from previous norms.

Funding

The research leading to these results has received support from the Medical Research Council Mental Health Data Pathfinder Award to King's College London. RS and MB are part-funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at the South London and Maudsley NHS Foundation Trust and King's College London; RS is additionally part-funded by: i) a Medical Research Council (MRC) Mental Health Data Pathfinder Award to King's College London; ii) an NIHR Senior Investigator Award; iii) the National Institute for Health Research (NIHR) Applied Research Collaboration South London (NIHR ARC South London) at King's College Hospital NHS Foundation Trust. The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

References

- (1) Stewart R, Martin E, Broadbent M. Mental health service activity during COVID-19 lockdown: South London and Maudsley data on working age community and home treatment team services and mortality from February to mid-May 2020. medRxiv 2020;doi: 10.1101/2020.06.13.20130419.
- (2) Stewart R, Soremekun M, Perera G, Broadbent M, Callard F, Denis M, et al. The South London and Maudsley NHS Foundation Trust Biomedical Research Centre (SLaM BRC) Case Register: development and descriptive data. BMC Psychiatry 2009;9:51.
- (3) Perera G, Broadbent M, Callard F, Chang C-K, Downs J, Dutta R, et al. Cohort profile of the South London and Maudsley NHS Foundation Trust Biomedical Research Centre (SLaM BRC) Case Register: current status and recent enhancement of an Electronic Mental Health Record derived data resource. BMJ Open 2016;6:e008721.
- (4) Fernandes AC, Cloete D, Broadbent MTM, Hayes RD, Chang C-K, Roberts A, et al. Development and evaluation of a de-identification procedure for a case register sourced from mental health electronic records. BMC Med Inform Decis Mak 2013 Jan 1;13:71.
- (5) CRIS NLP Service. Library of production-ready applications, Version 1.2. <https://www.maudsleybrc.nihr.ac.uk/media/313767/nlp-applications-library-v12.pdf>; 2020.

Table 1. Crisis days attributable to current/recent, past, and new mental health service users at the commencement of the 12-month follow-up period

Year	Cohort	N	Service use in the 12 months from 16 th March (person-days)					
			All crisis days (inpatient plus HTT)		Inpatient days		HTT days*	
2019	Recent	3105	254377	72.8%	220908	78.5%	36706	50.4%
	Past	1035	44124	12.6%	27422	9.7%	17780	24.4%
	New	1403	50780	14.5%	33242	11.8%	18367	25.2%
	Total		349281		281572	80.6%	72853	20.9%
2018	Recent	3054	260593	73.8%	226798	79.1%	36848	51.9%
	Past	992	39793	11.3%	23866	8.3%	16655	23.5%
	New	1360	52875	15.0%	36225	12.6%	17449	24.6%
	Total		353261		286889	81.2%	70952	20.1%
2017	Recent	3074	262578	75.3%	230060	80.4%	36139	53.3%
	Past	921	37720	10.8%	23528	8.2%	15201	22.4%
	New	1280	48198	13.8%	32487	11.4%	16426	24.2%
	Total		348496		286075	82.1%	67766	19.4%
2016	Recent	3102	276472	76.1%	241020	80.5%	40137	57.1%
	Past	823	34641	9.5%	22198	7.4%	13309	18.9%
	New	1246	52127	14.4%	36198	12.1%	16878	24.0%
	Total		363240		299416	82.4%	70324	19.4%
2015	Recent	3085	277800	76.6%	248261	80.9%	33455	54.7%
	Past	865	36138	10.0%	24108	7.9%	12810	20.9%
	New	1214	48726	13.4%	34557	11.3%	14917	24.4%
	Total		362664		306926	84.6%	61182	16.9%

HTT: home treatment team

*Excluding concurrent inpatient and HTT days

Figure 1. Crisis (home treatment team [HTT] and inpatient care) person-days receipt over 12 month period, for patients with current/recent mental healthcare at the commencement of that period, displaying total days and days by quarters of the follow-up period.

Figure 2. Crisis service use for the 12-month period following 16th March in patients with current/recent mental healthcare at that point, displayed by year and presence or not of certain previously recorded diagnoses¹

¹ SMI – serious mental illness; SUD – alcohol/substance use disorder; PD – personality disorder. Diagnoses based on any received up to 15th March of the year in question.

Table 2. Service contact changes 31 days before and after March 16th in current/recent patients.²

Year	Number	<i>Person-days as inpatient</i>			
		1 month before Mar 16th	1 month after Mar 16th	Difference	
2020	50487	23450	17313	-6137	-26.2%
2019	46606	24762	23455	-1307	-5.3%
2018	44362	24085	22998	-1087	-4.5%
2017	43283	24302	23774	-528	-2.2%
2016	42367	26008	25033	-975	-3.7%
2015	40972	26023	25093	-930	-3.6%

<i>Number of face-to-face contacts</i>					
2020	50487	36218	12803	-23415	-64.7%
2019	46606	42899	34180	-8719	-20.3%
2018	44362	41014	31269	-9745	-23.8%
2017	43283	41975	33877	-8098	-19.3%
2016	42367	39634	34183	-5451	-13.8%
2015	40972	36100	33105	-2995	-8.3%

<i>Number of virtual contacts</i>					
2020	50487	11622	25207	13585	116.9%
2019	46606	11351	9581	-1770	-15.6%
2018	44362	10026	7825	-2201	-22.0%
2017	43283	9269	7765	-1504	-16.2%
2016	42367	8277	7313	-964	-11.6%
2015	40972	6758	6556	-202	-3.0%

<i>Total number of contacts</i>					
2020	50487	47840	38010	-9830	-20.5%
2019	46606	54250	43761	-10489	-19.3%
2018	44362	51040	39094	-11946	-23.4%
2017	43283	51244	41642	-9602	-18.7%
2016	42367	47911	41496	-6415	-13.4%
2015	40972	42858	39661	-3197	-7.5%

² Those with any active SLaM referral within 3 calendar months prior to that date.

Figure 3. Proportional service contact changes³ from 31 days before to 31 days from/after March 16th in current/recent patients, displayed by year and presence of previous recorded diagnosis⁴.

³ Changes are expressed as percentages of the activity level in the 31 days prior to the 16th March index date

⁴ SMI – serious mental illness; SUD – alcohol/substance use disorder; PD – personality disorder. Diagnoses based on any received up to 15th March of the year in question.