Epidemic Trend Analysis of SARS-CoV-2 in SAARC Countries Using Modified SIR (M-SIR) Predictive Model

Samrat Kumar Dey¹, Md. Mahbubur Rahman^{*2}, Kabid Hassan Shibly³, Umme Raihan Siddiqi,⁴ and Arpita Howlader⁵

¹Department of Computer Science and Engineering (CSE), Dhaka International University (DIU), Dhaka-1205, Bangladesh, email: sopnil.samrat@gmail.com

²Department of Computer Science and Engineering (CSE), Military Institute of Science and Technology (MIST), Mirpur Cantonment, Dhaka-1216, Bangladesh, email: mahbub.rahman.cse@gmail.com

³Department of Computer Science and Engineering (CSE), Dhaka International University (DIU), Dhaka-1205, Bangladesh, email: khshibly00@gmail.com

⁴Department of Physiology, Shaheed Suhrawardy Medical College (ShSMC), Dhaka-1207, Bangladesh, email: u.raihan.siddiqi@gmail.com

⁵Department of Computer and Communication Engineering (CCE), Patuakhali Science and Technology University (PSTU), Dumki-8602, Patuakhali, Bangladesh, email: arpita.cce.pstu@gmail.com

*** Corresponding Author:** Dr. Md. Mahbubur Rahman,
Present address: Department of Computer Science and Engineering (CSE),
Military Institute of Science and Technology (MIST), Mirpur Cantonment,
Dhaka-1216, Bangladesh.
Email: mahbub.rahman.cse@gmail.com

Word count: 2097 (Excluding Abstract, Figures, Tables, and Reference)

Abstract

A novel coronavirus causing the severe and fatal respiratory syndrome was identified in China, is now producing outbreaks in more than two hundred countries around the world, and became pandemic by the time. In this article, a modified version of the well known mathematical epidemic model Susceptible (S)- Infected (I)- Recovered (R) is used to analyze the epidemic's course of COVID-19 in eight different countries of the South Asian Association for Regional Cooperation (SAARC). To achieve this goal, the parameters of the SIR model are identified by using publicly available data for the corresponding countries: Afghanistan, Bangladesh, Bhutan, India, the Maldives, Nepal, Pakistan and Sri Lanka. Based on the prediction model we estimated the epidemic trend of COVID-19 outbreak in SAARC countries for 20 days, 90 days, and 180 days respectively. An SML (short-mid-long) term prediction model has been designed to understand the early dynamics of COVID-19 Epidemic in the south-east Asian region. The maximum and minimum basic reproduction number ($R_0 = 1.33$ and 1.07) for SAARC countries are predicted to be in Pakistan and Bhutan. We equate simulation results with real data in the SAARC countries on the COVID-19 outbreak, and model potential countermeasure implementation scenarios. Our results should provide policymakers with a method for evaluating the impacts of possible interventions, including lockdown and social distancing, as well as testing and contact tracking.

Keywords

SARS-CoV-2; COVID-19; epidemics; SAARC; modified susceptible-infected-recovered; prediction model;

1. Introduction

A novel coronavirus (SARS-CoV-2), called COVID-19, caused an outbreak in the city of Wuhan, Hubei Province, China that is linked to the Huanan Seafood Wholesale Market^{1,2} in late December 2019. Till now, more than 200 countries around the world have been infected by a novel coronavirus. As of May 30, 2020, according to the World Health Organization (WHO), globally 5817385 confirmed cases have reported with a death count of 362705³. Among them, the south-east Asia region has confirmed 4.2% of cases while the case fatality ratio is 2.86%.

The very first COVID-19 patient was treated for being coronavirus positive in Wuhan, China, at the beginning of December 2019. In the south-east Asia region, the very first coronavirus infected patient was detected in Pakistan on February 26, 2020⁴. On 30 January 30, 2020, WHO officially declared this outbreak of COVID-19 as global pandemic⁵. To mitigate the spread of COVID-19, the affected countries of the world have taken various measures, including citywide lockdown, social distancing, traffic halt, community management, and information on health education knowledge. More importantly, the outbreak of COVID-19 possessed a massive threat to global health and economics all over the world. One of the significant feature of novel coronavirus unlike other infectious diseases like SARS (Severe Acute Respiratory Syndrome), and MERS (Middle East Respesoritory Syndrom), it causes asymptomatic infections (symptoms are very mild)^{6,7, 8}. In this circumstances, the rate of transmission among a large number of people can increase within no time. According to the latest World Health Organization survey, only 87.9% of COVID-19 patients have a fever, and 67.7% have dry cough⁹. Therefore, this is highly crucial to estimate the intensity of the COVID-19 epidemic and predict the time course, peak time, total duration, and so on. Our study focuses on the COVID-19 case prediction using a modified SIR model (M-SIR) in the countries of the South Asian Association for Regional Cooperation (SAARC). SAARC is considered as an intergovernmental organization and geopolitical union of states in South Asia. Its member states are Afghanistan, Bangladesh, Bhutan, India, the Maldives, Nepal, Pakistan and Sri Lanka. Our aim is to develop a prediction model for the SAARC countries to understand the epidemiological trend of novel coronavirus outbreak in these countries. Here we explore a modified version based on the Susceptible (S)- Infected (I)- Recovered (R) epidemic model to predict the short term (20 days), mid-term (90 days) and long term (180 days) evaluation of COVID-19 situation in these countries of SAARC regions.

2. Methodology

2.1 Dataset Analysis

For this paper, we have used datasets from various sources for our analysis and building the model. We have used four different sources of the dataset including, COVID-19 Data Repository by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University

(January - May 2020)¹⁰, COVID-19 Dataset (January - May 2020)¹¹, COVID-19 Open Research Dataset (CORD-19)¹² and population by country 2020 dataset¹³. **Table 1** provides insight on each dataset and their respective data files with their column description.

Table 1: Tabular representation of different data sources of COVID-19

Dataset	Description	Columns
Novel Coronavirus 2019 Dataset		
COVID19_line_list_data.csv	This file is an aggregated version of the Novel Coronavirus dataset collected by Johns Hopkins University.	Id, case_in_country, reporting date, summary, location, country, gender, age, symptom_onset, If_onset_approximate, hosp_visit_date, exposure_start, exposure_end, visiting Wuhan, from Wuhan, death, recovered, symptom, source
covid_19_data.csv	Daily level information on the number of COVID-19 affected cases across the globe	Sno, Observation Date, Province/State, Country/Region, Last Update, Confirmed, Deaths, Recovered
time_series_covid_19_confirmed.csv	Time series data on the number of confirmed cases	Province/State, Country/Region, Lat, Long, 1/22/20...5/6/20
time_series_covid_19_deaths.csv	Time series data on the number of death cases	Province/State, Country/Region, Lat, Long, 1/22/20...5/6/20
time_series_covid_19_recovered.csv	Time series data on the number of recovered cases	Province/State, Country/Region, Lat, Long, 1/22/20...5/6/20
COVID-19 (nCoV-19) Coronavirus Spread Dataset		
covid_19_clean_complete.csv	The file contains the cumulative count of confirmed, death and recovered cases of COVID-19 from different countries from January 2020	Province/State, Country/Region, Lat, Long, Date, Confirmed, Deaths, Recovered
COVID-19 Open Research Dataset		
metadata.csv	This dataset contains metadata for 59k articles on COVID-19	Cord_uid, sha, source_x, title, doi, pmcid, license, abstract, publish_time, authors, journal
Population by country 2020 dataset		
population_by_country_2020.csv	This dataset contains the information from 235 countries along with their population till 2020.	Country (or dependency), population (2020), Yearly Change, Net Change, Density (P/Km ²), Land Area (Km ²), Migrants (net), Fert. Rate, Med. Age, Urban Pop %, World Share

We also developed a custom dataset to develop and evaluate our model. The following **Table 2** is developed based on collected data from different sources.

Table 2: Columns description of custom build COVID-19 dataset

Field Name	Field Data Description
Sno.	Serial number
Province / State	Province or state of the observation
Country/ Region	Country of observation
Date	Date and time of the observation in MM/DD/YYYY HH:MM: SS
Confirmed	Number of confirmed cases
Deaths	Number of deaths
Recovered	Number of recovered cases
Active	Number of active cases
1/22/20	First reporting date
5/18/2020	Latest reporting date

2.2 Modified SIR (M-SIR) Model

Predictive mathematical disease models are important for understanding the trajectory of the outbreak and for preparing successful response strategies. One commonly used model is the human-to-human transmission SIR model, which defines people's flow through three mutually exclusive stages of infection: Susceptible (S), Infected (I), and Recovered (R). Most epidemic models are based on dividing the population into a small number of sections. Each person is identical in terms of their status with the considered disease. The SIR model based on three sections: Susceptible (S) is the class for those who are susceptible to infection. This can include passive immune systems as soon as they lose their immunity. In the Infected (I) class, the parasite level within the host is large enough, and there is a possibility of spreading the infection to other susceptible people. The Recovered (R) class includes all infected and recovered individuals. This epidemiological model captures the dynamics of acute infections that, after recovery, confer lifelong immunity. In general, the overall size of the population is considered constant $N=S+I+R$. The two cases should be examined and characterized by the inclusion or exclusion of demographic factors. Let's assume in the SIR model; there is a natural host lifespan of $1/\mu$ years. Then the rate at which individuals in an epidemiological compartment suffer from natural mortality is given by μ . It is important to emphasize that this factor is independent of the

disease and should not reflect the pathogenicity of the infectious agent. Diachronically, it can be said,

$$\frac{dS}{dt} + \frac{dI}{dt} + \frac{dR}{dt} = 0 \quad (1)$$

So, the SIR model can be defined as,

$$\begin{cases} \frac{dS}{dt} = \mu - \beta SI - \mu S \\ \frac{dI}{dt} = \beta SI - \gamma I - \mu I \\ \frac{dR}{dt} = \gamma I - \mu R \end{cases} \quad (2)$$

Here, the initial conditions $S(0) > 0$, $I(0) \geq 0$ and $R(0) \geq 0$. It is important to enter the expression of basic reproduction number R_0 for this model. The basic reproduction number R_0 is the parameter that estimates whether a disease has spread to the population or not. If the estimated $R_0 < 1$, we can assume that the disease will die out, and if $R_0 = 1$, the disease remains in the system and is stable. But if $R_0 > 1$, the disease will spread and cause an outbreak. The higher the value of R_0 , the more difficult it is to control.

2.3 Proposed Model and Algorithm

In this exploration, we have used a susceptible (S)-infected (I)-recovered (R) epidemic model. In a general SIR model, transmission rate (β) and recovery rate (γ) are considered as two time-invariant variables. Moreover, several research studies have shown that a SIR model works much better in presenting the information contained in the confirmed case data than an SEIR model¹⁴. Therefore we have developed a model that can dynamically adjust the crucial parameters while working on time-varying data, which is also treated as a modified SIR (M-SIR) model¹⁵. However, in a basic SIR model, the reproduction number (R_0) is a simple division of transmission and recovery rates, as shown in Equation (3).

$$R_0 = \frac{\beta}{\gamma + \mu} \quad (3)$$

For building the model, we modified the primary reproduction number, R_0 with respect to time (t). Equation (4) represents the changes that happened depending on time (t).

(4)

We also considered detectable () and non-detectable () infected persons for building our model effectively, as shown in Equation (5).

(5)

In general, detectable persons contain a lower transmission rate than non-detectable persons. Therefore we calculated the transmission rate (β) and recovery rate (γ) for each country of the SAARC region. For instance, Bangladesh has 165 million people¹⁶, and the first confirmed case reported in the country on March 7, 2020. As of May 30, 2020, a total of 42,844 COVID-19 infected people detected in the country³. Depending on the modified SIR model, Bangladesh contains a transmission rate of 0.63 with a recovery rate rate of 0.49. The initial reproduction number for the country is 1.27. Based on available data sources of every SAARC country, we have calculated these parameters for the prediction model, which is shown in **Table 3**. **Algorithm 1** represents the working procedure of our proposed (M-SIR) prediction model.

Figure 1: Modified SIR (M-SIR) model with the time-varying transmission

Algorithm 1: M-SIR prediction model

1. Evaluate the transmission rate , recovery rate within the dataset time period, T.
2. Train multiple regression or ridge regression
4. Estimate number of an infected person after the dataset time period, T
5. Estimate number of a recovered person after the dataset time period, T
6. **while** dataset time period, T respective time, t estimated time for prediction **do**

7. Estimate predicted transmission rate and recovery rate
 8. Predict the number of affected persons, the number of recovered persons, the number of active cases.
 9. end while
-

3. Results

We performed our simulations and tabulated the predicted results (short term, midterm, and long term) with all countries from the SAARC region. We also explored 3D parameter space using gradient descent to minimize the error. However, a lag parameter is also used in this experiment to reduce the gap of first confirmed cases of SAARC countries' pandemic situation. The following **Table 3** represents the transmission rate (β), recovery rate (γ), and the reproduction numbers (R_0) of all countries in the SAARC regions. Bangladesh contains the highest transmission rate of ($\beta = \sim 0.63$) in the SAARC region, whereas India shows a much lower transmission rate of only ~ 0.51 comparing with other countries. Also, data analysis visualization tool have been employed (**Figure 2**) to understand the current (as of May 30, 2020) scenario of the countries (Afghanistan, Bangladesh, Bhutan, India, Maldives, Nepal, Pakistan, and, Sri Lanka) in terms of their confirmed and death cases ratio.

Table 3. Transmission rate, Recovery rate, and Reproduction number of SAARC countries.

Countries	Transmission rate (β)	Recovery rate (γ)	Reproduction Number (R_0)
Afghanistan	0.5997691	0.4889147	1.22673566575
Bangladesh	0.6286615	0.4932285	1.2745847006
Bhutan	0.5334682	0.4982385	1.07070850607
India	0.5060977	0.3955793	1.27938367857
Maldives	0.5774223	0.4920300	1.17355100299
Nepal	0.5406592	0.4954823	1.09117762632
Pakistan	0.5254487	0.3947732	1.33101411139
Sri Lanka	0.5595021	0.4933888	1.13399838018

Figure 2: This graphical representation provides a comprehensive overview of the total number of confirmed and death cases in SAARC regions. Country-wise COVID-19 confirmed and death cases along with the number of cases per million illustrated in 2(A), 2(B), 2(C), and 2(D) till May 30, 2020. Based on the illustration on 2(A) and 2(C), India has confirmed the highest number of infected cases (173763) along with the most number of death cases (5164) in the region.

Surprisingly the slow growth of infections in the South-Asian region could be a result of a lower number of testing and testing strategy¹⁷. Initially, the testing was limited to specific individuals who have come from high-risk countries. Even their immediate contacts were also ignored primarily. The initial growth rate of COVID-19 infection in the SAARC region is comparatively lower than countries like the US, France, Germany, Spain, China, Italy, and so on. However, based on the current scenario in the SAARC region (as of May 30, 2020), India confirmed the highest number of COVID-19 reported cases. A short term model prediction for the next 20 days (till June 19, 2020) is illustrated in **Figure 3** for all the countries of SAARC regions.

Similarly, we also predicted the epidemic curve of SAARC regions for the midterm (90 days) and long term (180 days) COVID-19 cases. After predicting the 20 days of the epidemic, we noticed that India, Bangladesh, Sri Lanka, and Pakistan are increasing gradually by minimizing the active cases, and their recovery rate is also rising over time. However, in this short period, the confirmed cases of these countries are also increasing till June 19, 2020.

Figure 3: Short term prediction model for the next 20 days using M-SIR approach. This prediction provides a depiction of all SAARC countries evaluation of COVID-19 till Mid June. All the countries prediction curve are denoted here from (A-H) respectively.

4. Discussion

The US, Italy, Spain, Germany, China are the top most affected countries in the world due to the pandemic of COVID-19. Though the novel coronavirus originated and started to spread from China but, countries from the South-Asian region are also infected faster than other countries globally. The overall population density of the countries in the SAARC regions is also too high. According to **Table 4**, we can observe the various parameters (population, land area, density, and world share) and their impact on analyzing the spread of novel coronavirus in these regions.

Table 4. Population, total area and density of SAARC countries with their individual world share¹³

Countries	Population	Land Area (Km ²)	Density (P/Km ²)	World Share %
Afghanistan	38742911	652860	60	0.50
Bangladesh	164354176	130170	1265	2.11
Bhutan	769867	38117	20	0.01
India	1377233523	2973190	464	17.70
Maldives	538558	300	1802	0.01
Nepal	29027347	143350	203	0.37
Pakistan	219992900	770880	287	2.83
Sri Lanka	21395196	62710	341	0.27

We further extended our prediction model for the next 90 days and 180 days, respectively. Besides, the total number of the population of a country is also parameterized because the total population of a country cannot be infected. Therefore, if we consider the total population as infected, then the probable number of infected persons remains unknown. Based on the prediction model, we tabulated the model predicted results of all SAARC countries for short term, mid term and long term prediction respectively (**Table 5**).

Table 5. Predicted data based on M-SIR model for the countries of SAARC region

Countries		Short term prediction (20 days)	Mid-term prediction (90 days)	Long term prediction (180 days)	Remarks
Afghanistan	Confirmed	35286	297264	1092714	<ul style="list-style-type: none"> Confirmed, recovery and active cases will increase at each predictive term In the long-term forecast, the number of death cases will decrease comparing with the mid-term forecast.
	Deaths	652	5498	3521	
	Recovered	2555	8521	13416	
	Active	32078	283245	1075775	
Bangladesh	Confirmed	100037	451840	867700	<ul style="list-style-type: none"> significant rise will be observed in confirmed and active cases The number of recovered cases after mid-term prediction will not increase much in the long-term prediction.
	Deaths	1346	6080	11676	
	Recovered	17724	29418	30476	
	Active	80966	416341	825547	
Bhutan	Confirmed	61	372	1683	<ul style="list-style-type: none"> Large number of confirmed cases but not enough recovery. The number of deaths will remain zero.
	Deaths	0	0	0	
	Recovered	11	32	79	
	Active	50	340	1603	
India	Confirmed	378657	1717831	3709965	<ul style="list-style-type: none"> The highest number of confirmed, deaths, recovered and active cases will be observed in the region. It will show significant improvement in recovery cases
	Deaths	10648	48307	104329	
	Recovered	217681	1269646	2609589	
	Active	150327	399877	996047	
Maldives	Confirmed	2873	5970	7081	<ul style="list-style-type: none"> Both the confirmed and recovered cases will increase with a similar trend. Death cases will also fall to zero as active cases.
	Deaths	14	0	0	
	Recovered	1139	5970	7081	
	Active	1719	0	0	
Nepal	Confirmed	6295	185528	1089821	Confirmed cases will notably increase.
	Deaths	0	0	2955	

	Recovered	1783	99292	658568	<ul style="list-style-type: none"> However, the death cases will jump to 2955 from zero in the long-term prediction.
	Active	4512	86236	428297	
Pakistan	Confirmed	118775	332528	497800	<ul style="list-style-type: none"> Confirmed, deaths, recovered and active cases will increase gradually. recovered cases will show almost double in the long-term prediction.
	Deaths	2534	7096	10623	
	Recovered	49434	254510	487176	
	Active	66806	70921	588234	
Sri Lanka	Confirmed	2496	8832	23151	<ul style="list-style-type: none"> In the mid-term and long-term forecast, active cases will become zero
	Deaths	0	40	106	
	Recovered	1840	8791	23045	
	Active	655	0	0	

A midterm (90 Days) prediction model is further assessed until August 31, 2020, and the nature of the epidemic curve has been depicted in **Figure 4**. For all the countries of the SAARC regions, we have predicted the total number of confirmed, death, recovery, and active cases.

Figure 4: Based on the M-SIR prediction model, prediction statistics for the upcoming 90 days for all the countries in the SAARC regions has depicted here. Different predicted case scenarios (confirmed, death, recovered, and active) till August 31, 2020, helps to understand the epidemiological nature of COVID-19 in the south Asian region. The predicted model indicates that in the next 90 days prediction curve will increase sharply for India (in terms of confirmed and death cases). Moreover, India will also show a substantial increase in its recovery rate. However, the number of active cases will increase in Pakistan till August 31, 2020.

Lastly, a long term prediction model for the next 180 days (till November 30 2020) situation has been presented in **Figure 5**. According to the prediction model, active cases will fall to zero at the end of June in Sri Lanka, and at the beginning of August in India (**Figure 5**). However, Bangladesh, Maldives, and Pakistan will take more few months to reduce its active cases. On the other hand, countries like Afghanistan, Nepal and Bhutan will show a steep increase in their active cases.

Figure 5: Long term prediction model for the next 180 days using M-SIR approach. This prediction provides a depiction of all SAARC countries evaluation of COVID-19 till Mid November. All the countries prediction curve are denoted here from (A-H) respectively.

5. Conclusion

The COVID-19 epidemic has brought unprecedented health concern for the community of all around the globe. The pandemic data of COVID-19 of SAARC countries have analyzed by using the M-SIR model to predict the confirmed, deaths, recovered and active cases with respect to the time in days. We predicted the epidemic data in the basis of short term, midterm and long term situation. To the best of our knowledge, this study is the very first COVID-19 prediction model which focused on the countries of SAARC regions. This epidemic modelling can be a helpful tool for estimating and predicting the scale and time course of COVID-19, evaluation of the effectiveness of public health interventions, and information on public health policies in SAARC countries. In future machine learning tools can be further used to identify and optimize time profile for the confinement.

Author Contribution

All authors conceptualized and designed the study. SKD, KHS and MR had the idea for and designed the study and had full access to all the data in the study and take the responsibility for the predictive model and accuracy of the epidemic trend analysis with their visualization. URS and AH and contributed to the writing of the article. MR contributed to the critical revision of the report. All the visualization and data presentation methods developed by SKD, KHS and MR. All authors contributed to data acquisition, data analysis, and reviewed and approved the final version.

Funding

None

Declaration of Interests

The authors declare that there are no conflicts of interest.

Ethical Approval

Not required

References

- [1] Wu, Joseph T., Kathy Leung, and Gabriel M. Leung. "Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study." *The Lancet* 395, no. 10225 (2020): 689-697.
- [2] Bogoch, Isaac I., Alexander Watts, Andrea Thomas-Bachli, Carmen Huber, Moritz UG Kraemer, and Kamran Khan. "Pneumonia of Unknown Etiology in Wuhan, China: Potential for International Spread Via Commercial Air Travel." *Journal of Travel Medicine* (2020).
- [3] Coronavirus disease (COVID-19) Situation Report – 131. World Health Organization (WHO). https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200530-covid-19-sitrep-131.pdf?sfvrsn=d31ba4b3_2. Accessed 30 May 2020.
- [4] Coronavirus disease (COVID-19) Situation Report – 38. World Health Organization (WHO). https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200227-sitrep-38-covid-19.pdf?sfvrsn=2db7a09b_4. Accessed 30 May 2020.
- [5] WHO Timeline - COVID-19. World Health Organization (WHO). <https://www.who.int/news-room/detail/27-04-2020-who-timeline---covid-19>. Accessed 17 May 2020.
- [6] Guan, W.J., Ni, Z.Y., Hu, Y., Liang, W.H., Ou, CQ, He, J.X., Liu, L., Shan, H., Lei, C.L., Hui, D.S. and Du, B., 2020. Clinical characteristics of coronavirus disease 2019 in China. *New England journal of medicine*, 382(18), pp.1708-1720.
- [7] Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X. and Cheng, Z., 2020. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The lancet*, 395(10223), pp.497-506.
- [8] Mahase, E., 2020. Coronavirus: covid-19 has killed more people than SARS and MERS combined, despite lower case fatality rate.
- [9] Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19). World Health Organization (WHO). <https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf>. Accessed 12 May 2020.
- [10] COVID-19 Data Repository by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (Available from: <https://github.com/CSSEGISandData/COVID-19>)
- [11] COVID-19 Dataset: Number of Confirmed, Death and Recovered cases every day across the globe, (Available from: <https://www.kaggle.com/imdevskp/corona-virus-report>). Accessed 30 May 2020.

- [12] COVID-19 Open Research Dataset Challenge (CORD-19), An AI challenge with AI2, CZI, MSR, Georgetown, NIH & The White House, (Available from: <https://www.kaggle.com/allen-institute-for-ai/CORD-19-research-challenge>). Accessed 30 May 2020.
- [13] T. N. Prabhu "Population by Country - 2020, Version 3." Retrieved June 13, 2020 from <https://www.kaggle.com/tanuprabhu/population-by-country-2020>. Accessed 30 May 2020.
- [14] Roda, W.C., Varughese, M.B., Han, D. and Li, M.Y., 2020. Why is it difficult to accurately predict the COVID-19 epidemic?. Infectious Disease Modelling.
- [15] I. Nesteruk, "Statistics based predictions of coronavirus 2019-ncov spreading in mainland china," MedRxiv, 2020.
- [16] Report on Bangladesh Sample Vital Statistics 2018. BANGLADESH BUREAU OF STATISTICS.
http://bbs.portal.gov.bd/sites/default/files/files/bbs.portal.gov.bd/page/6a40a397_6ef7_48a3_80b3_78b8d1223e3f/SVRS_Report_2018_29-05-2019%28Final%29.pdf. Accessed 14 May 2020.
- [17] Gupta, R. and Pal, S.K., 2020. Trend Analysis and Forecasting of COVID-19 outbreak in India. medRxiv.