

1 **Modeling the Stability of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-**  
2 **2) on Skin, Currency, and Clothing**

3

4 **Authors:** David E. Harbourt\*<sup>1</sup>, Andrew D. Haddow<sup>2</sup>, Ashley E. Piper<sup>3</sup>, Holly Bloomfield<sup>4</sup>,  
5 Brian J. Kearney<sup>4</sup>, David Fetterer<sup>5</sup>, Kathleen Gibson<sup>4</sup> and Timothy Minogue<sup>6</sup>

6

7 **Affiliations:**

8

9 <sup>1</sup>Biosafety Division, United States Army Medical Research Institute of Infectious Diseases, 1425  
10 Porter Street, Ft. Detrick Maryland, 21702

11 <sup>2</sup>General Dynamics Health Solutions in support of USAMRIID, 1425 Porter Street, Ft. Detrick,  
12 Maryland, 21702

13 <sup>3</sup>Oak Ridge Institute of Science and Education, 1425 Porter Street, Ft. Detrick, Maryland, 21702

14 <sup>4</sup>Core Laboratory Services Directorate, United States Army Medical Research Institute of  
15 Infectious Diseases, 1425 Porter Street, Ft. Detrick Maryland, 21702

16 <sup>5</sup>ICON Global Public Health Solutions, 1425 Porter Street, Ft. Detrick, Maryland, 21702

17 <sup>6</sup>Diagnostic Systems Division, United States Army Medical Research Institute of Infectious  
18 Diseases, 1425 Porter Street, Ft. Detrick Maryland, 21702

19

20 \*Correspondence to: David.e.harbourt.civ@mail.mil

21

22

23

24

25

26

27

28

29

30

31

## Abstract

32  
33  
34 A new coronavirus (SARS-CoV-2) emerged in the winter of 2019 in Wuhan, China, and rapidly  
35 spread around the world. The extent and efficiency of SARS-CoV-2 pandemic is far greater than  
36 previous coronaviruses that emerged in the 21<sup>st</sup> Century. Here, we modeled stability of SARS-  
37 CoV-2 on skin, paper currency, and clothing to determine if these surfaces may factor in the  
38 fomite transmission dynamics of SARS-CoV-2. Skin, currency, and clothing samples were  
39 exposed to SARS-CoV-2 under laboratory conditions and incubated at three different  
40 temperatures ( $4^{\circ}\text{C} \pm 2^{\circ}\text{C}$ ,  $22^{\circ}\text{C} \pm 2^{\circ}\text{C}$ , and  $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ ). Stability was evaluated at 0 hours (h), 4  
41 h, 8 h, 24 h, 72 h, 96 h, 7 days, and 14 days post-exposure. SARS-CoV-2 was shown to be stable  
42 on skin through the duration of the experiment at  $4^{\circ}\text{C}$  (14 days). Virus remained stable on skin  
43 for at least 96 h at  $22^{\circ}\text{C}$  and for at least 8h at  $37^{\circ}\text{C}$ . There were minimal differences between the  
44 tested currency samples. The virus remained stable on the \$1 U.S.A. Bank Note for at least 96 h  
45 at  $4^{\circ}\text{C}$  while viable virus was not detected on the \$20 U.S.A. Bank Note samples beyond 72 h.  
46 The virus remained stable on both Bank Notes for at least 8 h at  $22^{\circ}\text{C}$  and 4 h at  $37^{\circ}\text{C}$ . Clothing  
47 samples were similar in stability to the currency with the virus being detected for at least 96 h at  
48  $4^{\circ}\text{C}$  and at least 4 h at  $22^{\circ}\text{C}$ . No viable virus was detected on clothing samples at  $37^{\circ}\text{C}$  after  
49 initial exposure. This study confirms the inverse relationship between virus stability and  
50 temperature. Furthermore, virus stability on skin demonstrates the need for continued hand  
51 hygiene practices to minimize fomite transmission both in the general population as well as  
52 workplaces where close contact is common.

53

54

55 **Background**

56

57 The emergence of SARS-CoV-2 represents the third major outbreak of a new human coronavirus  
58 disease over the past twenty years. This novel coronavirus (SARS-CoV-2) that initially emerged  
59 in Wuhan, China in late 2019 resulted in a global pandemic that, as of April 22, 2020, has  
60 officially resulted in more than 2.5 million cases and 175,000 deaths (1-3). The rapid and  
61 extensive spread of the virus could be indicative of both aerosol and fomite transmission which  
62 has been seen in previous coronavirus outbreaks (4). Previous studies have shown that SARS-  
63 CoV-2 is stable at room temperature on stainless steel for approximately 24 hours (h) and on  
64 cardboard for up to three days (5-6). Additional studies have shown an inverse relationship  
65 between surface temperature and stability which is consistent with previous reporting on stability  
66 of human coronaviruses (5).

67

68 While some aspects of SARS-CoV-2 have been reported, there have been only limited  
69 investigations into stability on skin specimens and paper currency (7). Despite limited evidence,  
70 some countries have taken measures to limit the spread of the virus by either burning or  
71 disinfecting paper currency or discouraging the use of cash during transactions (8). It would be  
72 expected that the stability of SARS-CoV-2 on currency would be similar to cardboard given that  
73 both surfaces are porous, but the effect of ink and toner on the virus remains unknown (6).  
74 Furthermore, it was not known how long SARS-CoV-2 could remain viable on human or animal  
75 skin as no similar studies had been performed to date. Handwashing and hand hygiene have been  
76 a key part of mitigation efforts, but fomite transmission likely remains a contributing factor to

77 the speed and extent of the pandemic (9-11). Herein, we model the stability of SARS-CoV-2  
78 across animal skin, paper currency, and clothing.

79

## 80 **Methods and Materials**

81

82 **Virus isolate.** We utilized the USA-WA1/2020 strain of SARS-CoV-2 isolated from a human  
83 patient in Washington State, USA, in January 2020 (GenBank accession no. MN985325.1). This  
84 isolate was selected due to its use in related studies (6). A working virus stock was prepared by  
85 adding virus to Grivet (*Chlorocebus aethiops*) Vero 76 kidney cells (ATCC, Manassas, VA;  
86 #CRL-1587) at a multiplicity of infection (MOI) of 0.01. Cells were incubated for 1 h for virus  
87 adsorption and maintained in Eagles Minimal Essential Media (EMEM) with 10% fetal bovine  
88 serum at 5% CO<sub>2</sub>. The cell supernatant was harvested 50 h post-inoculation. The supernatant was  
89 clarified at 10,000 x g for 10 minutes at 4°C and stored at -70°C until use.

90

91 **Virus surface stability.** We evaluated the surface stability of SARS-CoV-2 on four common  
92 surfaces (approx. 6.3 mm<sup>2</sup>). These included: swine skin (*Sus scrofa*) with the hair removed  
93 (acquired from a local butcher); uncirculated United States of America \$1 and \$20 Federal  
94 Reserve notes comprised of 25% linen and 75% cotton with red and blue security fibers (United  
95 States Secret Service, Washington, DC, USA); and unused scrub fabric consisting of 35% cotton  
96 and 65% polyester (Labforce, Swedesboro, NJ, USA). Using a pipette, we deposited 50 µL of  
97 virus onto the surface of each material in triplicate. Individual groups of samples were incubated  
98 for 0 h, 4 h, 8 h, 24 h, 72 h, 96 h, 7 days, and 14 days post-exposure across three temperatures;  
99 4°C ± 2°C, 22°C ± 2°C, and 37°C ± 2°C at a relative humidity of 40-50%. Following incubation,

100 samples were transferred to 2-mL CryoS sure tubes (Caplugs Evergreen, Buffalo, NY, USA)  
101 containing 1 mL of media minimum essential media (MEM: Corning, catalog 10-010-CM),  
102 supplemented with 4.0 µg/mL gentamicin (GIBCO, Carlsbad, CA, USA), 2% Penicillin (Sigma-  
103 Aldrich, St. Louis, MO, USA), 2% streptomycin (Sigma-Aldrich), and 2.5 mg/mL of  
104 amphotericin B (Sigma-Aldrich, St. Louis, MO, USA) using featherweight forceps (BioQuip  
105 Products, Inc, Rancho Doming o, CA, USA) to reduce the potential for sample damage. Forceps  
106 were disinfected using 5% Microchem Plus™ followed by 70% ETOH between samples.  
107 Samples were stored at -80°C prior to virus quantification.

108  
109 **Detection and quantification of infectious virus.** Confluent cultures (90-95% confluency) of  
110 ATCC Vero 76 cells in 6-well plates were utilized for all assays. Samples were thawed at  
111 ambient temperature and diluted by performing a series of 1:10 dilutions in MEM + 5% Heat  
112 Inactivated (HI) FBS + 2% penicillin, 2% streptomycin + 0.5% fungizone (MEM Complete). All  
113 samples were assayed as undiluted and up to three additional ten-fold dilutions. Media was  
114 removed from plates and cells were infected with 100 µL of sample in triplicate. Cells were  
115 incubated at 37°C and 5% CO<sub>2</sub> for one hour, with rocking approximately every 15 minutes.  
116 Following incubation, media-agarose overlay (2mL of a 1:1 mixture of 1.0% agarose and 2X  
117 EBME + 10% HI FBS + 2% penicillin, 2% streptomycin + 1% fungizone (2X EBME  
118 Complete)) was added to each well. Once overlay solidified, plates were incubated at 37°C and  
119 5% CO<sub>2</sub> for 48 h ± 4 h. Following incubation, a second media agarose overlay containing 4%  
120 neutral red in a 1:1 mixture of 1.0% agarose and 2X EBME complete was added to each well.  
121 Once overlay solidified, plates were incubated at 37°C and 5% CO<sub>2</sub> overnight. Following  
122 incubation, plaques were counted and the virus yield (plaque forming unit, PFU/mL) for each

123 sample calculated, with a lower limit of quantification (LLOQ) of 2.0 log<sub>10</sub> PFU/mL. The limit  
124 of detection (LOD) for virus isolation attempts was 0.1 log<sub>10</sub> PFU/mL.

125

126 **Statistical analyses.** The half-life was estimated by fitting, to each trial, a log-linear Poisson  
127 regression model of the form

$$128 \quad \log(\text{PFU}) = m \cdot \text{time} + b + \log(\text{Dilution} * \text{Volume Sampled})$$

129 such that log( Dilution \* Volume Sampled ) was the offset. The half-life was estimated as  
130  $-\log(2)/m$ . This analysis was implemented in SAS/PROC GENMOD. The log of the half-lives  
131 so estimated were entered into a two-way ANOVA, as implemented in SAS/PROC MIXED.  
132 The mixed model procedure was used to allow for a heterogeneous variance structure (11).  
133 Denominator degrees of freedom were estimated by Satterthwaite's method, and LS-mean  
134 differences between temperatures and surfaces were evaluated. For the purpose of comparing  
135 half-lives, one outlier was removed from the clothing surface at 22°C, pursuant to a Dixon gap  
136 test applied to the log transformed half-life. Analysis was implemented in SAS version 9.4 (SAS  
137 Institute, Cary, NC), with the exception of the gap test, which was implemented in R (12)  
138 package outliers version 0.14. No adjustment for multiplicity has been applied to the reported p-  
139 values.

140

## 141 **Results**

142

143 SARS-CoV-2 remained stable on skin at 4°C for the duration of the experiment (Fig 1 and Fig  
144 2). The virus exhibited similar initial decay profiles at 4°C across all surfaces. Initially, the virus  
145 exhibited a loss of 1-2 log<sub>10</sub> PFU in the first 8 h across all surfaces. However, after 8 h, the virus

146 appeared to stabilize to varying degrees for the remainder of the experiment. Clothing and \$1  
 147 U.S.A. Bank Note samples remained viable until 96 h with \$20 U.S.A. Bank Note samples  
 148 remaining detectable for 7 days. While the virus continued to lose the remaining 2-3 log<sub>10</sub> PFU  
 149 over the remaining time points of the experiment, the decay rate at 4°C was slower on skin than  
 150 any other tested surface.


151

	Positive Samples by Hour Post-Exposure <sup>†</sup>							
	0	4	8	24	72	96	168	336
Skin	+++	+++	+++	++	+	++	+	+
Clothing	+++	+++	+	+	+	+		
\$1 U.S.A. Bank Note	+++	+++	++	+	+	+	+	
\$20 U.S.A. Bank Note	+++	+++	+++	++	+	+	+	

152

153 Fig 1. Recovery of infectious virus.  
 154 Limit of detection was one plaque forming unit  
 155 <sup>†</sup>Not tested: Skin (22°C at 336 h post-exposure, 37°C at 168 and 336 h post-exposure); and  
 156 Cloth, \$1 U.S.A. Bank Note, and \$20 U.S.A. Bank Note (22°C and 37°C, at 168 h and 336 h  
 157 post-exposure)  
 158 Positive samples at 4°C are represented by a blue +  
 159 Positive samples at 22°C are represented by a green +  
 160 Positive samples at 37°C are represented by a red +

161  
 162  
 163  
 164  
 165  
 166  
 167  
 168  
 169  
 170  
 171  
 172  
 173  
 174  
 175



176

177 Fig 2. Quantification of infectious virus. Viral concentrations were determined as described in  
178 METHODS. Viral concentrations are expressed as mean  $\pm$  SD log<sub>10</sub> PFU/mL.

179 †Not tested: Skin (22°C at 336 h post-exposure, 37°C at 168 h and 336 h post-exposure); and  
180 Cloth, \$1 U.S.A. Bank Note, and \$20 U.S.A. Bank Note (22°C and 37°C, at 168 h and 336 h  
181 post-exposure)

182 \*Lower limit of quantification (LLOQ) of 2.0 log<sub>10</sub> PFU/mL

183

184 At 22°C, the virus appeared to lose approximately 2-3 log<sub>10</sub> PFU within the first 8 h across all  
185 surfaces (Fig 2). Virus was not isolated from the \$1 U.S.A. Bank Note or clothing beyond 8 h at  
186 room temperature. Viable virus was detected on the \$20 U.S.A. Bank Note at 24 h but all  
187 subsequent samples were negative for infectious virus (Fig 1 and Fig 2). Approximately 3 log<sub>10</sub>  
188 PFU of viable virus was detectable on the skin samples at 24 h at 22°C. Virus was isolated at 96  
189 h but all other tested samples were below the LOD.

190  
191 SARS-CoV-2 remained viable at 37°C on skin samples for up to 8 h (Fig 1 and Fig 2). No  
192 viable, infectious virus was detected on clothing samples after four hours at 37°C (Fig 2). There  
193 were minor differences in stability at 37°C between the \$1 U.S.A. Bank Note and the \$20 U.S.A.  
194 Bank Note, but those differences were not statistically significant. There was a discernable  
195 difference in virus stability across each of the temperatures with the 4°C conditions being the  
196 most hospitable conditions for virus stability even at 50% RH.

197  
198 Statistical analysis indicated that skin samples had the longest half-life at each of the tested  
199 temperatures compared to the other surfaces (Table 1). Significant differences in virus stability  
200 were noted when each temperature condition was compared (Fig S1). Significant differences  
201 were observed in virus stability between the skin samples and all other tested surfaces (Fig S2).  
202 No other significant differences were observed during direct comparisons of the other tested  
203 surfaces.

204

205 Table 1: Estimates of Geometric Mean Half-Life by Temperature and Surface

	Temperature		
	4°C	22°C	37°C
Skin	46.8 ( 65.7 , 33.3 )	3.5 ( 14.6 , 0.9 )	0.6 ( 0.9 , 0.4 )
Clothing	33.7 ( 87.6 , 12.9 )	1.0 ( 1.6 , 0.6 )	0.2 ( 0.7 , 0.1 )
\$1 U.S.A. Bank Note	33.2 ( 77.1 , 14.3 )	1.3 ( 5.8 , 0.3 )	0.4 ( 2.8 , 0.1 )
\$20 U.S.A Bank Note	15.9 ( 79.1 , 3.2 )	1.1 ( 1.8 , 0.7 )	0.6 ( 1.3 , 0.3 )

Values indicate geometric mean (95%CL) of the half-life, in hours.

206

207

208

## Discussion

209

210 While the outbreak of SARS-CoV in 2002-2003 resulted in approximately 8000 confirmed cases  
211 and the 2012 MERS-CoV outbreak has culminated in less than 2500 cases, the current SARS-  
212 CoV-2 pandemic has resulted in over 2.5 million cases as of this writing and has spread to nearly  
213 every country on Earth (1-3). There are likely a number of reasons for the rapid emergence of  
214 SARS-CoV-2. Possibilities for enhanced spread include a different human infection case profile,  
215 virus transmission through asymptomatic carriers, or potentially enhanced fomite or aerosol  
216 transmission capability (2-4, 6, 7). Previous reports indicate that SARS-CoV-2 is more stable on  
217 cardboard and plastic compared to SARS-CoV, but aerosol decay rates are similar under  
218 laboratory conditions (6).

219

220 We have found that the skin samples were most hospitable for SARS-CoV-2, especially under  
221 refrigerated conditions. There are noticeable differences in virus decay rates at increasing  
222 temperatures which aligns with previous literature (13). Similar stability profiles were observed  
223 in both the currency and clothing samples. All three samples retained viable virus out to at least  
224 96 h at 4°C with no recoverable virus beyond 8 h at 22°C in any of the three surfaces. There  
225 were small differences in viral concentration between the \$1 U.S.A. Bank Note and \$20 U.S.A.  
226 Bank Note samples, but they were not statistically significant. It is possible that differences in  
227 ink type, concentration, or both, affected virus stability and could warrant further study.

228

229 This is the first report to our knowledge modeling the stability of SARS-CoV-2 on skin. We note  
230 that even at 22°C, SARS-CoV-2 remained infectious on skin samples for 96 h post-exposure.

231 While we understand real-world conditions cannot be replicated in the laboratory setting, this  
232 observation indicates the potential for fomite transmission in indoor environments in the absence  
233 of good hand hygiene practices, given that even trained medical students have been observed  
234 touching their faces approximately 23 times per hour (14). Furthermore, while we did use swine  
235 skin samples as a substitute for human skin, swine skin has enough similarities to human skin  
236 that it has been used for human allograft transplantation especially in burn victims (15). Based on  
237 this, it is expected that if this experiment were ever replicated using human skin samples, it  
238 would generate similar results.

239  
240 We report the results of SARS-CoV-2 stability on animal skin as a series of outbreaks have been  
241 reported in the United States meat packing industry. Since most meat packing and processing  
242 procedures are carried out between 4-8°C, it is likely that any viral shedding from either  
243 symptomatic or asymptomatic workers in the absence of appropriate PPE would remain viable  
244 for an extended period of time on the surface of meat products or other surfaces (16-17).

245 However, even with extensive cleaning, transmission could still occur in the presence of  
246 asymptomatic, undiagnosed workers due to both the enhanced stability of the virus and the high  
247 viral loads even asymptomatic cases maintain in the nasal passages (18). Without an extensive  
248 testing and contact tracing program, transmission around meat packing plants will likely  
249 continue to be an issue until herd immunity is reached either by infection or through the  
250 administration of an efficacious vaccine.

251  
252 It is important to note the limitations of this study when analyzing the results. The variability in  
253 the decay curves between earlier studies and our results could have been due to differences in

254 exposure doses (4, 6, 7). In addition, inherent variability with a study where virus is recovered  
255 from a tested surface can potentially confound some of the results, particularly at virus  
256 concentrations close to the limit of detection (LOD) or lower limit of quantification (LLOQ)  
257 (Fig. 1).

258  
259 The results in this study demonstrate the continued inverse relationship between virus stability  
260 and temperature which is seen both in the laboratory and in the field when evaluating different  
261 transmission rates of SARS-CoV-2 in different parts of the world (19). While both fomite and  
262 aerosol transmission could be significant factors, due to the stability of SARS-CoV-2 on skin  
263 there is a continued need to reinforce proper hand hygiene practices and social distancing  
264 guidelines to minimize ongoing transmission potential.

265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276

277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299

### **Acknowledgements**

The authors are indebted to Mr. Brian Sauerbry who assisted with material procurement, Dr. Pamela Glass for technical assistance and Ms. Sarah Norris for assistance with the statistical analysis. The views expressed in this article are those of the authors and do not reflect the official policy or position of the U.S. Department of Defense or the U.S. Army.

300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322

## References

1. Coronavirus disease (COVID-2019) situation reports. 2020. (Accessed 22<sup>nd</sup> of April, 2020, at <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/>.)
2. Zhu, N. et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med*, <https://doi.org/10.1056/NEJMoa2001017> (2020).
3. Wang C, Horby PW, Hayden FG, Gao GF. A novel coronavirus outbreak of global health concern. *Lancet*, [https://doi.org/10.1016/S0140-6736\(20\)30185-9](https://doi.org/10.1016/S0140-6736(20)30185-9)
4. Otter JA, Donskey C, Yezli S, Douthwaite S, Goldenberg SD, Weber DJ. Transmission of SARS and MERS coronaviruses and influenza virus in healthcare settings: the possible role of dry surface contamination. *J Hosp Infect* 2016;92:235-50.
5. Kampf G, Todt D, Pfaender S, Steinmann E. Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *J Hosp Infect*. 2020 Mar;104(3):246-251.
6. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, Tamin A, Harcourt JL, Thornburg NJ, Gerber SI, Lloyd-Smith JO, de Wit E, Munster VJ. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N Engl J Med*. 2020 Apr 16;382(16):1564-1567

- 323
- 324 7. Chin AWH, Chu JTS, Mahen RA, Perera K, Hui PY, Hui-Ling YM, Chan CW, Peiris M,  
325 Poon LLM. Stability of SARS-CoV-2 in different environmental conditions.  
326 <https://doi.org/10.1101/2020.03.15.20036673>.
- 327
- 328 8. Keyes D. Business Insider. 9 Mar 2020 [cited 16 Mar 2020] In: Business Insider  
329 [Internet]. Available from: [https://www.businessinsider.com/south-korea-burning-cash-](https://www.businessinsider.com/south-korea-burning-cash-amid-coronavirus-outbreak-2020-3)  
330 [amid-coronavirus-outbreak-2020-3](https://www.businessinsider.com/south-korea-burning-cash-amid-coronavirus-outbreak-2020-3).
- 331
- 332 9. Rabenau HF, Cinatl J, Morgenstern B, Bauer G, Preiser W, Doerr HW. Stability and  
333 inactivation of SARS coronavirus. *Med Microbiol Immunol* (2005) 194: 1–6
- 334
- 335 10. Geller C, Varbanov M and Duval RE. Human Coronaviruses: Insights into  
336 Environmental Resistance and Its Influence on the Development of New Antiseptic  
337 Strategies. *Viruses* 2012, 4, 3044-3068.
- 338
- 339 11. SAS Institute Inc. Example 59.7: Influence in Heterogeneous Variance Model. in  
340 *SAS/STAT 14.1 User's Guide: The Mixed Procedure*, Cary, NC, SAS Institute Inc., 2015,  
341 pp. 5048-5056.
- 342
- 343 12. R Core Team (2019). R: A language and environment for statistical computing. R  
344 Foundation for Statistical Computing., Vienna, Austria, 2019.
- 345
- 346

- 347 13. Casanova LM, Jeon S, Rutala WA, Weber DJ, Sobsey MD. Effects of air temperature  
348 and relative humidity on coronavirus survival on surfaces. *Appl Environ Microbiol.* 2010  
349 May;76(9):2712-7.  
350
- 351 14. Kwok YL, Gralton J, McLaws ML. Face touching: a frequent habit that has implications  
352 for hand hygiene. *Am J Infect Control.* 2015;43(2):112-4. doi:  
353 10.1016/j.ajic.2014.10.015.  
354
- 355 15. Yamamoto T, Iwase H, King TW, Hara H, Cooper DKC. Skin xenotransplantation:  
356 Historical review and clinical potential. *Burns.* 2018;44(7):1738-1749.  
357
- 358 16. Krug PW, Davis T, O'Brien C, LaRocco M, Rodriguez LL. Disinfection of transboundary  
359 animal disease viruses on surfaces used in pork packing plants. *Vet Microbiol.* 2018  
360 Jun;219:219-225.  
361
- 362 17. Sun XD and Holley, RA. Antimicrobial and Antioxidative Strategies to Reduce  
363 Pathogens and Extend the Shelf Life of Fresh Red Meats. *Comprehensive Reviews in*  
364 *Food Science and Food Safety.* Vol.11, 2012. doi: 10.1111/j.1541-4337.2012.00188.x.  
365
- 366 18. Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 Viral Load in  
367 Upper Respiratory Specimens of Infected Patients. *N Engl J Med.* 2020;382(12):1177-  
368 1179. doi: 10.1056/NEJMc2001737.  
369

370 19. Sil, A. Does weather affect the growth rate of COVID-19, a study to comprehend  
371 transmission dynamics on human health. BioRxiv [Preprint]. [posted 29 April 2020;  
372 cited 6 May 2020] <https://doi.org/10.1101/2020.04.29.20085795>.

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

## Supplementary Figures

394

395 Supplementary Figure 1. Geometric mean ratio of half-lives, averaged across all three  
396 temperatures.

397

398

399 Supplementary Figure 2. Geometric mean ratio of half-lives, averaged across all four surfaces.