Predicting Eating Disorder and Anxiety Symptoms Using Anorexia Nervosa and Obsessive-Compulsive Disorder Polygenic Scores

Zeynep Yilmaz, PhD;^{1,2} Katherine Schaumberg, PhD;^{1,3} Matt Halvorsen, PhD;² Erica L. Goodman, MA;⁴ Leigh C. Brosf, MS;⁵ James J. Crowley, PhD;^{1,2,6} Anorexia Nervosa Genetics Initiative; Eating Disorders Working Group of the Psychiatric Genomics Consortium; Tourette Syndrome/Obsessive-Compulsive Disorder Working Group of the Psychiatric Genomics Consortium; Carol A. Mathews, MD;⁷ Manuel Mattheisen, MD;^{6,8,9,10} Gerome Breen, PhD;^{11,12} Cynthia M. Bulik, PhD;^{1,13,14} Nadia Micali, MD;^{15,16,17*} Stephanie C. Zerwas; PhD^{1*}

¹ Department of Psychiatry, University of North Carolina at Chapel Hill, Neurosciences Hospital, 101 Manning Drive, Campus Box #7160, Chapel Hill, NC, USA 27599-7160

² Department of Genetics, University of North Carolina at Chapel Hill, 120 Mason Farm Road, 5000 D, Genetic Medicine Building, Campus Box #7264, Chapel Hill, NC, USA 27599-7264

³ Department of Psychiatry, University of Wisconsin – Madison, 6001 Research Park Blvd, Madison, WI, USA, 53719

⁴ Department of Psychology, University of North Dakota, Columbia Hall, 501 North Columbia Road Stop 8380, Grand Forks, ND, USA, 58202-8380

⁵ Department of Psychological and Brain Sciences, University of Louisville, 317 Life Sciences Building, Louisville, KY, USA, 40292

⁶ Department of Clinical Neuroscience, Karolinska Institutet, Tomtebodavägen 18A, 5th floor, Stockholm, Sweden, 171 77

⁷ Department of Psychiatry, Genetics Institute, University of Florida, PO Box 100256, 1149 Newell Drive, L4-100, Gainesville, FL, USA, 32611

⁸ Department of Biomedicine, Aarhus University, Høegh-Guldborgs Gade 10, Aarhus, Denmark, 8000

⁹ The Lundbeck Foundation Initiative of Integrative Psychiatric Research (iPSYCH), Fuglesangs Allé 26, Aarhus, Denmark, 8210

¹⁰ Department of Psychiatry, Psychosomatics and Psychotherapy, University of Würzburg, Anstalt des öffentlichen Rechts, Josef-Schneider-Straße 2, Würzburg, Germany, 97080

¹¹ Institute of Psychiatry, Psychology and Neuroscience, MRC Social, Genetic and Developmental Psychiatry (SGDP) Centre, King's College London, PO80, De Crespigny Park, Denmark Hill, London, United Kingdom, SE5 8AF

¹² National Institute for Health Research Biomedical Research Centre, South London and Maudsley National Health Service Trust, London, UK

¹³ Department of Nutrition, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, 135 Dauer Drive, Campus Box #7461, Chapel Hill, NC, USA 27599-7461

¹⁴ Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, PO Box 281, Stockholm, Sweden, 171-77

¹⁵ Department of Psychiatry, Faculty of Medicine, University of Geneva, HUG, 24 rue du Général-Dufour, Geneva, Switzerland, 1211

¹⁶ Institute of Child Health, University College London, 30 Guilford Street, London, United Kingdom, WC1N-1EH

¹⁷ Department of Paediatrics, Gynecology and Obstetrics, Faculty of Medicine, University of Geneva, HUG, 24 rue du Général-Dufour, Geneva, Switzerland, 1211

* Co-senior authors

Corresponding author: Nadia Micali, MD. Department of Psychiatry, Faculty of Medicine, University of Geneva, HUG, Geneva, Switzerland; telephone: + 41 22 372 89 55; e-mail: Nadia.Micali@unige.ch.

Abstract

Clinical, epidemiological, and genetic findings support an overlap between eating disorders, obsessive-compulsive disorder (OCD), and anxiety symptoms. However, little research has examined the role of genetic factors in the expression of eating disorders and OCD/anxiety phenotypes. We examined whether the anorexia nervosa (AN), OCD, or AN/OCD transdiagnostic polygenic scores (PGS) predict eating disorders, OCD, and anxiety symptoms in a large population-based developmental cohort. Using summary statistics files from the Psychiatric Genomics Consortium Freeze 2 AN and Freeze 1 OCD GWAS, we first conducted an AN/OCD transdiagnostic GWAS meta-analysis and then calculated PGS for AN, OCD, and AN/OCD in participants from the Avon Longitudinal Study of Parents and Children with available genetic and phenotype data on eating disorder, OCD, and anxiety diagnoses and symptoms (sample size 3,212–5,369 per phenotype). We observed sex differences in the PGS prediction of eating disorder, OCD, and anxiety-related phenotypes, with AN genetic risk manifesting at an earlier age and playing a more prominent role in eating disorder phenotypes in boys than in girls. Compulsive exercise was the only phenotype predicted by all three PGS (e.g., $P_{AN(boys)}=0.0141$ at age 14; $P_{OCD(girls)}=0.0070$ at age 16; $P_{AN/OCD(all)}=0.0297$ at age 14). Our results suggest that earlier detection of eating disorder, OCD, and anxiety-related symptoms could be made possible by including measurement of genetic risk for these psychiatric conditions while being mindful of sex differences.

Keywords: ALSPAC, eating disorders, anorexia nervosa, obsessive-compulsive disorder, anxiety, symptom dimensions, polygenic score, sex differences, developmental, population study.

INTRODUCTION

Eating disorders and obsessive-compulsive disorder (OCD) are serious psychiatric conditions with high social, psychological, and physical impact.¹⁻³ Clinical, epidemiological, and genetic findings support an overlap between eating disorders, particularly anorexia nervosa (AN), and OCD.⁴⁻¹⁵ Whilst research on eating disorders and OCD comorbidity has primarily focused on diagnoses, many symptoms and behaviors are common to both diagnoses, spanning diagnostic categories, and their presence often precedes disorder onset.^{16,17} Little research has examined these associations—or symptom phenotypes—in a developmental context. Premorbid OCD symptoms, and anxiety disorders or symptoms are common in patients with AN.¹⁸ Childhood anxiety may precede eating disorder symptoms and AN in adolescence,¹⁹ and shared genetic and environmental influences play a role in anxiety and disordered eating symptoms.²⁰ Though no longer classified as an anxiety disorder,¹ OCD is highly comorbid with anxiety disorders and includes anxiety symptoms,²¹ especially in children.²² An improved understanding of the overlap among eating disorders and OCD, and intermediate phenotypes such as anxiety symptoms could aid in conceptualizing mechanisms and processes contributing to the clinical and genetic overlap between these disorders. Additionally, symptom dimensions may transmute over development, shifting from childhood obsessive-compulsive symptoms to adolescent eating disorders^{23,24} and vice versa. Thus, shared and unique risk factors may contribute to symptoms of OCD and eating disorders across development.

Genome-wide association studies (GWAS) of AN¹⁴ and OCD²⁵ have provided important insights into the highly polygenic architecture of these disorders and their positive genetic correlation.¹⁴ Application of polygenic scores (PGS)—the weighted sum of common risk variants per individual—examine the genetic architecture of complex traits using evidence for association

from variants below the stringent threshold for genome-wide significance.²⁶ The use of PGS has been validated across psychiatric diagnoses and symptom-level measures,²⁷⁻³³ demonstrating that genetic variants associated with risk are often shared across diagnostic categories.³³ Moreover, transdiagnostic PGS (determined by either AN or OCD case status) of genetically correlated disorders may enhance predictive power for either disorder.³⁴

Sex differences in the prevalence and presentation of eating disorders, anxiety disorders, and OCD warrant sex-specific examination of risk factors. While the majority of AN cases are female,^{35,36} AN in males often has an earlier age of onset and is likely to be more severe.³⁷⁻³⁹ Similarly, the lifetime prevalence of eating disorders is much higher in females than males,^{35,36} possibly with the exception of subthreshold binge eating.³⁵ The lifetime prevalence of anxiety disorders is up to 60% higher in women than in men.⁴⁰ In the case of OCD, childhood onset is more common among males and adolescent onset is more common among females.⁴¹ Importantly, sex differences in the presentation of symptoms such as restraint and weight and shape concern in eating disorders³⁷ and contamination/cleaning and sexual/religious symptoms in OCD⁴² have also been reported. Given these discrepancies, we could expect: a) notable sex differences in the role of genetic risk and eating disorders, OCD, and anxiety symptom phenotypes; and b) that genetic risk may be more impactful and predictive for boys, especially in the case of eating disorders.

This study examined whether the AN, OCD, or AN/OCD PGS predicts eating disorders, OCD, and anxiety symptom dimensions or diagnoses using a developmental framework in male and female participants from a population-based cohort. Our main hypothesis was that AN/OCD PGS would demonstrate better statistical power than AN or OCD PGS, and the transdiagnostic PGS would evidence the most benefit compared with single-trait PGS when predicting intermediate phenotypes *shared* across the two disorders, such as generalized anxiety or worrying.

We also hypothesized that symptom dimensions specific to each disorder would be predicted by disorder-specific PGS (e.g., thin ideal internalization by AN, or symmetry/checking behavior by OCD).⁴³ Finally, in light of the differences in lifetime prevalence and/or age of onset of eating disorders, OCD, and anxiety disorders between sexes, we hypothesized that there would be significant sex differences in whether AN, OCD, and AN/OCD PGS predicted eating disorders and anxiety phenotypes, and high AN genetic risk would play a larger role in predicting eating disorder symptoms in boys than girls.

MATERIALS AND METHODS

Participants

The Avon Longitudinal Study of Parents and Children (ALSPAC) is a longitudinal, population-based study of women and their children.⁴⁴ All pregnant women living in Avon, United Kingdom who were expected to deliver between April 1, 1991, and December 31, 1992 were invited to participate. Children from 14,541 pregnancies were enrolled, 13988 of whom were alive at one year. An additional 713 children were enrolled at or after age 7.⁴⁴ Please note that the study website contains details of all the data that is available through a fully searchable data dictionary and variable search tool (<http://www.bristol.ac.uk/alspac/researchers/our-data/>). Ethical approval was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees.

For genetic analyses, we used post-quality control (QC) dosage files for 7,977 unrelated participants,^{45,46} 7,779 of whom passed additional QC performed as a part of this study (*Supplementary Information*). The final number of participants with phenotype and genotype data was 3,212–5,369 (see *Tables 2–4* for phenotype-specific sample sizes).

Measures

Table 1 provides a list of all measures, assessment timepoints, and methods of administration.

Eating disorder symptoms for the previous year were evaluated at ages 14 and 16 using questions adapted from the Youth Risk Behavior Surveillance System Questionnaire,⁴⁷ validated in a population-based study of adolescents.⁴⁸ Binge-eating, purging, fasting, and compulsive exercise were characterized and categorized as described previously (**Supplementary Information**).^{49,50} Eating disorder diagnoses at ages 14 and 16 were derived using DSM-5 criteria¹ as detailed in a previous publication by our group.¹⁹ Eating disorder cognitions, including: body image distortion, emotional eating, external eating, body dissatisfaction, thin ideal internalization, dietary restraint, weight concern, and shape concern were assessed by validated, age appropriate self-report measurements (**Supplementary Information**).

OCD and anxiety symptoms at age 7, 10, 13, and 15 were collected using the Development and Wellbeing Assessment (DAWBA; **Supplementary Information**).^{51,52} Probabilities of anxiety disorder diagnoses at ages 7 (specific phobia and separation anxiety), 10 (OCD), 13 (OCD, social phobia, and generalized anxiety disorder), and 15 (generalized anxiety disorder) were determined using computer-generated DAWBA band variables,⁵² which assign the probability of the participant meeting DSM-IV criteria for an anxiety disorder. However, due to a very small number of participants scoring high on these variables, we dichotomized them to capture whether the likelihood of a diagnosis was <50% versus ≥50%. We also defined five latent OCD or anxiety factors for ages 10 and 13: (1) OCD-symmetry; (2) OCD-dirt/germs; (3) physical anxiety; (4) worrying; and (5) social phobia (**Supplementary Information**).¹⁹

Data Analysis

We calculated AN, OCD, and AN/OCD PGS to predict 27 eating disorder, 6 OCD, and 11 anxiety phenotypes in the ALSPAC target sample using PRSice v2.2.⁵³ AN PGS was constructed using the Anorexia Nervosa Genetics Initiative & Psychiatric Genomics Consortium (PGC) Eating Disorder Working Group Freeze 2 AN GWAS,¹⁴ and OCD PGS was calculated using the Freeze 1 PGC OCD GWAS.²⁵ AN/OCD summary statistics files were obtained from a GWAS conducted by combining these datasets (see *Supplementary Information, Table S1*, and *Figure S1*). We first examined how well AN, OCD, and AN/OCD PGS predicted eating disorder, OCD, and anxiety symptom phenotypes in the combined sample and then investigated sex-specific differences. Due to insufficient power, only binary phenotypes with ≥ 50 cases are reported. Significance threshold was set at empirical $P < 0.05$ following 10,000 permutations per target phenotype. Corresponding odds ratios and 95% confidence intervals for binary phenotypes are in *Table S2*.

RESULTS

Eating disorder symptom phenotypes and diagnoses

Figures 1a and *1b* show the significant results for eating disorder phenotypes. In the combined sample, AN PGS predicted compulsive exercise ($P=0.0448$) and body dissatisfaction at age 14 ($P=0.0410$), whereas OCD PGS predicted eating disorder not otherwise specified/purging disorder ($P=0.0410$), the presence of a threshold or subthreshold eating disorder ($P=0.0111$), and compulsive exercise ($P<0.0042$) at age 16 (*Table 2*). In addition to predicting compulsive exercise ($P=0.0297$) and body dissatisfaction at age 14 ($P=0.0200$), AN/OCD PGS uniquely predicted higher restraint at age 14 ($P=0.0147$).

In girls, none of the eating disorder-related constructs were predicted by AN PGS (**Table 3**). OCD PGS predicted higher thin-ideal internalization at age 14 ($P=0.0361$) as well as presence of a threshold or subthreshold eating disorder ($P=0.0232$), binge eating ($P=0.0264$), and compulsive exercise at age 16 ($P=0.0070$). AN/OCD PGS predicted increased thin-ideal internalization at age 14 ($P=0.0185$).

In boys, AN PGS predicted eating disorder not otherwise specified/purging disorder ($P=0.0400$), increased compulsive exercise ($P=0.0141$), and greater internal pressure to lose weight ($P=0.0432$) at age 14 (**Table 3**). OCD PGS predicted less internal pressure to lose weight at age 14 ($P=0.0051$), and the presence of any eating disorder or subthreshold eating disorder at age 16 ($P=0.0479$). AN/OCD PGS predicted greater internal pressure to lose weight ($P=0.0122$) and higher restraint at age 14 ($P=0.013$).

OCD and anxiety symptom phenotypes and diagnoses

Figures 1c and **1d** show the significant results for OCD and anxiety phenotypes. In the combined sample, AN PGS predicted an increased likelihood of separation anxiety at age 7 ($P=0.0132$), higher scores for the latent factors OCD dirt/germs ($P=0.0319$), worrying at age 10 ($P=0.0163$), and social phobia ($P=0.0380$) at age 13 (**Table 2**). None of the anxiety phenotypes were predicted by OCD PGS. AN/OCD PGS predicted an increased likelihood of specific phobia at age 7 ($P=0.0181$).

In girls, AN PGS predicted an increased likelihood of separation anxiety ($P=0.0035$) at age 7 (**Table 3**). OCD PGS predicted lower scores for the latent factors worrying ($P=0.0307$) and social phobia at age 13 ($P=0.0072$). AN/OCD PGS predicted an increased likelihood of separation anxiety ($P=0.0134$) and specific phobia at age 7 ($P=0.0378$).

In boys, AN PGS predicted a higher worrying score at age 10 ($P=0.0143$), whereas neither OCD nor AN/OCD PGS predicted any anxiety-related phenotypes (*Table 4*).

DISCUSSION

We were able to predict eating disorder, OCD, and anxiety phenotypes using AN, OCD, and AN/OCD PGS during different developmental points in a large population sample. In contrast to our hypothesis, the majority of symptom phenotypes were predicted by either the AN or OCD PGS and not by the AN/OCD PGS, suggesting that the genetic risk associated with these measures during development may be more disorder-specific than a transdiagnostic common factor. Alternatively, multi-trait PGS may not have reached the point at which it could enhance predictive power, especially when the contribution by one of the two disorder-specific PGS is more subtle and requires much larger sample sizes. This could also be driven by the notably smaller sample size of the OCD GWAS compared to the AN GWAS (2,688 vs 16,992 cases), therefore limiting the contribution of OCD genetic risk to the transdiagnostic PGS.

Significant PGS predictions did not fall cleanly in accordance with hypothesized disorder-specific symptom phenotypes. For instance, AN—but not OCD—PGS predicted higher scores for OCD-dirt/germs and worrying at age 10. While contamination fears are often associated with OCD, they are not unique to OCD and have a cross-disorder component. In fact, it is not uncommon for individuals with AN to present with food-related contamination fears.⁵⁴ In our ALSPAC previous study, we found that the latent factor worrying significantly predicts eating disorder symptoms at ages 14 and 16 as well as AN diagnosis at age 16.¹⁹ This may suggest that uncontrolled worrying may be an underlying early symptom of AN and disordered eating that precedes the manifestation of an eating disorder. In contrast, OCD PGS predicted less worrying at age 13, which may be due to qualitative differences in worry and rumination between general

anxiety and OCD.⁵⁵ Additionally, OCD—but not AN—PGS predicted increased risk for binge eating, eating disorders not otherwise specified/purging disorder, and the presence of any eating disorder at age 16. This may be due to a larger obsessive and/or compulsive genetic component to eating disorders other than AN, which genetically has a key metabolic component.¹⁴ Additionally, purging often has a significant compulsive component.⁵⁶ These results corroborate findings from our team of a contribution of OCD genetic risk to bulimia nervosa.⁵⁷ While currently AN is the only eating disorder for which GWAS data are available, future studies on the molecular genetics of other eating disorders could shed light on their potential genetic overlap with OCD.

Our findings further emphasize the potential for sex differences in the biological pathways and vulnerabilities leading to these symptom phenotypes. As per our hypothesis, our results suggest that AN genetics may play a more prominent role in risk for eating disorder and related phenotypes in boys, as compared to girls, in *early development*. In fact, significant eating disorder symptom phenotypes at age 14 were predicted predominantly by AN PGS in boys, whereas it was the OCD PGS that predicted significant eating disorder phenotypes at age 16 in both sexes. Twin studies in female participants have shown changes in the contribution of genetic and environmental risk factors for disordered eating during different stages of adolescence—with a higher genetic risk being present during later teen years^{58,59}—whereas the twin-based heritability estimate for disordered eating may be much higher in boys than girls prior to puberty (0.52 in boys versus 0 in girls),⁶⁰ further supporting our hypothesis that AN genetic load may manifest itself earlier during adolescence in boys. Similarly, since OCD often precedes eating disorder onset temporally, it is possible that OCD genetic risk loading might need longer environmental exposure to lead to eating disorders, which could also explain why many eating disorder phenotypes were successfully predicted by OCD PGS later during development in both sexes.

Notably, AN PGS did not predict any eating disorder phenotypes in girls. It is possible that the manifestation of eating disorder phenotypes may be more heavily influenced by the modern sociocultural environment in girls than boys, therefore reducing the contribution of genetic risk. For boys, genetic risk may be of increased importance as weight-related societal pressure is less pronounced; whereas for girls, a lower genetic risk load coupled with sociocultural factors may suffice for the expression of eating disorder-related symptoms. Additionally, it is unclear how much the genetic etiology of AN overlaps with other eating disorder diagnoses or symptoms. Twin studies have shown disordered eating symptoms to be moderately heritable (i.e., a heritability estimate of up to 0.65),⁵⁹⁻⁶² but while these symptom phenotypes have a notable twin-based genetic correlation with eating disorders other than AN, their genetic correlation with AN is much weaker (0.52 for other eating disorders versus 0.26 for AN).⁶² Although SNP-based heritability data for non-AN eating disorder phenotypes are limited and based on small sample sizes, a recent GWAS by our group did not find a significant genetic correlation between AN and bulimia factor score,⁶³ despite both phenotypes showing significant SNP-based heritability estimates.⁶⁴ While these studies were conducted in female participants only and cannot be necessarily generalized to men, currently little evidence exists at a genetic level on whether these intermediate phenotypes truly lie on a continuum with AN diagnosis.

We also observed sex differences in the genetic prediction of anxiety symptoms and diagnoses. AN PGS predicted separation anxiety at age 7 in girls and increased worrying at age 10 in boys. Epidemiological studies show over a 10-fold increase in AN risk among girls with separation anxiety disorder,⁶⁵ and a twin-based study reported a shared genetic effect influencing liability to AN, separation anxiety, and childhood overanxious disorder (which is very similar to generalized anxiety disorder in adults) during different stages of development,⁶⁶ supporting our

findings about the presence of a shared genetic pathway between anxiety and AN. We unexpectedly observed that *lower* OCD-specific genetic risk predicted higher anxiety symptoms in girls (e.g., worrying and social phobia latent factors at age 13). As worrying and social phobia are symptoms that are shared among many anxiety disorders and may be non-specific, enhanced risk for these symptoms may arise through more general pathways. Additionally, while anxiety symptoms are common in patients with OCD, our results suggest that OCD may be distinct from anxiety disorders at a genetic level, especially for men. Replication of these associations is crucial to better understand the nature of this relationship and the importance of sex differences in the biological pathways associated with anxiety risk.

Compulsive exercise was the only intermediate phenotype that was positively predicted by more than one PGS for both sexes, strongly suggesting it may be one intermediate phenotype that, although commonly associated with eating disorders, is truly transdiagnostic, not sex-specific, and influenced by genetic risk for both AN and OCD. Together with evidence for shared genetic risk between a broad AN phenotype and general propensity for physical activity,¹⁴ this finding suggests that genetic factors may be particularly relevant to understanding the development of compulsive exercise in eating disorders.

Compulsive exercise encompasses many of the hallmark symptoms of AN (e.g., weight and shape concern) and OCD (e.g., compulsive behavior).⁶⁷ Furthermore, comorbid OCD symptoms are especially pronounced in the subpopulation of AN patients with compulsive exercise,⁶⁷⁻⁷¹ which has significant clinical relevance since the presence of compulsive exercise in AN is an established predictor of negative treatment outcomes, including higher pathology at discharge from inpatient treatment,⁷² relapse,⁷³ and greater energy requirements for weight gain.⁷⁴ Treatments for this symptom are currently lacking, and our results point to the need for additional

investigation of the habitual and compulsive nature of exercise behavior, which may lead to targeted intervention development for this symptom that derives from a modern biobehavioral understanding of both eating disorders and OCD.

Our study has notable strengths that merit consideration. This is the first study to use AN, OCD, or AN/OCD PGS to predict eating disorder and anxiety intermediate phenotypes, and to examine sex differences in how genetic risk manifests in the presence of these intermediate phenotypes during different stages of development in a large population sample. We augmented the diagnostic approach by including intermediate phenotypes measured continuously to capture the full range of these underlying traits in the general population. Furthermore, studying these associations in the general population allows a finer understanding of intermediate phenotypes and broader psychopathology as treatment-seeking individuals might have notable differences from the general population (e.g., increased comorbid psychopathology).

Limitations of our study include reliance on self- or parent-report symptoms instead of clinical diagnoses, phenotype data being available for only a subset of participants with the potential for response bias, and potential Type 2 error due to lack of statistical power for PGS in the prediction of genetic risk. Of note, few participants met diagnostic criteria for AN, bulimia nervosa, or binge-eating disorder, whereas we had better statistical power for the non-specific eating disorder statuses, which likely explains why AN PGS did not predict AN diagnosis. Similarly, a high probability (50% or higher) for anxiety disorder diagnoses was uncommon despite our attempt to increase power through dichotomizing these items, and even with dichotomizing, we did not have enough cases to include OCD diagnosis in our outcomes. Additionally, we did not address the presence of comorbid psychiatric diagnoses, therefore we cannot account for the role of comorbidities or the genetic risk associated with these additional

diagnoses. However, as comorbidity is the norm and not the exception, our results are likely to capture associations that are more likely to be present in clinical and population settings, as pure forms of eating disorders and OCD are not common. From a genetic perspective, whether all of the eating disorder-related symptom phenotypes examined as a part of our study actually fall on an etiological continuum with AN is not clear.⁶² Finally, the AN PGS was constructed using a much larger GWAS than the OCD GWAS, which may have translated to OCD PGS being underpowered and the AN/OCD transdiagnostic GWAS being more heavily skewed by AN PGS than OCD PGS.

Taken together, results of our study point to significant sex differences in the manifestation of genetic risk for eating disorder and anxiety symptoms. Genetic risk associated with AN may be a stronger predictor of eating disorder symptoms in boys, whereas OCD genetic risk may play a bigger role in symptomatology in these domains for girls, and may increase in effect across adolescence, which is in line with previous research findings that genetic effects may play a more prominent role post-puberty for eating disorder risk in girls.⁷⁵ These findings could have important implications for earlier detection of disordered eating and anxiety-related symptoms while being mindful of sex differences. In clinical settings, men with AN often have an earlier onset, present with lower BMI and fat mass, and score higher on weight suppression than women with AN,³⁷⁻³⁹ raising the possibility of a stronger genetic role in the etiology of AN in men. Our genetic results provide support for this hypothesis, especially in the context of a developmental framework. Apart from sex differences, one significant observation was that compulsive exercise may be an intermediate phenotype or clinical manifestation of shared genetic risk factors for both AN and OCD in both sexes. It is possible that compulsive exercise might be a distinct AN subphenotype, and clinical research should continue to explore habitual and compulsive processes associated with

this symptom. Finally, this study opens up new avenues for a clearer understanding of biology of behaviors and intermediate phenotypes in eating disorders.

ACKNOWLEDGEMENTS

ZY was responsible for genetic study design, quality control of genotype data, genetic analyses, and manuscript preparation. KS, ELG, and LCB was responsible for study design, quality control and preparation of phenotype data, statistical analysis of phenotype data, and manuscript preparation. MH was responsible for carrying out the transdiagnostic AN/OCD GWAS. MH, JJC, MM, CAM, and GB were responsible for genetic study design, oversight of genetic analyses, and manuscript preparation. CMB oversaw and contributed to development of the research question, study design, and manuscript preparation. NM and SCZ were responsible for development of the research question and provided oversight for all aspects of the study.

We are deeply grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. The UK Medical Research Council and the Wellcome Trust (Grant ref: 217065/Z/19/Z) and the University of Bristol provide core support for ALSPAC. This publication is the work of the authors and Dr. Micali will serve as guarantor for the contents of this paper.

Dr. Micali had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Funding

This research was specifically funded by NIH 5R01MH073842 and R21MH109917. ZY is funded by the National Institute of Mental Health (K01MH109782; R01MH105500) and a Brain and Behavior Research Foundation NARSAD Young Investigator Award (grant # 28799). KS is supported by NIH K01MH123914 and NIH L30MH120619. SCZ is supported by NIH

K01MH100435. CMB acknowledges funding from the National Institute of Mental Health (R01MH120170; R01MH119084; R21MH115397; U01 MH109528), the Swedish Research Council (VR Dnr: 538-2013-8864), and the Klarman Family Foundation. NM acknowledges grant funding from the Medical Research Council (MR/R004803/1), NIH (R01MH108595; R21MH115397), the Swiss National Fund (320030_182484), and a Brain and Behavior Research Foundation Independent Investigator Award (grant #24608). JJC acknowledges grant funding from NIH (R01MH105500; R01MH110427).

The UK Medical Research Council and the Wellcome Trust (Grant ref: 217065/Z/19/Z) and the University of Bristol provide core support for ALSPAC. This publication is the work of the authors and they will serve as guarantors for the contents of this paper. A comprehensive list of grant funding is available on the ALSPAC website.

Conflict of Interest

GB received grant funding and consultancy fees in preclinical genetics from Eli Lilly, consultancy fees from Otsuka, and has received honoraria from Illumina. CMB served on Shire Scientific Advisory Boards, is a consultant for Idorsia, and receives author royalties from Pearson. All other authors have no conflicts of interest to disclose.

CONSORTIA CO-AUTHORS

Anorexia Nervosa Genetics Initiative

Jessica H. Baker, Andrew W. Bergen, Andreas Birgegård, Joseph M. Boden, Harry Brandt, Cynthia M. Bulik, Steven Crawford, Laramie E. Duncan, Scott Gordon, Jakob Grove, Katherine A. Halmi, Anjali K. Henders, L. John Horwood, Craig Johnson, Jennifer Jordan, Anders Juréus, Allan S. Kaplan, Walter Kaye, Martin Kennedy, Katherine M. Kirk, Mikael Landén, Janne T. Larsen, Virpi M. Leppä, Paul Lichtenstein, Nicholas G. Martin, Manuel Mattheisen, James Mitchell, Grant W. Montgomery, Preben Bo Mortensen, Melissa A. Munn-Chernoff, Claes Norring, Catherine M. Olsen, Richard Parker, John F. Pearson, Nancy L. Pedersen, Liselotte Petersen, Michael Strober, Patrick F. Sullivan, Laura M. Thornton, Tracey D. Wade, Hunna J. Watson, Thomas Werge, David C. Whiteman, D. Blake Woodside, Zeynep Yilmaz

Eating Disorders Working Group of the PGC

Hunna J. Watson, Zeynep Yilmaz, Laura M. Thornton, Christopher Hübel, Jonathan R. I. Coleman, Hélène A. Gaspar, Julien Bryois, Anke Hinney, Virpi M. Leppä, Manuel Mattheisen, Sarah E. Medland, Stephan Ripke, Shuyang Yao, Paola Giusti-Rodríguez, Ken B. Hanscombe, Kirstin L. Purves, Roger A. H. Adan, Lars Alfredsson, Tetsuya Ando, Ole A. Andreassen, Jessica H. Baker, Wade H. Berrettini, Ilka Boehm, Claudette Boni, Vesna Boraska Perica, Katharina Buehren, Roland Burghardt, Matteo Cassina, Sven Cichon, Maurizio Clementi, Roger D. Cone, Philippe Courtet, Scott Crow, James J. Crowley, Unna N. Danner, Oliver S. P. Davis, Martina de Zwaan, George Dedoussis, Daniela Degortes, Janiece E. DeSocio, Danielle M. Dick, Dimitris Dikeos, Christian Dina, Monika Dmitrzak-Weglarz, Elisa Docampo, Laramie E. Duncan, Karin Egberts, Stefan Ehrlich, Geòrgia Escaramís, Tõnu Esko, Xavier Estivill, Anne Farmer, Angela

Favaro, Fernando Fernández-Aranda, Manfred M. Fichter, Krista Fischer, Manuel Föcker, Lenka Foretova, Andreas J. Forstner, Monica Forzan, Christopher S. Franklin, Steven Gallinger, Ina Giegling, Johanna Giuranna, Fragiskos Gonidakis, Philip Gorwood, Monica Gratacos Mayora, Sébastien Guillaume, Yiran Guo, Hakon Hakonarson, Konstantinos Hatzikotoulas, Joanna Hauser, Johannes Hebebrand, Sietske G. Helder, Stefan Herms, Beate Herpertz-Dahlmann, Wolfgang Herzog, Laura M. Huckins, James I. Hudson, Hartmut Imgart, Hidetoshi Inoko, Vladimir Janout, Susana Jiménez-Murcia, Antonio Julià, Gursharan Kalsi, Deborah Kaminská, Jaakko Kaprio, Leila Karhunen, Andreas Karwautz, Martien J. H. Kas, James L. Kennedy, Anna Keski-Rahkonen, Kirsty Kiezebrink, Youl-Ri Kim, Lars Klareskog, Kelly L. Klump, Gun Peggy S. Knudsen, Maria C. La Via1, Stephanie Le Hellard, Robert D. Levitan, Dong Li, Lisa Lilienfeld, Bochao Danae Lin, Jolanta Lissowska, Jurjen Luykx, Pierre J. Magistretti, Mario Maj, Katrin Mannik, Sara Marsal, Christian R. Marshall, Morten Mattingsdal, Sara McDevitt, Peter McGuffin, Andres Metspalu, Ingrid Meulenbelt, Nadia Micali, Karen Mitchell, Alessio Maria Monteleone, Palmiero Monteleone, Melissa A. Munn-Chernoff, Benedetta Nacmias, Marie Navratilova, Ioanna Ntalla, Julie K. O'Toole, Roel A. Ophoff, Leonid Padyukov, Aarno Palotie, Jacques Pantel, Hana Papezova, Dalila Pinto, Raquel Rabionet, Anu Raevuori, Nicolas Ramoz, Ted Reichborn-Kjennerud, Valdo Ricca, Samuli Ripatti, Franziska Ritschel, Marion Roberts, Alessandro Rotondo, Dan Rujescu, Filip Rybakowski, Paolo Santonastaso, André Scherag, Stephen W. Scherer, Ulrike Schmidt, Nicholas J. Schork, Alexandra Schosser, Jochen Seitz, Lenka Slachtova, P. Eline Slagboom, Margarita C. T. Slof-Op 't Landt, Agnieszka Slopian, Sandro Sorbi, Beata Świątkowska, Jin P. Szatkiewicz, Ioanna Tachmazidou, Elena Tenconi, Alfonso Tortorella, Federica Tozzi, Janet Treasure, Artemis Tsitsika, Marta Tyszkiewicz-Nwafor, Konstantinos Tziouvas, Annemarie A. van Elburg, Eric F. van Furth, Gudrun Wagner, Esther Walton, Elisabeth

Widen, Eleftheria Zeggini, Stephanie Zerwas, Stephan Zipfel, Andrew W. Bergen, Joseph M. Boden, Harry Brandt, Steven Crawford, Katherine A. Halmi, L. John Horwood, Craig Johnson, Allan S. Kaplan, Walter H. Kaye, James E. Mitchell, Catherine M. Olsen, John F. Pearson, Nancy L. Pedersen, Michael Strober, Thomas Werge, David C. Whiteman, D. Blake Woodside, Garret D. Stuber, Scott Gordon, Jakob Grove, Anjali K. Henders, Anders Juréus, Katherine M. Kirk, Janne T. Larsen, Richard Parker, Liselotte Petersen, Jennifer Jordan, Martin Kennedy, Grant W. Montgomery, Tracey D. Wade, Andreas Birgegård, Paul Lichtenstein, Claes Norring, Mikael Landén, Nicholas G. Martin, Preben Bo Mortensen, Patrick F. Sullivan, Gerome Breen, Cynthia M. Bulik.

Obsessive Compulsive Disorder Working Group of the PGC

Paul D. Arnold, Kathleen D. Askland, Cristina Barlassina, Laura Bellodi, O. J. Bienvenu, Donald Black, Michael Bloch, Helena Brentani, Christie L. Burton, Beatriz Camarena, Carolina Cappi, Danielle Cath, Maria Cavallini, David Conti, Edwin Cook, Vladimir Coric, Bernadette A. Cullen, Danielle Cusi, Lea K. Davis, Richard Delorme, Damiaan Denys, Eske Derks, Valsamma Eapen, Christopher Edlund, Lauren Erdman, Peter Falkai, Martijn Figee, Abigail J. Fyer, Daniel A Geller, Fernando S. Goes, Hans Grabe, Marcos A. Grados, Benjamin D. Greenberg, Edna Grünblatt, Wei Guo, Gregory L. Hanna, Sian Hemmings, Ana G. Hounie, Michael Jenicke, Clare Keenan, James Kennedy, Ekaterina A. Khramtsova, Anuar Konkashbaev, James A. Knowles, Janice Krasnow, Cristophe Lange, Nuria Lanzagorta, Marion Leboyer, Leonhard Lennertz, Bingbin Li, K-Y Liang, Christine Lochner, Fabio Macciardi, Brion Maher, Wolfgang Maier, Maurizio Marconi, Carol A. Mathews, Manuel Mattheisen, James T. McCracken, Nicole C. McLaughlin, Euripedes C. Miguel, Rainald Moessner, Dennis L. Murphy, Benjamin Neale, Gerald Nestadt, Paul Nestadt, Humberto

Nicolini, Ericka Nurmi, Lisa Osiecki, David L. Pauls, John Piacentini, Danielle Posthuma, Ann E. Pulver, H-D Qin, Steven A. Rasmussen, Scott Rauch, Margaret A. Richter, Mark A. Riddle, Stephan Ripke, Stephan Ruhrmann, Aline S. Sampaio, Jack F. Samuels, Jeremiah M. Scharf, Yin Yao Shugart, Jan Smit, Daniel Stein, S. Evelyn Stewart, Maurizio Turiel, Homero Vallada, Jeremy Veenstra-VanderWeele, Michael Wagner, Susanne Walitza, Y. Wang, Jens Wendland, Nienke Vulink, Dongmei Yu, Gwyneth Zai.

REFERENCES

1. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders, fifth edition*, (Author, Washington DC, 2013).
2. Keshaviah, A. *et al.* Re-examining premature mortality in anorexia nervosa: a meta-analysis redux. *Compr Psychiatry* **55**, 1773-84 (2014).
3. World Health Organization. The Global Burden of Disease: 2004 Update. (2008).
4. Kaye, W.H., Bulik, C.M., Thornton, L., Barbarich, N. & Masters, K. Comorbidity of anxiety disorders with anorexia and bulimia nervosa. *Am J Psychiatry* **161**, 2215-21 (2004).
5. Godart, N.T., Flament, M.F., Perdereau, F. & Jeammet, P. Comorbidity between eating disorders and anxiety disorders: a review. *Int J Eat Disord* **32**, 253-70 (2002).
6. Swinbourne, J.M. & Touyz, S.W. The co-morbidity of eating disorders and anxiety disorders: a review. *Eur Eat Disord Rev* **15**, 253-74 (2007).
7. du Toit, P.L., van Kradenburg, J., Niehaus, D. & Stein, D.J. Comparison of obsessive-compulsive disorder patients with and without comorbid putative obsessive-compulsive spectrum disorders using a structured clinical interview. *Compr Psychiatry* **42**, 291-300 (2001).
8. Rubenstein, C.S., Pigott, T.A., L'Heureux, F., Hill, J.L. & Murphy, D.L. A preliminary investigation of the lifetime prevalence of anorexia and bulimia nervosa in patients with obsessive compulsive disorder. *J Clin Psychiatry* **53**, 309-14 (1992).
9. Cederlof, M. *et al.* Etiological overlap between obsessive-compulsive disorder and anorexia nervosa: a longitudinal cohort, multigenerational family and twin study. *World Psychiatry* **14**, 333-8 (2015).

10. Lilenfeld, L.R. *et al.* A controlled family study of anorexia nervosa and bulimia nervosa: psychiatric disorders in first-degree relatives and effects of proband comorbidity. *Arch Gen Psychiatry* **55**, 603-10 (1998).
11. Strober, M., Freeman, R., Lampert, C. & Diamond, J. The association of anxiety disorders and obsessive compulsive personality disorder with anorexia nervosa: evidence from a family study with discussion of nosological and neurodevelopmental implications. *Int J Eat Disord* **40 Suppl**, S46-51 (2007).
12. Meier, S.M. *et al.* Diagnosed Anxiety Disorders and the Risk of Subsequent Anorexia Nervosa: A Danish Population Register Study. *Eur Eat Disord Rev* **23**, 524-30 (2015).
13. Yilmaz, Z. *et al.* Examination of the shared genetic basis of anorexia nervosa and obsessive-compulsive disorder. *Mol Psychiatry* (2018).
14. Watson, H.J. *et al.* Genome-wide association study identifies eight risk loci and implicates metabo-psychiatric origins for anorexia nervosa. *Nat Genet* **51**, 1207-1214 (2019).
15. Anttila, V. *et al.* Analysis of shared heritability in common disorders of the brain. *Science* **360**(2018).
16. Stice, E. Interactive and Meditational Etiologic Models of Eating Disorder Onset: Evidence from Prospective Studies. *Annu Rev Clin Psychol* **12**, 359-81 (2016).
17. Nolen-Hoeksema, S. & Watkins, E.R. A Heuristic for Developing Transdiagnostic Models of Psychopathology: Explaining Multifinality and Divergent Trajectories. *Perspect Psychol Sci* **6**, 589-609 (2011).

18. Udo, T. & Grilo, C.M. Psychiatric and medical correlates of DSM-5 eating disorders in a nationally representative sample of adults in the United States. *Int J Eat Disord* **52**, 42-50 (2019).
19. Schaumberg, K. *et al.* Anxiety disorder symptoms at age 10 predict eating disorder symptoms and diagnoses in adolescence. *J Child Psychol Psychiatry* **60**, 686-696 (2019).
20. Fairweather-Schmidt, A.K. & Wade, T.D. Common genetic architecture and environmental risk factors underpin the anxiety-disordered eating relationship: Findings from an adolescent twin cohort. *Int J Eat Disord* **53**, 52-60 (2020).
21. Torres, A.R. *et al.* Comorbidity variation in patients with obsessive-compulsive disorder according to symptom dimensions: Results from a large multicentre clinical sample. *J Affect Disord* **190**, 508-516 (2016).
22. Anagnostopoulos, D.C. *et al.* Comorbid psychopathology and clinical symptomatology in children and adolescents with obsessive-compulsive disorder. *Psychiatriki* **27**, 27-36 (2016).
23. Anderluh, M.B., Tchanturia, K., Rabe-Hesketh, S. & Treasure, J. Childhood obsessive-compulsive personality traits in adult women with eating disorders: defining a broader eating disorder phenotype. *Am J Psychiatry* **160**, 242-7 (2003).
24. Micali, N. *et al.* Is childhood OCD a risk factor for eating disorders later in life? A longitudinal study. *Psychol Med* **41**, 2507-13 (2011).
25. International Obsessive Compulsive Disorder Foundation Genetics Collaborative (IOCDF-GC) and OCD Collaborative Genetics Association Studies (OC GAS). Revealing the complex genetic architecture of obsessive-compulsive disorder using meta-analysis. *Mol Psychiatry* **23**, 1181-1188 (2018).

26. Wray, N.R. *et al.* Research review: Polygenic methods and their application to psychiatric traits. *J Child Psychol Psychiatry* **55**, 1068-87 (2014).
27. Lee, S.H. *et al.* Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. *Nat Genet* **45**, 984-94 (2013).
28. Cross-Disorder Group of the Psychiatric Genomics Consortium *et al.* Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet* **381**, 1371 (2013).
29. Ripke, S. *et al.* Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511**, 421 (2014).
30. Ripke, S. *et al.* Genome-wide association analysis identifies 13 new risk loci for schizophrenia. *Nat Genet* **45**, 1150-9 (2013).
31. Psychiatric GWAS Consortium Bipolar Disorder Working Group. Large-scale genome-wide association analysis of bipolar disorder identifies a new susceptibility locus near ODZ4. *Nat Genet* **43**, 977-83 (2011).
32. Axelrud, L.K. *et al.* Polygenic Risk Score for Alzheimer's Disease: Implications for Memory Performance and Hippocampal Volumes in Early Life. *Am J Psychiatry* **175**, 555-563 (2018).
33. Mistry, S., Harrison, J.R., Smith, D.J., Escott-Price, V. & Zammit, S. The use of polygenic risk scores to identify phenotypes associated with genetic risk of bipolar disorder and depression: A systematic review. *J Affect Disord* **234**, 148-155 (2018).
34. Maier, R. *et al.* Joint analysis of psychiatric disorders increases accuracy of risk prediction for schizophrenia, bipolar disorder, and major depressive disorder. *Am J Hum Genet* **96**, 283-94 (2015).

35. Hudson, J.I., Hiripi, E., Pope, H.G., Jr. & Kessler, R.C. The prevalence and correlates of eating disorders in the National Comorbidity Survey Replication. *Biol Psychiatry* **61**, 348-58 (2007).
36. Swanson, S.A., Crow, S.J., Le Grange, D., Swendsen, J. & Merikangas, K.R. Prevalence and correlates of eating disorders in adolescents. Results from the national comorbidity survey replication adolescent supplement. *Arch Gen Psychiatry* **68**, 714-23 (2011).
37. Kinasz, K., Accurso, E.C., Kass, A.E. & Le Grange, D. Does Sex Matter in the Clinical Presentation of Eating Disorders in Youth? *J Adolesc Health* **58**, 410-416 (2016).
38. Voderholzer, U. *et al.* Are there differences between men and women in outcome of intensive inpatient treatment for anorexia nervosa? An analysis of routine data. *Eur Eat Disord Rev* **27**, 59-66 (2019).
39. El Ghoch, M., Calugi, S., Milanese, C., Bazzani, P.V. & Dalle Grave, R. Body composition in men with anorexia nervosa: Longitudinal study. *Int J Eat Disord* **50**, 856-860 (2017).
40. Kessler, R.C., Chiu, W.T., Demler, O., Merikangas, K.R. & Walters, E.E. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* **62**, 617-27 (2005).
41. Ruscio, A.M., Stein, D.J., Chiu, W.T. & Kessler, R.C. The epidemiology of obsessive-compulsive disorder in the National Comorbidity Survey Replication. *Mol Psychiatry* **15**, 53-63 (2010).
42. Torresan, R.C. *et al.* Symptom dimensions, clinical course and comorbidity in men and women with obsessive-compulsive disorder. *Psychiatry Res* **209**, 186-95 (2013).

43. Bipolar Disorder and Schizophrenia Working Group of the Psychiatric Genomics Consortium. Genomic Dissection of Bipolar Disorder and Schizophrenia, Including 28 Subphenotypes. *Cell* **173**, 1705-1715.e16 (2018).
44. Boyd, A. *et al.* Cohort Profile: the 'children of the 90s'--the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol* **42**, 111-27 (2013).
45. Paternoster, L. *et al.* Genome-wide association study of three-dimensional facial morphology identifies a variant in PAX3 associated with nasion position. *Am J Hum Genet* **90**, 478-85 (2012).
46. Martin, J., Hamshere, M.L., Stergiakouli, E., O'Donovan, M.C. & Thapar, A. Genetic risk for attention-deficit/hyperactivity disorder contributes to neurodevelopmental traits in the general population. *Biol Psychiatry* **76**, 664-71 (2014).
47. Kann, L. *et al.* Youth risk behavior surveillance--United States, 1995. *J Sch Health* **66**, 365-77 (1996).
48. Field, A.E., Taylor, C.B., Celio, A. & Colditz, G.A. Comparison of self-report to interview assessment of bulimic behaviors among preadolescent and adolescent girls and boys. *Int J Eat Disord* **35**, 86-92 (2004).
49. Micali, N., Daniel, R.M., Ploubidis, G.B. & De Stavola, B.L. Maternal Prepregnancy Weight Status and Adolescent Eating Disorder Behaviors: A Longitudinal Study of Risk Pathways. *Epidemiology* **29**, 579-589 (2018).
50. Micali, N. *et al.* Adolescent eating disorder behaviours and cognitions: gender-specific effects of child, maternal and family risk factors. *Br J Psychiatry* **207**, 320-7 (2015).

51. Goodman, R., Ford, T., Richards, H., Gatward, R. & Meltzer, H. The Development and Well-Being Assessment: description and initial validation of an integrated assessment of child and adolescent psychopathology. *J Child Psychol Psychiatry* **41**, 645-55 (2000).
52. Goodman, A., Heiervang, E., Collishaw, S. & Goodman, R. The 'DAWBA bands' as an ordered-categorical measure of child mental health: description and validation in British and Norwegian samples. *Soc Psychiatry Psychiatr Epidemiol* **46**, 521-32 (2011).
53. Choi, S.W. & O'Reilly, P.F. PRSice-2: Polygenic Risk Score software for biobank-scale data. *Gigascience* **8**(2019).
54. Drummond, L.M. & Kolb, P. Obsessive–Compulsive Contamination Fears and Anorexia Nervosa: The Application of the New Psycho-Educational Treatment of Danger Ideation Reduction Therapy (DIRT). *Behav Change* **25**, 44-50 (2008).
55. Dar, K.A. & Iqbal, N. Worry and Rumination in Generalized Anxiety Disorder and Obsessive Compulsive Disorder. *J Psychol* **149**, 866-80 (2015).
56. Thomas, M. & Lovell, A. Anxiety and compulsion patterns in the maintenance of bingeing/purging behaviours by individuals with bulimia nervosa. *J Psychiatr Ment Health Nurs* **22**, 20-9 (2015).
57. Hübel, C. *et al.* Binge-eating disorder, anorexia nervosa, and constitutional thinness differ in their associations with anthropometric and psychiatric polygenic scores. *medRxiv* **2020.03.24.20042648**(2020).
58. Fairweather-Schmidt, A.K. & Wade, T.D. Changes in genetic and environmental influences on disordered eating between early and late adolescence: a longitudinal twin study. *Psychol Med* **45**, 3249-58 (2015).

59. O'Connor, S.M., Culbert, K.M., Mayhall, L.A., Burt, S.A. & Klump, K.L. Differences in genetic and environmental influences on body weight and shape concerns across pubertal development in females. *J Psychiatr Res* **121**, 39-46 (2020).
60. Klump, K.L. *et al.* The effects of puberty on genetic risk for disordered eating: evidence for a sex difference. *Psychol Med* **42**, 627-37 (2012).
61. Wade, T. *et al.* The structure of genetic and environmental risk factors for three measures of disordered eating. *Psychol Med* **29**, 925-34 (1999).
62. Dinkler, L. *et al.* Association of etiological factors across the extreme end and continuous variation in disordered eating in female Swedish twins. *Psychol Med*, 1-11 (2019).
63. Wade, T.D. *et al.* Genetic variants associated with disordered eating. *Int J Eat Disord* **46**, 594-608 (2013).
64. Munn-Chernoff, M.A. *et al.* Shared Genetic Risk between Eating Disorder- and Substance-Use-Related Phenotypes: Evidence from Genome-Wide Association Studies. *bioRxiv*, 741512 (2019).
65. Bulik, C.M., Sullivan, P.F., Fear, J.L. & Joyce, P.R. Eating disorders and antecedent anxiety disorders: a controlled study. *Acta Psychiatr Scand* **96**, 101-7 (1997).
66. Silberg, J.L. & Bulik, C.M. The developmental association between eating disorders symptoms and symptoms of depression and anxiety in juvenile twin girls. *J Child Psychol Psychiatry* **46**, 1317-26 (2005).
67. Davis, C. & Kaptein, S. Anorexia nervosa with excessive exercise: a phenotype with close links to obsessive-compulsive disorder. *Psychiatry Res* **142**, 209-17 (2006).
68. Błachno, M. *et al.* Obsessive-compulsive symptoms and physical activity in patients with anorexia nervosa—possible relationships. *Psychiatria Polska* **50**, 55-64 (2016).

69. Davis, C. & Claridge, G. The eating disorders as addiction: a psychobiological perspective. *Addict Behav* **23**, 463-75 (1998).
70. Davis, C., Katzman, D.K. & Kirsh, C. Compulsive physical activity in adolescents with anorexia nervosa: a psychobehavioral spiral of pathology. *J Nerv Ment Dis* **187**, 336-42 (1999).
71. Naylor, H., Mountford, V. & Brown, G. Beliefs about excessive exercise in eating disorders: the role of obsessions and compulsions. *Eur Eat Disord Rev* **19**, 226-36 (2011).
72. Dalle Grave, R., Calugi, S. & Marchesini, G. Compulsive exercise to control shape or weight in eating disorders: prevalence, associated features, and treatment outcome. *Compr Psychiatry* **49**, 346-52 (2008).
73. Carter, J.C., Blackmore, E., Sutandar-Pinnock, K. & Woodside, D.B. Relapse in anorexia nervosa: a survival analysis. *Psychol Med* **34**, 671-9 (2004).
74. Kaye, W.H., Gwirtsman, H.E., Obarzanek, E. & George, D.T. Relative importance of calorie intake needed to gain weight and level of physical activity in anorexia nervosa. *Am J Clin Nutr* **47**, 989-94 (1988).
75. Klump, K.L., Burt, S.A., McGue, M. & Iacono, W.G. Changes in genetic and environmental influences on disordered eating across adolescence: a longitudinal twin study. *Arch Gen Psychiatry* **64**, 1409-15 (2007).

Table 1. Eating disorder, obsessive-compulsive disorder, and anxiety diagnostic and symptom-based constructs.

Eating Disorder			
Age(s)	Construct	Scale	Report
10	Body image distortion	Stunkard Figure Rating Scale	Self
14 and/or 16	Anorexia nervosa	Youth Risk Behavior Surveillance System Questionnaire	Self
	Bulimia nervosa or subthreshold bulimia nervosa		
	Binge-eating disorder or subthreshold binge-eating disorder		
	Eating disorders not otherwise specified or purging disorder		
	Any threshold/subthreshold eating disorder		
	Fasting		
	Purging		
	Binge eating		
	Fear of weight gain		
	Compulsive exercise		
	Pressure to lose weight	Perceived Sociocultural Pressure Scale	Self
	Restraint	Dutch Eating Behavior Questionnaire	Self
	Emotional eating		
	External eating		
	Thin-ideal internalization	Ideal-Body Stereotype Scale-Revised	Self
Body dissatisfaction	Perceived Sociocultural Pressure Scale	Self	
Weight and shape concern	McKnight Risk Factor Survey	Self	
Obsessive-Compulsive Disorder			
10 & 13	Obsessive-compulsive disorder	Development and Wellbeing Assessment	Parent
	Obsessive-compulsive disorder latent factor symmetry		
	Obsessive-compulsive disorder latent factor dirt/germs		
Anxiety			
Age(s)	Construct	Scale	Report
7	Separation anxiety	Development and Wellbeing Assessment	Parent
10 & 13	Social phobia		
	Latent factor physical anxiety		
	Latent factor worrying		
	Latent factor social phobia		
13	Generalized anxiety disorder	Development and Wellbeing Assessment	Parent
15		Development and Wellbeing Assessment	Self

Table 2. Prediction of eating disorder, obsessive-compulsive disorder, and anxiety symptom dimensions and diagnoses using polygenic scores in all participants^a

Eating Disorder Symptom Dimensions and Diagnoses							
Phenotype	Sample size	PGS	P_T	N SNPs	R^2	β (SE)	Empirical P
Body image distortion at age 10	4758	AN	0.4	57861	0.0002	-0.0084 (0.0098)	0.7889
		OCD	0.2	38302	0.0004	-0.0139 (0.0097)	0.4322
		AN/OCD	0.05	18540	0.0001	-0.0069 (0.0098)	0.8724
AN at age 14 ^b	106 cases, 4129 controls	AN	0.05	15063	0.0044	0.1948 (0.0986)	0.1545
		OCD	0.001	601	0.0019	0.1271 (0.0979)	0.5193
		AN/OCD	0.3	54054	0.0017	0.1223 (0.0992)	0.5160
Bulimia nervosa or subthreshold bulimia nervosa at age 14 ^b	63 cases, 4172 controls	AN	0.001	1081	0.0012	0.1080 (0.1257)	0.7847
		OCD	0.001	601	0.0012	-0.1095 (0.1279)	0.8268
		AN/OCD	0.001	1388	0.0026	0.1596 (0.1272)	0.5025
Binge-eating disorder or subthreshold binge-eating disorder at age 14 ^{b,c}	25 cases, 4210 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Eating disorders not otherwise specified or purging disorder at age 14 ^b	593 cases, 3642 controls	AN	0.01	5029	0.0012	0.0755 (0.0445)	0.2588
		OCD	0.001	601	0.0004	-0.0418 (0.0444)	0.7729
		AN/OCD	0.05	18540	0.0010	0.0694 (0.0445)	0.3181
Any threshold/subthreshold eating disorder at age 14 ^b	787 cases, 3448 controls	AN	0.01	5029	0.0015	0.0780 (0.0396)	0.1471
		OCD	0.01	3915	0.0001	-0.0229 (0.0396)	0.9556
		AN/OCD	0.01	6379	0.0013	0.0737 (0.0398)	0.1775
Fasting at age 14 ^b	271 cases, 3861 controls	AN	0.3	48847	0.0006	0.0633 (0.0641)	0.7006
		OCD	0.4	60239	0.0007	-0.0653 (0.0628)	0.7066
		AN/OCD	0.5	68009	0.0009	0.0761 (0.0640)	0.5449
Purging at age 14	4132	AN	0.05	15063	0.0006	0.0063 (0.0040)	0.3280
		OCD	0.001	601	0.0008	-0.0071 (0.0040)	0.2441
		AN/OCD	0.1	28747	0.0007	0.0070 (0.0040)	0.2333
Binge eating at age 14	4167	AN	0.01	5029	0.0001	0.0050 (0.0064)	0.8335
		OCD	0.1	23472	0.0007	0.0109 (0.0064)	0.2751
		AN/OCD	0.001	1388	0.0002	0.0062 (0.0064)	0.6942
Fear of weight gain at age 14	4127	AN	0.01	5029	0.0003	0.0128 (0.0107)	0.5470
		OCD	0.05	13919	0.0002	0.0092 (0.0107)	0.8251
		AN/OCD	1	82076	0.0002	0.0087 (0.0108)	0.8113
Compulsive exercise at age 14	4027	AN	0.05	15063	0.0015	0.0247 (0.0099)	0.0448*
		OCD	0.001	601	0.0004	-0.0131 (0.0099)	0.5130
		AN/OCD	0.01	6379	0.0018	0.0265 (0.0100)	0.0297*
Pressure to lose weight at age 14	4116	AN	0.001	1081	0.0009	0.0524 (0.0279)	0.1802
		OCD	0.001	601	0.0007	-0.0494 (0.0281)	0.2460
		AN/OCD	0.001	1388	0.0010	0.0573 (0.0283)	0.1322
Restraint at age 14	4074	AN	0.3	48847	0.0014	0.0434 (0.0179)	0.0557
		OCD	0.05	13919	0.0002	0.0140 (0.0178)	0.8611
		AN/OCD	0.2	43351	0.0020	0.0518 (0.0179)	0.0147*
Emotional eating at age 14	3927	AN	0.001	1081	0.0001	0.0515 (0.0888)	0.9393
		OCD	0.001	601	0.0004	-0.1166 (0.0890)	0.5123
		AN/OCD	0.05	18540	0.0001	0.0552 (0.0893)	0.9126

External eating at age 14	3608	AN	0.001	1081	0.0001	0.0403 (0.0551)	0.8617
		OCD	0.05	13919	0.0005	0.0775 (0.0554)	0.4512
		AN/OCD	0.001	1388	0.0003	0.0593 (0.0558)	0.6353
Thin ideal internalization at age 14	4053	AN	0.001	1081	0.0010	0.0864 (0.0424)	0.1351
		OCD	0.1	23472	0.0008	0.0784 (0.0426)	0.2164
		AN/OCD	0.001	1388	0.0006	0.0659 (0.0430)	0.3325
Body dissatisfaction at age 14	4169	AN	0.1	24182	0.0015	0.3044 (0.1204)	0.0410⁺
		OCD	0.001	601	0.0003	−0.1294 (0.1202)	0.6720
		AN/OCD	0.2	43351	0.0018	0.3329 (0.1209)	0.0200⁺
Weight and shape concern at age 14	4164	AN	0.001	1081	0.0002	0.0269 (0.0284)	0.7268
		OCD	0.05	13919	0.0002	0.0248 (0.0285)	0.8105
		AN/OCD	1	82076	0.0002	0.0268 (0.0289)	0.7250
AN at age 16 ^b	71 cases, 3511 controls	AN	0.001	1081	0.0060	−0.2354 (0.1213)	0.1643
		OCD	0.01	3915	0.0008	0.0824 (0.1182)	0.9028
		AN/OCD	0.1	28747	0.0037	−0.1825 (0.1189)	0.3203
Bulimia nervosa at age 16 ^b	146 cases, 3436 controls	AN	0.001	1081	0.0004	0.0555 (0.0845)	0.9059
		OCD	0.1	23472	0.0005	0.0629 (0.0843)	0.8835
		AN/OCD	0.001	1388	0.0017	0.1120 (0.0839)	0.4432
Binge-eating disorder at age 16 ^b	58 cases, 3524 controls	AN	0.3	48847	0.0026	0.1583 (0.1336)	0.5600
		OCD	0.1	23472	0.0033	0.1779 (0.1320)	0.4711
		AN/OCD	0.001	1388	0.0034	0.1806 (0.1315)	0.4135
Eating disorders not otherwise specified or purging disorder at age 16 ^b	1079 cases, 2503 controls	AN	0.001	1081	0.0004	−0.0370 (0.0365)	0.6728
		OCD	0.05	13919	0.0026	0.0935 (0.0366)	0.0410⁺
		AN/OCD	0.05	18540	0.0001	−0.0157 (0.0363)	0.9776
Any threshold/subthreshold eating disorder at age 16 ^b	1354 cases, 2228 controls	AN	0.001	1081	0.0005	−0.0384 (0.0345)	0.6112
		OCD	0.1	23472	0.0035	0.1045 (0.0345)	0.0111⁺
		AN/OCD	0.2	43351	< 0.0001	0.0068 (0.0343)	0.9993
Fasting at age 16	3379	AN	0.001	1081	0.0002	−0.0101 (0.0114)	0.7675
		OCD	0.05	13919	0.0014	0.0243 (0.0112)	0.1070
		AN/OCD	1	82076	0.0001	−0.0075 (0.0113)	0.8968
Purging at age 16	3402	AN	0.1	24182	0.0008	0.0142 (0.0085)	0.2536
		OCD	0.01	3915	0.0007	0.0125 (0.0084)	0.3789
		AN/OCD	0.05	18540	0.0012	0.0176 (0.0084)	0.1045
Binge eating at age 16	2929	AN	0.3	48847	0.0002	0.0102 (0.0122)	0.7994
		OCD	0.1	23472	0.0011	0.0219 (0.0120)	0.2197
		AN/OCD	0.001	1388	0.0012	0.0230 (0.0122)	0.1771
Compulsive exercise at age 16	3186	AN	0.1	24182	0.0002	0.0147 (0.0172)	0.7860
		OCD	0.05	13919	0.0034	0.0555 (0.0170)	0.0042⁺
		AN/OCD	0.4	61888	0.0006	0.0227 (0.0171)	0.4465
Obsessive-Compulsive Disorder Symptom Dimensions and Diagnosis							
Phenotype	Sample size	PGS	P _T	N SNPs	R ²	β (SE)	Empirical P
OCD at age 10 ^{b,c}	18 cases, 5198 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
OCD latent factor – symmetry, checking at age 10	5197	AN	0.001	1081	0.0009	0.0151 (0.0069)	0.0949
		OCD	0.2	38302	0.0001	−0.0060 (0.0070)	0.8131
		AN/OCD	0.2	43351	0.0007	0.0132 (0.0070)	0.1781

OCD latent factor – dirt/germs at age 10	5197	AN	0.001	1081	0.0013	0.0157 (0.0060)	0.0319[*]
		OCD	0.05	13919	0.0001	0.0032 (0.0060)	0.9645
		AN/OCD	0.2	43351	0.0002	0.0068 (0.0060)	0.5745
OCD at age 13 ^{b,c}	10 cases 4841 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
OCD latent factor – symmetry, checking at age 13	4832	AN	1	85364	0.0003	0.0125 (0.0109)	0.5815
		OCD	0.001	601	0.0002	−0.0106 (0.0107)	0.7313
		AN/OCD	0.01	6379	< 0.0001	0.0052 (0.0109)	0.9669
OCD latent factor – dirt/germs at age 13	4832	AN	0.05	15063	0.0001	−0.0064 (0.0093)	0.8866
		OCD	0.05	13919	0.0009	−0.0196 (0.0092)	0.1208
		AN/OCD	0.001	1388	< 0.0001	−0.0045 (0.0093)	0.9646
Anxiety Symptom Dimensions and Diagnoses							
Phenotype	Sample size	PGS	P_T	N SNPs	R^2	β (SE)	Empirical P
Separation anxiety at age 7 ^b	148 cases, 5214 controls	AN	0.3	48847	0.0072	0.2475 (0.0841)	0.0132[*]
		OCD	0.05	13919	0.0001	0.0308 (0.0831)	0.9954
		AN/OCD	0.2	43351	0.0039	0.1812 (0.0843)	0.0982
Specific phobia at age 7 ^b	104 cases, 5265 controls	AN	0.01	5029	0.0063	0.2432 (0.1001)	0.0501
		OCD	0.001	601	0.0010	−0.0955 (0.0988)	0.7517
		AN/OCD	0.01	6379	0.0085	0.2809 (0.0999)	0.0181⁺
Latent factor – physical anxiety at age 10	5197	AN	0.01	5029	0.0009	0.0221 (0.0102)	0.0945
		OCD	0.2	38302	0.0001	−0.0065 (0.0102)	0.9302
		AN/OCD	0.1	28747	0.0007	0.0195 (0.0102)	0.1675
Latent factor – worrying at age 10	5197	AN	0.01	5029	0.0016	0.0283 (0.0099)	0.0163[*]
		OCD	0.1	23472	0.0003	0.0131 (0.0099)	0.5107
		AN/OCD	0.1	28747	0.0004	0.0150 (0.0099)	0.3376
Latent factor – social phobia at age 10	5197	AN	0.001	1081	0.0005	0.0140 (0.0085)	0.2810
		OCD	0.01	3915	0.0009	−0.0184 (0.0085)	0.1102
		AN/OCD	0.01	6379	0.0002	0.0083 (0.0086)	0.7001
Social phobia at age 13 ^b	55 cases, 4795 controls	AN	0.3	48847	0.0015	0.1254 (0.1366)	0.7521
		OCD	0.01	3915	0.0088	0.3036 (0.1359)	0.0959
		AN/OCD	0.001	1388	0.0015	0.1258 (0.1371)	0.7412
Generalized anxiety disorder at age 13 ^b	154 cases, 4687 controls	AN	0.01	5029	0.0029	0.1544 (0.0826)	0.1916
		OCD	1	91297	0.0019	0.1248 (0.0823)	0.3813
		AN/OCD	0.01	6379	0.0025	0.1422 (0.0830)	0.2363
Latent factor – physical anxiety at age 13	4832	AN	1	85364	0.0003	−0.0118 (0.0096)	0.5315
		OCD	0.01	3915	0.0001	−0.0067 (0.0095)	0.9022
		AN/OCD	0.2	43351	0.0002	−0.0098 (0.0096)	0.6668
Latent factor – worrying at age 13	4832	AN	0.3	48847	0.0006	0.0197 (0.0114)	0.2414
		OCD	0.001	601	0.0003	−0.0131 (0.0112)	0.6147
		AN/OCD	0.2	43351	0.0003	0.0134 (0.0114)	0.5571
Latent factor – social phobia at age 13	4832	AN	0.3	48847	0.0014	0.0278 (0.0108)	0.0380⁺
		OCD	0.001	601	0.0005	−0.0168 (0.0107)	0.3497
		AN/OCD	0.2	43351	0.0006	0.0179 (0.0108)	0.2637
Generalized anxiety disorder at age 15 ^b	196 cases, 3712 controls	AN	0.001	1081	0.0019	0.1136 (0.0723)	0.3184
		OCD	0.1	23472	0.0006	0.0657 (0.0738)	0.8034
		AN/OCD	0.001	1388	0.0003	0.0453 (0.0742)	0.9199

^a Principal components 1–5 were used as covariates.

^b Binary phenotype.

^c Due to insufficient statistical power, any binary measure with less than 50 cases is not included in the final analysis.

* (also bolded) Statistically significant after P for regression was corrected for multiple testing via 10,000 permutations (significance threshold=0.05).

Abbreviations: PGS=polygenic score; P_T =best discovery P threshold; SNP=single nucleotide polymorphism; R^2 =Nagelkerke's pseudo- R^2 for total variance explained in target phenotype; β =standardized beta regression coefficient; SE =standard error; AN=anorexia nervosa; OCD=obsessive-compulsive disorder; AN/OCD=anorexia nervosa/obsessive-compulsive transdiagnostic phenotype.

Table 3. Prediction of eating disorder, obsessive-compulsive disorder, and anxiety symptom dimensions and diagnoses using polygenic scores in girls^a

Eating Disorder Symptom Dimensions and Diagnoses							
Phenotype	Sample size	PGS	P_T	N SNPs	R^2	β (SE)	Empirical P
Body image distortion at age 10	2512	AN	0.001	1079	0.0005	-0.0158 (0.0135)	0.5689
		OCD	0.001	602	0.0006	0.0165 (0.0135)	0.5702
		AN/OCD	0.001	1385	0.0002	-0.0095 (0.0136)	0.8783
AN at age 14 ^b	78 cases, 2242 controls	AN	0.05	15037	0.0062	0.2254 (0.1173)	0.1731
		OCD	0.001	602	0.0009	0.0863 (0.1149)	0.8775
		AN/OCD	0.001	1385	0.0059	0.2167 (0.1163)	0.1865
Bulimia nervosa or subthreshold bulimia nervosa at age 14 ^{b,c}	44 cases, 2276 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Binge-eating disorder or subthreshold binge-eating disorder at age 14 ^{b,c}	17 cases, 2303 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Eating disorders not otherwise specified or purging disorder at age 14 ^b	419 cases, 1901 controls	AN	0.3	48846	0.0007	0.0531 (0.0543)	0.7027
		OCD	0.01	3917	0.0011	-0.0680 (0.0545)	0.5441
		AN/OCD	0.4	61950	0.0014	0.0754 (0.0542)	0.4122
Any threshold/subthreshold eating disorder at age 14 ^b	558 cases, 1762 controls	AN	0.05	15037	0.0006	0.0490 (0.0492)	0.6948
		OCD	0.01	3917	0.0002	-0.0294 (0.0486)	0.9452
		AN/OCD	0.3	54071	0.0017	0.0801 (0.0488)	0.2844
Fasting at age 14 ^b	227 cases, 2047 controls	AN	1	85306	0.0015	0.0890 (0.0706)	0.4959
		OCD	0.5	68461	0.0005	-0.0533 (0.0697)	0.8726
		AN/OCD	0.5	68072	0.0024	0.1145 (0.0709)	0.2870
Purging at age 14	2278	AN	0.4	57808	0.0013	0.0116 (0.0067)	0.2441
		OCD	0.001	602	0.0011	-0.0106 (0.0067)	0.3376
		AN/OCD	0.05	18537	0.0016	0.0129 (0.0067)	0.1606
Binge eating at age 14	2289	AN	0.05	15037	0.0008	-0.0124 (0.0094)	0.4818
		OCD	0.1	23463	0.0010	0.0143 (0.0093)	0.3599
		AN/OCD	0.05	18537	0.0004	-0.0095 (0.0094)	0.6817
Fear of weight gain at age 14	2277	AN	0.4	57808	0.0003	0.0124 (0.0161)	0.8368
		OCD	0.001	602	0.0011	0.0251 (0.0161)	0.3431
		AN/OCD	0.5	68072	0.0003	0.0134 (0.0161)	0.7958
Compulsive exercise at age 14	2240	AN	0.05	15037	0.0016	0.0266 (0.0142)	0.1809
		OCD	0.001	602	0.0001	-0.0053 (0.0141)	0.9943
		AN/OCD	0.01	6370	0.0021	0.0306 (0.0141)	0.0955
Pressure to lose weight at age 14	2263	AN	0.001	1079	0.0003	0.0334 (0.0420)	0.8185
		OCD	0.3	50361	0.0001	0.0205 (0.0422)	0.9768
		AN/OCD	0.001	1385	0.0005	0.0454 (0.0422)	0.6297
Restraint at age 14	2257	AN	0.3	48846	0.0011	0.0423 (0.0265)	0.3096
		OCD	0.001	602	0.0002	0.0194 (0.0263)	0.8930
		AN/OCD	0.1	28752	0.0015	0.0489 (0.0265)	0.1852
Emotional eating at age 14	2167	AN	0.4	57808	0.0006	-0.1482 (0.1276)	0.5808
		OCD	0.3	50361	0.0015	0.2301 (0.1261)	0.2207
		AN/OCD	0.4	61950	0.0005	-0.1370 (0.1276)	0.6266

External eating at age 14	2068	AN	0.05	15037	0.0008	−0.0926 (0.0722)	0.4879
		OCD	0.1	23463	0.0007	0.0848 (0.0711)	0.5969
		AN/OCD	0.2	43327	0.0001	0.0277 (0.0724)	0.9895
Thin ideal internalization at age 14	2283	AN	0.001	1079	0.0018	0.1010 (0.0495)	0.1303
		OCD	0.1	23463	0.0030	0.1287 (0.0492)	0.0361*
		AN/OCD	0.001	1385	0.0034	0.1394 (0.0497)	0.0185*
Body dissatisfaction at age 14	2297	AN	0.2	37940	0.0020	0.3528 (0.1654)	0.1050
		OCD	0.05	13923	0.0006	0.1886 (0.1640)	0.6161
		AN/OCD	0.2	43327	0.0020	0.3521 (0.1650)	0.1015
Weight and shape concern at age 14	2299	AN	0.1	24166	0.0001	−0.0233 (0.0412)	0.9453
		OCD	0.001	602	0.0008	0.0562 (0.0410)	0.4725
		AN/OCD	1	82082	0.0001	0.0216 (0.0409)	0.9551
AN at age 16 ^b	56 cases, 2035 controls	AN	0.001	1079	0.0021	−0.1342 (0.1380)	0.6993
		OCD	0.05	13923	0.0017	0.1183 (0.1356)	0.8054
		AN/OCD	1	82082	0.0020	−0.1295 (0.1348)	0.7006
Bulimia nervosa at age 16 ^b	121 cases, 1970 controls	AN	0.01	5029	0.0012	−0.0908 (0.0943)	0.7122
		OCD	0.1	23463	0.0017	0.1057 (0.0930)	0.6295
		AN/OCD	0.001	1385	0.0018	0.1089 (0.0932)	0.5619
Binge-eating disorder at age 16 ^{b,c}	48 cases, 2043 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Eating disorders not otherwise specified or purging disorder at age 16 ^b	843 cases, 1248 controls	AN	0.001	1079	0.0003	−0.0326 (0.0449)	0.8643
		OCD	0.05	13923	0.0018	0.0736 (0.0447)	0.3027
		AN/OCD	0.01	6370	0.0003	−0.0288 (0.0448)	0.9038
Any threshold/subthreshold eating disorder at age 16 ^b	1068 cases, 1023 controls	AN	0.001	1079	0.0005	−0.0369 (0.0439)	0.7914
		OCD	0.1	23463	0.0049	0.1206 (0.0434)	0.0232*
		AN/OCD	0.3	54071	0.0003	−0.0306 (0.0431)	0.8700
Fasting at age 16	1972	AN	0.2	37940	0.0005	−0.0181 (0.0180)	0.6891
		OCD	0.05	13923	0.0025	0.0395 (0.0178)	0.0951
		AN/OCD	1	82082	0.0004	−0.0158 (0.0178)	0.7573
Purging at age 16	1990	AN	0.1	24166	0.0010	0.0196 (0.0139)	0.4144
		OCD	0.001	602	0.0009	0.0180 (0.0137)	0.5050
		AN/OCD	0.05	18537	0.0021	0.0281 (0.0138)	0.1312
Binge eating at age 16	1715	AN	0.3	48846	0.0003	0.0129 (0.0184)	0.8860
		OCD	0.1	23463	0.0043	0.0489 (0.0181)	0.0264*
		AN/OCD	0.001	1385	0.0023	0.0362 (0.0183)	0.1468
Compulsive exercise at age 16	1840	AN	1	85306	0.0004	0.0210 (0.0245)	0.7928
		OCD	0.1	23463	0.0056	0.0776 (0.0241)	0.0070*
		AN/OCD	0.4	61950	0.0010	0.0327 (0.0244)	0.4451
Obsessive-Compulsive Disorder Symptom Dimensions and Diagnosis							
Phenotype	Sample size	PGS	P _T	N SNPs	R ²	β (SE)	Empirical P
OCD at age 10 ^{b,c}	< 5 cases	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
OCD latent factor – symmetry, checking at age 10	2590	AN	0.01	5029	0.0008	0.0136 (0.0092)	0.3740
		OCD	0.05	13923	0.0006	0.0118 (0.0092)	0.5210
		AN/OCD	0.2	43327	0.0007	0.0125 (0.0093)	0.4549

OCD latent factor – dirt/germs at age 10	2590	AN	0.001	1079	0.0006	0.0099 (0.0081)	0.5269
		OCD	0.05	13923	0.0005	0.0094 (0.0081)	0.6212
		AN/OCD	0.2	43327	0.0003	0.0071 (0.0082)	0.7768
OCD at age 13 ^{b,c}	< 5 cases	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
OCD latent factor – symmetry, checking at age 13	2421	AN	0.05	15037	0.0011	−0.0250 (0.0150)	0.2739
		OCD	0.1	23463	0.0007	0.0196 (0.0148)	0.4988
		AN/OCD	0.01	6370	0.0003	−0.0127 (0.0151)	0.7972
OCD latent factor – dirt/germs at age 13	2421	AN	1	85306	0.0006	0.0160 (0.0128)	0.5192
		OCD	1	91289	0.0012	−0.0216 (0.0127)	0.2652
		AN/OCD	1	82082	0.0006	0.0161 (0.0129)	0.5044
Anxiety Symptom Dimensions and Diagnoses							
Phenotype	Sample size	PGS	P_T	N SNPs	R^2	β (SE)	Empirical P
Separation anxiety at age 7 ^b	68 cases, 2544 controls	AN	0.2	37940	0.0203	0.4229 (0.1256)	0.0035 ⁺
		OCD	1	91289	0.0018	0.1243 (0.1226)	0.7173
		AN/OCD	0.1	28752	0.0153	0.3655 (0.1251)	0.0134 ⁺
Specific phobia at age 7 ^b	53 cases, 2564 controls	AN	0.01	5029	0.0033	0.1746 (0.1396)	0.5141
		OCD	1	91289	0.0028	0.1590 (0.1374)	0.6183
		AN/OCD	0.01	6370	0.0138	0.3562 (0.1401)	0.0378 ⁺
Latent factor – physical anxiety at age 10	2590	AN	0.001	1079	0.0010	0.0234 (0.0143)	0.2819
		OCD	0.001	602	0.0012	−0.0254 (0.0145)	0.2500
		AN/OCD	0.3	54071	0.0009	0.0227 (0.0145)	0.3147
Latent factor – worrying at age 10	2590	AN	0.001	1079	0.0010	0.0228 (0.0144)	0.3139
		OCD	0.05	13923	0.0006	0.0181 (0.0145)	0.5495
		AN/OCD	1	82082	0.0003	0.0125 (0.0147)	0.7854
Latent factor – social phobia at age 10	2590	AN	0.01	5029	0.0003	0.0106 (0.0117)	0.7604
		OCD	0.001	602	0.0013	−0.0219 (0.0118)	0.2121
		AN/OCD	0.05	18537	0.0003	-0.0099 (0.0118)	0.8044
Social phobia at age 13 ^{b,c}	30 cases, 2396 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Generalized anxiety disorder at age 13 ^b	100 cases, 2317 controls	AN	0.001	1079	0.0039	0.1701 (0.1017)	0.2682
		OCD	1	91289	0.0044	0.1795 (0.1022)	0.2504
		AN/OCD	0.01	6370	0.0056	0.2078 (0.1044)	0.1474
Latent factor – physical anxiety at age 13	2421	AN	1	85306	0.0002	−0.0101 (0.0131)	0.8391
		OCD	0.001	602	0.0004	−0.0130 (0.0129)	0.7252
		AN/OCD	1	82082	0.0006	−0.0159 (0.0131)	0.5329
Latent factor – worrying at age 13	2421	AN	0.2	37940	0.0008	0.0227 (0.0166)	0.4388
		OCD	0.001	602	0.0029	−0.0430 (0.0161)	0.0307 ⁺
		AN/OCD	0.2	43327	0.0016	0.0329 (0.0165)	0.1440
Latent factor – social phobia at age 13	2421	AN	0.3	48846	0.0011	0.0259 (0.0155)	0.2680
		OCD	0.001	602	0.0039	−0.0471 (0.0152)	0.0072 ⁺
		AN/OCD	0.2	43327	0.0015	0.0295 (0.0156)	0.1769
Generalized anxiety disorder at age 15 ^{b,c}	154 cases, 1899 controls	AN	0.1	24166	0.0036	0.1486 (0.0849)	0.2354
		OCD	0.1	23463	0.0013	0.0880 (0.0840)	0.6895
		AN/OCD	0.1	28752	0.0028	0.1304 (0.0840)	0.3313

^a Principal components 1–5 were used as covariates.

^b Binary phenotype.

^c Due to insufficient statistical power, any binary measure with less than 50 cases is not included in the final analysis.

* (also bolded) Statistically significant after *P* for regression was corrected for multiple testing via 10,000 permutations (significance threshold=0.05).

Abbreviations: PGS=polygenic score; P_T =best discovery *P* threshold; SNP=single nucleotide polymorphism; R^2 =Nagelkerke's pseudo- R^2 for total variance explained in target phenotype; β =standardized beta regression coefficient; *SE*=standard error; AN=anorexia nervosa; OCD=obsessive-compulsive disorder; AN/OCD=anorexia nervosa/obsessive-compulsive transdiagnostic phenotype.

Table 4. Prediction of eating disorder, obsessive-compulsive disorder, and anxiety symptom dimensions and diagnoses using polygenic scores in boys^a

Eating Disorder Symptom Dimensions and Diagnoses							
Phenotype	Sample size	PGS	P_T	N SNPs	R^2	β (SE)	Empirical P
Body image distortion at age 10	2246	AN	0.001	1080	0.0005	0.0141 (0.0136)	0.6621
		OCD	0.05	13924	0.0028	-0.0342 (0.0136)	0.0500
		AN/OCD	0.4	61850	0.0001	-0.0053 (0.0138)	0.9868
AN at age 14 ^{b,c}	28 cases, 1887 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Bulimia nervosa or subthreshold bulimia nervosa at age 14 ^{b,c}	19 cases, 1896 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Binge-eating disorder or subthreshold binge-eating disorder at age 14 ^{b,c}	8 cases, 1907 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Eating disorders not otherwise specified or purging disorder at age 14 ^b	174 cases, 1741 controls	AN	0.001	1080	0.0075	0.2019 (0.0787)	0.0400[*]
		OCD	0.001	603	0.0006	-0.0595 (0.0803)	0.8915
		AN/OCD	0.001	1388	0.0031	0.1334 (0.0805)	0.2667
Any threshold/subthreshold eating disorder at age 14 ^b	229 cases, 1686 controls	AN	0.01	5032	0.0058	0.1701 (0.0710)	0.0614
		OCD	0.001	603	0.0001	-0.0218 (0.0710)	0.9973
		AN/OCD	0.01	6385	0.0024	0.1108 (0.0719)	0.3190
Fasting at age 14 ^{b,c}	44 cases, 1814 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Purging at age 14	1854	AN	0.01	5032	0.0001	0.0016 (0.0034)	0.9773
		OCD	0.1	23487	0.0011	0.0049 (0.0034)	0.4181
		AN/OCD	0.01	6385	0.0001	0.0015 (0.0034)	0.9777
Binge eating at age 14	1878	AN	0.2	37905	0.0014	0.0137 (0.0083)	0.2825
		OCD	1	91248	0.0006	0.0090 (0.0083)	0.6657
		AN/OCD	0.01	6385	0.0020	0.0160 (0.0083)	0.1631
Fear of weight gain at age 14	1850	AN	0.01	5032	0.0016	0.0193 (0.0113)	0.2553
		OCD	0.001	603	0.0007	-0.0133 (0.0114)	0.5988
		AN/OCD	0.001	1388	0.0009	0.0152 (0.0115)	0.4509
Compulsive exercise at age 14	1787	AN	0.001	1080	0.0046	0.0376 (0.0130)	0.0141[*]
		OCD	0.05	13924	0.0014	0.0203 (0.0131)	0.3586
		AN/OCD	0.001	1388	0.0026	0.0287 (0.0134)	0.0967
Pressure to lose weight at age 14	1853	AN	0.001	1080	0.0034	0.0804 (0.0321)	0.0432[*]
		OCD	0.001	603	0.0058	-0.1069 (0.0324)	0.0051[*]
		AN/OCD	0.001	1388	0.0047	0.0974 (0.0329)	0.0122[*]
Restraint at age 14	1817	AN	0.2	37905	0.0038	0.0547 (0.0207)	0.0312[*]
		OCD	0.05	13924	0.0010	0.0279 (0.0206)	0.4822
		AN/OCD	0.3	54002	0.0046	0.0605 (0.0208)	0.0148[*]
Emotional eating at age 14	1760	AN	0.5	65261	0.0028	0.2485 (0.1119)	0.0890
		OCD	0.001	603	0.0017	-0.1911 (0.1103)	0.2695
		AN/OCD	0.5	67970	0.00191	0.2054 (0.1119)	0.1857

External eating at age 14	1540	AN	0.001	1080	0.0006	0.0854 (0.0863)	0.6987
		OCD	0.01	3913	0.0014	0.1299 (0.0877)	0.3993
		AN/OCD	0.001	1388	0.0007	0.0949 (0.0887)	0.6259
Thin ideal internalization at age 14	1770	AN	0.001	1080	0.0006	0.0736 (0.0699)	0.6471
		OCD	0.001	603	0.0017	−0.1216 (0.0707)	0.2694
		AN/OCD	0.01	6385	0.0011	−0.1013 (0.0711)	0.3837
Body dissatisfaction at age 14	1872	AN	0.01	5032	0.0031	0.3898 (0.1616)	0.0534
		OCD	0.001	603	0.0015	−0.273 (0.1629)	0.2809
		AN/OCD	0.3	54002	0.0022	0.3312 (0.1636)	0.1324
Weight and shape concern at age 14	1865	AN	0.01	5032	0.0018	0.0614 (0.0338)	0.2026
		OCD	0.001	603	0.0006	−0.0366 (0.0339)	0.6651
		AN/OCD	0.5	67970	0.0009	0.0442 (0.0344)	0.4775
AN at age 16 ^{b,c}	15 cases, 1476 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Bulimia nervosa at age 16 ^{b,c}	25 cases, 1466 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Binge-eating disorder at age 16 ^{b,c}	10 cases, 1481 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Eating disorders not otherwise specified or purging disorder at age 16 ^b	236 cases, 1255 controls	AN	0.2	37905	0.0009	0.0648 (0.0723)	0.7575
		OCD	0.1	23487	0.0074	0.1821 (0.0721)	0.0479*
		AN/OCD	0.001	1388	0.0008	0.0599 (0.0711)	0.7776
Any threshold/subthreshold eating disorder at age 16 ^b	286 cases, 1205 controls	AN	0.2	37905	0.0008	0.0596 (0.0670)	0.7633
		OCD	1	91248	0.0028	0.1060 (0.0659)	0.3226
		AN/OCD	0.2	43348	0.0012	0.0708 (0.0660)	0.6129
Fasting at age 16	1407	AN	0.2	37905	0.0017	0.0133 (0.0085)	0.3145
		OCD	0.001	603	0.0008	−0.0088 (0.0082)	0.6764
		AN/OCD	0.001	1388	0.0007	−0.0085 (0.0084)	0.6552
Purging at age 16	1412	AN	0.2	37905	0.0013	0.0063 (0.0047)	0.4416
		OCD	0.01	3913	0.0005	0.0040 (0.0046)	0.8002
		AN/OCD	0.01	6385	0.0009	0.0052 (0.0046)	0.5648
Binge eating at age 16	1214	AN	0.05	15049	0.0010	0.0143 (0.0128)	0.6055
		OCD	0.2	38311	0.0023	−0.0210 (0.0126)	0.2894
		AN/OCD	0.05	18541	0.0025	0.0223 (0.0127)	0.2211
Compulsive exercise at age 16	1346	AN	0.01	5032	0.0005	0.0154 (0.0193)	0.8224
		OCD	0.05	13924	0.0014	0.0268 (0.0195)	0.4685
		AN/OCD	0.001	1388	0.0008	0.0202 (0.0199)	0.6562
Obsessive-Compulsive Disorder Symptom Dimensions and Diagnosis							
Phenotype	Sample size	PGS	P _T	N SNPs	R ²	β (SE)	Empirical P
OCD at age 10 ^{b,c}	15 cases, 2610 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
OCD latent factor – symmetry, checking at age 10	2607	AN	0.3	48867	0.0015	0.0207 (0.0104)	0.1395
		OCD	0.1	23487	0.0005	−0.0123 (0.0105)	0.6101
		AN/OCD	0.1	28755	0.0013	0.0191 (0.0104)	0.1879

OCD latent factor – dirt/germs at age 10	2607	AN	0.001	1080	0.0023	0.0214 (0.0088)	0.0503
		OCD	0.01	3913	0.0002	−0.0055 (0.0088)	0.9378
		AN/OCD	0.05	18541	0.0007	0.0120 (0.0088)	0.4186
OCD at age 13 ^{b,c}	6 cases, 2419 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
OCD latent factor – symmetry, checking at age 13	2411	AN	0.05	15049	0.0018	0.0301 (0.0145)	0.1221
		OCD	0.05	13924	0.0015	−0.0275 (0.0144)	0.1961
		AN/OCD	0.01	6385	0.0011	0.0239 (0.0145)	0.2754
OCD latent factor – dirt/germs at age 13	2411	AN	0.1	24168	0.0005	−0.0144 (0.0132)	0.6217
		OCD	0.05	13924	0.0012	−0.0221 (0.0132)	0.2860
		AN/OCD	0.01	6385	0.0007	−0.0175 (0.0133)	0.4556
Anxiety Symptom Dimensions and Diagnoses							
Phenotype	Sample size	PGS	P_T	N SNPs	R^2	β (SE)	Empirical P
Separation anxiety at age 7 ^b	80 cases, 2670 controls	AN	0.1	24168	0.0025	0.1443 (0.1141)	0.5109
		OCD	0.2	38311	0.0028	−0.1515 (0.1138)	0.5005
		AN/OCD	0.001	1388	0.0037	0.1798 (0.1164)	0.3234
Specific phobia at age 7 ^b	51 cases, 2701 controls	AN	0.01	5032	0.0121	0.3405 (0.1434)	0.0567
		OCD	0.1	23487	0.0025	−0.1518 (0.1418)	0.6784
		AN/OCD	0.01	6385	0.0043	0.2010 (0.1427)	0.3931
Latent factor – physical anxiety at age 10	2607	AN	0.4	57882	0.0022	0.0341 (0.0143)	0.0626
		OCD	0.001	603	0.0005	0.0158 (0.0143)	0.6611
		AN/OCD	0.01	6385	0.0008	0.0203 (0.0145)	0.4087
Latent factor – worrying at age 10	2607	AN	0.5	65261	0.0034	0.0396 (0.0133)	0.0143*
		OCD	0.5	68431	0.0006	0.0163 (0.0132)	0.5582
		AN/OCD	0.1	28755	0.0011	0.0220 (0.0133)	0.2615
Latent factor – social phobia at age 10	2607	AN	0.001	1080	0.0011	0.0214 (0.0124)	0.2491
		OCD	0.01	3913	0.0021	−0.0288 (0.0124)	0.0768
		AN/OCD	0.001	1388	0.0008	0.0180 (0.0127)	0.3922
Social phobia at age 13 ^{b,c}	25 cases, 2399 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--
Generalized anxiety disorder at age 13 ^b	54 cases, 2370 controls	AN	0.05	15049	0.0060	0.2298 (0.1368)	0.2606
		OCD	0.01	3913	0.0064	0.2370 (0.1376)	0.2663
		AN/OCD	0.1	28755	0.0019	0.1277 (0.1371)	0.7181
Latent factor – physical anxiety at age 13	2411	AN	0.2	37905	0.0006	−0.0166 (0.0141)	0.5694
		OCD	0.01	3913	0.0005	−0.0154 (0.0139)	0.6580
		AN/OCD	0.001	1388	0.0004	0.0137 (0.0142)	0.7044
Latent factor – worrying at age 13	2411	AN	0.3	48867	0.0004	0.0159 (0.0154)	0.6721
		OCD	0.001	603	0.0003	0.0134 (0.0153)	0.8070
		AN/OCD	0.01	6385	0.0001	−0.0056 (0.0154)	0.9894
Latent factor – social phobia at age 13	2411	AN	0.3	48867	0.0015	0.0283 (0.0148)	0.1660
		OCD	0.05	13924	0.0007	−0.0189 (0.0146)	0.5196
		AN/OCD	0.1	28755	0.0001	0.0084 (0.0146)	0.9319
Generalized anxiety disorder at age 15 ^{b,c}	42 cases, 1813 controls	AN	--	--	--	--	--
		OCD	--	--	--	--	--
		AN/OCD	--	--	--	--	--

^a Principal components 1–5 were used as covariates.

^b Binary phenotype.

^c Due to insufficient statistical power, any binary measure with less than 50 cases is not included in the final analysis.

* (also bolded) Statistically significant after *P* for regression was corrected for multiple testing via 10,000 permutations (significance threshold=0.05).

Abbreviations: PGS=polygenic score; P_T =best discovery *P* threshold; SNP=single nucleotide polymorphism; R^2 =Nagelkerke's pseudo- R^2 for total variance explained in target phenotype; β =standardized beta regression coefficient; *SE*=standard error; AN=anorexia nervosa; OCD=obsessive-compulsive disorder; AN/OCD=anorexia nervosa/obsessive-compulsive transdiagnostic phenotype.

Figure 1. Significant results for the prediction of **(a)** eating disorder symptoms; **(b)** eating disorder diagnoses; **(c)** obsessive-compulsive disorder and anxiety-related symptoms; and **(d)** obsessive-compulsive disorder and anxiety disorder diagnoses by polygenic scores in all participants (combined and by sex) during different developmental timepoints. Significant findings (threshold set at $p > 0.05$ following 10,000 permutations per target phenotype) are indicated with an asterisk. The bars depict standardized beta regression coefficients, and the lines depict standard errors.