Health literacy in the general population in the context of epidemic or pandemic coronavirus outbreak situations: rapid scoping review

Uwe Mattered¹, Nina Egger³, Jana Tempes³, Christina Tischer¹, Jonas Lander², Marie-Luise Dierks², Eva-Maria Bitzer³, Christian Apfelbacher^{1,4}

¹Institute of Social Medicine and Health Systems Research, Medical Faculty, Otto von Guericke University, Magdeburg, Germany

²Institute for Epidemiology, Social Medicine and Health Systems Research, Hannover Medical School, Hannover, Germany

³Department of Public Health and Health Education, Freiburg University of Education, Freiburg, Germany

⁴Family Medicine and Primary Care, Lee Kong Chian School of Medicine, Nanyang Technological University Singapore, Singapore

Summary

BACKGROUND: Authorities responded with contact restrictions and other measures to the global spread of SARS-CoV-2. Health literacy (HL) has been linked to health outcomes and refers to the ability to access, understand, appraise and apply health information in order to make good health decisions. When restrictions are gradually lifted, individual HL becomes essential to control the pandemic and to prevent the resumption of these restriction, should infection numbers surge again. The aim of this rapid scoping review, for which only studies from the general population were considered, was to describe the extent of existing research on HL in the context of previous coronavirus outbreaks (SARS-CoV-1, MERS-CoV and SARS-CoV-2). Facets of HL that were of particular interest were: type of assessment of HL (theory-based versus proxy assessment; validated instrument versus ad hoc assessment), domains of HL, interventions aiming to improve HL during outbreak situations, and HL surveillance during outbreak.

METHODS: We searched two major databases and included publications of quantitative and qualitative studies in English and German on any type of research on the functional, critical and communicative domains of HL conducted in the context of the three outbreaks in the general population. We extracted and tabulated relevant data and narratively reported where and when the study was conducted, the design and method used, and how HL was measured.

FINDINGS: 72 studies were included. Three investigated HL or explicitly referred to the concept of HL, 14 were guided by health behaviour theory. We did not find any study designed to develop or psychometrically evaluate pandemic HL instruments, or relate pandemic or general HL to a pandemic outcome, or any controlled intervention study. Type of assessment of the domains of HL varied widely.

INTERPRETATION: Theory-driven observational studies as well as interventions, examining whether pandemic-related HL can be improved are needed. In addition, the development and validation of instruments that measure pandemic-related HL is desirable.

Introduction

In late 2019 an outbreak of a new viral disease occurred in Wuhan, China and later spread to almost all countries of the world¹. It is caused by a novel beta coronavirus, the Severe Acute Respiratory Syndrome – Coronavirus – 2 (SARS-CoV-2), which causes Coronavirus Disease (COVID) – 19². The clinical epidemiology of COVID-19 is currently being investigated intensely³. Course of disease may be very mild, asymptomatic to very severe with respiratory and systemic damage and requiring mechanical ventilation⁴. Responses of governments to the COVID-19 pandemic have been multi-faceted including outbreak management (suppression versus mitigation), provision of adequate clinical treatment facilities for severe cases and measures to alleviate the economic and psychosocial impact of the pandemic and the measures taken to manage it². Public health measures implemented in many countries across the globe encompass contact restrictions and physical distancing, hygiene rules (i.e. frequent and thorough handwashing or disinfection), mask wearing, eye protection and recommendations about how to sneeze and cough^{3,5}. Some of these measures, particularly contact restrictions, have been law enforced in many countries⁶. Relaxing regulations and re-organising social life requires people to voluntarily adhere to the named measures in order to avoid exponential growth of SARS-CoV-2 to reoccur. Further, people who contract SARS-CoV-2 need to know when and how to seek health care and/or be tested. Those who suffer from severe COVID-19 and survive will have to seek health care to mitigate the potentially long-lasting physical and psychological sequelae such as kidney damage⁷ or post-traumatic stress disorder⁸. In all these and other different scenarios, the concept of health literacy (HL) becomes a vital public health concept that is essential to counterpart on the individual level the social restrictions enforced by law. It “entails the motivation, knowledge and competencies to access, understand, appraise and apply health information in order to make judgements and take decisions in everyday life concerning healthcare, disease prevention and health promotion to maintain or improve quality of life throughout the course of life”⁹. When restrictions are gradually lifted, the role of individual level HL increases in order to prevent the resumption of these restriction, should infection numbers surge again.

In other words, what is necessary beyond governmental regulations and policy, is an increase in the levels of COVID-19 related health literacy^{10,11}. We not only need to monitor the pandemic’s epidemiology during the course of the pandemic including the creation of herd immunity but also HL and health behavioural responses related to the pandemic in the population¹¹. HL is considered a major determinant of a person’s health^{12,13}, a factor that contributes to health inequalities¹², and a person’s health behaviour, for instance, healthy diet adherence or non-smoking¹³ and health care utilisation¹⁴. There is evidence that lower HL is consistently associated with mortality¹⁴ or lower self-rated health status¹⁵.

Research suggests that adequate HL may not be as prevalent among populations as might be necessary in order to navigate the increasingly complex healthcare landscape^{13,14,16}. Synthesised evidence

suggests a relationship between levels of HL and infectious disease prevention in non-pandemic contexts. Inadequate HL was found to be associated with reduced adoption of protective behaviours such as vaccination uptake and poor understanding of antibiotics. Large research gaps were found in relation to infectious diseases with a high clinical and societal impact, such as tuberculosis and malaria¹⁷. For instance, it was emphasised that critical HL, which focuses on supporting effective political and social action, was not considered in any of the reviewed studies¹⁷. The strengths of this relationship may be exponentially higher under pandemic circumstances, but no synthesised information on this topic appears to exist to date. Further, the importance of individual HL in pandemic control has been emphasised more urgently^{10,11}.

Therefore, the aim of this rapid scoping review, for which only studies from the general population were considered, was to describe the extent of existing research on HL in the context of previous coronavirus outbreaks (SARS-CoV-1, MERS-CoV and SARS-CoV-2). Facets of HL that were of particular interest were: type of assessment of HL (theory-based versus proxy assessment; validated instrument versus ad hoc assessment), interventions aiming to improve HL during outbreak situations, or HL surveillance during outbreak.

Method

Overview

This scoping review was performed according to the methodological framework as outlined by Khalil et al.¹⁸. Their guidelines regarding scoping reviews build on the work of Arksey and O'Malley's five-stage scoping review framework¹⁹, complemented with the Joanna Briggs Institute methodology²⁰, in order to (1) identify the research questions, (2) identify relevant studies, (3) select studies, (4) chart the data, and (5) collate and summarise the data. A scoping review's objective is to identify the nature and extent of the existing evidence. Unlike other types of review, it does not endeavour to systematically evaluate the quality of available research, but rather seek to identify the contribution of existing literature to an area of interest²¹. Our methodology was also guided by the rapid review approach which inevitable uses less rigor as necessary in a traditional systematic review due to the need for production within a short time-frame using limited resources²². The protocol for this rapid review was registered at OSFREGISTRIES on 06/04/2020²³.

Search strategy, selection criteria, extraction strategy and data analysis

Two authors (UM, NE) ran the search strategy on PubMed (MEDLINE®) and PsycINFO® on 20th April 2020. Citations were downloaded to Citavi (Swiss Academic Software). We included publications in English and German of quantitative and qualitative studies. The same authors evaluated titles and abstracts excluding any irrelevant ones. Full texts of the remaining citations were obtained, and two authors (UM, NE) reviewed these, excluding any, which did not meet the inclusion criteria. Finally, reference lists of remaining papers were hand-searched for additional relevant studies.

We then compared results from full text screening; there were only minor discrepancies, which were resolved through discussion with the whole team. Data extraction was carried out by five authors (UM, NE, JT, CT, JL) in independent pairs of two. Consensus was achieved through discussion and arbitration within the team. The search strategy was informed by HL theory (derivation of search terms) and is displayed in appendix 1. Inclusion criteria were:

We included reports on any type of research on the functional, critical and communicative domains of HL²⁴ conducted in the context of SARS-CoV-2, SARS-CoV-1 and MERS-CoV in the general population. This was a rational decision as an initial search using HL as the chief search term in conjunction with the aforementioned coronavirus outbreaks resulted in very few hits. We used the following definitions / concepts of functional, communicative and critical HL: Functional HL is broadly compatible with the narrow definition of ‘health literacy’ which can be considered to consist of health-related knowledge, attitudes, motivation, behavioural intentions, personal skills, or self-efficacy²⁴. Communicative HL means to be able ...’to derive meaning from different forms of communication’..., while the ability to critically analyse information is referred to as critical HL²⁴.

The following data were extracted from the included studies: authors, publication year, country of study, type of epidemic or pandemic outbreak (SARS-CoV-2, SARS-CoV-1, MERS-CoV), participants (including sample size), design, method, and instruments, and measured constructs including how they were measured (only if applicable e.g. not in qualitative studies). Findings were synthesized quantitatively and narratively and reporting followed the guidelines as proposed in PRISMA-ScR²⁵. A critical appraisal of the quality of the included studies was not within the scope of this review. We do however, comment on major methodological issues regarding the studies.

There was no funding source for this study.

Results

The search in PubMed (MEDLINE®) and PsycInfo® yielded 3394 references, two were obtained from colleagues^{26,27}, leading to 2766 references after removal of duplicates. Title and abstract screening resulted in exclusion of 2652 articles. Full-texts of the remaining 114 references were assessed for eligibility leading to inclusion of 77 publications pertaining to 72 studies. Details of the selection stages are provided in Figure 1. SARS-CoV-2 was investigated in 10, MERS-CoV in 26 and SARS-CoV-1 in 36 studies. Only three studies investigated HL or explicitly referred to the concept of HL²⁷⁻²⁹. 14 studies were conducted in the context of health behaviour theory, seven of another theory (Appendix 2). All studies, while mainly not explicitly investigating HL, measured one or more components of HL (Appendix 2). Most studies were observational or short longitudinal (58 cross-sectional, eight pre-post) and six qualitative. All SARS-CoV-two studies were conducted during, of the MERS-CoV studies 27 during, one during (first wave), eight after, of the MERS-CoV studies 24 during, one after and one both during and after the pandemic/outbreak. 66 studies used questionnaires,

two focus group discussion, four other qualitative methods (e.g. interviews) for data collection. 49 studies studied convenience or opportunity samples and 23 representative samples drawn from general populations. Sample size ranged from 19 – 222,599 participants.

Functional, critical, communicative HL and individual behaviour in quantitative studies

Within the nine quantitative SARS-CoV-2 studies knowledge was measured in seven, attitude in seven, risk perceptions in four, SE in three, critical HL in five, communicative HL in three, Health-information seeking behaviour (HISB) in two, and behavioural aspects in four studies. Only one study²⁶ measured functional, critical and communicative HL as well as a behavioural outcome. All others assessed only some aspects of HL. While seven studies reported on knowledge, most studies asked only about knowledge of symptoms, no study undertook a comprehensive assessment covering a broad range of SARS-CoV-2 related knowledge. Within the knowledge domain, symptoms were most often assessed. (Table 1). Wearing a mask was the most frequently assessed behaviour (Table 1).

31 quantitative studies were conducted in the context of SARS-CoV-1. 25 measured knowledge, 28 attitude, 20 risk perception, eight SE, 11 critical HL, 12 communicative HL, and 18 behaviour. One study³⁰ reported all six HL aspects, the others one to five aspects. Within knowledge, transmission mode was most often measured. Although 25 studies reported knowledge assessment, most studies did not comprehensively assess knowledge (Table 1). Handwashing was the most frequently measured behaviour (Table 1).

Of the 26 quantitative MERS-CoV studies 21 measured knowledge, 17 attitude, 18 risk perceptions, 3 SE, 4 critical HL, 17 communicative HL, and 10 behaviour. Two studies assessed five of the six HL aspects^{31,32}, the remainder one to four. Within knowledge, transmission mode was most often assessed. Again, most studies did not comprehensively assess knowledge. Handwashing was the most frequently assessed behaviour (Table 1).

The reported measured depth within the domains of HL varied widely among the studies (results not shown). For instance, the number of knowledge components ranged from one to at least eight.

Health-information seeking behaviour (HISB) in quantitative studies

HISB was measured in two (SARS-CoV-2), four (SARS-CoV-1), two (MERS-CoV) quantitative studies.

Pandemic HL measurement and relationship to pandemic outcome (quantitative and qualitative studies)

Our search failed to come across any studies designed to develop or psychometrically evaluate pandemic HL instruments or relate pandemic or general HL to a pandemic outcome.

Other aspects of HL measurement in pandemic contexts (quantitative and qualitative studies)

The number of items per HL aspect varied widely (data not shown), hardly any study reported on psychometric properties, two studies from three publications^{28,33,34} were the notable exception (Appendix 3) and a clear distinction between knowledge, attitude, or risk perceptions was sometimes absent. For instance, perceived vulnerability was reported as an attitude³⁵.

Qualitative studies

Six studies explored domains of HL in the context of SARS-CoV-2, SARS-CoV-1, and MERS-CoV. One study³⁶ reported low risk perceptions and a lack of seeking relevant health information in relation to SARS-CoV-2. Two studies^{37,38} explored risk perceptions and preventive behaviour in relation to SARS-CoV-1, another³⁹ explored individual experiences during quarantine. One study reported low knowledge about SARS-CoV-1 and its prevention⁴⁰. Another study in the context of SARS-CoV-1 concluded that attitudes towards mask wearing had substantially changed in the post-SARS-CoV-1 period⁴¹.

Discussion

While individual HL is recognised as an increasingly important construct in public health⁴², it is of note that only three studies emerged from our extensive search, which explicitly referred to the construct of HL in the context of any of the three coronavirus outbreaks. One used the Newest Vital Sign (NVS), a test measuring nutrition label information processing ability²⁹, another study²⁸ administered a short form of the HLS-EU-Q47, an HL instrument rooted in testable theory⁴³ and the third²⁷ study used a version of the HLS-EU-Q47 adapted to SARS-CoV-2. However, the latter provided no evidence on the psychometric properties of the adapted instrument. Hence, at present there seems to be no tested instrument designed to measure coronavirus pandemic-related HL. There is, however, one HL instrument assessing print and multimedia literacy in respect to respiratory diseases⁴⁴.

Most of the other included studies were not theory-based. It is important to highlight that these studies did not purport to measure HL, but were included in this review because the search strategy was based on a pragmatic application of suggested HL components within domains^{24,45}. Of those that were theory-driven, the majority employed health-behaviour theory as conceptualised by social cognition models. There is substantial overlap between socio-cognitive predictors of health behaviour and HL. For instance, attitude and self-efficacy (defined as behavioural control) are part of the theory of planned behaviour⁴⁶, risk perceptions part of the health belief model⁴⁷ or knowledge part of protection motivation theory⁴⁸. Theory-based research allows the formulation of testable a priori hypotheses, and if necessary revision of the theory. Nonetheless, the measures obtained from those studies lacking an explicit theoretical foundation can be considered proxies of HL because they constitute or at least contribute to one or more HL domains.

While there appears to be no evidence linking validly measured (epidemic or pandemic) HL to coronavirus outbreak/pandemic outcomes there is evidence that HL can be linked to other epidemic outbreaks, e.g. the 2014–2016 Ebola epidemic outbreak in West Africa resulted among other factors from low health literacy⁴⁹. A Center for Disease Control and Prevention campaign, with input from partners, helped increase HL⁵⁰. HL has also been shown to be associated with health and health behaviour in general. Hence, one would expect that this association would hold in coronavirus outbreak situations.

Communicative HL included the measurement of access to different sources of information. Whether this had anything to do with better decisions about health in relation to any of the three outbreaks, remained unclear. Knowledge items were generally devised by the authors, and very few reported to have items checked against guidelines. This and the lack of an objective standard for cut-offs make knowledge assessment arbitrary as it cannot be established whether knowledge items reflect current and correct evidenced knowledge. Similarly, while risk perceptions generally pertain to perceptions of vulnerability/susceptibility to and severity of a disease, they were not always measured accordingly or sometimes subsumed under the term attitudes or knowledge.

We also observed very little evidence about the psychometric properties of instruments used to measure the socio-cognitive variables attitude, risk perceptions, and self-efficacy. It is desirable to know whether measures are reliable and valid, and sensitive to change if the aim is to reflect the effects of health literacy interventions by e.g. education (responsiveness).

Even if knowledge, attitudinal constructs, risk perceptions or self-efficacy were composed in a clear-cut and unequivocal way and psychometrically sound, uncertainty as to whether HL in its broader definition^{9,51} as a composite/compound construct was measured, would still prevail. HL was proposed to be a latent construct⁴³ thus indicators for its measurement are necessary. There is the need for the development of adequate measurement models.

The present review cannot ascertain, whether established instruments such as TOFHLA (Test of Functional Health Literacy)⁵³, or the broader dimension based instruments, for instance the HLQ (Health Literacy Questionnaire)⁵⁴ could be used to predict a pattern of association between HL and epidemic or pandemic outcomes (and antecedents such as favourable behaviours and practices), because no such investigations appear to have been carried out, yet. The study²⁸ that used a short form of the HLS-EU-Q did not investigate the relationship between HL and pandemic outcome/preventive behaviour but coping responses to the outbreak (depression, quality of life). Okan et al.²⁷ reported individuals' subjective perceptions about how well they could access, understand, appraise and apply information in the SARS-CoV-2 context but did not test the actual level of what these skills pertain to and whether they are related to better/more favourable behaviour/practices.

Further, it also not possible to state at present whether pandemic outbreaks require a specific HL instrument, that is able to explain variance in relevant behaviour and practices over and above that of general instruments (i.e. latent trait/construct measured by discrete manifest cognitive antecedents of behaviour).

In this rapid review, it was not possible to adhere to the methodological rigour that is expected from a standard scoping review. As this review was conducted as a scoping review, we did not look at the strengths of any reported associations between HL aspects and behavioural aspects. Further, it is beyond the scope of this review to assess the quality of the reviewed studies according to standard guidelines for observational studies.

At present HL in the context of coronavirus outbreaks is at an early stage to inform public health/educational strategies aimed at improving the public's HL in order to contain the spread of pandemics. One study²⁶ appears to be able to shed light on the question whether HL related aspects change over the course of the pandemic as its survey is conducted in weekly intervals.

We recommend future research be guided theory from HL research^{9,45} in the much needed work on HL in pandemic outbreak situations. Consequently, assessment of HL should be based on the ability to access, understand, critically appraise and eventually apply information to make better choices about one's health in pandemic outbreak situations when viewed as a set of meta-cognitive skills or a latent trait⁴³. Nevertheless, operationalisations at the manifest level, for example, knowledge, or attitudes (which influence critical appraisal) need to be considered, as latent constructs cannot be directly measured. Nevertheless, in the interim, public health communication could benefit from what is generally known from HL research. Health information should be clear so that all members of the public can access needed health information for routine and critical decisions⁵⁵.

Beside theory-driven observational studies, we also need interventions, examining whether coronavirus pandemic-related HL can be improved. In addition, research should also attempt to develop HL instruments that measure coronavirus pandemic-related HL and test the reliability, validity and responsiveness to change. The latter is of particular importance, if we want to be able to examine change during the stages of a pandemic.

Contributors

CA conceived the study. CA & UM developed the protocol. UM, NE, CT, JT, JL, CA & EMB contributed to the search strategy, screened the search results, extracted data, and interpreted the findings. UM, CA & EMB wrote the paper. All authors reviewed the paper for important intellectual content.

Declaration of interest

We declare no competing interests.

Acknowledgments

There was no funding source for this study.

References

- 1 World Health Organisation. Novel Coronavirus(2019-nCoV)SITUATION REPORT-121 JANUARY 2020. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200121-sitrep-1-2019-ncov.pdf?sfvrsn=20a99c10_4 (accessed May 20, 2020).
- 2 Wilder-Smith A, Chiew CJ, Lee VJ. Can we contain the COVID-19 outbreak with the same measures as for SARS? *Lancet Infect Dis* 2020; **20**: e102-e107. [https://doi.org/10.1016/S1473-3099\(20\)30129-8](https://doi.org/10.1016/S1473-3099(20)30129-8).
- 3 Lai C-C, Shih T-P, Ko W-C, Tang H-J, Hsueh P-R. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The epidemic and the challenges. *Int J Antimicrob Agents* 2020; **55**: 105924. <https://doi.org/10.1016/j.ijantimicag.2020.105924>.
- 4 Remuzzi A, Remuzzi G. COVID-19 and Italy: what next? *Lancet* 2020; **395**: 1225–28. [https://doi.org/10.1016/S0140-6736\(20\)30627-9](https://doi.org/10.1016/S0140-6736(20)30627-9).
- 5 Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schünemann HJ. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet* 2020. [https://doi.org/10.1016/S0140-6736\(20\)31142-9](https://doi.org/10.1016/S0140-6736(20)31142-9).
- 6 Mattioli AV, Ballerini Puviani M, Nasi M, Farinetti A. COVID-19 pandemic: the effects of quarantine on cardiovascular risk. *Eur J Clin Nutr* 2020; **74**: 852–55. <https://doi.org/10.1038/s41430-020-0646-z>.
- 7 Cheng Y, Luo R, Wang K, et al. Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney Int* 2020; **97**: 829–38. <https://doi.org/10.1016/j.kint.2020.03.005>.
- 8 Röhr S, Müller F, Jung F, Apfelbacher C, Seidler A, Riedel-Heller SG. Psychosocial Impact of Quarantine Measures During Serious Coronavirus Outbreaks: A Rapid Review. *Psychiatr Prax* 2020; **47**: 179–89. <https://doi.org/10.1055/a-1159-5562>.
- 9 Sorensen K, van den Broucke S, Fullam J, et al. Health literacy and public health: a systematic review and integration of definitions and models. *BMC Public Health* 2012; **12**. <https://doi.org/10.1186/1471-2458-12-80>.
- 10 Abel T, McQueen D. Critical health literacy and the COVID-19 crisis. *Health Promot Int* 2020. <https://doi.org/10.1093/heapro/daaa040>.
- 11 Paakkari L, Okan O. COVID-19: health literacy is an underestimated problem. *Lancet Public Health* 2020; **5**: e249-e250. [https://doi.org/10.1016/S2468-2667\(20\)30086-4](https://doi.org/10.1016/S2468-2667(20)30086-4).
- 12 Rikard RV, Thompson MS, McKinney J, Beauchamp A. Examining health literacy disparities in the United States: a third look at the National Assessment of Adult Literacy (NAAL). *BMC Public Health* 2016; **16**: 975. <https://doi.org/10.1186/s12889-016-3621-9>.
- 13 Wagner C von, Knight K, Steptoe A, Wardle J. Functional health literacy and health-promoting behaviour in a national sample of British adults. *J Epidemiol Community Health* 2007; **61**: 1086–90. <https://doi.org/10.1136/jech.2006.053967>.
- 14 Eichler K, Wieser S, Brugger U. The costs of limited health literacy: a systematic review. *Int J Public Health* 2009; **54**: 313–24. <https://doi.org/10.1007/s00038-009-0058-2>.
- 15 van der Heide I, Wang J, Droomers M, Spreeuwenberg P, Rademakers J, Uiters E. The Relationship Between Health, Education, and Health Literacy: Results From the Dutch Adult Literacy and Life Skills Survey. *Journal of Health Communication* 2013; **18**: 172–84. <https://doi.org/10.1080/10810730.2013.825668>.

- 16 Paasche-Orlow MK, Parker RM, Gazmararian JA, Nielsen-Bohlman LT, Rudd RR. The prevalence of limited health literacy. *J Gen Intern Med* 2005; **20**: 175–84. <https://doi.org/10.1111/j.1525-1497.2005.40245.x>.
- 17 Castro-Sánchez E, Chang PWS, Vila-Candel R, Escobedo AA, Holmes AH. Health literacy and infectious diseases: why does it matter? *International Journal of Infectious Diseases* 2016; **43**: 103–10. <https://doi.org/10.1016/j.ijid.2015.12.019>.
- 18 Khalil H, Peters M, Godfrey CM, McInerney P, Soares CB, Parker D. An Evidence-Based Approach to Scoping Reviews. *Worldviews Evid Based Nurs* 2016; **13**: 118–23. <https://doi.org/10.1111/wvn.12144>.
- 19 Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *International Journal of Social Research Methodology* 2005; **8**: 19–32. <https://doi.org/10.1080/1364557032000119616>.
- 20 Peters MDJ, Godfrey C, McInerney P, Munn Z, Tricco AC, Khalil, H. Chapter 11: Scoping Reviews (2020 version). In: Aromataris E MZ, ed. Joanna Briggs Institute Reviewer's Manual, JBI, 2020.
- 21 Grant MJ, Booth A. A typology of reviews: an analysis of 14 review types and associated methodologies. *Health Info Libr J* 2009; **26**: 91–108. <https://doi.org/10.1111/j.1471-1842.2009.00848.x>.
- 22 Khangura S, Konnyu K, Cushman R, Grimshaw J, Moher D. Evidence summaries: the evolution of a rapid review approach. *Syst Rev* 2012; **1**: 10. <https://doi.org/10.1186/2046-4053-1-10>.
- 23 Matterne, U., Egger, N., Apfelbacher, C., Lander, J., Bitzer, E. M., & Dierks, M. L. Health literacy in the context of pandemic and epidemic (outbreak): rapid scoping review. <https://osf.io/sgw76/> (accessed Apr 06, 2020).
- 24 Nutbeam D. Health literacy as a public health goal: a challenge for contemporary health education and communication strategies into the 21st century. *Health Promot Int* 2000; **15**: 259–67. <https://doi.org/10.1093/heapro/15.3.259>.
- 25 Tricco AC, Lillie E, Zarin W, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med* 2018; **169**: 467–73. <https://doi.org/10.7326/M18-0850>.
- 26 Betsch C, Wieler L, Bosnjak M, et al. Germany COVID-19 Snapshot MOonitoring (COSMO Germany): Monitoring knowledge, risk perceptions, preventive behaviours, and public trust in the current coronavirus outbreak in Germany. <https://www.psycharchives.org/handle/20.500.12034/2501> (accessed May 08, 2020).
- 27 Okan O, de Sombre S, Hurrelmann K, Berens EM, Bauer U, Schaeffer D. Gesundheitskompetenz der Bevölkerung im Umgang mit der Coronavirus-Pandemie (accessed May 07, 2020).
- 28 Nguyen HC, Nguyen MH, Do BN, et al. People with Suspected COVID-19 Symptoms Were More Likely Depressed and Had Lower Health-Related Quality of Life: The Potential Benefit of Health Literacy. *J Clin Med* 2020; **9**. <https://doi.org/10.3390/jcm9040965>.
- 29 Wolf MS, Serper M, Opsasnick L, et al. Awareness, Attitudes, and Actions Related to COVID-19 Among Adults With Chronic Conditions at the Onset of the U.S. Outbreak: A Cross-sectional Survey. *Ann Intern Med* 2020. <https://doi.org/10.7326/M20-1239>.
- 30 Voeten, Helene A C M, Zwart O de, Veldhuijzen IK, et al. Sources of information and health beliefs related to SARS and avian influenza among Chinese communities in the United Kingdom and The Netherlands, compared to the general population in these countries. *Int J Behav Med* 2009; **16**: 49–57. <https://doi.org/10.1007/s12529-008-9006-4>.
- 31 Jang K, Park N. The Effects of Repetitive Information Communication through Multiple Channels on Prevention Behavior during the 2015 MERS Outbreak in South Korea. *J Health Commun* 2018; **23**: 670–78. <https://doi.org/10.1080/10810730.2018.1501440>.

- 32 Kim S, Kim S. Exploring the Determinants of Perceived Risk of Middle East Respiratory Syndrome (MERS) in Korea. *Int J Environ Res Public Health* 2018; **15**.
<https://doi.org/10.3390/ijerph15061168>.
- 33 Wang C, Pan R, Wan X, et al. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *Int J Environ Res Public Health* 2020; **17**.
<https://doi.org/10.3390/ijerph17051729>.
- 34 Wang C, Pan R, Wan X, et al. A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. *Brain Behav Immun* 2020.
<https://doi.org/10.1016/j.bbi.2020.04.028>.
- 35 Alhomoud F, Alhomoud F. "Your Health Essential for Your Hajj": Muslim pilgrims' knowledge, attitudes and practices regarding Middle East respiratory syndrome coronavirus (MERS-CoV) during Hajj season. *J Infect Chemother* 2017; **23**: 286–92.
<https://doi.org/10.1016/j.jiac.2017.01.006>.
- 36 Ma T, Heywood A, MacIntyre CR. Travel health risk perceptions of Chinese international students in Australia - Implications for COVID-19. *Infect Dis Health* 2020.
<https://doi.org/10.1016/j.idh.2020.03.002>.
- 37 Wills BSH, Morse JM. Responses of Chinese elderly to the threat of severe acute respiratory syndrome (SARS) in a Canadian community. *Public Health Nurs* 2008; **25**: 57–68.
<https://doi.org/10.1111/j.1525-1446.2008.00680.x>.
- 38 Jiang X, Elam G, Yuen C, et al. The perceived threat of SARS and its impact on precautionary actions and adverse consequences: a qualitative study among Chinese communities in the United Kingdom and the Netherlands. *Int J Behav Med* 2009; **16**: 58–67. <https://doi.org/10.1007/s12529-008-9005-5>.
- 39 Cava MA, Fay KE, Beanlands HJ, McCay EA, Wignall R. The Experience of Quarantine for Individuals Affected by SARS in Toronto. *Public Health Nurs* 2005; **22**: 398–406.
<https://doi.org/10.1111/j.0737-1209.2005.220504.x>.
- 40 Tse MMY, Pun SPY, Benzie IFF. Experiencing SARS: perspectives of the elderly residents and health care professionals in a Hong Kong nursing home. *Geriatr Nurs* 2003; **24**: 266–69.
[https://doi.org/10.1016/s0197-4572\(03\)00251-9](https://doi.org/10.1016/s0197-4572(03)00251-9).
- 41 Siu JY-M. Qualitative study on the shifting sociocultural meanings of the facemask in Hong Kong since the severe acute respiratory syndrome (SARS) outbreak: implications for infection control in the post-SARS era. *Int J Equity Health* 2016; **15**: 73. <https://doi.org/10.1186/s12939-016-0358-0>.
- 42 Nielsen-Bohlman L, Panzer AM, Kindig DA. Health Literacy: A Prescription to End Confusion. Washington (DC), 2004.
- 43 Sørensen K, van den Broucke S, Pelikan JM, et al. Measuring health literacy in populations: illuminating the design and development process of the European Health Literacy Survey Questionnaire (HLS-EU-Q). *BMC Public Health* 2013; **13**: 948. <https://doi.org/10.1186/1471-2458-13-948>.
- 44 Sun X, Chen J, Shi Y, et al. Measuring health literacy regarding infectious respiratory diseases: a new skills-based instrument. *PLoS One* 2013; **8**: e64153.
<https://doi.org/10.1371/journal.pone.0064153>.
- 45 Nutbeam D, Kickbusch I. Advancing health literacy: a global challenge for the 21st century. *Health Promot Int* 2000; **15**: 183–84. <https://doi.org/10.1093/heapro/15.3.183>.
- 46 Ajzen I. The theory of planned behaviour: Reactions and reflections. *Psychol Health* 2011; **26**: 1113–27. <https://doi.org/10.1080/08870446.2011.613995>.

- 47 Rosenstock IM. The health belief model: Explaining health behavior through expectancies. In: Health behavior and health education: Theory, research, and practice. San Francisco, CA, US: Jossey-Bass, 1990: 39–62.
- 48 Rogers RW, Prentice-Dunn S. Protection motivation theory. In: Handbook of health behavior research 1: Personal and social determinants. New York, NY, US: Plenum Press, 1997: 113–32.
- 49 Fowler RA, Fletcher T, Fischer WA2, et al. Caring for critically ill patients with ebola virus disease. Perspectives from West Africa. *Am J Respir Crit Care Med* 2014; **190**: 733–37. <https://doi.org/10.1164/rccm.201408-1514CP>.
- 50 Bedrosian SR, Young CE, Smith LA, et al. Lessons of Risk Communication and Health Promotion - West Africa and United States. *MMWR Suppl* 2016; **65**: 68–74. <https://doi.org/10.15585/mmwr.su6503a10>.
- 51 Pleasant A, Kuruvilla S. A tale of two health literacies: public health and clinical approaches to health literacy. *Health Promot Int* 2008; **23**: 152–59. <https://doi.org/10.1093/heapro/dan001>.
- 52 Baker DW. The meaning and the measure of health literacy. *J Gen Intern Med* 2006; **21**: 878–83. <https://doi.org/10.1111/j.1525-1497.2006.00540.x>.
- 53 Parker RM, Baker DW, Williams MV, Nurss JR. The test of functional health literacy in adults: a new instrument for measuring patients' literacy skills. *J Gen Intern Med* 1995; **10**: 537–41. <https://doi.org/10.1007/BF02640361>.
- 54 Osborne RH, Batterham RW, Elsworth GR, Hawkins M, Buchbinder R. The grounded psychometric development and initial validation of the Health Literacy Questionnaire (HLQ). *BMC Public Health* 2013; **13**: 658. <https://doi.org/10.1186/1471-2458-13-658>.
- 55 Rudd R, Baur C. Health literacy and early insights during a pandemic. *Journal of Communication in Healthcare* 2020: 1–4. <https://doi.org/10.1080/17538068.2020.1760622>.
- 56 Geldsetzer P. Knowledge and Perceptions of COVID-19 Among the General Public in the United States and the United Kingdom: A Cross-sectional Online Survey. *Ann Intern Med* 2020. <https://doi.org/10.7326/M20-0912>.
- 57 Geldsetzer P. Use of Rapid Online Surveys to Assess People's Perceptions During Infectious Disease Outbreaks: A Cross-sectional Survey on COVID-19. *J Med Internet Res* 2020; **22**: e18790. <https://doi.org/10.2196/18790>.
- 58 Khan S, Khan M, Maqsood K, Hussain T, Noor-Ul-Huda, Zeeshan M. Is Pakistan prepared for the COVID-19 epidemic? A questionnaire-based survey. *J Med Virol* 2020. <https://doi.org/10.1002/jmv.25814>.
- 59 Roy D, Tripathy S, Kar SK, Sharma N, Verma SK, Kaushal V. Study of knowledge, attitude, anxiety & perceived mental healthcare need in Indian population during COVID-19 pandemic. *Asian J Psychiatr* 2020; **51**: 102083. <https://doi.org/10.1016/j.ajp.2020.102083>.
- 60 Zhong B-L, Luo W, Li H-M, et al. Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional survey. *Int J Biol Sci* 2020; **16**: 1745–52. <https://doi.org/10.7150/ijbs.45221>.
- 61 Bener A, Al-Khal A. Knowledge, attitude and practice towards SARS. *J R Soc Promot Health* 2004; **124**: 167–70. <https://doi.org/10.1177/146642400412400408>.
- 62 Bergeron SL, Sanchez AL. Media effects on students during SARS outbreak. *Emerg Infect Dis* 2005; **11**: 732–34. <https://doi.org/10.3201/eid1105.040512>.
- 63 Blendon RJ, Benson JM, DesRoches CM, Raleigh E, Taylor-Clark K. The public's response to severe acute respiratory syndrome in Toronto and the United States. *Clin Infect Dis* 2004; **38**: 925–31. <https://doi.org/10.1086/382355>.
- 64 Brug J, Aro AR, Oenema A, Zwart O de, Richardus JH, Bishop GD. SARS risk perception, knowledge, precautions, and information sources, the Netherlands. *Emerg Infect Dis* 2004; **10**: 1486–89. <https://doi.org/10.3201/eid1008.040283>.

- 65 Chan SSC, So WKW, Wong DCN, Lee ACK, Tiwari A. Improving older adults' knowledge and practice of preventive measures through a telephone health education during the SARS epidemic in Hong Kong: a pilot study. *Int J Nurs Stud* 2007; **44**: 1120–27. <https://doi.org/10.1016/j.ijnurstu.2006.04.019>.
- 66 Cheng C, Ng A-K. Psychosocial Factors Predicting SARS-Preventive Behaviors in Four Major SARS-Affected Regions. *Journal of Applied Social Psychology* 2006; **36**: 222–47. <https://doi.org/10.1111/j.0021-9029.2006.00059.x>.
- 67 Chuo H-Y. Restaurant diners' self-protective behavior in response to an epidemic crisis. *Int J Hosp Manag* 2014; **38**: 74–83. <https://doi.org/10.1016/j.ijhm.2014.01.004>.
- 68 Des Jarlais DC, Galea S, Tracy M, Tross S, Vlahov D. Stigmatization of newly emerging infectious diseases: AIDS and SARS. *Am J Public Health* 2006; **96**: 561–67. <https://doi.org/10.2105/AJPH.2004.054742>.
- 69 Des Jarlais DC, Stuber J, Tracy M, Tross S, Galea S. Social factors associated with AIDS and SARS. *Emerg Infect Dis* 2005; **11**: 1767–69. <https://doi.org/10.3201/eid1111.050424>.
- 70 Deurenberg-Yap M, Foo LL, Low YY, Chan SP, Vijaya K, Lee M. The Singaporean response to the SARS outbreak: knowledge sufficiency versus public trust. *Health Promot Int* 2005; **20**: 320–26. <https://doi.org/10.1093/heapro/dai010>.
- 71 Hazreen AM, Myint Myint S, Farizah H, et al. An evaluation of information dissemination during the severe acute respiratory syndrome (SARS) outbreak among selected rural communities in Kuala Kangsar. *Med J Malaysia* 2005; **60**: 180–87.
- 72 Ishizaki T, Imanaka Y, Hirose M, et al. Estimation of the impact of providing outpatients with information about SARS infection control on their intention of outpatient visit. *Health Policy* 2004; **69**: 293–303. <https://doi.org/10.1016/j.healthpol.2004.04.008>.
- 73 Lau JTF, Yang X, Tsui H, Kim JH. Monitoring community responses to the SARS epidemic in Hong Kong: from day 10 to day 62. *J Epidemiol Community Health* 2003; **57**: 864–70. <https://doi.org/10.1136/jech.57.11.864>.
- 74 Lau JTF, Yang X, Pang E, Tsui HY, Wong E, Wing YK. SARS-related perceptions in Hong Kong. *Emerg Infect Dis* 2005; **11**: 417–24. <https://doi.org/10.3201/eid1103.040675>.
- 75 Lau JTF, Yang X, Tsui H, Pang E, Kim JH. SARS preventive and risk behaviours of Hong Kong air travellers. *Epidemiol Infect* 2004; **132**: 727–36. <https://doi.org/10.1017/s0950268804002225>.
- 76 Leung GM, Lam T-H, Ho L-M, et al. The impact of community psychological responses on outbreak control for severe acute respiratory syndrome in Hong Kong. *J Epidemiol Community Health* 2003; **57**: 857–63. <https://doi.org/10.1136/jech.57.11.857>.
- 77 Leung GM, Ho L-M, Chan SKK, et al. Longitudinal assessment of community psychobehavioral responses during and after the 2003 outbreak of severe acute respiratory syndrome in Hong Kong. *Clin Infect Dis* 2005; **40**: 1713–20. <https://doi.org/10.1086/429923>.
- 78 Leung GM, Quah S, Ho L-M, et al. A tale of two cities: community psychobehavioral surveillance and related impact on outbreak control in Hong Kong and Singapore during the severe acute respiratory syndrome epidemic. *Infect Control Hosp Epidemiol* 2004; **25**: 1033–41. <https://doi.org/10.1086/502340>.
- 79 Leung GM, Quah S, Ho LM, et al. Community psycho-behavioural surveillance and related impact on outbreak control in Hong Kong and Singapore during the SARS epidemic. *Hong Kong Med J* 2009; **15 Suppl 9**: 30–34.
- 80 Lim VKG. War with SARS: an empirical study of knowledge of SARS transmission and effects of SARS on work and the organisations. *Singapore Med J* 2003; **44**: 457–63.
- 81 Peng EY-C, Lee M-B, Tsai S-T, et al. Population-based post-crisis psychological distress: an example from the SARS outbreak in Taiwan. *J Formos Med Assoc* 2010; **109**: 524–32. [https://doi.org/10.1016/S0929-6646\(10\)60087-3](https://doi.org/10.1016/S0929-6646(10)60087-3).

- 82 Quah SR, Hin-Peng L. Crisis prevention and management during SARS outbreak, Singapore. *Emerg Infect Dis* 2004; **10**: 364–68. <https://doi.org/10.3201/eid1002.030418>.
- 83 Seng SL, Lim PS, Ng MY, Wong HB, Emmanuel SC. A study on SARS awareness and health-seeking behaviour - findings from a sampled population attending National Healthcare Group Polyclinics. *Ann Acad Med Singapore* 2004; **33**: 623–29.
- 84 So WKW, Chan SSC, Lee ACK, Tiwari AFY. The knowledge level and precautionary measures taken by older adults during the SARS outbreak in Hong Kong. *Int J Nurs Stud* 2004; **41**: 901–09. <https://doi.org/10.1016/j.ijnurstu.2004.04.004>.
- 85 Tan X, Li S, Wang C, Chen X, Wu X. Severe acute respiratory syndrome epidemic and change of people's health behavior in China. *Health Educ Res* 2004; **19**: 576–80. <https://doi.org/10.1093/her/cyg074>.
- 86 Tang CSK, Wong C-Y. An outbreak of the severe acute respiratory syndrome: predictors of health behaviors and effect of community prevention measures in Hong Kong, China. *Am J Public Health* 2003; **93**: 1887–88. <https://doi.org/10.2105/ajph.93.11.1887>.
- 87 Tang CS-K, Wong C-Y. Psychosocial Factors Influencing the Practice of Preventive Behaviors Against the Severe Acute Respiratory Syndrome Among Older Chinese in Hong Kong. *J Aging Health* 2005; **17**: 490–506. <https://doi.org/10.1177/0898264305277966>.
- 88 Tracy CS, Rea E, Upshur REG. Public perceptions of quarantine: community-based telephone survey following an infectious disease outbreak. *BMC Public Health* 2009; **9**: 470. <https://doi.org/10.1186/1471-2458-9-470>.
- 89 Vartti A-M, Oenema A, Schreck M, et al. SARS knowledge, perceptions, and behaviors: A comparison between Finns and the Dutch during the SARS outbreak in 2003. *Int J Behav Med* 2009; **16**: 41–48. <https://doi.org/10.1007/s12529-008-9004-6>.
- 90 Wong C-Y, Tang CS-K. Practice of habitual and volitional health behaviors to prevent severe acute respiratory syndrome among Chinese adolescents in Hong Kong. *J Adolesc Health* 2005; **36**: 193–200. <https://doi.org/10.1016/j.jadohealth.2004.02.024>.
- 91 Yip HK, Tsang PCS, Samaranayake LP, Li AHP. Knowledge of and attitudes toward severe acute respiratory syndrome among a cohort of dental patients in Hong Kong following a major local outbreak. *Community Dent Health* 2007; **24**: 43–48.
- 92 Zwart O de, Veldhuijzen IK, Elam G, et al. Perceived threat, risk perception, and efficacy beliefs related to SARS and other (emerging) infectious diseases: Results of an international survey. *Int J Behav Med* 2009; **16**: 30–40. <https://doi.org/10.1007/s12529-008-9008-2>.
- 93 Al-Hazmi A, Gosadi I, Somily A, Alsubaie S, Bin Saeed A. Knowledge, attitude and practice of secondary schools and university students toward Middle East Respiratory Syndrome epidemic in Saudi Arabia: A cross-sectional study. *Saudi J Biol Sci* 2018; **25**: 572–77. <https://doi.org/10.1016/j.sjbs.2016.01.032>.
- 94 Al-Mohaissen M. Awareness among a Saudi Arabian university community of Middle East respiratory syndrome coronavirus following an outbreak. *East Mediterr Health J* 2017; **23**: 351–60. <https://doi.org/10.26719/2017.23.5.351>.
- 95 Almutairi KM, Al Helih EM, Moussa M, et al. Awareness, Attitudes, and Practices Related to Coronavirus Pandemic Among Public in Saudi Arabia. *Fam Community Health* 2015; **38**: 332–40. <https://doi.org/10.1097/FCH.000000000000082>.
- 96 Alotaibi MS, Alsubaie AM, Almohaime KA, et al. To what extent are Arab pilgrims to Makkah aware of the middle east respiratory syndrome coronavirus and the precautions against it? *J Family Community Med* 2017; **24**: 91–96.
- 97 Alqahtani AS, Wiley KE, Mushta SM, et al. Association between Australian Hajj Pilgrims' awareness of MERS-CoV, and their compliance with preventive measures and exposure to camels. *J Travel Med* 2016; **23**. <https://doi.org/10.1093/jtm/taw046>.

- 98 Alqahtani AS, Wiley KE, Tashani M, et al. Camel exposure and knowledge about MERS-CoV among Australian Hajj pilgrims in 2014. *Virol Sin* 2016; **31**: 89–93. <https://doi.org/10.1007/s12250-015-3669-1>.
- 99 Alqahtani AS, Rashid H, Basyouni MH, Alhawassi TM, BinDhim NF. Public response to MERS-CoV in the Middle East: iPhone survey in six countries. *J Infect Public Health* 2017; **10**: 534–40. <https://doi.org/10.1016/j.jiph.2016.11.015>.
- 100 Althobaity HM, Alharthi RAS, Altowairqi MH, et al. Knowledge and awareness of Middle East respiratory syndrome coronavirus among Saudi and Non-Saudi Arabian pilgrims. *Int J Health Sci (Qassim)* 2017; **11**: 20–25.
- 101 Ashok N, Rodrigues JC, Azouni K, et al. Knowledge and Apprehension of Dental Patients about MERS-A Questionnaire Survey. *J Clin Diagn Res* 2016; **10**: ZC58-62. <https://doi.org/10.7860/JCDR/2016/17519.7790>.
- 102 Bawazir A, Al-Mazroo E, Jradi H, Ahmed A, Badri M. MERS-CoV infection: Mind the public knowledge gap. *J Infect Public Health* 2018; **11**: 89–93. <https://doi.org/10.1016/j.jiph.2017.05.003>.
- 103 Gautret P, Benkouiten S, Salaheddine I, et al. Hajj pilgrims knowledge about Middle East respiratory syndrome coronavirus, August to September 2013. *Euro Surveill* 2013; **18**: 20604. <https://doi.org/10.2807/1560-7917.es2013.18.41.20604>.
- 104 Hoda J. Identification of information types and sources by the public for promoting awareness of Middle East respiratory syndrome coronavirus in Saudi Arabia. *Health Educ Res* 2016; **31**: 12–23.
- 105 Hou Y'a, Tan Y-R, Lim WY, et al. Adequacy of public health communications on H7N9 and MERS in Singapore: insights from a community based cross-sectional study. *BMC Public Health* 2018; **18**: 436. <https://doi.org/10.1186/s12889-018-5340-x>.
- 106 Kamau E, Ongus J, Gitau G, et al. Knowledge and practices regarding Middle East Respiratory Syndrome Coronavirus among camel handlers in a Slaughterhouse, Kenya, 2015. *Zoonoses Public Health* 2019; **66**: 169–73. <https://doi.org/10.1111/zph.12524>.
- 107 Lee M, Ju Y, You M. The effects of social determinants on public health emergency preparedness mediated by health communication: The 2015 mers outbreak in south korea. *Health Commun* 2019. <https://doi.org/10.1080/10410236.2019.1636342>.
- 108 Lee SY, Yang HJ, Kim G, Cheong H-K, Choi BY. Preventive behaviors by the level of perceived infection sensitivity during the Korea outbreak of Middle East Respiratory Syndrome in 2015. *Epidemiol Health* 2016; **38**: e2016051. <https://doi.org/10.4178/epih.e2016051>.
- 109 Lin L, McCloud RF, Bigman CA, Viswanath K. Tuning in and catching on? Examining the relationship between pandemic communication and awareness and knowledge of MERS in the USA. *J Public Health (Oxf)* 2017; **39**: 282–89. <https://doi.org/10.1093/pubmed/fdw028>.
- 110 Migault C, Kanagaratnam L, Hentzien M, et al. Effectiveness of an education health programme about Middle East respiratory syndrome coronavirus tested during travel consultations. *Public Health* 2019; **173**: 29–32. <https://doi.org/10.1016/j.puhe.2019.05.017>.
- 111 Nooh HZ, Alshammary RH, Alenezy JM, et al. Public awareness of coronavirus in Al-Jouf region, Saudi Arabia. *Z Gesundh Wiss* 2020: 1–8. <https://doi.org/10.1007/s10389-020-01209-y>.
- 112 Sahin MK, Aker S, Kaynar Tuncel E. Knowledge, attitudes and practices concerning Middle East respiratory syndrome among Umrah and Hajj pilgrims in Samsun, Turkey, 2015. *Euro Surveill* 2015; **20**. <https://doi.org/10.2807/1560-7917.ES.2015.20.38.30023>.
- 113 Tashani M, Alfelali M, Barasheed O, et al. Australian Hajj pilgrims' knowledge about MERS-CoV and other respiratory infections. *Virol Sin* 2014; **29**: 318–20. <https://doi.org/10.1007/s12250-014-3506-y>.

- 114 Yang S, Cho S-I. Middle East respiratory syndrome risk perception among students at a university in South Korea, 2015. *Am J Infect Control* 2017; **45**: e53-e60. <https://doi.org/10.1016/j.ajic.2017.02.013>.
- 115 Yang J, Park E-C, Lee SA, Lee SG. Associations Between Hand Hygiene Education and Self-Reported Hand-Washing Behaviors Among Korean Adults During MERS-CoV Outbreak. *Health Educ Behav* 2019; **46**: 157–64. <https://doi.org/10.1177/1090198118783829>.
- 116 Yoo W, Choi D-H, Park K. The effects of SNS communication: How expressing and receiving information predict MERS-preventive behavioral intentions in South Korea. *Comput Human Behav* 2016; **62**: 34–43. <https://doi.org/10.1016/j.chb.2016.03.058>.
- 117 Betsch C, Wieler L, Bosnjak M, et al. Germany COVID-19 Snapshot MOonitoring (COSMO Germany): Monitoring knowledge, risk perceptions, preventive behaviours, and public trust in the current coronavirus outbreak in Germany. <https://www.psycharchives.org/handle/20.500.12034/2501> (accessed May 08, 2020).
- 118 Okan O, de Sombre S, Hurrelmann K, Berens EM, Bauer U, Schaeffer D. Gesundheitskompetenz der Bevölkerung im Umgang mit der Coronavirus-Pandemie (accessed May 07, 2020).
- 119 Alotaibi MS, Alsubaie AM, Almohaimede KA, et al. To what extent are Arab pilgrims to Makkah aware of the middle east respiratory syndrome coronavirus and the precautions against it? *J Family Community Med* 2017; **24**: 91–96.
- 120 Hoda J. Identification of information types and sources by the public for promoting awareness of Middle East respiratory syndrome coronavirus in Saudi Arabia. *Health Educ Res* 2016; **31**: 12–23.

PRISMA 2009 Flow Diagram²⁵

Appendix 1: search strategy

PubMed (MEDLINE®)

(health literacy[MeSH Terms] OR (health[Title/Abstract] AND competence[Title/Abstract]) OR literacy[Title/Abstract] OR knowledge[Title/Abstract] OR attitude[Title/Abstract] OR motivation* OR intention* OR skills OR self-efficacy[MeSH] OR organisation*[Title/Abstract] OR community[Title/Abstract]) AND ((2019-nCoV OR 2019nCoV OR COVID-19 OR SARS-CoV-2 OR ((wuhan AND coronavirus) AND 2019/12[PDAT]:2030[PDAT])) OR ("Severe Acute Respiratory Syndrome"[MeSH Terms] OR SARS) OR ("middle east respiratory syndrome coronavirus"[MeSH Terms] OR MERS)) NOT (animals [mh] NOT humans [mh])

PsycINFO®

(health literacy or health education or health knowledge or health information or health understanding) AND (pandemic* or epidemic* or outbreak or covid-19 or coronavirus OR 2019-ncov or sars OR sars-cov-2 OR mers) NOT (hiv or aids or acquired human immunodeficiency syndrome or human immunodeficiency virus)

Appendix 2: Summary of study characteristics

Study ID	Country	n	When conducted	Design	Method	Sampling method	HL specific	Other theory guided	Type of theory
SARS-CoV-2									
Betsch 2020 ²⁶	GER	1000	during	obs cross	Quest.	repr.	no	no	n.a.
Geldsetzer 2020a 2020b ^{56,57}	UK, USA	5974	during	obs cross	Quest.	Conv.	no	no	n.a.
Khan 2020 ⁵⁸	PAK	302	during	obs cross	Quest.	Conv.	no	no	n.a.
Ma 2020 ³⁶	AUS	28	during	qual.	Focus groups	Conv.	no	no	n.a.
Nguyen 2020 ²⁸	VNM	3974	during	obs cross	Quest.	Conv.	yes	no	n.a.
Okan 2020 ²⁷	GER	1000	during	obs cross	Quest.	repr.	yes	no	n.a.
Roy 2020 ⁵⁹	IND	662	during	obs cross	Quest.	Conv.	no	no	n.a.
Wang 2020a 2020b ^{33,34}	CHN	1738	during	obs Pp	Quest.	Conv.	no	no	n.a.
Wolf 2020 ²⁹	USA	630	during	obs cross	Quest.	Conv.	yes	no	n.a.
Zhong 2020 ⁶⁰	CHN	6919	during	obs cross	Quest.	Conv.	no	no	n.a.
SARS-CoV-1									
Bener 2004 ⁶¹	QAT	1386	during	obs cross	Quest.	Conv.	no	no	n.a.
Bergeron 2005 ⁶²	CAN	300	post	obs cross	Quest.	Conv.	no	no	n.a.
Blendon 2004 ⁶³	CAN, USA	1500	during	obs cross	Quest.	repr.	no	no	n.a.
Brug 2004 ⁶⁴	NLD	373	during	obs cross	Quest.	unclear	no	no	n.a.
Cava 2005 ³⁹	CAN	21	during	qual.	Interview	Conv.	no	no	n.a.
Chan 2007 ⁶⁵	HKG	296/122	during	obs Pp Intervention	Quest.	Conv.	no	no	n.a.
Cheng 2006 ⁶⁶	CHN, SGP, HKG, CAN	300	during	obs Pp	Quest.	Conv.	no	yes	THB
Chuo 2014 ⁶⁷	TWN	294/88	partly during (first survey wave)	obs Pp	Quest.	Conv.	no	yes	THB
Des Jarlais 2006/2005 ^{68,69}	USA	928	during	obs cross	Quest.	repr.	no	yes	OM
Deurenberg-Yap 2005 ⁷⁰	SGP	853	during	obs cross	Quest.	repr.	no	yes	OM
Hazreen 2005 ⁷¹	MYS	201 house holds	during	obs cross	Quest.	repr.	no	no	n.a.
Ishizaki 2004 ⁷²	JPN	821	during	obs cross	Quest.	Conv.	no	yes	THB
Jiang 2009 ³⁸	UK, NLD	164	post	Qual.	Focus groups	Conv.	no	yes	GT
Lau 2003 ⁷³	HKG	1397	during	obs Pp	Quest.	repr.	no	yes	THB
Lau 2005 ⁷⁴	HKG	820	during	obs cross	Quest.	repr.	no	no	n.a.
Lau 2004 ⁷⁵	HKG	820	during	obs cross	Quest.	Conv.	no	yes	THB
Leung 2003 ⁷⁶	HKG	1115	during	obs cross	Quest.	repr.	no	no	n.a.
Leung 2005 ⁷⁷	HKG	4481	post	obs Pp	Quest.	repr.	no	no	n.a.
Leung 2004/2009 ^{78,79}	HKG, SGP	1906	during	obs cross	Quest.	repr.	no	no	n.a.
Lim 2003 ⁸⁰	SGP	101	during	obs cross	Quest.	Conv.	no	no	n.a.
Peng 2010 ⁸¹	TWN	1278	post	obs cross	Quest.	repr.	no	no	n.a.
Quah 2004 ⁸²	SGP	1201	during	obs cross	Quest.	repr.	no	no	n.a.
Seng 2004 ⁸³	SGP	593	post	obs cross	Quest.	Conv.	no	no	n.a.
Siu 2016 ⁴¹	HKG	40	post	qual.	interview	Conv.	no	no	n.a.
So 2004 ⁸⁴	HKG	163/112	during	obs cross	Quest.	Conv.	no	no	n.a.
Tan 2004 ⁸⁵	CHN	1807	during	obs retros	Quest.	Conv.	no	no	n.a.
Tang 2003 ⁸⁶	HKG	2331	during	obs cross	Quest.	Conv.	no	yes	THB
Tang 2005 ⁸⁷	HKG	354	during	obs cross	Quest.	Conv.	no	yes	THB
Tracy 2009 ⁸⁸	CHN	500	post	obs cross	Quest.	Conv.	no	yes	PHM
Tse 2003 ⁴⁰	HKG	40	during	qual.	interview	Conv.	no	no	n.a.
Vartti 2009 ⁸⁹	NLD, FIN	681	during	obs cross	Quest.	Conv.	no	yes	MISR
Voeten 2009 ³⁰	UK, NLD	300	post	obs cross	Quest.	Conv.	no	yes	THB
Wills 2008 ³⁷	CAN	19	during	qual.	interview	Conv.	no	no	n.a.
Wong 2005 ⁹⁰	HKG	230	during	obs cross	Quest.	repr.	no	yes	THB
Yip 2007 ⁹¹	HKG	463	during	obs cross	Quest.	Conv.	no	no	n.a.

Zwart 2009 ⁹²	DNK, NLD, UK, ESP, POL, SGP, CHN, HKG	3436	during	obs cross	Quest.	repr.	no	yes	THB
MERS-CoV									
Al-Hazmi 2018 ⁹³	SAU	1109	during	obs cross	Quest.	Conv.	no	no	n.a.
Al-Mohaisen 2017 ⁹⁴	SAU	1541	during	obs cross	Quest.	Conv.	no	no	n.a.
Alhomoud 2017 ⁹⁵	SAU	292/257	during	obs cross	Quest.	Conv.	no	no	n.a.
Almutairi 2015 ⁹⁶	SAU	1147	during	obs cross	Quest.	Conv.	no	no	n.a.
Alotaibi 2017 ⁹⁶	SAU	417	during	obs cross	Quest.	Conv.	no	no	n.a.
Alqahtani 2016a/2016b ^{97,98}	AUS	421	during	obs Pp	Quest.	Conv.	no	yes	THB
Alqahtani 2017 ⁹⁹	SAU, KWT, UAE, QAT, BHR, OM	1812	during	obs cross	Quest.	Conv.	no	yes	THB
Althobaity 2017 ¹⁰⁰	SAU	2120	during	obs cross	Quest.	Conv.	no	no	n.a.
Ashok 2016 ¹⁰¹	SAU	404	during	obs cross	Quest.	Conv.	no	no	n.a.
Bawazir 2018 ¹⁰²	SAU	676	during	obs cross	Quest.	Conv.	no	no	n.a.
Gautret 2013 ¹⁰³	FRA	360	during	obs cross	Quest.	Conv.	no	no	n.a.
Hoda 2016 ¹⁰⁴	SAU	854/658	during	obs cross	Quest.	repr.	no	yes	CPM
Hou 2018 ¹⁰⁵	SGP	2969	during	obs cross	Quest.	Conv.	no	yes	THB
Jang 2018 ³¹	KOR	1036	during	obs cross	Quest.	repr.	no	yes	THB
Kamau 2019 ¹⁰⁶	KEN	22	during	obs cross	Quest.	Conv.	no	no	n.a.
Kim 2018 ³²	KOR	814	during	obs cross	Quest.	Conv.	no	yes	VBM
Lee 2019 ¹⁰⁷	KOR	855	post	obs cross	Quest.	repr.	no	no	n.a.
Lee 2016 ¹⁰⁸	KOR	6739	during	obs cross	Quest.	repr.	no	no	n.a.
Lin 2017 ¹⁰⁹	USA	627	both	obs cross	Quest.	repr.	no	no	n.a.
Migault 2019 ¹¹⁰	FRA	82	during	obs Pp Intervention	Quest.	Conv.	no	no	n.a.
Nooh 2010 ¹¹¹	SAU	384	during	obs cross	Quest.	repr.	no	no	n.a.
Sahin 2015 ¹¹²	TUR	381	during	obs cross	Quest.	Conv.	no	no	n.a.
Tashani 2014 ¹¹³	AUS	119	during	obs cross	Quest.	Conv.	no	no	n.a.
Yang 2017 ¹¹⁴	KOR	1470	during	obs cross	Quest.	Conv.	no	no	n.a.
Yang 2019 ¹¹⁵	KOR	222,599	during	obs cross	Quest.	repr.	no	no	n.a.
Yoo 2016 ¹¹⁶	KOR	1000	during	obs cross	Quest.	repr.	no	no	n.a.

CPM: communication-persuasion model; Conv.: convenience or opportunity sample; GT: grounded theory; HL specific: Do the authors specifically refer to health literacy and/or a health literacy concept?; MISR: Model of Illness Perceptions and Self-Regulation; n.a.: not applicable; obs cross: observational cross-sectional; obs Pp: observational pre-post; OM: other model; THB: Theory of health behaviour; PHM: Public health model; repr.: representative; VBM: value-based model; Quest.: Questionnaire

Appendix 3: Health literacy (HL) dimensions and their components and behaviour/practice measured in studies

	Health Literacy						HISB	Behavior/practice	Instrument validated
	Functional				Critical	Communicative			
	Knowledge	Attitude	Risk perception†	Skills‡					
SARS-CoV-2									
Betsch 2020 ¹¹⁷	●	●	●	●	●	●	●	●	n.r.
Geldsetzer 2020a 2020b ^{56,57}	●	●	●	○	○	○	○	○	n.r.
Khan 2020 ⁵⁸	●	●	○	○	○	●	○	●	n.r.
Ma 2020 ³⁶	●	●	●	○	○	●	●	●	n.a.
Nguyen 2020 ²⁸	○	○	○	○	○	○	○	●	partly
Okan 2020 ¹¹⁸	○	○	○	●	●	●	○	○	n.r.
Roy 2020 ⁵⁹	●	●	○	○	○	○	○	○	n.r.
Wang 2020a 2020b ^{33,34}	●	●	●	○	●	○	○	○	partly
Wolf 2020 ²⁹	●	●	●	●	●	○	●	●	n.r.
Zhong 2020 ⁶⁰	●	●	○	○	●	○	○	○	n.r.
SARS-1									
Bener 2004 ⁶¹	●	●	○	○	●	●	●	●	n.r.
Bergeron 2005 ⁶²	●	●	○	○	○	●	○	○	n.r.
Blendon 2004 ⁶³	●	●	○	○	○	●	●	●	n.r.
Brug 2004 ⁶⁴	●	○	●	●	●	●	○	●	n.r.
Cava 2005 ³⁹	○	○	○	○	○	○	○	●	n.r.
Chan 2007 ⁶⁵	●	●	●	○	○	●	○	●	n.r.
Cheng 2006 ⁶⁶	●	●	○	●	○	○	●	●	n.r.
Chuo 2014 ⁶⁷	○	●	○	○	○	○	○	○	n.r.
Des Jarlais 2006/2005 ^{68,69}	●	●	○	○	○	○	○	○	n.r.
Deurenberg-Yap 2005 ⁷⁰	●	●	●	○	●	○	○	○	n.r.
Hazreen 2005 ⁷¹	●	●	●	○	●	●	○	●	n.r.
Ishizaki 2004 ⁷²	●	●	●	○	●	●	○	○	n.r.
Jiang 2009 ³⁸	●	●	●	●	○	●	●	●	n.a.
Lau 2003 ⁷³	●	●	●	○	○	●	○	●	n.r.
Lau 2005 ⁷⁴	●	●	●	○	○	○	○	○	n.r.
Lau 2004 ⁷⁵	○	●	●	○	○	○	○	●	n.r.
Leung 2003 ⁷⁶	●	●	●	○	○	○	○	●	n.r.
Leung 2005 ⁷⁷	●	●	●	○	○	○	○	●	n.r.
Leung 2004/2009 ^{78,79}	●	●	●	○	○	○	○	●	n.r.
Lim 2003 ⁸⁰	●	●	○	○	○	○	○	○	n.r.
Peng 2010 ⁸¹	○	●	●	●	●	●	○	○	n.r.
Quah 2004 ⁸²	●	●	●	○	●	○	○	●	n.r.
Seng 2004 ⁸³	●	○	○	○	○	●	○	○	n.r.
Siu 2016 ⁴¹	○	●	○	○	○	○	○	●	n.a.
So 2004 ⁸⁴	●	●	○	○	○	○	○	●	n.r.
Tan 2004 ⁸⁵	●	○	○	○	○	○	○	●	n.r.
Tang 2003 ⁸⁶	●	●	●	●	●	○	○	●	n.r.
Tang 2005 ⁸⁷	○	●	●	●	○	○	○	●	n.r.
Tracy 2009 ⁸⁸	○	●	○	○	○	○	○	○	n.r.
Tse 2003 ⁴⁰	●	○	○	○	○	●	○	○	n.a.
Vartti 2009 ⁸⁹	●	●	●	○	●	●	●	●	n.r.
Voeten 2009 ³⁰	●	●	●	●	●	●	○	○	n.r.
Wills 2008 ³⁷	○	●	●	○	○	●	●	●	n.a.
Wong 2005 ⁹⁰	●	●	●	●	○	○	○	●	n.r.
Yip 2007 ⁹¹	○	●	○	○	○	○	○	○	n.r.
Zwart 2009 ⁹²	●	●	●	●	●	○	○	○	n.r.
MERS									
Al-Hazmi 2018 ⁹³	●	●	●	○	○	●	○	○	n.r.
Al-Mohaissen 2017 ⁹⁴	●	●	●	○	○	○	○	●	n.r.
Alhomoud 2017 ³⁵	●	●	●	○	○	○	○	○	n.r.
Almutairi 2015 ⁹⁵	●	●	●	○	○	○	○	●	n.r.
Alotaibi 2017 ¹¹⁹	●	●	●	○	○	●	○	○	n.r.
Alqahtani 2016a/2016b ^{97,98}	●	●	●	○	○	●	○	●	n.r.

	Health Literacy					HISB	Behavior/practice	Instrument validated
	Functional			Critical	Communicative			
	Knowledge	Attitude	Risk perception [†]	Skills [‡]				
Alqahtani 2017 ⁹⁹	●	●	●	○	○	●	○	n.r.
Althobaity 2017 ¹⁰⁰	●	○	●	○	○	●	○	n.r.
Ashok 2016 ¹⁰¹	●	●	●	○	○	○	○	n.r.
Bawazir 2018 ¹⁰²	●	●	●	○	○	●	○	n.r.
Gautret 2013 ¹⁰³	●	●	○	○	○	●	○	n.r.
Hoda 2016 ¹²⁰	●	●	●	○	○	●	○	n.r.
Hou 2018 ¹⁰⁵	●	●	●	○	○	●	○	n.r.
Jang 2018 ³¹	●	●	●	●	○	●	●	n.r.
Kamau 2019 ¹⁰⁶	●	○	○	○	○	○	○	n.r.
Kim 2018 ³²	●	●	●	●	●	○	○	n.r.
Lee 2019 ¹⁰⁷	○	○	○	○	○	●	●	n.r.
Lee 2016 ¹⁰⁸	○	●	○	○	●	○	●	n.r.
Lin 2017 ¹⁰⁹	●	○	○	○	○	●	○	n.r.
Migault 2019 ¹¹⁰	●	○	●	○	○	●	○	n.r.
Nooh 2010 ¹¹¹	●	○	●	○	○	●	○	n.r.
Sahin 2015 ¹¹²	●	●	○	○	○	○	○	n.r.
Tashani 2014 ¹¹³	●	○	○	○	●	●	○	n.r.
Yang 2017 ¹¹⁴	●	●	●	○	●	○	○	n.r.
Yang 2019 ¹¹⁵	○	○	○	○	○	●	○	n.r.
Yoo 2016 ¹¹⁶	○	○	●	●	○	●	○	n.r.

● yes; ○ no; n.a.: not applicable; HISB: health information seeking behaviour; n.r.: not reported; * includes related concepts (e.g. outcome expectancies, response efficacy); † perceived vulnerability and/or severity; ‡ Skills (self-efficacy, skills, preparedness)

Table 1: Health literacy (HL) and behavioural dimensions assessment in quantitative studies

		SARS-CoV-2 9 studies	SARS-CoV-1 31 studies	MERS-CoV 26 studies
Functional HL	Knowledge dimension	7 26,29,33,34,56-60	25 30,61-66,68-74,76-80,82-86,89,90,92	21 31,35,93-106,109-114
	Awareness	2 58,59	3 64,83,89	9 97,98,100,101,104-106,109,110,113
	Nature of disease	2 26,59	10 30,61-64,66,71,85,89,92	9 35,95,96,100,102,105,110,111,114
	Transmission mode	4 26,33,34,58,60	18 61-63,65,66,70-74,76-80,82-85	18 31,35,93-96,99-102,104-106,109-111,113,114
	Symptoms	6 26,29,56-60	8 30,61,62,66,70,85,89,92	13 31,35,93-100,102,110,111,114
	Course	2 26,29	4 66,73,74,89	7 35,94-96,101,111,114
	Treatment	2 26,33,34	5 62,63,70,74,89	8 31,35,94,95,100,109,111,114
	Risk groups	1 56	0	3 35,94,100
	Fatality rate/# infections	2 33,34,56,57	1 64	3 35,103,111
	Measures of infection control	0	1 83	1 100
	Prevention	4 29,56,57,59,60	3 70,71,74	8 35,93,94,96,101,102,110,114
	Attitude	7 26,33,34,56-60	28 30,61-63,65-82,84,86-92	17 31,32,35,93-99,101-105,108,112,114
	Risk perception	4 26,33,34,56,57	20 30,64,65,70-79,81,82,84,86,87,89,90,92	18 31,32,35,93-102,104,105,110,111,114,116
	Self-efficacy/skills/preparedness	3 26,27,29	8 30,64,66,81,86,87,90,92	3 31,32,116
Critical HL		5 26,27,29,33,34,60	11 30,61,64,70-72,81,82,86,89,92	4 32,108,113,114
Communicative HL		3 26,27,58	12 30,61-65,71-73,81,83,89	17 31,93,96-100,102-105,107,109-111,113,115,116
HISB		2 26,29	4 61,63,66,89	2 31,107
Behaviour/practices		4 26,29,33,34,60	18 61,63-66,71,73,75-79,82,84-87,89,90	10 31,35,95,97-99,106-108,114,115
	Wash hands	1 33,34	13 64-66,73,76-79,82,84,85,87,89,90	9 31,35,95,97-99,106-108,115
	Wear mask	3 26,33,34,60	15 64-66,73,75-79,82,84-87,89,90	7 31,35,95,97-99,106,108
	Avoid touching face	0	3 76-79	4 35,95,99,108
	Cough/Sneeze hygiene	1 33,34	7 65,76-79,82,84,90	6 31,35,95,97-99,108
	Correct tissue disposal	0	0	3 35,95,97,98
	Self-impose quarantine if experiencing symptoms	0	1 75	2 35,95
	Consult doctor if experiencing symptoms	0	5 63,66,75,87,89	1 35
	Avoid sharing personal items	1 33,34	6 65,76-79,82,84	0
	Avoid travelling	0	3 61,63,89	0
	Avoid travelling to high-risk areas	0	2 71,89	0
	Avoid public transport	0	4 64,73,75,89	0
	Avoid crowded places	2 26,60	5 63,64,71,73,89	1 108
	Avoid eating out	0	3 64,66,89	0
	Avoid people with flu-like symptoms	0	2 61,66	2 99,108
	General personal self-care (pro-active HB)	0	6 64,66,85,87,89,90	0
	Home hygiene	0	7 63,64,66,73,85,87,90	1 99
	Avoid camel exposure and products	0	0	3 97-99,106
	Other behavioural responses	2 26,29	7 63,65,66,73,84,85,89	3 106-108

HB: health behaviour; HISB: health-information seeking behaviour; note: cited references do not correspond to number of studies but publications