Clinical, Behavioral and Social Factors Associated with Racial Disparities in Hospitalized and Ambulatory COVID-19 Patients from an Integrated Health Care System in Georgia

Felipe Lobelo, MD PhD^{1,2*}; Alan Bienvenida, MPH¹; Serena Leung, MPH¹; Armand Mbanya, MD Msc²; Elizabeth J Leslie, MA¹; Kate E Koplan, MD MPH¹; S. Ryan Shin, MD MA¹

¹Department of Quality and Patient Safety, The Southeast Permanente Medical Group, Kaiser Permanente Georgia and ²Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, Georgia.

Please address correspondence to:

Felipe Lobelo, MD PhD
Physician Program Director Epidemiology, Public Health and Preparedness
and Senior Physician Consultant, Population Health Research
Department of Quality and Patient Safety
The Southeast Permanente Medical Group; Kaiser Permanente Georgia
3495 Piedmont Road NE; 9 Piedmont Center, 3rd floor
Atlanta GA 30305-1736
P: (470) 825-6846
Felipe.lobelo@kp.org ; felipelobelo@emory.edu

Abstract

Introduction: Racial and ethnic minorities have shouldered a disproportional burden of coronavirus disease 2019 (COVID-19) infection to date in the US, but data on the various drivers of these disparities is limited.

Objectives: To describe the characteristics and outcomes of COVID-19 patients and explore factors associated with hospitalization risk by race.

Methods: Case series of 448 consecutive patients with confirmed COVID-19 seen at Kaiser Permanente Georgia (KPGA), an integrated health care system serving the Atlanta metropolitan area, from March 3 to May 12, 2020. KPGA members with laboratory-confirmed COVID-19. Multivariable analyses for hospitalization risk also included an additional 3489 persons under investigation (PUI) with suspected infection. COVID-19 treatment and outcomes, underlying comorbidities and quality of care management metrics, socio-demographic and other individual and community-level social determinants of health (SDOH) indicators.

Results: Of 448 COVID-19 positive members, 68,3% was non-Hispanic Black (n=306), 18% non-Hispanic White (n=81) and 13,7% Other race (n=61). Median age was 54 [IQR 43-63] years. Overall, 224 patients were hospitalized, median age 60 (50-69) years. Black race was a significant factor in the Confirmed + PUI, female and male models (ORs from 1.98 to 2.19). Obesity was associated with higher hospitalization odds in the confirmed, confirmed + PUI, Black and male models (ORs from 1.78 to 2.77). Chronic disease control metrics (diabetes, hypertension, hyperlipidemia) were associated with lower odds of hospitalization ranging from 48% to 35% in the confirmed + PUI and Black models. Self-reported physical inactivity was associated with 50% higher hospitalization odds in the Black and Female models. Residence in

the Northeast region of Atlanta was associated with lower hospitalization odds in the Confirmed + PUI, White and female models (ORs from 0.22 to 0.64)

Conclusions: We found that non-Hispanic Black KPGA members had a disproportionately higher risk of infection and, after adjusting for covariates, twice the risk of hospitalization compared to other race groups. We found no significant differences in clinical outcomes or mortality across race/ethnicity groups. In addition to age, sex and comorbidity burden, pre-pandemic self-reported exercise, metrics on quality of care and control of underlying cardio-metabolic diseases, and location of residence in Atlanta were significantly associated with hospitalization risk by race groups. Beyond well-known physiologic and clinical factors, individual and community-level social indicators and health behaviors must be considered as interventions designed to reduce COVID-19 disparities and the systemic effects of racism are implemented.

Manuscript Text

Introduction

As of July 2020, despite having 4.25% of the global population, the United States (US) has contributed a quarter of the more than 10 million cases and 500,000 deaths recorded globally due to coronavirus disease 2019 (COVID-19).(1) In the US — the current epicenter of the pandemic — it has been widely reported that Black/African Americans (AA) and other racial/ethnic minorities, particularly those living in large and diverse urban centers, shoulder a disproportionate burden of COVID-19 infection risk and associated adverse health outcomes.(2-6)

Earlier descriptive studies from patients admitted during March/April 2020 in Georgia showed an over-representation of COVID-19 hospitalizations and death rates among the Black/AA population.(7, 8) Subsequent reports from two large health care systems in Louisiana and California, and from the Veterans Affairs health system(9) also found racial disparities in COVID-19 outcomes and clinical risk factors for hospitalization. These reports also theorized that chronic disease control, health behaviors, social and other factors may contribute to such disparities.(3, 5),(7),(8) However, limited availability of quality of care history and social determinants of health metrics in most medical health records has precluded a comprehensive, broader analyses of potential drivers of these racial disparities.

The U.S. Census Bureau reports the racial/ethnic demographic distribution of Georgia as 58.3% White alone, 31.6% Black/AA, 9.7% Latino or Hispanic, and 4.1% Asian.(10) As of May 12th, the Georgia Department of Public Health (DPH) reported confirmed COVID-19 cases by race/ethnicity as 35.1% Black/AA, 30% white, 10.1% Hispanic or Latino and 1.4% Asian(11). This overrepresentation of Black/AA and other minority populations has been observed in other Georgia counties.(11) Kaiser Permanente Georgia (KPGA) is a regional integrated health care system serving over 300,000 members in 32 counties located in the Atlanta Metropolitan Area and the Northeast region of the state. As of April 2020, KPGA membership is 43% Black/AA, 30% White, 5% Asian, 4% Hispanic or Latino (18% Unknown/other), which more closely resembles metro Atlanta than overall Georgia.(12) In this study, we conducted a descriptive analysis of KPGA members with suspected and laboratory-confirmed COVID-19. KPGA's robust electronic health record (EHR) data enabled analyses throughout the continuum of care including pre-pandemic underlying disease control, COVID-19 outpatient/inpatient management and post-discharge, with a particular focus on racial/ethnic comparisons. In addition, we

conducted multivariable analyses for hospitalization risk based on demographics, comorbidities, quality of care metrics, lifestyle behaviors and other available individual and community-level social determinants of health indicators.

Methods

For this retrospective cohort study, we performed an EHR review of KPGA members seen with COVID-19 related symptoms between March 3, 2020 and May 12, 2020. Patients were screened according to the Centers for Disease Control (CDC) and Georgia DPH guidelines.(13, 14) Patients who met criteria were tested for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) by polymerase chain reaction (PCR). Due to limits on the testing capacity in Georgia during this period, patients with symptoms or exposures consistent with SARS-CoV-2 were categorized as having been tested or as a person under investigation (PUI). Patients who received tests were further categorized as confirmed or ruled-out. At the start of the epidemic in our region, KPGA prioritized testing among symptomatic health care workers and/or with relevant exposures and symptomatic KPGA members requiring hospital admission. After April 22, testing was progressively expanded to include clinical dispositions dependent on test results (pre-operative clearance, dialysis pending, skilled nurse facility or hospice placement), high risk symptomatic patients based on clinical criteria (>65 years, immunocompromise, chronic obstructive pulmonary disease (COPD), moderate-to-severe asthma, serious heart condition, Body Mass Index (BMI)>40, diabetes, chronic kidney disease (CKD), liver disease, pregnancy) and symptomatic patients with public health implications (non-KPGA healthcare workers, first responders, jail and elder care employees, etc.)

Patient Demographics

We characterized confirmed SARS-CoV-2 and PUI patients by age, sex, self-reported race/ethnicity, insurance type, and area of residence. Race/ethnicity was categorized as non-Hispanic Black/AA (confirmed n=306; 68,3%), non-Hispanic White (n=81; 18%), and Other (n=61; 13,7%), which included Hispanic or Latino (n=16), Asian (n=15), Native American (n=1) and unknown/declined to report (n=29).

We obtained patient's location of residence from the EHR and categorized it into four different regions of metro Atlanta: Northeast, Northwest, Southeast, and Southwest. Residence location was also linked to the neighborhood deprivation index (NDI), a composite SDOH measure including income, education, employment and housing quality.(15, 16) The higher the NDI value, the higher the level of deprivation in the neighborhood.(15, 16) We also utilized ESRI® Business Analyst data, a comprehensive demographic and lifestyle database which provides data to help interpolate patient's socioeconomic status(17). Specifically, we linked patients' places of residency with ESRI's® zip code level classifications of median household income, occupation, and educational attainment. We used this data to cross-reference median household income with the government-defined poverty line.(18)

Comorbidities

Existing comorbidities of SARS-CoV-2 confirmed patients were obtained from the patient's EHR as classified by the International Statistical Classification of Diseases and Related Health Problems codes (ICD-10)(19). The Charlson Comorbidity Index (CCI) was used as a continuous measure of total comorbidity burden.(20) We used pharmacy dispensing data to compile the frequency of outpatient medications used by patients.

We used Healthcare Effectiveness Data and Information Set (HEDIS)(21) as a marker of hypertension (blood pressure reading lower than <140/90mm Hg) and diabetes control (glycated hemoglobin HbA1c < 8%) within a minimum rolling 12-month period.

Using KPGA's Exercise Vital Sign (EVS) data, patient's physical activity levels were classified as inactive, insufficiently active, and sufficiently active for those self-reporting ≤10 minutes, 11-149 minutes, and ≥150 minutes of exercise/week, respectively. The EVS has been previously validated(22) and is considered a clinically relevant screening tool for physical activity behaviors in the health care setting(23, 24).

Clinical Outcomes

Hospitalized patients with confirmed COVID-19 were characterized by hospital length of stay (LOS), ICU LOS, invasive mechanical ventilation initiation and length of use, hospital discharge, readmission, currently hospitalized (n=20; 8.9%), and deceased. Instances of admission and discharge on the same date were defined as LOS of one day. Mechanical ventilation data was compiled using an ICD-10 code flagging instances of emergency endotracheal intubation during hospital stay. Once identified, a manual chart review was conducted for each eligible patient to calculate the length of mechanical ventilation, from intubation to extubation or death. Readmissions were defined as instances of subsequent admission within 30 days to a hospital after recent discharge. We conducted manual record reviews to distinguish between encounters of readmission and patient transfers from a hospital to another medical facility.

Statistical Analysis

We report numbers (percentages) for binary and categorical variables and medians (interquartile ranges, IQR) for continuous variables. Chi-square tests ANOVAs and two sample t-tests were used to determine significant differences between groups. For two sample t-tests with statistically unequal variances, the Satterthwaite method was applied and reported.

Multivariable logistic regression was used to explore factors associated with having a COVID-19 related hospitalization in seven different models: COVID-19 confirmed cases only, confirmed cases plus PUIs, and confirmed cases plus PUIs stratified by race/ethnicity (Black/AA, White, Other) and by sex (Male and Female). All multivariable logistic regression models included age, gender and race/ethnicity as independent variables and hospitalization as the dependent variable. All additional independent variables were assessed using a bivariate analysis, either chi-squared or two sample t-test, and only the variables showing evidence of a statistically significant ($\alpha=0.05$) relationship with the dependent variable were considered for entry into the models. A subset of the dependent variables was considered for the confirmed cases model due to the reduced sample size of the population. Stepwise selection method was used for final dependent variable selection with effect entry and effect remain significance levels of 0.05. All data analysis was conducted using SAS 9.4 software.

The KPGA institutional review board approved this study with a waiver of informed consent.

Results

Epidemiologic Characteristics

Within the study period we screened 6,568 patients, tested 2,920 (44.5%) and 448 (15.3% of tested) patients were positive for SARS-CoV-2. The median age of confirmed positive patients was 54 [IQR, 43-63] years old. Black/AA patients resided in neighborhoods with the highest rate under the federal poverty level (14.2%), unfavorable NDI (0.45), and the highest mean percentage of frontline (35.7%) and healthcare workers (7.5%). (Table 1). The highest percentage of the KPGA members with confirmed SARS-COV-2 resided in the Northeast Metro Atlanta area (31.5%). However, different areas of metro Atlanta showed varying prevalence of KPGA members with confirmed SARS-COV-2 when stratified by race/ethnicity. More Black/AA and Other race patients lived in the Southern areas of metro Atlanta which visible correlates with more socially deprived neighborhoods. (Figure)

Clinical Characteristics

Black/AA patients had higher rates of obesity (67.3%), hypertension (54.9%), and 2 or more comorbidities (66.3%). White patients presented with higher rates of hyperlipidemia (50.6%), congestive heart failure (CHF; 24.7%), coronary artery disease (CAD; 13.6%), arrhythmia (13.8%), CKD (11.1%) and overall CCI Scores (3.2 [2.2]) (all $p < 0.001$) (Table 2).

Compared to other race/ethnicity groups, White patients had the highest rate of diabetes control (14.8%) but Black/AA patients had higher rates of blood pressure control (27.5%) as measured by HEDIS measures. Black/AA patients self-reported the least mean [SD] average weekly exercise minutes in all race/ethnicity groups (61.9 [88.1]; $p < 0.001$). The prevalence of

physical inactivity was higher for both Black/AA and White females compared to Other race females (38.7% vs. 41.5% vs 7.4%; all $p < 0.001$, respectively).

Clinical Outcomes of Hospitalization

Overall, 224 patients with laboratory confirmed COVID-19 were hospitalized with 248 hospital stays, a median age of 60 (50-69) and a median length of stay of 6 (3-11.3) days. (Exhibit 3) There were no significant differences between Black/AA and White patients in ICU admission, ICU LOS, invasive mechanical ventilation and death (8.1% vs. 14.6%). Black/AA females were hospitalized on average 2.4 days longer than white females (95% CI 0.11 to 4.6; $p \leq 0.05$). White females had higher 30-day readmission rates than Black/AA females (17.9% vs. 4%; $p \leq 0.05$). Other race females showed significantly higher rates of invasive mechanical ventilation compared to Black/AA and White females (50% vs. 17% vs 10.7%, $p \leq 0.05$; respectively) (Table 3).

Multivariable Analysis and Factors Associated with Hospitalization

Increasing age was a significant risk factor in all models and females had lower hospitalization odds in the confirmed, confirmed + PUI, Black/AA and Other race models (ORs ranging from 0.33 to 0.51) (Table 4). Black/AA race was a significant factor in the Confirmed + PUI, female and male models (ORs ranging from 1.98 to 2.19). Obesity was associated with higher hospitalization odds in the confirmed, confirmed + PUI, Black/AA and male models (ORs ranging from 1.78 to 2.77). Every point increase in the CCI Index showed increased hospitalization odds in the White model (OR 1.35 95% CI 1.15 to 1.59; $p < 0.001$) while patients with 2 or more Comorbidities had higher hospitalization odds in the Female model (OR 2.38

95% CI 1.43 to 3.94; $p < 0.001$). Cardio-metabolic disease management and control metrics (diabetes, hypertension, hyperlipidemia) were associated with lower odds of hospitalization ranging from 48% to 35% in the confirmed + PUI and Black/AA models.

Self-reported physical inactivity was associated with 50% higher hospitalization odds in the Black/AA and Female models. Residence in the Northeast region of Atlanta was associated with lower hospitalization odds in the Confirmed + PUI, White and female models (ORs ranging from 0.22 to 0.64)

Discussion

This study shows an over-representation of Black/AA populations and other minorities in both the outpatient and inpatient phases of care for COVID-19 in an integrated care system, similar to previous reports (4, 6-9) Although a higher number of Black/AA KPGA members with COVID-19 were hospitalized (69.8%) than other races, there were no significant differences between racial/ethnic groups in ICU admission and duration, invasive ventilation and duration, 30-day readmissions, and mortality. However, further stratification by sex showed Black/AA females were hospitalized on average 2.4 days longer than white females, white females had higher 30-day readmission rates than Black/AA females (17.9% vs. 4%; $p \leq 0.05$) and Other race females had higher rates of invasive mechanical ventilation compared to both Black/AA and White females (50% vs. 17% vs 10.7%, $p \leq 0.05$; respectively). Previous studies reported similar clinical outcomes between Black/AA and non-Black hospitalized COVID-19 patients in Georgia(7, 8) and some but not all previous report have also showed no difference in clinical outcomes between racial/ethnic groups.(9)

Compared to White KPGA members, a higher percentage of Black/AA members with COVID-19 were female, younger, and more likely to reside in neighborhoods with median household income less than \$75,000. Furthermore, Black/AA patients also reside at a higher proportion in neighborhoods with the highest rate of households below the federal poverty level (14.2%), positive neighborhood deprivation index (0.45), and the highest percentage of frontline (35.7%) and healthcare workers (7.5%) compared to other racial groups. All of these factors were associated with an increased risk of COVID-19 infection in our findings. In addition, Black/AA patients had the highest prevalence of obesity, hypertension, and presence of 2 or more comorbidities, all associated with increased disease severity in our analysis, as was found in previous reports.(4) However, we found the comorbidity burden was somewhat different by race, with White patients in our sample being on average older and showing higher CCI scores compared to Black/AA patients, and a different mix of specific underlying conditions (hyperlipidemia, CAD, CHF, arrhythmia, CKD). Although there is a high prevalence of obesity, diabetes and other chronic diseases in the US population,(25-27) these findings suggest that a different comorbidity profile may influence COVID-19 disease severity across racial groups.

Similar to previous reports, our multivariable analysis revealed males were consistently more likely to be hospitalized while increasing age, obesity and hypertension were predominant factors associated with higher odds of hospitalization in all models.(7, 8) Black/AA race, diabetes, COPD, CHF, and CKD were also significant factors in different models. Of note, in the confirmed + PUIs model we detected a significant protective effect of outpatient hypertension and lipid management. Furthermore, Black/AA patients with adequate HEDIS blood pressure and diabetes control were significantly less likely to be hospitalized. Overall, these findings confirm that although the presence of various comorbidities is associated with COVID-19

admission, emphasis on providing adequate clinical management of baseline cardio-metabolic diseases could help ameliorate hospitalization rates. As the pandemic progresses over time, enhanced measures to ensure high quality of care for patients with multiple comorbidities should be reinforced, including those that leverage novel avenues of care including telemedicine and patient-generated actionable data, as well as sustainable linkages with community resources.(28, 29)

Beyond demographic and underlying comorbidity burden and management, our analyses also took into account the potential role of additional SDOH, including indicators of education, economic stability, neighborhood and physical environment and lifestyle behaviors. Of these metrics, we found that residence in the Northeast area of metro Atlanta was one of the most powerful protective factors for hospitalization in the Confirmed + PUI model, particularly for White and female KPGA members. Counties in the NE region have consistently higher levels of safety, quality housing, green space, education and income and have a lower prevalence of obesity compared to the southern regions of KPGA's catchment area(30, 31).

Furthermore, self-reported physical inactivity — engaging in less than 10 minutes of moderate to vigorous exercise/week — increased by 50% the odds of hospitalization among Black/AA and female populations. Several biologic mechanisms may explain this novel association. Physical inactivity is a consistent risk factor for a plethora of chronic diseases shown to also increase COVID-19 severity.(32) Increased inactivity and sedentary time and related comorbidities are also associated with an increased low-grade chronic inflammatory state,(33) which may contribute to the known increased systemic inflammatory effects of SARS-COV-2. In addition to being a modulator of inflammation, regular moderate exercise is also an important immunomodulator, particularly of the virus-fighting cytotoxic immune response.(34) This is

reinforced by epidemiologic studies showing a link between moderate-to-vigorous regular exercise and a lower risk of upper respiratory tract viral infections – including influenza and pneumonia – as well as improved vaccine responses.(35) Although previous reports have shown that self-reported exercise is a predictor of clinical outcomes(24), it is noteworthy that physical inactivity remained a significant correlate of hospitalization risk in our study population, after adjusting for traditional “hard” risk factors such as age, BMI, comorbidity burden and therapeutic management. This reinforces the clinical value of promoting fitness and an active lifestyle, preferably outdoors, to reduce the risk of infection and disease severity of a novel infectious agent such as SARS-COV-2.(36)

This study has some limitations. Limited testing availability in the early stages of the pandemic — globally and in Georgia — led to prioritizing those with the most symptomatic and severe disease requiring admission, as well as testing healthcare workers to prevent further nosocomial infection. For this reason, we included in our analysis not only laboratory-confirmed but also persons under investigation seeking care with COVID-like symptoms. However, we acknowledge that not all PUIs would necessarily have SARS-COV-2. The target population in this analysis included only KPGA members that by definition have insurance and ready access to health care services. However, our analysis showed a diverse socioeconomic background of KPGA members. Merging racial/ethnic groups with low sample sizes into a combined “Other” race category was necessary for statistical power reasons but limits the interpretation of findings for this group. Finally, despite having some SDOH indicators in our member’s EHR, we also included neighborhood level data to extrapolate additional SDOH metrics. Ongoing investigation of drivers in COVID-19 disparities will benefit from more individual level SDOH data. Despite these limitations, by integrating underlying chronic disease management history, outpatient

information, hospitalization, clinical outcomes and post-discharge follow-up data, this study provides a comprehensive longitudinal assessment of COVID-19 patients in relation to racial/ethnic disparities.

To our knowledge, this investigation is the first COVID-19 retrospective cohort to include a multivariate analysis on multiple measures of SDOH and pre-pandemic comorbidity management. Our study suggests that, within our sample of KPGA members with ready access to insurance and high quality of care within an integrated health system, Black/AA members were still being disproportionately affected by COVID-19 risk of infection and hospitalization. However, we found no significant differences in clinical outcomes such as ICU length of stay or mortality across race/ethnicity groups. Location of residence, a proxy for the overall community context of our patients, appears to be a factor strongly associated with increased infection risk among Black/AA and other minorities. SDOH have also shown to contribute to a more unfavorable baseline health status and therefore, can indirectly impact COVID-19 risk of hospitalization and severity.(6) In addition to age, sex, location of residence and presence of comorbidities, pre-pandemic self-reported exercise levels and underlying cardio-metabolic disease control may also significantly impact hospitalization risk in different race groups. Therefore, beyond well-known physiologic and clinical factors, individual and community-level social factors and health behaviors must be considered by clinicians, health care systems(37) and public health stakeholders (6) as interventions designed to reduce COVID-19 disparities and the systemic effects of racism(38) are implemented.

Acknowledgements: Special thanks to all the clinicians, providers and staff of the Southeast Permanente Medical Group and Kaiser Permanente Georgia.

References

1. World Health Organization. Coronavirus disease (COVID-19) Situation Report – 164 July 2nd 2020 [Available from: https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200702-covid-19-sitrep-164.pdf?sfvrsn=ac074f58_2.
2. Price-Haywood EG, Burton J, Fort D, Seoane L. Hospitalization and Mortality among Black Patients and White Patients with Covid-19. N Engl J Med 2020;382(26):2534-2543 doi:101056/NEJMsa2011686. 2020.
3. Azar KMJ, Shen Z, Romanelli RJ, Lockhart SH, Smits K, Robinson S, et al. Disparities In Outcomes Among COVID-19 Patients In A Large Health Care System In California. [published online ahead of print, 2020 May 21]. Health Aff (Millwood) 2020;101377hlthaff202000598 doi:101377/hlthaff202000598. 2020.
4. Suleyman G, Fadel RA, Malette KM, Hammond C, Abdulla H, Entz A, et al. Clinical Characteristics and Morbidity Associated With Coronavirus Disease 2019 in a Series of Patients in Metropolitan Detroit. JAMA Netw Open. 2020;3(6):e2012270.
5. Millett GA, Jones AT, Benkeser D, Baral S, Mercer L, Beyrer C, et al. Assessing Differential Impacts of COVID-19 on Black Communities [published online ahead of print, 2020 May 14]. Ann Epidemiol 2020;101016/jannepidem202005003 doi:101016/jannepidem202005003. 2020.
6. Centers for Disease Control and Prevention. National Center for Immunization and Respiratory Diseases (NCIRD) DoVD. COVID-19 in Racial and Ethnic Minority Groups

- Atlanta, GA2020 [Available from: <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/racial-ethnic-minorities.html>.]
7. Gold JA WK, Szablewski CM. Characteristics and Clinical Outcomes of Adult Patients Hospitalized with COVID-19 — Georgia, March 2020. MMWR Morb Mortal Wkly Rep 2020. March 2020;69(18):545-50.
8. Killerby ME L-GR, Haight SC. Characteristics Associated with Hospitalization Among Patients with COVID-19 — Metropolitan Atlanta, Georgia, March–April 2020. MMWR Morb Mortal Wkly Rep 2020. 2020.
9. Rentsch CT, Kidwai-Khan F, Tate JP, Park LS, King Jr JT, Skanderson M, et al. Covid-19 by Race and Ethnicity: A National Cohort Study of 6 Million United States Veterans. Preprint. medRxiv 2020;2020051220099135 Published 2020 May 18
doi:101101/2020051220099135. 2020.
10. United States Census Bureau. Georgia 2020 [Available from: <https://data.census.gov/cedsci/profile?q=Georgia&g=0400000US13&tid=ACSDP1Y2018.DP05>.]
11. Georgia Department of Public Health. Georgia Department of Public Health Daily Status Report 2020 [updated 6/6/2020. Available from: <https://dph.georgia.gov/covid-19-daily-status-report>.]
12. Metro Atlanta Chamber. Profile of Metro Atlanta 2020 [Available from: <https://www.metroatlantachamber.com/resources/reports-and-information/executive-profile>.]

13. National Center for Immunization and Respiratory Diseases (NCIRD) Division of Viral Diseases. Standard Operating Procedure (SOP) for Triage of Suspected COVID-19 Patients in non-US Healthcare Settings: Early Identification and Prevention of Transmission during Triage: Centers for Disease Control and Prevention; 2020 [updated May 28th 2020. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/non-us-settings/sop-triage-prevent-transmission.html>.
14. Georgia Department of Public Health. COVID-19: Guidance for Healthcare Professionals 2020 [updated April 16, 2020. Available from: <https://dph.georgia.gov/covid-19-guidance-healthcare-professionals>.
15. Andrews MR, Tamura K, Claudel SE, Xu S, Ceasar JN, Collins BS, et al. Geospatial analysis of neighborhood deprivation index (NDI) for the United States by county. Journal of Maps. 2020;16(1):101-12.
16. Messer LC, Laraia BA, Kaufman JS, Eyser J, Holzman C, Culhane J, et al. The development of a standardized neighborhood deprivation index. J Urban Health. 2006;83(6):1041-62.
17. ArcGIS. Esri Demographics - Tapestry Segmentation: ArcGIS; 2019 [updated 2019. Available from: <https://doc.arcgis.com/en/esri-demographics/data/tapestry-segmentation.htm>.
18. Office of the Assistant Secretary for Planning and Evaluation. U.S. Federal Poverty Guidelines Used to Determine Financial Eligibility for Certain Federal Programs: U.S Department of Health & Human Services; 2020 [updated January 17th 2020. Available from: <https://aspe.hhs.gov/poverty-guidelines>.

- 405 19. World Health Organization. The ICD-10 classification of mental and behavioural
406 disorders: diagnostic criteria for research 1993.
- 407 20. Deyo RA, Cherkin DC, Ciol MA. Adapting a clinical comorbidity index for use with
408 ICD-9-CM administrative databases. Journal of clinical epidemiology. 1992;45(6):613-9.
- 409 21. National Committee for Quality Assurance (NCQA). HEDIS Measures and Technical
410 Resources 2020 [Available from: <https://www.ncqa.org/hedis/measures/>].
- 411 22. Coleman KJ, Ngor E, Reynolds K, Quinn VP, Koebnick C, Young DR, et al. Initial
412 validation of an exercise “vital sign” in electronic medical records. Medicine & Science in Sports
413 & Exercise. 2012;44(11):2071-6.
- 414 23. Young DR, Coleman KJ, Ngor E, Reynolds K, Sidell M, Sallis RE. Associations between
415 physical activity and cardiometabolic risk factors assessed in a Southern California health care
416 system, 2010-2012. Preventing chronic disease. 2014;11:E219.
- 417 24. Lobelo F, Rohm Young D, Sallis R, Garber MD, Billinger SA, Duperly J, et al. Routine
418 Assessment and Promotion of Physical Activity in Healthcare Settings: A Scientific Statement
419 From the American Heart Association. Circulation. 2018;137(18):e495-e522.
- 420 25. Hales CM, Carroll MD, Fryar CD, Ogden CL. Prevalence of obesity and severe obesity
421 among adults: United States, 2017–2018. NCHS Data Brief, no 360 Hyattsville, MD: National
422 Center for Health Statistics 2020. 2020.
- 423 26. Centers for Disease Control and Prevention. National Diabetes Statistics Report A, GA:
424 Centers for Disease Control and Prevention, U.S. Dept of Health and Human Services; 2020. . . .

27. Clark A, Jit M, Warren-Gash C, Guthrie B, Wang HHX, Mercer SW, et al. Global, regional, and national estimates of the population at increased risk of severe COVID-19 due to underlying health conditions in 2020: a modelling study. The Lancet Global health. 2020;published online ahead of print, 2020 Jun 15]. Lancet Glob Health. 2020;S2214-109X(20)30264-3. doi:10.1016/S2214-109X(20)30264-3.
28. Ceriello A, Schnell O. COVID-19: Considerations of Diabetes and Cardiovascular Disease Management. Journal of diabetes science and technology. 2020:1932296820930025.
29. Shabto JM, Loerinc L, O'Keefe GA, O'Keefe J. Characteristics and outcomes of COVID-19 positive patients with diabetes managed as outpatients. Diabetes research and clinical practice. 2020;164:108229.
30. Emory University. COVID-19 Health Equity Interactive Dashboard 2020 [Available from: <https://covid19.emory.edu/>].
31. County Health Rankings & Roadmaps. 2020 Georgia Report 2020 [Available from: <https://www.countyhealthrankings.org/reports/state-reports/2020-georgia-report>].
32. Powell KE, King AC, Buchner DM, Campbell WW, DiPietro L, Erickson KI, et al. The Scientific Foundation for the Physical Activity Guidelines for Americans, 2nd Edition. J Phys Act Health. 2018:1-11.
33. Henson J, Yates T, Edwardson CL, Khunti K, Talbot D, Gray LJ, et al. Sedentary time and markers of chronic low-grade inflammation in a high risk population. PLoS One. 2013;8(10):e78350.

34. Nieman DC, Wentz LM. The compelling link between physical activity and the body's defense system. *Journal of sport and health science*. 2019;8(3):201-17.
35. Song Y, Ren F, Sun D, Wang M, Baker JS, Istvan B, et al. Benefits of Exercise on Influenza or Pneumonia in Older Adults: A Systematic Review. *Int J Environ Res Public Health*. 2020;17(8).
36. Sallis J, Pratt M. A Call to Action: Physical Activity and COVID-19 Exercise is Medicine2020 [updated April 3, 2020. Available from: https://www.exerciseismedicine.org/support_page.php/stories/?b=896.
37. Parodi S CB, Young S, Bellows J, Grossman D, Liu VX. Kaiser Permanente's system capabilities to suppress Covid-19. *NEJM Catal* Published June 9, 2020 doi:101056/CAT200187.
38. Bryan A D-GJ, Davis NJ, Chokshi DA, Galea S. Moving From The Five Whys To Five Hows: Addressing Racial Inequities In COVID-19 Infection And Death. *Health Aff (Millwood)* 2020; 101377/hblog20200625389260

Table 1: Socio-economic and demographic characteristics of KPGA members with confirmed SARS-COV-2 seen from March 3 to May 12, 2020

Patients Stratified by Race ^a No. (%)					
	All (N= 448)	Black/AA (n= 306)	White (n= 81)	Other (n= 61)	p-value
Age, median [IQR], years	54 [43-63]	54 [43-62]	58 [49-73]	51 [37-61]	<.001
Age range, years					<.001
18-49	169 (37.7)	116 (37.9)	23 (28.4)	30 (49.2)	
50-64	177 (39.5)	132 (43.1)	24 (29.6)	21 (34.4)	
65 and above	102 (22.8)	58 (19.0)	34 (42.0)	10 (16.4)	
Gender					<.001
Male	176 (39.3)	102 (33.3)	40 (49.4)	34 (55.7)	
Female	272 (60.7)	204 (66.7)	41 (50.6)	27 (44.3)	
Insurance					
Commercial	282 (62.9)	194 (63.4)	43 (53.1)	45 (73.8)	0.11
Medicare	94 (21)	61 (19.9)	26 (32.1)	7 (11.5)	
Self-pay	44 (9.8)	31 (10.1)	7 (8.6)	6 (9.8)	
Other ^b	28 (6.3)	20 (6.6)	5 (6.2)	3 (4.9)	
Median Household Income ^c No.	439	299	80	60	
25k-50k	95 (21.6)	74 (24.7)	9 (11.3)	12 (20)	<.001
50k-75k	230 (52.4)	178 (59.5)	27 (33.8)	25 (41.7)	
75k-100k	88 (20)	43 (14.4)	29 (36.3)	16 (26.7)	
100k+	26 (5.9)	4 (1.3)	15 (18.8)	7 (11.7)	
Households Under Poverty Level, % ^d	13.1	14.2	10.0	12.0	<.001
Residential Region, No. (%) ^e	308	208	51	49	
Northeast	97 (31.5)	55 (26.4)	15 (29.4)	27 (55.1)	<.001
Northwest	46 (14.9)	23 (11.1)	19 (37.3)	4 (8.2)	
Southeast	84 (27.3)	65 (31.3)	6 (11.8)	13 (26.5)	
Southwest	81 (26.3)	65 (31.3)	11 (21.6)	5 (10.2)	
Neighborhood Deprivation Index ^f	0.21	0.45	-0.48	-0.07	<.001
Occupation ^g, mean % ^h					
Frontline Workers	34.4	35.7	29.6	34.3	<.001
Healthcare Workers	7.3	7.5	6.9	6.8	<.001
Other Workers	58.3	56.8	63.5	58.9	<.001
Education, mean % ^h					
Some High School	6.8	7.2	5.5	6.5	<.001
High School	22.9	24.4	18.9	20.9	<.001
Associates Degree	8.4	8.6	7.8	8.3	<.001
Some College	21	22	18.7	19.3	<.001
Bachelors	21.1	19.2	26.3	23.4	<.001
Graduate	12.4	11.4	15.6	13.4	<.001

Abbreviations: IQR, Interquartile Range; COVID-19, Coronavirus Disease 2019; KPGA, Kaiser Permanente Georgia; AA, African American.

^a Other is defined as all other racial/ethnic groups (Asian, Hispanic/Latino), Unknown, and those who declined to report their race.

^b Other Insurances include military Health Maintenance Organization (HMO) or Preferred Provider Organization (PPO).

^c Based on ESRI® Business Analyst dataset showing median household income by zip code and then linked to individual patients based on their recorded residence.¹⁹

^d Poverty line was defined by the Federal poverty level ¹⁸

^e The Atlanta metro area was divided up by county in four sub-regions

- Northwest: Cobb, Cherokee, Paulding, Bartow, Pickens, Polk, Troup, Habersham.
- Northeast: Dekalb, Gwinnett, Forsyth, Hall, Barrow, Jackson, Butts, Gilmer, Pike, Gordon, Jasper, Monroe.
- Southwest: Fulton, Douglas, Fayette, Coweta, Carroll, Meriwether, Heard, Dawson, Madison, Lumpkin.
- Southeast: Clayton, Henry, Rockdale, Walton, Clarke, Spalding, Oconee, Muscogee, Brooks, Town.

^f The Neighborhood Deprivation Index (NDI) is a composite measure of social and economic factors such as income, education, employment and housing quality that reflect neighborhood deprivation.^{13,14} The higher the index value, the higher the level of deprivation in the neighborhood.

^g Based on ESRI® Business Analyst data.¹⁶ Occupation Breakdown:

- Frontline workers included community/social services, protective services, food preparation/serving related services, building/grounds cleaning/maintenance services, construction/extraction services, installation/maintenance/repair services, production services and transportation/material moving services.
- Healthcare workers included healthcare practitioners/technicians and healthcare support staff.
- Other workers included personal care/service workers, sales and sales related workers, office/administrative support workers, farming/fishing/forestry workers, management/business/financial workers, computer/mathematical service workers, architecture/engineering workers, life/physical/social science workers, community/social service workers, legal workers, education/training/library workers and arts/design/entertainment/sports/media workers.

^h Based on ESRI® Business Analyst data. It is expressed in mean percentage and provides the counts of individual education attainment and occupation by category within each zip code (denominator).¹⁹

Table 2: Comorbidities, outpatient medication, history of disease control, & exercise frequency of KPGA members with confirmed SARS-COV-2 by race & sex

	Race/Ethnicity ^a									
	All	Black/AA No. (%)			White No. (%)			Other No. (%)		
	(N=448)	Total (n=306)	Male (n=102)	Female (n=204)	Total (n=81)	Male (n=40)	Female (n=41)	Total (n=61)	Male (n=34)	Female (n=27)
Comorbidities										
Hypertension	226 (50.5)	168 (54.9) *†, *°°	61 (59.8) *‡‡	107(52.5) *EE	37(45.7) *†	21 (52.5)	16 (39) *†, *°°	21 (34.4) *†, *°°	13 (38.2) *‡‡	8 (29.6) *EE
Diabetes	116 (25.9)	86 (28.1) *°°	40 (39.2) §, *‡‡	46 (22.5) §	21(25.9)	14 (35) *‡‡‡‡	7 (17.1) *°°	9 (14.8) *°°	5 (14.7) *‡‡, *‡‡‡‡	4 (14.8)
Obesity (BMI>30)	273 (60.9)	207 (67.3) **†, *°, *°°	70(68.6) *‡‡	137(67.2) *E	39(48.1) **†, *°	21 (52.5)	18 (43.9) *E	28 (45.9) **†, *°°	15 (44.1) *‡‡	13 (48.1)
Hyperlipidemia	150 (33.5)	95 (31) **†, *°	41 (40.2) §, *‡	54 (52.9) §, *E	41(50.6) **†, *°, *°°°	23 (57.5) *‡, *‡‡‡‡	18 (43.9) *E, *EEE	14 (23) **†, *°°°	9 (26.5) *‡‡‡‡	5 (18.5) *EEE
CAD	34 (7.6)	19 (6.2) *°	11 (10.8) §	8 (3.9) §, *E	11(13.6) *°	6 (15)	5 (12.2) *E	4 (6.6)	3 (8.8)	1 (3.7)
CHF	61 (13.6)	38 (12.4) *†, *°	17 (16.7) *‡‡	21 (10.3) *E	20(24.7) *†, *°, *°°°	10 (25)	10 (24.4) *E	3 (4.9) *†, *°°°	1 (2.9) *‡‡	2 (7.4)
Asthma	47 (10.5)	37 (11.8) *†, *°	10 (9.8)	26 (12.7)	2(2.5) *†, *°, *°°°	1 (2.5) *‡‡‡‡	1 (2.4) *EEE	9 (14.8) *†, *°°°	2 (5.9) ¶, *‡‡‡‡	7 (25.9) ¶, *EEE
COPD	28 (6.3)	13 (4.2) *†, **°	5 (4.9)	8 (3.9) **E	12(14.8) *†, **°	2 (5) #	10 (24.4) #, **E	3 (4.9) *†	1 (2.9)	2 (7.4)
Arrhythmia	29 (6.6)	16 (5.3) *†, *°	7 (6.9) *‡	9 (4.4)	11(13.8) *†, *°, *°°°	8 (20) *‡, *‡‡‡‡	3 (7.3)	2 (3.4) *†, *°°°	1 (2.9) *‡‡‡‡	1 (3.7)
ESRD ^b	1 (0.2)	1 (0.3)	1 (1)	0 (0)	0(0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
HIV	6 (1.3)	5 (1.6)	4 (3.9) §	1 (0.5) §	1(1.2)	1 (2.5)	0 (0)	0 (0)	0 (0)	0 (0)
Depression	69 (15.4)	45 (14.7)	5 (4.9) §	40 (19.6) §	15(18.5)	3 (7.5) #	12 (29.3) #	9 (14.8)	5 (14.7)	4 (14.8)
CKD ^c	25 (5.6)	15 (4.9) *†, *°	6 (5.9)	9 (4.4) *E	9(11.1) *†, *°, *°°°	3 (7.5)	6 (14.6) *E	1 (1.6) *†, *°°°	0 (0)	1 (3.7)
Cancer	15 (3.4)	12 (4)	7 (6.9)	5 (2.5)	2(2.5)	0 (0)	2 (4.9)	1 (1.6)	1 (2.9)	0 (0)
2+ Comorbidities ^d	281 (62.7)	203 (66.3) *†, *°°	72 (70.6) *‡‡	131(64.2)	50(61.7) *†	26 (65)	24 (58.5)	28 (45.9) *†, *°°	15 (44.1) *‡‡	13 (48.2)
3+ Comorbidities ^d	179 (40)	124 (40.5)	48 (47.1)	76 (37.3)	38(46.9) *°°°	20 (50)	18 (43.9)	17 (27.9) *°°°	11 (32.3)	6 (22.2)
CCI, mean [SD] ^e	2.4 [1.8]	2.3 [1.6] **†, *°, *°°	2.7 [1.8]	2.1 [1.5] *E	3.2[2.2] **†, *°, *°°°°	3 [1.8] *‡‡‡‡	3.4 [2.5] *E, *EEE	1.5 [1] **†, *°, *°°°°	1.6 [1.0] *‡‡‡‡	1.36 [1.1] *EEE

511

512

Table 2: Comorbidities, outpatient medication, history of disease control, & exercise frequency of KPGA members with confirmed SARS-COV-2 by race & sex

	Race/Ethnicity ^a									
	All	Black/AA No. (%)			White No. (%)			Other No. (%)		
	(N=448)	Total (n=306)	Male (n=102)	Female (n=204)	Total (n=81)	Male (n=40)	Female (n=41)	Total (n=61)	Male (n=34)	Female (n=27)
Outpatient Medication, No. (%)										
Anti-Rheumatic	2 (0.4)	1 (0.3)	1 (1)	0 (0) ***	0(0)	0 (0)	0 (0)	1 (1.6)	0 (0)	1 (3.7) **
Anti-Hypertensive	129 (28.8)	91 (29.7)	34 (33.3)	57 (27.9)	24(29.6)	15 (37.5)	9 (22)	14 (23)	7 (20.6)	7 (25.9)
Anti-Asthmatic	103 (23)	72 (23.5)	27 (26.5)	45 (22.1) ***	17(21)	9 (22.5)	8 (19.5) ****	14 (23)	4 (11.8) ¶	10 (37) ¶, **, ****
Anti-Hyperlipidemic	125 (27.9)	81 (26.5)	40 (39.2)	41 (20.1)	29(35.8)	17 (42.5)	12 (29.3)	15 (24.6)	9 (26.5)	6 (22.2)
Corticosteroids	151 (33.7)	112 (36.6)	37 (36.3)	75 (36.8)	22(27.2)	9 (22.5)	13 (31.7)	17 (27.9)	7 (20.6)	10 (37)
Anti-malarial	6 (1.3)	4 (1.3)	0 (0)	4 (2)	1(1.2)	0 (0)	1 (2.4)	1 (1.6)	0 (0)	1 (3.7)
HEDIS Measures ^f										
Blood Pressure Control ^g	113 (25.2)	84 (27.5) *°	35 (34.3)	49 (24)	18(22.2) *°, ****	11 (27.5)	7 (17.1)	11 (18) ****	5 (14.7)	6 (22.2)
Diabetes Control HbA1c < 8% ^h	55 (12.3)	37 (12.1) *†, *°	14 (13.7) *¶	23 (11.3)	12(14.8) *†, *°	7 (17.5) *¶	5 (12.2)	6 (9.8) *†	3 (8.8)	3 (11.1)
Weekly Exercise, mean [SD], mins ⁱ										
	70.1 [101.8]	61.9 [88.1] *†, ***	86.7 [104.5]	49.5 [76.1] ***	75.7 [118.5] *†	76.4 [97.74]	75.2 [136.2]	108.4 [134.9] *†, ***	98.4 [126.8]	123 [148.9] ***
Exercise Vital Sign Category, No. (%) ^j										
Inactive	151 (33.7)	106 (34.6) **†, ***	27 (26.5)	79 (38.7) ***	32(39.5) **†, ****	15 (37.5)	17 (41.5) ****	13 (21.3) **†, **, ****	11 (32.3)	2 (7.4) ***, ****
Insufficiently Active	144 (32.1)	103 (33.7) **†	34 (33.3)	69 (33.8)	23(28.4) **†	11 (27.5)	12 (29.3)	18 (29.5) **†	7 (20.6)	11 (40.8)
Sufficiently Active	64 (14.3)	37 (12.1) **†	21 (20.6)	16 (7.8) **	16(19.8) **†	8 (20)	8 (19.5) **	11 (18) **†	7 (20.6)	4 (14.8)
No information	89 (19.9)	60 (19.6) **†, ***	20 (19.6)	40 (19.7) ***	10(12.3) **†, ****	6 (15)	4 (9.7) ****	19 (31.2) **†, **, ****	9 (26.5)	10 (37) ***, ****

513

514

515

516

517

518

Abbreviations: AA, African American; HbA1c, Glycated Hemoglobin; BMI, Body Mass Index; CAD, Coronary Artery Disease; CCI, Charlson Comorbidity Index; CHF, Congestive Heart Failure; CKD, Chronic Kidney Disease; COPD, Chronic Obstructive Pulmonary Disease; ESRD, End Stage Renal Disease;

Significance Levels * $P \leq 0.05$, ** $P \leq 0.001$

Significance Tests

† Across Race Groups

519 ° Black/AA vs. White, °° Black/AA vs. Other, °°° White vs. Other
 520 § Black/AA Male vs. Black/AA Female
 521 # White Male vs. White Female
 522 ¶ Other Male vs. Other Female
 523 º Black/AA Female vs. White Female, ºº Black/AA Female vs. Other Female, ººº White Female vs Other Female
 524 º Black/AA Male vs. White Male, ºº Black/AA Male vs. Other Male, ººº White Male vs. Other Male
 525
 526 º Other Race is defined as all other racial/ethnic groups (Asian, Hispanic/Latino), Unknown, and those who declined to report their race.
 527 º ESRD classified based on ICD-10 in patient’s medical history.²⁰
 528 º CKD classified based on diagnosis reported by the ICD-10 code in patient’s medical history.²⁰
 529 º Comorbidities are medical diagnoses included in medical history as ICD-10 codes. These include but are not limited to those presented in the table.
 530 º Charlson Comorbidity Index. The lowest score of 0 corresponds to a 98% estimated 10-year survival rate.²¹
 531 º Healthcare Effectiveness Data and Information Set (HEDIS), is a performance improvement tool used by healthcare organizations in the United States.²² For this study, this tool was used to
 532 identify the proportion of COVID-19 confirmed patients with their chronic disease successfully controlled. Of note, the total number of people with diabetes (n=116) included in this analysis is
 533 larger than the total number of diabetics reported by HEDIS measure (n=55). This discrepancy is also observed when comparing total number of people with hypertension in this analysis (n=226)
 534 with total number of hypertensive patients reported by HEDIS measure (n=116). Explanations for these differences in sample size are related to lack of continuous membership enrollment and
 535 the likelihood of no interactions with patient within the last year. These reasons could explain the lack of documentation of HEDIS disease control.
 536 º HEDIS measure for blood pressure control (<140/90mm HG) for KPGA members with a 12-month rolling enrollment.²²
 537 º HEDIS measure for diabetes control based on a HbA1C < 8% for KPGA members with a 12-month rolling enrollment.²²
 538 º Average exercise was collected from self-reported data during a clinical encounter in the last year.
 539 º Exercise Vital Sign is based on patient reported weekly exercise minutes. Three categories are coded: Inactive (< 10 mins/week), insufficiently active (11-149 minutes/week) and sufficiently active
 540 (150 or more mins/week)²⁴
 541
 542
 543
 544
 545
 546
 547
 548
 549
 550
 551
 552

553 **Table 3: Clinical outcomes of hospitalized KPGA members with confirmed COVID-19 by race & age groups**

	Race/Ethnicity ^b									Age group (years)			
	No. (%)									No. (%)			
	All No. (%)	Black/AA			White			Other			18-49	50-64	65+
		Total	Male	Female	Total	Male	Female	Total	Male	Female			
Hospitalization Characteristics	N = 248 ^c	n=173	n=79	n=94	n=48	n=20	n=28	n=27	n=17	n=10	n=62	n=97	n=89
Age of Hospitalized Patients, median (IQR) ^d , years	60 (50.0-69.0)	59 (49.5-66.0)	59 (49.5-66.0)	59 (49.5-66.0)	67 (54.0-75.5)	66.5 (52.5-74.5)	68 (55.5-77.0)	60.5 (39.5-65.5)	62 (37.0-66.0)	52 (41.0-60.0)	n/a	n/a	n/a
Hospital Length of Stay, median (IQR), days	6 (3.0-11.3)	6 (3.0-11.0)	7 (3.0-12.0)	6 (3.0-11.0) * ^e	6 (2.8-10.3)	7.5 (3.0-14.5)	4 (2.0-8.5)	5 (2.0 - 17.0)	4 (2.0-14.0)	12 (2.0-20)	5 (3.0-8.0) * [¶] ,	6 (3.0-12.0) * [¶] , * ^{◊◊}	6 (3.0-14.0) * [¶] , * [◊]
Admitted to ICU	104 (41.9)	69 (39.9)	33 (41.8)	36 (38.3)	19 (39.6)	9 (45.0)	10 (35.7)	16 (59.3)	9 (52.9)	7 (70.0)	20 (32.3)	44 (45.4)	40 (44.9)
ICU Length of Stay, median (IQR), days	7 (2.0-13.0)	6 (2.0-13.3)	6 (3.0-14.0)	7 (1.0-13.0)	8 (1.0-10.5)	10 (8.0-13.0)	5.5 (1.0-8.0)	9 (4.0-18.0)	8.5 (2.0-18.5)	9 (5.0-19.0)	4 (1.0-16.0)	6 (3.0-12.0)	9 (3.5-16.0)
Invasive Mechanical Ventilation	47 (19.0)	32 (18.5)	16 (20.3)	16 (17.0)	6 (12.5)	3 (15.0)	3 (10.7)	9 (33.3) * ^{°°°}	4 (23.5)	5 (50.0) * ^{°°°} , * ^{°°°°}	5 (8.1)	20 (20.6)	22 (24.7)
Ventilator duration, median (IQR), days	13 (8.0-16.5)	13 (8.0-15.5)	11.5 (6.0-15.5)	13 (9.5-16.0)	10.5 (6.0-16.0)	13 (4.0-24.0)	8 (6.0-16.0)	11 (8.0-14.0)	12.5 (9.5-16.5)	9 (8.0-14.0)	14 (13.0-15.0) * [¶]	10.5 (6.5-13.0) * [¶]	15.0 (8.0-19.0) * [¶] , * ^{◊◊◊}
Outcomes													
Discharged alive	206 (83.0)	149 (86.1)	65 (82.3)	84 (89.4)	37 (77.1)	16 (80.0)	21 (75.0)	20 (74.1)	12 (70.6)	8 (80.0)	60 (96.8) ** [¶]	86 (88.7) ** [¶]	60 (67.4) ** [¶]
Still Hospitalized	20 (8.1)	10 (5.8) * [†]	6 (7.6)	4 (4.3)	4 (8.3) * [†]	1 (5.0)	3 (10.7)	6 (22.2) * [†]	5(29.4) * [‡] , * ^{‡‡‡}	1 (10.0)	2 (3.2)	7 (7.2)	11 (12.4)
30-day Readmission Rate	13 (5.2)	7 (4.0)	4 (5.1)	3 (3.2)	5 (10.4)	0 (0.0)	5 (17.9) * ^ε	1 (3.7)	1 (5.9)	0 (0.0)	2 (3.2)	2 (2.1)	6 (6.7)
Deceased	22 (8.9)	14 (8.1)	8 (10.1)	6 (6.4)	7 (14.6)	3 (15.0)	4 (14.3)	1 (3.7)	0 (0.0)	1 (10.0)	0 (0.0) ** [¶]	4 (4.1) ** [¶]	18 (20.2) ** [¶]

554 **Abbreviations:** ICU, Intensive Care Unit; IQR, interquartile range; n/a = not applicable; AA, African American

555 ^a Significance Levels * P ≤ 0.05, ** P ≤ 0.001; Significance Tests (Categorical: Chi-Squared Test of Significance, Continuous: ANOVA Test of Significance, Two-Sample T-Test of Means)

556 ^b Other is defined as all other racial/ethnic groups (Asian, Hispanic/Latino), Unknown, and those who declined to report their race.

557 ^c N=248 includes patients at level of hospital stay. Thus, participants who were readmitted, or transferred, are accounted for more than once.

558 ^d Median age represents all unique hospitalized patients n=224 (Black n=160, White n=40, Other n=24); ^e In this table, column percentages are provided for categorical variables.

559 [†] Across Race Groups, [°] Other vs. Black, ^{°°°} White vs. Other, [¶] Across Age Groups, [◊] Age 18-49 vs. 50-64, ^{◊◊} Age 18-49 vs. 65+, ^{◊◊◊} Age 50-64 vs. 65+

560 ^ε Black/AA Female vs. White Female, ^{°°} Black/AA Female vs. Other Female, ^{°°°°} White Female vs Other Female; [‡] Black/AA Male vs. Other Male, ^{‡‡‡} White Male vs. Other Male

561 **Table 4: Multivariable logistic regression model odds ratios for hospitalization by COVID-19 status, Rrce, and sex**

Sample Population	SARS-COV-2 Status		Race/Ethnicity Groups			Sex Groups	
	Confirmed	Confirmed & PUI	Black/AA	White	Other ^a	Female	Male
Total sample size	n=448	n=3,937	n=2,156	n=981	n=800	n=2,536	n=1,401
Variables, OR (95% CI)							
Age	1.03**(1.02,1.05)	1.05**(1.04,1.06)	1.04**(1.03,1.06)	1.05**(1.02,1.08)	1.06**(1.03,1.09)	1.04**(1.03,1.06)	1.06**(1.05,1.08)
Race: Black		1.98**(1.44,2.72)	n/a ^b	n/a ^b	n/a ^b	2.19**(1.40,3.42)	2.03*(1.31,3.13)
Female	0.42**(0.27,0.65)	0.51**(0.38,0.68)	0.52**(0.37,0.73)		0.33*(0.14,0.77)	n/a ^c	n/a ^c
Obesity (BMI ≥30)	1.87*(1.19,2.93)	1.80**(1.31,2.47)	1.78*(1.21,2.64)				2.77**(1.78,4.32)
CCI ^d				1.35**(1.15,1.59)			
CHF ^e		1.83*(1.19,2.80)	1.99*(1.20,3.31)			2.65**(1.54,4.52)	
Hypertension ^e	2.09*(1.33,3.29)	1.76*(1.21,2.55)	2.02*(1.26,3.21)				
CKD ^e	11.04*(1.39,87.60)						
Diabetes ^e		1.76*(1.23,2.51)	2.00*(1.27,3.13)				
Physically Inactive ^f			1.50*(1.05,2.13)			1.49*(1.00,2.20)	
COPD ^e	5.37*(1.45,19.81)	1.72*(1.02,2.90)					
2+ Comorbidities ^g						2.38**(1.43,3.94)	
Antihypertensive		0.65*(0.46,0.94)					
Antihyperlipidemic		0.65*(0.45,0.95)					
HEDIS ^h BP Control ⁱ			0.62*(0.41,0.94)				
HEDIS ^h Diabetes Control ^j			0.52*(0.29,0.93)				
NE County Residence ^k		0.64*(0.47,0.89)		0.22*(0.07,0.66)		0.61*(0.38,0.95)	

562 **Abbreviations:** OR, Odds Ratio; CI, Confidence Interval; PUI, Persons Under Investigation; n/a, not applicable; BMI, Body Mass Index; AA, African American; CCI, Charlson Comorbidity Index; CHF, Congestive Heart Failure; CKD, Chronic Kidney Disease; COPD, Chronic Obstructive Pulmonary Disease; HEDIS, Healthcare Effectiveness Data and Information Set; BP: blood pressure
563
564 Significance Levels * P ≤ 0.05, ** P ≤ 0.001

565 Odds Ratios (OR) represent yes vs no for all variables except Age (per year) and Charlson Comorbidity Index (per point)
566 ^a Other is defined as all other racial/ethnic groups (Asian, Hispanic/Latino), Unknown, and those who declined to state race
567 ^b Race Black/AA is not available as independent variable for race stratified models
568 ^c Gender Female not available as an independent variable for the gender stratified models
569 ^d Charlson Comorbidity Index ²¹
570 ^e CHF, Hypertension, CKD, Diabetes, and COPD classified based on ICD-10 diagnosis in member's electronic medical record²⁰
571 ^f Physically Inactive defined as self-reported exercise < 10 minutes/week.²⁴
572 ^g Comorbidities here are medical diagnoses included in medical history as ICD-10 codes.²⁰
573 ^h HEDIS is a performance improvement tool used by healthcare organizations in the United States.²² For this study, this tool was used to identify the proportion of members with their chronic
574 conditions successfully controlled
575 ⁱ HEDIS measure for blood pressure control (<140/90mm HG) for KPGA members with a 12-month rolling enrollment.²²
576 ^j HEDIS measure for diabetes control based on a glycated hemoglobin HbA1c < 8% for KPGA members with a 12-month rolling enrollment.²²
577 ^k NE County Area of Residence, extracted from member's electronic medical record, includes Dekalb, Gwinnett, Forsyth, Hall, Barrow, Jackson, Butts, Gilmer, Pike, Gordon, Jasper, Monroe
578 counties

579 **Figure 1. Heat Map of SARS-COV-2 Confirmed Positive Patients By Race/ethnicity Groups Across Metro Atlanta's KPGA Catchment Area**

580

581

582 Panel A: non-Hispanic Black/African American; Panel B: non-Hispanic White; Panel C: Other race group (Hispanic/Latino, Asian, American Indian,

583 unknown/declined to report); Panel D: Metro Atlanta Neighborhood Deprivation Index

584