Effects of high intensity interval exercise on cerebrovascular function: A systematic review

Alicen A Whitaker^{1¶}, Mohammed Alwatban^{1&}, Andrea Freemyer^{1&}, Jaime Perales-Puchalt^{2,3&} and Sandra A Billinger^{1,3,4,¶*}

¹ Department of Physical Therapy and Rehabilitation Science, University of Kansas Medical Center, Kansas City, KS, United States

² University of Kansas Alzheimer's Disease Center, Fairway, KS, United States

³ Department of Neurology, University of Kansas Medical Center, Kansas City, KS, United States

⁴ Department of Physical Medicine and Rehabilitation, University of Kansas Medical Center, Kansas City, KS, United States

*Corresponding Author:

Sandra A. Billinger

Email: sbillinger@kumc.edu

[¶]These authors contributed equally to this work.

[&]These authors also contributed equally to this work.

Short title: High intensity interval exercise and cerebrovascular function

Abstract

High intensity interval exercise (HIIE) improves aerobic fitness with decreased exercise time compared to moderate continuous exercise. A gap in knowledge exists regarding the effects of HIIE on cerebrovascular function such as cerebral blood velocity and autoregulation. The objective of this systematic review was to ascertain the effect of HIIE on cerebrovascular function in healthy individuals. We searched PubMed and the Cumulative Index to Nursing and Allied Health Literature databases with apriori key words. We followed the Preferred Reporting Items for Systematic Reviews. Twenty articles were screened and thirteen articles were excluded due to not meeting the apriori inclusion criteria. Seven articles were reviewed via the modified Sackett's quality evaluation. Outcomes included middle cerebral artery blood velocity (MCAv) (n=4), dynamic cerebral autoregulation (dCA) (n=2), cerebral de/oxygenated hemoglobin (n=2), cerebrovascular reactivity to carbon dioxide (CO₂) (n=2) and cerebrovascular conductance/resistance index (n=1). Quality review was moderate with 3/7 to 5/7 quality criteria met. HIIE acutely lowered exercise MCAv compared to moderate intensity. HIIE decreased dCA phase following acute and chronic exercise compared to rest. HIIE acutely increased de/oxygenated hemoglobin compared to rest. HIIE acutely decreased cerebrovascular reactivity to higher CO₂ compared to rest and moderate intensity. The acute and chronic effects of HIIE on cerebrovascular function vary depending on the outcomes measured. Therefore, future research is needed to confirm the effects of HIIE on cerebrovascular function in healthy individuals and better understand the effects in individuals with chronic conditions. In order to conduct rigorous systematic reviews in the future, we recommend assessing MCAv, dCA and CO₂ reactivity during and post HIIE.

Introduction

High intensity interval exercise (HIIE) has emerged at the forefront of exercise regimens due to the shorter activity time needed to benefit[1-3]. HIIE confers similar or significant increased aerobic fitness compared to conventional moderate intensity continuous exercise[1, 4-7]. While aerobic fitness is a measure of increased cardiovascular health, the entire vascular system (including the cerebral vascular system) may be improved following increased aerobic fitness[8]. A review and meta-analysis of HIIE in healthy adults has shown significant increases in aerobic fitness [1, 5, 6, 9]. However, the effects of HIIE on cerebrovascular function have not been systematically reviewed.

Cerebrovascular function is the ability of the cerebral blood vessels to deliver oxygen and nutrients for neuronal metabolism and maintain cerebral blood flow through dynamic autoregulation (dCA). dCA is the ability of the brain to sustain a constant cerebral blood flow despite large fluctuations in peripheral blood pressure[10] [11]. During resting conditions cerebral blood flow responds to arterial blood pressure fluctuations, neuronal metabolism, cortical activation, arterial blood gases and cardiac output [12]. Cerebral blood flow can be measured at rest using magnetic resonance imaging or transcranial Doppler ultrasound (TCD). Middle cerebral artery blood velocity (MCAv) measured by TCD is the only technique to measure cerebral blood flow during exercise, with high temporal resolution [13]. MCAv is linearly related to cerebral blood flow with the caveat that the MCA diameter remains unchanged[14].

A normal cerebrovascular response to submaximal moderate continuous exercise results in increased MCAv [15-17], increased cerebral oxygenation [18, 19] and sustained dCA[20, 21]. MCAv has been shown to concomitantly increase as exercise intensity increases, up to moderate intensity [12, 15, 22-25]. MCAv is affected differently during high intensity exercise. During continuous high intensity exercise and hyperventilation, MCAv is decreased due to a reduction in arterial carbon dioxide (CO₂)[26, 27] causing downstream arteriole constriction[12, 28]. Cerebrovascular reactivity is the ability of the small vessels in the brain to vasodilate and vasoconstrict in response to fluctuating CO₂ levels [29, 30]. The cerebrovascular response to HIIE may differ from continuous high intensity exercise due to the repetitive

short interval bouts that rapidly increase blood pressure which may cause cerebrovascular hyper-perfusion [31, 32]. If neuroprotective mechanisms of the brain such as dCA do not respond quickly to the repetitive and rapid increases in blood pressure, HIIE could elevate the risk for leakage within the blood brain barrier[31, 33].

Previous scientific statements and narrative reviews have recounted the molecular, hemodynamic and structural processes (i.e. CO₂, nitric oxide, systemic blood pressure, vessel compliance, glial cell integrity) associated with the cerebrovascular response that may occur during HIIE [33, 34]. However, these detailed narrative reviews[35, 36] did not report the statistical findings of previous studies showing cerebrovascular function during HIIE. To our knowledge, our current systematic review is the first to systematically search and report the results of the dynamic cerebrovascular response during HIIE. Reporting the cerebrovascular response during HIIE is important because it provides objective results to support the previously described narrative statements on hemodynamic processes during HIIE [35, 36]. The purpose of this systematic review was to address the gap in knowledge and report the various study results of HIIE on cerebrovascular function compared to moderate continuous exercise or rest conditions. We systematically examined the results of HIIE studies in healthy individuals based on the operationalization of cerebrovascular outcomes.

Methods

This review follows the guidelines for Preferred Reporting Items for Systematic Reviews [37]. Literature searches and reviews were performed using PubMed and the Cumulative Index to Nursing and Allied Health Literature (CINAHL) databases. The University of Kansas Medical Center Online Library system was used to access these databases in February, March, and June 2020. In this systematic review, we included peer-reviewed manuscripts written in English from January 2010 to June 2020.

Key words used to search the databases included “high intensity interval training”, “HIIT”, “high intensity interval exercise” “HIIE” AND “cerebral blood flow”, “cerebral blood velocity”, “dynamic autoregulation”.

We believe these key words primarily reflect the high intensity interval intervention and cerebrovascular function outcome measures. The main outcomes of this systematic review were MCAv and dCA. However, additional cerebrovascular measures were also included such as oxygenated hemoglobin, cerebrovascular reactivity, cerebrovascular conductance index and cerebrovascular resistance index. Cerebrovascular conductance index is a measure of the conductance of peripheral blood pressure to cerebral blood velocity and is calculated as MCAv/mean arterial pressure (MAP). Cerebrovascular resistance index (MAP/MCAv) measures the resistance of cerebral perfusion pressure to cerebral blood velocity.

The identified abstracts from PubMed and CINAHL were screened using the following inclusion criteria: 1) experimental or quasi-experimental, 2) aerobic exercise identified as the primary means of performing HIIE, 3) cerebrovascular measures were primary or secondary outcomes and 4) human subjects across the lifespan with no current disease. After the removal of duplicates, two researchers screened titles/abstracts for inclusion criteria (A.W. and M.A.). The full texts were examined, and data extracted (A.W. and M.A.). If the authors were unable to come to an agreement, a third author moderated incongruity (A.F.).

A quality review was performed for each article using the modified version of Sackett's 1981 criteria[38]. We critically analyzed each article's study design, population, HIIE protocols, cerebrovascular outcomes and results. If an article did not report enough information to determine sufficient quality criteria a "No" rating was given. Articles were rated based on the level of evidence including level I for large randomized control trials, level II for small randomized trial, level III for nonrandomized design, Level IV for case series and Level V for case reports [39].

Results

The search methods resulted in 67 articles. After removal of duplicates, 15[40-53] articles were identified in PubMed and 5[54-58] new articles in CINAHL. During the initial screening of titles/abstracts, 11 articles

were excluded due to HIIE not being the primary experimental protocol performed (n=6), studies not measuring cerebral arteries (n=4) and an animal study (n=1). Studies that combined other interventions with HIIE were excluded due to the confounding variables that could affect cerebrovascular outcomes. After the full text assessment, two articles were excluded due to not meeting experimental or quasi experimental criteria (n=2). See Fig 1 for flow diagram of article selection. [Insert Figure 1]

Fig 1. Flow Diagram of Article Selection.

We included seven articles describing cerebrovascular outcomes following HIIE within this review[41, 44, 47, 52, 56, 57, 59]. The full texts are described in Table 1. Of the articles reviewed six were small, randomized trials and one nonrandomized cross-over trial. All the studies involved healthy individuals although some only included men (n=1), women (n=1), or children (n= 1). Prior activity levels of participants ranged from inactive, recreationally active and endurance trained.

Table 1. Summary of reviewed articles.

Study	Design	Level of Evidence	Subjects	Intervention	Outcome measures	Results
Burma et al, March 2020	Small Randomized Cross-Over Trial	II	9 Young Adults	3 cycling conditions: - High intensity interval training (HIIT, 1-min interval at 85-90% predicted heart rate reserve with 1-min active recovery 15% power output for 10 intervals) - Moderate intensity continuous exercise (MICT, 50-60% predicted heart rate reserve for 45 min) - No-exercise control	Average Exercise MCAv dCA via Transfer Function Analysis during forced MAP oscillations (repeated squat-stand maneuver)	Significant increase in exercise MCAv during MICT compared to HIIT (p<0.05) and baseline (p<0.05). No change in exercise MCAv during HIIT compared to baseline (p>0.05). Significantly higher systolic gain/phase compared to diastolic/mean gain/phase at 0.05 and 0.10 Hz during control (p<0.05). Decreased systolic phase in 0.05 Hz immediately following HIIT (p>0.102) and MICT until hour 4

						(p>0.079). Decreased systolic phase in 0.10 Hz immediately following HIIT until hour 2 (p>.11) and immediately following MICT until hour 4 (p>0.079). No change in gain or coherence in 0.05 Hz or 0.10 Hz during HIIT or MICT.
Burma et al, June 2020	Secondary Analysis of the above Small Randomized Cross-Over Trial	II	Same as above	Same as above	Cerebrovascular reactivity to hypercapnia Cerebrovascular reactivity to hypocapnia	Significantly decreased absolute and relative MCA reactivity to hypercapnia immediately following HIIT up to hour 2 (p<0.018) compared to MICT and control (p<0.022). Significantly decreased relative MCA reactivity to hypercapnia immediately following MICT up to hour 1 (p<0.024). No significant differences in MCA reactivity to hypocapnia between conditions (p>0.31).
Coetsee et al, 2017	Small Randomized Controlled Trial	II	67 Inactive Adults	16-week intervention Treadmill 30 min, 3x/week 4 groups: - High intensity Interval training (HIIT, 4 min interval at 90-95% HRmax with 3 min active recovery 70% HRmax) - Moderate continuous training (MCT, 70-75% HRmax) - No-exercise control (CON).	Cerebrovascular measures taken during cortical activation (Stroop test): - Oxygenated Hemoglobin - Deoxygenated Hemoglobin - Total Hemoglobin Index	No significant differences in oxygenated (effect size=.45, p=.3), deoxygenated (effect size=0.67, p=.14), or total hemoglobin (effect size<.6, p>.18) after HIIT. Significant decrease in deoxygenated hemoglobin (effect size=1.14, p=.01) and total hemoglobin index (effect size=1.49, p<.01) after MCT. Significant increase in oxygenated hemoglobin

						in CON (effect size=.76, p=.03).
Drapeau et al, 2019	Small Randomized Clinical Trial	II	17 Endurance Trained Males	6-week intervention. Cycled until exhaustion, 3x/week 2 groups: - HIIT ₈₅ (1-7 min interval at 85% of maximal aerobic power, with active recovery of 50% of maximal aerobic power) - HIIT ₁₁₅ (30sec – 1min interval at 115% of maximal aerobic power, with active recovery of 50% of maximal aerobic power).	Resting MCAv Resting CVCi Resting CVRi dCA via Transfer Function Analysis during forced MAP oscillations (repeated squat-stand maneuver)	Significant decrease in phase at 0.10 Hz in HIIT ₈₅ and HIIT ₁₁₅ (p = .048) with no differences between intensity groups. No significant difference in power spectral density (p > .39), gain (p > .05), or coherence (p>.05) between time or intensity. No significant differences in MCAv (p = .4), CVCi (p = .87), or CVRi (p = .87).
Northey et al, 2019	Small Randomized Controlled Trial	II	17 Female Breast Cancer Survivors	12-week intervention Cycled 20-30 min 3x/week 3 groups: - High intensity interval training (HIIT, 30 sec intervals at ~90% maximal heart rate or ~105% peak power with 2 min active recovery) - Moderate intensity continuous exercise (MOD, 55-65% peak power) - No-exercise control (CON)	Resting MCAv, Cerebrovascular Reactivity to CO ₂	No significant differences in resting MCAv (p=.24) or cerebrovascular reactivity (p=.54) after HIIT compared to MOD. No significant differences in resting MCAv (p=.86) or cerebrovascular reactivity (p=.72) after HIIT compared to CON.
Tallon et al, 2019	Small Randomized Cross-over Trial	II	8 Prepubertal Children	2 Cycling conditions: - High intensity interval exercise (HIIE, 1 min interval at 90%max watt with 1 min	Exercise MCAv during each interval Immediate post-exercise MCAv 30-minutes post-	Significant decrease in exercise MCAv during the 6 th interval of HIIE compared to baseline (10.7%, p=.004). Significant decrease in exercise MCAv during

				active recovery at 20%max watt for 6 intervals) - Moderate-intensity steady-state exercise (MISS, 15 min at 44%max watt)	exercise MCAv	the 3 rd and 4 th intervals of HIIE compared to MISS (p=.001). Significant decrease in MCAv immediately post-exercise following HIIE and MISS (p<.001). No significant difference in MCAv at 30-minutes post-exercise following HIIE and MISS compared to baseline (p>.05). Significant increase in exercise MCAv during the 2 nd minute of MISS compared to baseline (5.8%, p=.004).
Monroe et al, 2015	Nonrandomized Cross-Over Trial	III	15 Recreationally Active Adults	2 cycling conditions: - Sprint Interval Cycling (SIC, 30 second all-out sprint interval with 4 min active recovery for 4 intervals) - Constant Resistance Cycling (CRC, 18 min at 70rpm with resistance set by matching total work performed during SIC)	Oxygenated Hemoglobin (HbO ₂) Deoxygenated Hemoglobin (HHb)	Significant increase in average HbO ₂ (effect size=.536, p=.001), minimum HbO ₂ during recovery (effect size=.392, p<.001) and maximum HbO ₂ during recovery (effect size=.588, p=.001) in SIC compared to CRC. Significant increase in average HHb during SIC compared to CRC (effect size=.386, p=.003).

MCAv = middle cerebral artery blood velocity, dCA = dynamic cerebral autoregulation, min = minute,

HRmax = maximum heart rate, CVCi = cerebrovascular conductance index, CVRi = cerebrovascular

resistance index, CO₂ = carbon dioxide

High intensity interval protocols

Methods of prescribing HIIE varied greatly between studies. HIIE protocols included 6- to 16-week

exercise interventions (n = 3) or one single bout of exercise(n=4). By examining 6- to 16-weeks of HIIE,

the long-term or chronic effects of this intervention were studied. By examining a single bout of HIIE, the

immediate or acute effects of the exercise were reported. In addition to the duration variability, we found that the mode of HIIE also differed across the included studies. One study used a treadmill as the mode of exercise with 4-minute intervals of 90-95% maximal heart rate for 30 minutes. The remaining six studies used cycling as the mode of exercise but differed in parameters ranging from 30 seconds to 7-minute intervals at 85% to 115% of maximal watts or ~ 85% to 90% maximal heart rate. A constant between all studies included an active (rather than passive) recovery interval between sprints. However, the intensity and duration of recovery intervals differed greatly.

Cerebrovascular outcome measures

The results of this review can be operationalized based on the outcome variables measured during HIIE such as MCAv (n=4), dCA (n=2), cerebral de/oxygenated hemoglobin (n=2), cerebrovascular reactivity to CO₂ (n=2) and cerebrovascular conductance/resistance index (n=1). Table 2 describes whether HIIE increased, decreased or had no influence on the operationalized cerebrovascular measures. A meta-analysis was not performed due to low number of studies (≤ 2) reporting each operationalized cerebrovascular measure.

Table 2. Summary of the effects of HIIE on operationalized cerebrovascular measures.

	Resting MCAv	Exercise MCAv	Post- Exercise MCAv	dCA phase	dCA Gain	dCA Coher ence	De/Oxy- genated Hemoglobin	Cerebro- vascular Conductance Resistance Index	Cerebro- vascular Reactivity to CO ₂
Burma et al, March 2020		↓ To moderate		↓	⊘	⊘			
Burma et al, June 2020									↓ To moderate and control
Coetsee et al, 2017							⊘ During cortical activation		
Drapeau et al, 2019	⊘			↓	⊘	⊘		⊘	
Northey	⊘								⊘

et al, 2019	
Tallon et al, 2019	<div> <div>↓ To moderate and rest</div> <div>↓ To rest</div> </div>
Monroe et al, 2015	<div> <div>↑ During HIIE</div> </div>

↓ = Decreased effect, ↑ = Increased effect, ⊘ = no effect

MCAv

Of the studies reporting MCAv outcomes, resting MCAv (n = 2), exercise MCAv (n = 2) and MCAv immediately post exercise (n = 1) were used. No significant differences were found for resting MCAv after 6- or 12-weeks of HIIE when compared to moderate continuous exercise or control[41, 44]. During HIIE, exercise MCAv was significantly decreased compared to moderate continuous exercise[56, 59]. Conflicting results were found between two studies comparing exercise MCAv to rest. Burma et al.[59] reported no significant difference between average exercise MCAv and rest in adults. However, rather than reporting average exercise MCAv of the entire HIIE bout, Tallon et al.[56] reported exercise MCAv for each 1-minute sprint interval of HIIE. During the 6th sprint interval of HIIE, Tallon et al.[56] reported significantly decreased exercise MCAv compared to rest which remained immediately following exercise[56].

dCA

Transfer function analysis of dCA was reported in the very low and low frequency bands (n = 2). Drapeau et al.[41] conducted a 6-week intervention of HIIE and reported a significant decrease in phase compared to rest with no significant change in coherence or gain. Burma et al.[59] conducted a single bout of HIIE and reported decreased MCAv systolic phase immediately following exercise that extended up to four hours later.

De/oxygenated hemoglobin

Oxygenated and deoxygenated hemoglobin were reported during a single bout of HIIE (n=1) and during a 16-week HIIE intervention (n=1). Monroe et al.[57] conducted a single bout of HIIE and reported an increase in oxygenated and deoxygenated hemoglobin during HIIE compared to moderate continuous exercise. Coetsee et al.[47] conducted a 16-week intervention of HIIE and reported no significant lasting changes in oxygenated or deoxygenated hemoglobin during cortical activation.

Cerebrovascular reactivity

Cerebrovascular reactivity to CO₂ were reported during a single bout of HIIE (n=1)[52] and during a 12-week HIIE intervention (n=1)[44]. After a single bout of HIIE, cerebrovascular reactivity to higher CO₂, or hypercapnia, was significantly decreased by 37% and remained an hour later. The reduced cerebrovascular reactivity to hypercapnia was also significantly different than moderate intensity and control. Cerebrovascular reactivity to lower CO₂, or hypocapnia, was not significantly different following a single bout of HIIE. Cerebrovascular reactivity to CO₂ was also not significantly different following 12 weeks of HIIE[44].

Cerebrovascular conductance and resistance

Cerebrovascular conductance index and cerebrovascular resistance index were only reported in a single study each. A 6-week HIIE intervention reported no significant changes in cerebrovascular conductance index or cerebrovascular resistance index[41].

Quality review

The quality review of each study is presented in Table 3. Out of seven total quality criteria, (n=2) reported five quality criteria, (n=1) reported four quality criteria and the remaining (n=4) reported three quality criteria. Therefore, the overall quality criteria results were moderately poor. All studies accounted for subjects and monitored the HIIE protocol parameters. No studies reported avoidance of contamination or co-intervention. No studies reported blinding of the outcome assessments. Only Burma et al.[52, 59] and Monroe et al.[57] reported their reliability via coefficient of reproducibility and intraclass coefficients of

their measures. And only Burma et al.[52, 59] and Drapeau et al.[41] reported validity of their respective cerebrovascular outcomes.

Table 3. Summary of quality review.

	Avoided Contamination and Co-Intervention	Random Assignment to Conditions	Blinded Assessment	Monitored Intervention	Accounted for All Subjects	Reported Reliability of Measures Used	Reported Validity of Measures Used	Total Number of Criteria Met
Burma et al, March 2020	No	Yes	No	Yes	Yes	Yes	Yes	5
Burma et al, June 2020	No	Yes	No	Yes	Yes	Yes	Yes	5
Coetsee et al, 2017	No	Yes	No	Yes	Yes	No	No	3
Drapeau et al, 2019	No	Yes	No	Yes	Yes	No	Yes	4
Northey et al, 2019	No	Yes	No	Yes	Yes	No	No	3
Tallon et al, 2019	No	Yes	No	Yes	Yes	No	No	3
Monroe et al, 2015	No	No	No	Yes	Yes	Yes	No	3

Discussion

This review met the objective of reporting the results of various HIIE studies and the effects on operationalized cerebrovascular function in healthy individuals. This review is the first to report the effects HIIE on cerebrovascular function compared to moderate continuous exercise and rest in healthy individuals. In general, we found that the acute and chronic effects of HIIE on cerebrovascular function vary largely depending on the methods and outcomes measured.

MCAv

In these studies, 6- to 12-week HIIE interventions had no effect on resting MCAv in healthy individuals. No significant change may be from too short of a HIIE intervention duration. Also a ceiling effect may be observed for young, healthy individuals and could explain no changes in resting MCAv[16]. During a single bout of HIIE, hyperventilation and downstream arteriole vasoconstriction may explain the acute decreases in exercise MCAv compared to moderate continuous exercise[12, 28, 60, 61]. Vasoconstriction may play a protective role during HIIE due to heightened peripheral blood pressure potentially causing hyper-perfusion or damage to the blood brain barrier[10, 33, 62]. During a single bout of HIIE, there is contradictory evidence comparing exercise MCAv to rest. One study reported no change in average exercise MCAv compared to resting. Another study reported decreased exercise MCAv after six sprint intervals of HIIE and remained decreased compared to rest immediately following HIIE[56]. The differences reported in exercise MCAv compared to rest could be due to age (adults versus prepubertal children) or due to the analysis of MCAv during HIIE (average over entire exercise versus separate sprint intervals). Decreases in exercise MCAv compared to rest may only occur in the late intervals of HIIE, during hyperventilation. Therefore, exercise MCAv should be reported for each interval of HIIE rather than an average of the entire exercise bout.

dCA

After a 6-week intervention and single bout of HIIE, dCA phase was decreased compared to rest. The chronic effects of HIIE on dCA phase may be due to elevated cardiorespiratory fitness in endurance trained individuals being associated with attenuated dCA [41, 63]. In healthy individuals, increased frequency within MCAv and MAP waveforms (that can occur with HIIE) may cause a reduction in phase due to dCA being a high-pass filter[64, 65]. Burma et al.[59] also suggests that systolic phase may reveal greater changes in dCA than both diastolic and mean phase. After a single bout of HIIE, reduction in systolic phase extended up to 4 hours and therefore the common approach of abstaining from exercise 12 hours before research studies[66-68] may be too conservative[59].

Although not included in this review due to the observational study design, contradictory evidence of sustained dCA during HIIE has been reported[45]. Differences in exercise parameters between HIIE may

be the cause to contradictory findings due to exhaustive exercise showing decreased dCA [31, 69]. More studies are needed to confirm the acute and chronic decreases in dCA following HIIE.

De/oxygenated hemoglobin

After a 16-week HIIE intervention, oxygenated and deoxygenated hemoglobin during cortical activation did not change. However, the 16-week HIIE intervention decreased reaction time during cortical activation and therefore may have increased efficiency of cortical oxygen use. Also, during a single bout of HIIE oxygenated and deoxygenated hemoglobin increased compared to moderate continuous exercise. The acute and chronic effects of HIIE on oxygenated and deoxygenated hemoglobin still needs further investigation due to each only being reported in a single study.

Cerebrovascular reactivity

A 12-week HIIE intervention did not significantly change cerebrovascular reactivity which could be due to vascular desensitization from chronic exposure to CO₂ during HIIE[33, 70]. Following a single bout of HIIE, cerebrovascular reactivity to hypercapnia was decreased showing the inability of the cerebrovascular system to maximally vasodilate. The maximal capacity for vasodilation after HIIE may be reduced following HIIE due to prolonged cerebrovascular vasoconstriction that occurs with hyperventilation during HIIE[33, 52]. Cerebrovascular reactivity to hypocapnia was not changed following a single bout of HIIE due to the ability of the vessels to vasoconstrict remaining intact[52]. The reduction in cerebrovascular reactivity to higher CO₂ remains an hour after HIIE. Therefore, the authors conclude again that the common approach of abstaining from exercise 12 hours before research studies[67, 71, 72] may be too conservative[52].

Cerebrovascular conductance and resistance

Cerebrovascular conductance and resistance were not significantly changed following a 6-week HIIE intervention. While a 6-week HIIE intervention significantly improved peripheral arterial conductance and resistance[4], this change in the peripheral arteries may not be demonstrated in the cerebrovascular

arterial conductance or resistance[35, 41]. However, due to the cerebrovascular conductance or resistance index being reported in only a single study, no conclusive effects of HIIE can be determined.

Article limitations

The individual study limitations have been previously acknowledged within the published articles or have been identified by authors (AW and MA) during the quality review. Most of the studies were small, randomized trials and included mostly young individuals. The small sample size available in males ($n = 17$), females ($n = 17$), or children ($n = 8$) limit the generalizability of results presented in this systematic review to a larger population. All studies implemented HIIE protocols that differed in time, repetitions and intensity levels, therefore making comparisons between studies difficult. These studies report potential limitations to the methods of measuring cerebrovascular function. TCD is the preeminent method of measuring MCAv during exercise but MCAv during HIIE may be underestimated[73]. However, some of the studies report the minor changes in expired CO_2 during HIIE were not likely to induce changes in MCA diameter[14, 56, 57]. The studies also report underestimation of cerebral oxygenation due to the two-channel near infrared spectrometer not measuring the motor, occipital, or parietal cortex [47, 57].

Review limitations

The authors acknowledge a risk of publication bias by only including peer-reviewed articles written in English and did not include grey literature. The cerebrovascular function measures included within this review vary greatly and have vast heterogeneity. While HIIE is not a new mode of exercise, studying cerebrovascular measures during HIIE is novel. Therefore, authors could only identify seven small studies with the oldest article dating back to 2015. The primary outcome of MCAv ($n=4$) and dCA ($n=2$) were reported in few studies with low power. Therefore, a meta-analysis could not be performed due to insufficient mathematical combination.

Conclusion

This review has provided preliminary information studying the effects of HIIE on cerebrovascular function. Although the amount of studies reporting cerebrovascular function during HIIE was small, we propose that a single bout of HIIE may acutely decrease MCAv, dCA and cerebrovascular reactivity to hypercapnia while increasing oxygenated hemoglobin, during or immediately following HIIE compared to moderate continuous exercise or rest conditions. No significant chronic effects were found for resting MCAv, cerebrovascular reactivity to CO₂, cerebrovascular conductance or resistance index after 6- or 12-week interventions of HIIE. Further research is warranted to confirm the acute and chronic effects of HIIE on MCAv, dCA, and cerebrovascular reactivity.

With increased interest in healthy brain aging and implementing interventions to maintain or improve brain health[74], studying the effects of HIIE interventions are critically needed[35, 36]. While this review only included healthy individuals, we provide an early reference to understanding “normal” physiological effects of HIIE on cerebrovascular function and the need to compare to clinical populations.

Future HIIE research is needed to evaluate cerebrovascular function not only during HIIE but also immediately following exercise. More studies such as randomized controlled trials with large sample sizes are needed to conduct a meta-analysis to combine and statistically analyze the summary results of HIIE on cerebrovascular function. Additionally, more studies are needed to determine the optimal interval parameters for HIIE and cerebrovascular function for humans.

References

1. Batacan RB, Jr., Duncan MJ, Dalbo VJ, Tucker PS, Fenning AS. Effects of high-intensity interval training on cardiometabolic health: a systematic review and meta-analysis of intervention studies. *Br J Sports Med.* 2017;51(6):494-503. Epub 2016/11/01. doi: 10.1136/bjsports-2015-095841. PubMed PMID: 27797726.
2. Costa EC, Hay JL, Kehler DS, Boreskie KF, Arora RC, Umpierre D, et al. Effects of High-Intensity Interval Training Versus Moderate-Intensity Continuous Training On Blood Pressure in Adults with Pre-to Established Hypertension: A Systematic Review and Meta-Analysis of Randomized Trials. *Sports Med.* 2018;48(9):2127-42. Epub 2018/06/28. doi: 10.1007/s40279-018-0944-y. PubMed PMID: 29949110.
3. Ramos JS, Dalleck LC, Tjonna AE, Beetham KS, Coombes JS. The impact of high-intensity interval training versus moderate-intensity continuous training on vascular function: a systematic review and meta-analysis. *Sports Med.* 2015;45(5):679-92. Epub 2015/03/17. doi: 10.1007/s40279-015-0321-z. PubMed PMID: 25771785.
4. Rakobowchuk M, Tanguay S, Burgomaster KA, Howarth KR, Gibala MJ, MacDonald MJ. Sprint interval and traditional endurance training induce similar improvements in peripheral arterial stiffness and flow-mediated dilation in healthy humans. *Am J Physiol Regul Integr Comp Physiol.* 2008;295(1):R236-42. Epub 2008/04/25. doi: 10.1152/ajpregu.00069.2008. PubMed PMID: 18434437; PubMed Central PMCID: PMC2494806.
5. Tabata I, Nishimura K, Kouzaki M, Hirai Y, Ogita F, Miyachi M, et al. Effects of moderate-intensity endurance and high-intensity intermittent training on anaerobic capacity and VO₂max. *Med Sci Sports Exerc.* 1996;28(10):1327-30. Epub 1996/10/01. doi: 10.1097/00005768-199610000-00018. PubMed PMID: 8897392.
6. Cunningham DA, McCrimmon D, Vlach LF. Cardiovascular response to interval and continuous training in women. *Eur J Appl Physiol Occup Physiol.* 1979;41(3):187-97. Epub 1979/07/02. doi: 10.1007/bf00430011. PubMed PMID: 488084.
7. Edge J, Bishop D, Goodman C. The effects of training intensity on muscle buffer capacity in females. *Eur J Appl Physiol.* 2006;96(1):97-105. Epub 2005/11/12. doi: 10.1007/s00421-005-0068-6. PubMed PMID: 16283370.
8. Ainslie PN, Cotter JD, George KP, Lucas S, Murrell C, Shave R, et al. Elevation in cerebral blood flow velocity with aerobic fitness throughout healthy human ageing. *J Physiol.* 2008;586(16):4005-10. Epub 2008/07/19. doi: 10.1113/jphysiol.2008.158279. PubMed PMID: 18635643; PubMed Central PMCID: PMC2538930.
9. Bayati M, Farzad B, Gharakhanlou R, Agha-Alinejad H. A practical model of low-volume high-intensity interval training induces performance and metabolic adaptations that resemble 'all-out' sprint interval training. *J Sports Sci Med.* 2011;10(3):571-6. Epub 2011/01/01. PubMed PMID: 24150635; PubMed Central PMCID: PMC3737829.
10. Aaslid R, Lindegaard KF, Sorteberg W, Nornes H. Cerebral autoregulation dynamics in humans. *Stroke.* 1989;20(1):45-52. Epub 1989/01/01. doi: 10.1161/01.str.20.1.45. PubMed PMID: 2492126.
11. Li Z, Zhang M, Xin Q, Luo S, Zhou W, Cui R, et al. Assessment of cerebral oxygenation oscillations in subjects with hypertension. *Microvasc Res.* 2013;88:32-41. Epub 2013/04/16. doi: 10.1016/j.mvr.2013.04.003. PubMed PMID: 23583904.
12. Smith KJ, Ainslie PN. Regulation of cerebral blood flow and metabolism during exercise. *Exp Physiol.* 2017;102(11):1356-71. Epub 2017/08/09. doi: 10.1113/EP086249. PubMed PMID: 28786150.
13. Brassard P, Ainslie PN, Secher NH. Cerebral oxygenation in health and disease. *Front Physiol.* 2014;5:458. Epub 2014/12/17. doi: 10.3389/fphys.2014.00458. PubMed PMID: 25505422; PubMed Central PMCID: PMC4241837.

14. Serrador JM, Picot PA, Rutt BK, Shoemaker JK, Bondar RL. MRI measures of middle cerebral artery diameter in conscious humans during simulated orthostasis. *Stroke*. 2000;31(7):1672-8. Epub 2000/07/08. doi: 10.1161/01.str.31.7.1672. PubMed PMID: 10884472.
15. Witte E, Liu Y, Ward JL, Kempf KS, Whitaker A, Vidoni ED, et al. Exercise intensity and middle cerebral artery dynamics in humans. *Respir Physiol Neurobiol*. 2019;262:32-9. Epub 2019/02/03. doi: 10.1016/j.resp.2019.01.013. PubMed PMID: 30710650; PubMed Central PMCID: PMC6393201.
16. Ward JL, Craig JC, Liu Y, Vidoni ED, Maletsky R, Poole DC, et al. Effect of healthy aging and sex on middle cerebral artery blood velocity dynamics during moderate-intensity exercise. *Am J Physiol Heart Circ Physiol*. 2018;315(3):H492-H501. Epub 2018/05/19. doi: 10.1152/ajpheart.00129.2018. PubMed PMID: 29775407; PubMed Central PMCID: PMC6172645.
17. Brugniaux JV, Marley CJ, Hodson DA, New KJ, Bailey DM. Acute exercise stress reveals cerebrovascular benefits associated with moderate gains in cardiorespiratory fitness. *J Cereb Blood Flow Metab*. 2014;34(12):1873-6. Epub 2014/10/02. doi: 10.1038/jcbfm.2014.142. PubMed PMID: 25269518; PubMed Central PMCID: PMC4269737.
18. Davenport MH, Hogan DB, Eskes GA, Longman RS, Poulin MJ. Cerebrovascular reserve: the link between fitness and cognitive function? *Exerc Sport Sci Rev*. 2012;40(3):153-8. Epub 2012/04/17. doi: 10.1097/JES.0b013e3182553430. PubMed PMID: 22504726.
19. Fisher JP, Hartwich D, Seifert T, Olesen ND, McNulty CL, Nielsen HB, et al. Cerebral perfusion, oxygenation and metabolism during exercise in young and elderly individuals. *J Physiol*. 2013;591(7):1859-70. Epub 2012/12/12. doi: 10.1113/jphysiol.2012.244905. PubMed PMID: 23230234; PubMed Central PMCID: PMC3624856.
20. Brys M, Brown CM, Marthol H, Franta R, Hilz MJ. Dynamic cerebral autoregulation remains stable during physical challenge in healthy persons. *Am J Physiol Heart Circ Physiol*. 2003;285(3):H1048-54. Epub 2003/08/14. doi: 10.1152/ajpheart.00062.2003. PubMed PMID: 12915389.
21. Ogoh S, Fadel PJ, Zhang R, Selmer C, Jans O, Secher NH, et al. Middle cerebral artery flow velocity and pulse pressure during dynamic exercise in humans. *Am J Physiol Heart Circ Physiol*. 2005;288(4):H1526-31. Epub 2004/12/14. doi: 10.1152/ajpheart.00979.2004. PubMed PMID: 15591094.
22. Steventon JJ, Hansen AB, Whittaker JR, Wildfong KW, Nowak-Fluck D, Tymko MM, et al. Cerebrovascular Function in the Large Arteries Is Maintained Following Moderate Intensity Exercise. *Front Physiol*. 2018;9:1657. Epub 2018/12/07. doi: 10.3389/fphys.2018.01657. PubMed PMID: 30519192; PubMed Central PMCID: PMC6258791.
23. Herholz K, Buskies W, Rist M, Pawlik G, Hollmann W, Heiss WD. Regional cerebral blood flow in man at rest and during exercise. *J Neurol*. 1987;234(1):9-13. Epub 1987/01/01. doi: 10.1007/bf00314001. PubMed PMID: 3102697.
24. Jorgensen LG, Perko M, Hanel B, Schroeder TV, Secher NH. Middle cerebral artery flow velocity and blood flow during exercise and muscle ischemia in humans. *J Appl Physiol* (1985). 1992;72(3):1123-32. Epub 1992/03/01. doi: 10.1152/jappl.1992.72.3.1123. PubMed PMID: 1568967.
25. Thomas SN, Schroeder T, Secher NH, Mitchell JH. Cerebral blood flow during submaximal and maximal dynamic exercise in humans. *J Appl Physiol* (1985). 1989;67(2):744-8. Epub 1989/08/01. doi: 10.1152/jappl.1989.67.2.744. PubMed PMID: 2507500.
26. Hellstrom G, Wahlgren NG. Physical exercise increases middle cerebral artery blood flow velocity. *Neurosurg Rev*. 1993;16(2):151-6. Epub 1993/01/01. doi: 10.1007/bf00258249. PubMed PMID: 8345909.
27. Moraine JJ, Lamotte M, Berre J, Niset G, Leduc A, Naeije R. Relationship of middle cerebral artery blood flow velocity to intensity during dynamic exercise in normal subjects. *Eur J Appl Physiol Occup Physiol*. 1993;67(1):35-8. Epub 1993/01/01. doi: 10.1007/bf00377701. PubMed PMID: 8375362.

28. Ogoh S, Ainslie PN. Regulatory mechanisms of cerebral blood flow during exercise: new concepts. *Exerc Sport Sci Rev.* 2009;37(3):123-9. Epub 2009/06/25. doi: 10.1097/JES.0b013e3181aa64d7. PubMed PMID: 19550203.
29. Maeda H, Matsumoto M, Handa N, Hougaku H, Ogawa S, Itoh T, et al. Reactivity of cerebral blood flow to carbon dioxide in various types of ischemic cerebrovascular disease: evaluation by the transcranial Doppler method. *Stroke.* 1993;24(5):670-5. Epub 1993/05/01. doi: 10.1161/01.str.24.5.670. PubMed PMID: 8488521.
30. Willie CK, Macleod DB, Shaw AD, Smith KJ, Tzeng YC, Eves ND, et al. Regional brain blood flow in man during acute changes in arterial blood gases. *J Physiol.* 2012;590(14):3261-75. Epub 2012/04/13. doi: 10.1113/jphysiol.2012.228551. PubMed PMID: 22495584; PubMed Central PMCID: PMC3459041.
31. Bailey DM, Evans KA, McEneny J, Young IS, Hullin DA, James PE, et al. Exercise-induced oxidative-nitrosative stress is associated with impaired dynamic cerebral autoregulation and blood-brain barrier leakage. *Exp Physiol.* 2011;96(11):1196-207. Epub 2011/08/16. doi: 10.1113/expphysiol.2011.060178. PubMed PMID: 21841038.
32. Phillips AA, Matin N, Jia M, Squair JW, Monga A, Zheng MMZ, et al. Transient Hypertension after Spinal Cord Injury Leads to Cerebrovascular Endothelial Dysfunction and Fibrosis. *J Neurotrauma.* 2018;35(3):573-81. Epub 2017/11/17. doi: 10.1089/neu.2017.5188. PubMed PMID: 29141501; PubMed Central PMCID: PMC6421994.
33. Lucas SJ, Cotter JD, Brassard P, Bailey DM. High-intensity interval exercise and cerebrovascular health: curiosity, cause, and consequence. *J Cereb Blood Flow Metab.* 2015;35(6):902-11. Epub 2015/04/03. doi: 10.1038/jcbfm.2015.49. PubMed PMID: 25833341; PubMed Central PMCID: PMC4640257.
34. Franklin BA, Thompson PD, Al-Zaiti SS, Albert CM, Hivert MF, Levine BD, et al. Exercise-Related Acute Cardiovascular Events and Potential Deleterious Adaptations Following Long-Term Exercise Training: Placing the Risks Into Perspective-An Update: A Scientific Statement From the American Heart Association. *Circulation.* 2020;141(13):e705-e36. Epub 2020/02/27. doi: 10.1161/CIR.0000000000000749. PubMed PMID: 32100573.
35. Lucas SJE, Cotter JD, Brassard P, Bailey DM. High-intensity interval exercise and cerebrovascular health: curiosity, cause, and consequence. *J Cerebr Blood F Met.* 2015;35(6):902-11. doi: 10.1038/jcbfm.2015.49. PubMed PMID: WOS:000355575600003.
36. Calverley TA, Ogoh S, Marley CJ, Steggall M, Marchi N, Brassard P, et al. HIITing the brain with exercise: mechanisms, consequences and practical recommendations. *J Physiol.* 2020;598(13):2513-30. Epub 2020/04/30. doi: 10.1113/JP275021. PubMed PMID: 32347544.
37. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ.* 2009;339:b2535. Epub 2009/07/23. doi: 10.1136/bmj.b2535. PubMed PMID: 19622551; PubMed Central PMCID: PMC2714657.
38. Sackett DL. How to read clinical journals. *Can Med Assoc J.* 1982;126(12):1373. Epub 1982/06/15. PubMed PMID: 20313761; PubMed Central PMCID: PMC1863136.
39. Sackett DL. Rules of evidence and clinical recommendations for the management of patients. *Can J Cardiol.* 1993;9(6):487-9. Epub 1993/07/01. PubMed PMID: 8221340.
40. Dupuy O, Tremblay J. Impact of Carbohydrate Ingestion on Cognitive Flexibility and Cerebral Oxygenation during High-Intensity Intermittent Exercise: A Comparison between Maple Products and Usual Carbohydrate Solutions. *Nutrients.* 2019;11(9). Epub 2019/08/31. doi: 10.3390/nu11092019. PubMed PMID: 31466305; PubMed Central PMCID: PMC6770040.
41. Drapeau A, Labrecque L, Imhoff S, Paquette M, Le Blanc O, Malenfant S, et al. Six weeks of high-intensity interval training to exhaustion attenuates dynamic cerebral autoregulation without influencing

resting cerebral blood velocity in young fit men. *Physiol Rep.* 2019;7(15):e14185. Epub 2019/08/03. doi: 10.14814/phy2.14185. PubMed PMID: 31373166; PubMed Central PMCID: PMC6675921.

42. Willis SJ, Borrani F, Millet GP. Leg- vs arm-cycling repeated sprints with blood flow restriction and systemic hypoxia. *Eur J Appl Physiol.* 2019;119(8):1819-28. Epub 2019/06/13. doi: 10.1007/s00421-019-04171-0. PubMed PMID: 31187281.

43. Klein T, Bailey TG, Abeln V, Schneider S, Askew CD. Cerebral Blood Flow during Interval and Continuous Exercise in Young and Old Men. *Med Sci Sports Exerc.* 2019;51(7):1523-31. Epub 2019/02/16. doi: 10.1249/MSS.0000000000001924. PubMed PMID: 30768552.

44. Northey JM, Pampa KL, Quinlan C, Ikin A, Toohey K, Smee DJ, et al. Cognition in breast cancer survivors: A pilot study of interval and continuous exercise. *J Sci Med Sport.* 2019;22(5):580-5. Epub 2018/12/18. doi: 10.1016/j.jsams.2018.11.026. PubMed PMID: 30554923.

45. Tsukamoto H, Hashimoto T, Olesen ND, Petersen LG, Sorensen H, Nielsen HB, et al. Dynamic Cerebral Autoregulation Is Maintained during High-Intensity Interval Exercise. *Med Sci Sports Exerc.* 2019;51(2):372-8. Epub 2018/09/27. doi: 10.1249/MSS.0000000000001792. PubMed PMID: 30252775.

46. Hanssen H, Minghetti A, Magon S, Rossmessl A, Rasenack M, Papadopoulou A, et al. Effects of different endurance exercise modalities on migraine days and cerebrovascular health in episodic migraineurs: A randomized controlled trial. *Scand J Med Sci Sports.* 2018;28(3):1103-12. Epub 2017/11/22. doi: 10.1111/sms.13023. PubMed PMID: 29161767.

47. Coetsee C, Terblanche E. Cerebral oxygenation during cortical activation: the differential influence of three exercise training modalities. A randomized controlled trial. *Eur J Appl Physiol.* 2017;117(8):1617-27. Epub 2017/06/02. doi: 10.1007/s00421-017-3651-8. PubMed PMID: 28567669.

48. Kirkham AA, Shave RE, Bland KA, Bovard JM, Eves ND, Gelmon KA, et al. Protective effects of acute exercise prior to doxorubicin on cardiac function of breast cancer patients: A proof-of-concept RCT. *Int J Cardiol.* 2017;245:263-70. Epub 2017/07/25. doi: 10.1016/j.ijcard.2017.07.037. PubMed PMID: 28735755.

49. Gjellesvik TI, Becker F, Tjonna AE, Indredavik B, Nilsen H, Brurok B, et al. Effects of High-Intensity Interval Training after Stroke (The HIIT-Stroke study) - A Multicenter Randomized Controlled Trial. *Arch Phys Med Rehabil.* 2020. Epub 2020/03/08. doi: 10.1016/j.apmr.2020.02.006. PubMed PMID: 32145280.

50. Scott JM, Tucker WJ, Martin D, Crowell JB, Goetchius E, Ozgur O, et al. Association of Exercise and Swimming Goggles With Modulation of Cerebro-ocular Hemodynamics and Pressures in a Model of Spaceflight-Associated Neuro-ocular Syndrome. *JAMA Ophthalmol.* 2019;137(6):652-9. Epub 2019/04/19. doi: 10.1001/jamaophthalmol.2019.0459. PubMed PMID: 30998818; PubMed Central PMCID: PMC6567831.

51. Saoi M, Percival M, Nemr C, Li A, Gibala M, Britz-McKibbin P. Characterization of the Human Skeletal Muscle Metabolome for Elucidating the Mechanisms of Bicarbonate Ingestion on Strenuous Interval Exercise. *Anal Chem.* 2019;91(7):4709-18. Epub 2019/03/06. doi: 10.1021/acs.analchem.9b00149. PubMed PMID: 30835436.

52. Burma JS, Macaulay A, Copeland P, Khatra O, Bouliane KJ, Smirl JD. Comparison of cerebrovascular reactivity recovery following high-intensity interval training and moderate-intensity continuous training. *Physiol Rep.* 2020;8(11):e14467. Epub 2020/06/09. doi: 10.14814/phy2.14467. PubMed PMID: 32506845; PubMed Central PMCID: PMC67276190.

53. Labrecque L, Drapeau A, Rahimaly K, Imhoff S, Billaut F, Brassard P. Comparable blood velocity changes in middle and posterior cerebral arteries during and following acute high-intensity exercise in young fit women. *Physiol Rep.* 2020;8(9):e14430. Epub 2020/04/29. doi: 10.14814/phy2.14430. PubMed PMID: 32342622; PubMed Central PMCID: PMC67186567.

54. Peyravi A, Yazdanpanahi N, Nayeri H, Hosseini SA. The effect of endurance training with crocin consumption on the levels of MFN2 and DRP1 gene expression and glucose and insulin indices in the

muscle tissue of diabetic rats. *J Food Biochem.* 2020;44(2):e13125. Epub 2019/12/19. doi: 10.1111/jfbc.13125. PubMed PMID: 31849103.

55. Komiyama T, Tanoue Y, Sudo M, Costello JT, Uehara Y, Higaki Y, et al. Cognitive Impairment during High-Intensity Exercise: Influence of Cerebral Blood Flow. *Med Sci Sports Exerc.* 2020;52(3):561-8. Epub 2019/10/15. doi: 10.1249/MSS.0000000000002183. PubMed PMID: 31609297.

56. Tallon CM, Simair RG, Koziol AV, Ainslie PN, McManus AM. Intracranial Vascular Responses to High-Intensity Interval Exercise and Moderate-Intensity Steady-State Exercise in Children. *Pediatr Exerc Sci.* 2019;31(3):290-5. Epub 2019/03/06. doi: 10.1123/pes.2018-0234. PubMed PMID: 30832540.

57. Monroe DC, Gist NH, Freese EC, O'Connor PJ, McCully KK, Dishman RK. Effects of Sprint Interval Cycling on Fatigue, Energy, and Cerebral Oxygenation. *Med Sci Sports Exerc.* 2016;48(4):615-24. Epub 2015/11/13. doi: 10.1249/MSS.0000000000000809. PubMed PMID: 26559448.

58. Hansen RK, Nielsen PS, Schelske MW, Secher NH, Volianitis S. CO₂ supplementation dissociates cerebral oxygenation and middle cerebral artery blood velocity during maximal cycling. *Scand J Med Sci Sports.* 2020;30(3):399-407. Epub 2019/10/28. doi: 10.1111/sms.13582. PubMed PMID: 31650627.

59. Burma JS, Copeland P, Macaulay A, Khatra O, Wright AD, Smirl JD. Dynamic cerebral autoregulation across the cardiac cycle during 8 hr of recovery from acute exercise. *Physiol Rep.* 2020;8(5):e14367. Epub 2020/03/13. doi: 10.14814/phy2.14367. PubMed PMID: 32163235; PubMed Central PMCID: PMC7066871.

60. Gonzalez-Alonso J, Dalsgaard MK, Osada T, Volianitis S, Dawson EA, Yoshiga CC, et al. Brain and central haemodynamics and oxygenation during maximal exercise in humans. *J Physiol.* 2004;557(Pt 1):331-42. Epub 2004/03/09. doi: 10.1113/jphysiol.2004.060574. PubMed PMID: 15004212; PubMed Central PMCID: PMC1665053.

61. Linkis P, Jorgensen LG, Olesen HL, Madsen PL, Lassen NA, Secher NH. Dynamic exercise enhances regional cerebral artery mean flow velocity. *J Appl Physiol* (1985). 1995;78(1):12-6. Epub 1995/01/01. doi: 10.1152/jappl.1995.78.1.12. PubMed PMID: 7713801.

62. Brassard P, Kim YS, van Lieshout J, Secher NH, Rosenmeier JB. Endotoxemia reduces cerebral perfusion but enhances dynamic cerebrovascular autoregulation at reduced arterial carbon dioxide tension. *Crit Care Med.* 2012;40(6):1873-8. Epub 2012/05/23. doi: 10.1097/CCM.0b013e3182474ca7. PubMed PMID: 22610190.

63. Labrecque L, Rahimaly K, Imhoff S, Paquette M, Le Blanc O, Malenfant S, et al. Diminished dynamic cerebral autoregulatory capacity with forced oscillations in mean arterial pressure with elevated cardiorespiratory fitness. *Physiol Rep.* 2017;5(21). Epub 2017/11/11. doi: 10.14814/phy2.13486. PubMed PMID: 29122957; PubMed Central PMCID: PMC5688778.

64. van Beek AH, Claassen JA, Rikkert MG, Jansen RW. Cerebral autoregulation: an overview of current concepts and methodology with special focus on the elderly. *J Cereb Blood Flow Metab.* 2008;28(6):1071-85. Epub 2008/03/20. doi: 10.1038/jcbfm.2008.13. PubMed PMID: 18349877.

65. Hamner JW, Cohen MA, Mukai S, Lipsitz LA, Taylor JA. Spectral indices of human cerebral blood flow control: responses to augmented blood pressure oscillations. *J Physiol.* 2004;559(Pt 3):965-73. Epub 2004/07/16. doi: 10.1113/jphysiol.2004.066969. PubMed PMID: 15254153; PubMed Central PMCID: PMC1665190.

66. Ainslie PN, Barach A, Murrell C, Hamlin M, Hellemans J, Ogoh S. Alterations in cerebral autoregulation and cerebral blood flow velocity during acute hypoxia: rest and exercise. *Am J Physiol Heart Circ Physiol.* 2007;292(2):H976-83. Epub 2006/10/03. doi: 10.1152/ajpheart.00639.2006. PubMed PMID: 17012355.

67. Ainslie PN, Hamlin M, Hellemans J, Rasmussen P, Ogoh S. Cerebral hypoperfusion during hypoxic exercise following two different hypoxic exposures: independence from changes in dynamic autoregulation and reactivity. *Am J Physiol Regul Integr Comp Physiol.* 2008;295(5):R1613-22. Epub 2008/09/05. doi: 10.1152/ajpregu.90420.2008. PubMed PMID: 18768767.

68. Smirl JD, Hoffman K, Tzeng YC, Hansen A, Ainslie PN. Relationship between blood pressure and cerebral blood flow during supine cycling: influence of aging. *J Appl Physiol* (1985). 2016;120(5):552-63. Epub 2015/11/21. doi: 10.1152/jappphysiol.00667.2015. PubMed PMID: 26586907; PubMed Central PMCID: PMC4773644.
69. Ogoh S, Dalsgaard MK, Yoshiga CC, Dawson EA, Keller DM, Raven PB, et al. Dynamic cerebral autoregulation during exhaustive exercise in humans. *Am J Physiol Heart Circ Physiol*. 2005;288(3):H1461-7. Epub 2004/10/23. doi: 10.1152/ajpheart.00948.2004. PubMed PMID: 15498819.
70. Thomas BP, Yezhuvath US, Tseng BY, Liu P, Levine BD, Zhang R, et al. Life-long aerobic exercise preserved baseline cerebral blood flow but reduced vascular reactivity to CO₂. *J Magn Reson Imaging*. 2013;38(5):1177-83. Epub 2013/03/26. doi: 10.1002/jmri.24090. PubMed PMID: 23526811; PubMed Central PMCID: PMC3695025.
71. Battisti-Charbonney A, Fisher J, Duffin J. The cerebrovascular response to carbon dioxide in humans. *J Physiol*. 2011;589(Pt 12):3039-48. Epub 2011/04/28. doi: 10.1113/jphysiol.2011.206052. PubMed PMID: 21521758; PubMed Central PMCID: PMC3139085.
72. Claassen JA, Zhang R, Fu Q, Witkowski S, Levine BD. Transcranial Doppler estimation of cerebral blood flow and cerebrovascular conductance during modified rebreathing. *J Appl Physiol* (1985). 2007;102(3):870-7. Epub 2006/11/18. doi: 10.1152/jappphysiol.00906.2006. PubMed PMID: 17110510.
73. Coverdale NS, Gati JS, Opalevych O, Perrotta A, Shoemaker JK. Cerebral blood flow velocity underestimates cerebral blood flow during modest hypercapnia and hypocapnia. *J Appl Physiol* (1985). 2014;117(10):1090-6. Epub 2014/07/12. doi: 10.1152/jappphysiol.00285.2014. PubMed PMID: 25012027.
74. Gorelick PB, Furie KL, Iadecola C, Smith EE, Waddy SP, Lloyd-Jones DM, et al. Defining Optimal Brain Health in Adults: A Presidential Advisory From the American Heart Association/American Stroke Association. *Stroke*. 2017;48(10):e284-e303. Epub 2017/09/09. doi: 10.1161/STR.000000000000148. PubMed PMID: 28883125; PubMed Central PMCID: PMC5654545.

Supporting Information

S1 Fig. Prisma flow diagram

S1 Table. Prisma checklist

Articles identified in database
searches of PubMed and
CINAHL
(n=67)

Articles screened by title/abstract
after duplicates were removed
(n=20)

Full text articles assessed for
eligibility
(n=9)

Articles included in review
(n=7)

Articles excluded (n=11)
- HIE not primary protocol (n=8)
- Not measuring cerebral arteries (n=4)
- Animal study (n=1)

Article excluded due to non-
experimental or quasi-experimental
study design
(n=2)