Immunohistochemical Detection of early and late Human Papillomaviruses (HPV) proteins in Retinoblastoma tumors

Nara Diniz¹, Thatiana C Melo², Rodrigo P Araldi³, Rodrigo F. Carvalho³, Willy Beçak³, Jonathan W de Medeiros¹, Giwellington S Albuquerque¹, Ana Pavla Gurgel^{4#}, Rocio Hassan^{5□}, Maria TC Muniz^{1*}, Rita de Cassia Stocco^{3□}

¹University of Pernambuco, Brazil;

²Federal University of São Paulo, Brazil;

³Butantan Institute, Brazil;

⁴Federal University of Paraíba, Brazil;

⁵National Câncer Institute (INCA), Brazil.

*Corresponding author: Maria Tereza Cartaxo Muniz, PhD. University of Pernambuco (UPE), Brazil. E-mail: tereza.cartaxo@upe.br.

#Alternative corresponding author: Ana Pavla Almeida Diniz Gurgel, PhD. Federal University of Paraíba (UFPB), Brazil. E-mail: anapavla@ccae.ufpb.br.

26 **Abstract**

27 In this study, we evaluated the presence of early and late Human
 28 Papillomavirus (HPV) proteins in retinoblastoma Brazilian patients. For this, 8
 29 formalin-fixed paraffin-embedded retinoblastoma tissue blocks were used. HPV
 30 DNA presence was determined by *in situ* hybridization (ISH).
 31 Immunohistochemistry were performed to verify HPV16/18 E6, E1[^]E4, and L1
 32 proteins. HPV was detected in all retinoblastoma tumors and viral DNA was
 33 labeled in tumor cells, retinal layers and optical nerve structures. In addition,
 34 E1[^]E4, E6 and L1 proteins were detected in all samples in the same areas
 35 where HPV DNA was detected. Our data showed the presence and expression
 36 of early and late HPV proteins in retinoblastoma tumors from Brazilian children.
 37 However, further studies should be performed to clarify the role of HPV infection
 38 in retinoblastoma tumor.

39

40 **Key words:** Human papillomavirus (HPV), retinoblastoma (RB), oncoproteins.

41

42 **Introduction**

43 Retinoblastoma is an intraocular tumor involving neuroectodermal cells of the
 44 retina, affecting 18,000-30,000 children per year (Antoneli et al., 2011; Chauhan
 45 et al., 2020; Orjuela et al., 2000; Palazzi et al., 2003; Ryoo et al., 2013;
 46 Sachdeva and O'Brien, 2012). Most retinoblastomas arise from inherited or
 47 acquired mutations in the *RB1* gene, leading to biallelic inactivation of this gene
 48 (Friend et al., 1986; Knudson, 1971). Tumors with intact *RB1* gene suggest
 49 other routes for their development. In this context, studies have demonstrated

50 that HPV may play a role in the RB development since the E7 oncoprotein can
51 inhibit action of pRb, destabilizing the cell cycle (Buitrago-Pérez et al., 2009;
52 Dyson et al., 1989; Moody and Laimins, 2010; Orjuela et al., 2000; Whyte et al.,
53 1988).

54 The prevalence of HPV in retinoblastoma has been shown with wide variations
55 in different populations (4.6% - 69.7%) with the use of molecular methods for
56 viruses detection (Antoneli et al., 2011; Bhuvaneswari et al., 2012; Espinoza et
57 al., 2005; Gillison et al., 2007; Javanmard et al., 2019; Mohan et al., 2009;
58 Montoya-Fuentes et al., 2003; Orjuela et al., 2000; Palazzi et al., 2003; Shetty
59 et al., 2012; Shukla et al., 2009). Additionally, techniques such as *in situ*
60 hybridization and immunohistochemistry have been performed and allowed
61 possible causal relationship between HPV and RB development (Antoneli et al.,
62 2011; Bhuvaneswari et al., 2012; Espinoza et al., 2005; Gillison et al., 2007;
63 Palazzi et al., 2003; Shetty et al., 2012).

64 Although this relationship is not always suggested (Palazzi et al., 2003; Ryoo et
65 al., 2013), child exposure to HPV during pregnancy and/or childbirth, indicates
66 vertical viral transmission. In this scenario, it is suggested that HPV could be a
67 factor in the development of oncogenic processes, supporting the idea that the
68 virus is related, in fact, with the etiopathogenesis of RB (Bhuvaneswari et al.,
69 2012; Lee et al., 2013; Skoczyński et al., 2014).

70 Evaluating the role of HPV in the development of retinoblastoma, we aimed to
71 assess not only the viral presence in the tumor, but also possible viral activity,
72 through detection of viral proteins in both retina and adjacent ocular tissues.

73 **Material and Methods**

74 Selected cases

75 The patients in this study were children diagnosed with retinoblastoma and
 76 treated at the "Center for Pediatric Oncohematology - University Hospital
 77 Oswaldo Cruz". Eight formalin-fixed paraffin-embedded were obtained by
 78 spontaneous demand. Parents or guardians consented to the release of the
 79 material for research in retinoblastoma although data on eventual evaluations of
 80 the patient relatives were not included in this report. This study was performed
 81 with approval of the ethics committee of the Oswaldo Cruz University Hospital
 82 (CAEE0015.0.000-11).

83 In situ hybridization

84 For the detection and typing of HPV in paraffin-embedded material, the *in situ*
 85 hybridization (ISH) technique was used. The detection was performed using the
 86 ZytoFast PLUS CISH Implementation Kit HRP-DAB (ZytoVision) with the
 87 ZytoFast HPV type 16/18/31/33/35 Probe (ZytoVision, T-1040-400), which
 88 identifies the 16, 18, 31, 33 and 35 types, according to the manufacturer's
 89 instructions.

90 Immunohistochemistry

91 The Immunohistochemistry technique was performed for the 8 paraffin-
 92 embedded cases, using EnVision TM + system-HRP (AEC) (Dako) with
 93 antibodies (Abcam, AbcamInc, Cambridge, London) against HPV 16 E1 ^ E4

94 proteins (TV G402), HPV 1, 6, 11, 16, 18 and 31 L1 protein (ab2417) and HPV
95 16 and 18 E6 proteins (ab70).

96

97 **Results**

98 RB tumor samples were analyzed for HPV presence by using *in situ*
99 hybridization. HPV DNA was detected in all samples (n=8). The labeled areas
100 included tumor cells, retinal layers and optical nerve structures (Figure 1 A, C
101 and E).

102 To analyze the possible HPV activity, viral proteins were analyzed by
103 immunohistochemistry. E1[^]E4, E6 and L1 proteins were visualized in
104 retinoblastoma samples in the same areas where HPV DNA was detected
105 (Figure 1B, D and F). These results showed early and late HPV genes
106 expression.

107

108 **Discussion**

109 The HPV relationship with the development of RB remains dubious, despite the
110 detection of high and low risk HPV sequences have been detected in tumor
111 samples (Antoneli et al., 2011; Shetty et al., 2012). In addition, HPV has been
112 detected in tumor segments, ocular surface squamous indicating that the virus
113 may be a risk factor for the development of other ocular tumors (Carreira et al.,
114 2013).

115 Although Gillison et al. (2007) (Gillison et al., 2007) have not reported the
116 presence of HPV in RB samples, suggesting that the virus has no role in the
117 development of these tumors, we have shown the presence of high-risk HPV.
118 High risk types have oncogenic potential by their oncoproteins ability to bind to
119 the tumor suppressor proteins such as p53 and pRb, destabilizing the cell cycle
120 and preventing apoptosis (McLaughlin-Drubin and Munger, 2009; Roman and
121 Munger, 2013).

122 Unlike Ryoo et al. (2013) (Ryoo et al., 2013), which did not detected no viral
123 DNA using *in situ* hybridization, the use of this technique allowed us to check
124 the exact location of DNA in biological samples. The labeled areas included
125 tumor cells, retinal layers and optical nerve structures. In addition,
126 immunohistochemistry was performed to verify the presence of viral proteins in
127 retinoblastoma samples. Thus, it was possible to verify the presence of DNA
128 and proteins not only in tumor cells but also in adjacent tissues to the tumor. It
129 was observed that the viral DNA is maintained in tumor tissues in advanced
130 stages, reaching in some cases the optic nerve, with expression of important
131 viral proteins, proteins that are related to the deregulation of the cell cycle (zur
132 Hausen, 2002).

133 We have identified viral DNA and proteins in nerve bundles of optic nerve.
134 Johnson et al. (2007) (Johnson et al., 2007) reported that retinoblastoma cells
135 are subjected to an intensive neuronal differentiation in early stages. Therefore,
136 according to these authors, the association between the presence of HPV and
137 the morphological differentiation of this tumor suggests a possible role of HPV
138 in the early stages of the RB development. Considering that the optic nerve

139 arises from embryonic cells of the retina, the results are relevant due to the
140 localization of virus and its proteins in verifying the viral location and activity in
141 tumor development. In according, the identification of viral DNA was confirmed
142 in fresh tumor.

143 In order to prove the presence and viral activity, we analyzed three different
144 HPV proteins, that demonstrated that HPV is not only present in the samples
145 analyzed as well as expressing its protein. The analysis of viral activity,
146 targeting E1[^]E4, E6 and L1 proteins in retinoblastoma samples, showed that
147 these proteins could be detected in the same areas where HPV DNA was
148 detected. The identification of HPV DNA and L1 protein in the same regions of
149 the eye tissue suggests this tissue as an organization site for productive
150 infection. In addition, proteins required for viral replication (E1 [^] E4) were
151 detected in the other regions, suggesting viral DNA replication. The presence of
152 the immunolabeled E6 protein also indicates the transforming action of HPV in
153 cells, since this oncoprotein can bind and degrade the p53 protein (McLaughlin-
154 Drubin and Munger, 2009). More studies are needed to identify other proteins
155 such as L2, E2 and E7, to verify the possible complete formation of the viral
156 particle, its integralization to the host genome and its transforming action on this
157 tumor stage and in these locations.

158 It is well described in the literature that the HPV complete its cycle in
159 differentiated, epithelial, cells (McLaughlin-Drubin and Munger, 2009) and
160 expression of structural genes is restricted to keratinocytes. Our results
161 suggest that the expression of viral proteins cannot depend solely on
162 differentiation of keratinocytes. We have identified the HPV proteins in tumor

163 cells, different layers of the retina, cornea epithelial tissue and nerve cells
164 constituting the optic nerve, as well as in inflammatory infiltrates.

165 Viral sources in these young individuals should be discussed. It is suggested
166 that HPV infects the child during childbirth (Bhuvaneswari et al., 2012), or
167 through vertical transmission, since HPV has been detected in the peripheral
168 blood of pregnant women, umbilical cord of newborns (Tseng et al., 1992) and
169 amniotic fluid (Armbruster-Moraes et al., 1994; Xu et al., 1998). Therefore, the
170 virus can induce retinoblastoma and other oncogenic processes (Tseng et al.,
171 1992). In addition, HPV has been detected in trophoblasts, through which
172 nutrient exchange occurs between the mother and the fetus. This infection has
173 been reported as the cause of some miscarriages (Hermonat et al., 1998).

174 Since the origin of HPV infection in children with retinoblastoma may be through
175 vertical transmission (Bhuvaneswari et al., 2012), reducing the incidence of this
176 virus in women may lead to a decrease in cancers linked to viral infection.
177 Efforts to reduce cases of HPV infection, lesions and/or cervical cancer are
178 extremely important for the reduction of cases of sporadic retinoblastomas
179 involving HPV. Like cervical cancer, the reduction of RB cases indirectly by the
180 monitoring of women, causes a decrease in the traumatic processes of the
181 children in their treatment, which often includes the surgical process of
182 enucleation, as well as chemotherapy and radiotherapy.

183

184 **Conclusions**

185 The results of this study showed the presence of HPV DNA in retinoblastoma
186 tumors. In addition, early and late HPV proteins were detected in all samples,
187 suggesting a viral activity in retinoblastoma tumors.

188

189 **Competing Interests**

190 The authors declare that they do not have anything to disclose regarding funding from
191 industries or conflict of interest with respect to this manuscript.

192

193 **Authors' Contributions**

194 ND and JWM carried out the molecular genetic studies, hybridization assays, and
195 drafted the manuscript. TCM and GSA participated in immunoassays. RPA participated
196 in hybridization assays. WB participated in the design of the study. MTCM, APG, RS
197 and participated in the design of the study, and helped to draft the manuscript. RCS
198 and conceived of the study, and participated in its design and coordination, RFC join
199 RCS in revision. All authors read and approved the final manuscript.

200

201 **Acknowledgments**

202 The authors acknowledge the CNPq (Conselho Nacional de Desenvolvimento Científico
203 e Tecnológico), and CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível
204 Superior) that supported this study.

205

206 **References**

207 Antoneli, C.B.G., Ribeiro, K.B., Sredni, S.T., Arias, V.E.A., Andreoli, M.A., de Camargo, B.,
208 Sobrinho, J.S., Prado, J.C.M., Soares, F.A., Villa, L.L., 2011. Low prevalence of HPV in
209 Brazilian children with retinoblastoma. J. Med. Virol. 83, 115–118.
210 <https://doi.org/10.1002/jmv.21925>

211 Armbruster-Moraes, E., Ioshimoto, L.M., Leão, E., Zugaib, M., 1994. Presence of Human
212 Papillomavirus DNA in Amniotic Fluids of Pregnant Women with Cervical Lesions.
213 Gynecol. Oncol. 54, 152–158. <https://doi.org/10.1006/gyno.1994.1185>
214 Bhuvaneswari, A., Pallavi, V.R., Jayshree, R.S., Kumar, R.V., 2012. Maternal transmission of
215 human papillomavirus in retinoblastoma: A possible route of transfer. Indian J. Med.
216 Paediatr. Oncol. Off. J. Indian Soc. Med. Paediatr. Oncol. 33, 210–215.
217 <https://doi.org/10.4103/0971-5851.107080>
218 Buitrago-Pérez, Á., Garaulet, G., Vázquez-Carballo, A., Paramio, J.M., García-Escudero, R., 2009.
219 Molecular Signature of HPV-Induced Carcinogenesis: pRb, p53 and Gene Expression
220 Profiling. Curr. Genomics 10, 26–34. <https://doi.org/10.2174/138920209787581235>
221 Carreira, H., Coutinho, F., Carrilho, C., Lunet, N., 2013. HIV and HPV infections and ocular
222 surface squamous neoplasia: systematic review and meta-analysis. Br. J. Cancer 109,
223 1981–1988. <https://doi.org/10.1038/bjc.2013.539>
224 Chauhan, S., Sen, S., Singh, N., Sharma, A., Chawla, B., Kashyap, S., 2020. Human
225 Papillomavirus Detection Strategies in Retinoblastoma. Pathol. Oncol. Res. 26, 1341–
226 1344. <https://doi.org/10.1007/s12253-018-00577-x>
227 Dyson, N., Howley, P.M., Münger, K., Harlow, E., 1989. The human papilloma virus-16 E7
228 oncoprotein is able to bind to the retinoblastoma gene product. Science 243, 934–937.
229 <https://doi.org/10.1126/science.2537532>
230 Espinoza, J.P.M., Cardenas, V.J.P., Luna, C.A., Fuentes, H.M., Camacho, G.V., Carrera, F.M.,
231 Garcia, J.R.G., 2005. Loss of 10p material in a child with human papillomavirus–positive
232 disseminated bilateral retinoblastoma. Cancer Genet. Cytogenet. 161, 146–150.
233 <https://doi.org/10.1016/j.cancergencyto.2005.01.012>
234 Friend, S.H., Bernards, R., Rogelj, S., Weinberg, R.A., Rapaport, J.M., Albert, D.M., Dryja, T.P.,
235 1986. A human DNA segment with properties of the gene that predisposes to
236 retinoblastoma and osteosarcoma. Nature 323, 643–646.
237 <https://doi.org/10.1038/323643a0>
238 Gillison, M.L., Chen, R., Goshu, E., Rushlow, D., Chen, N., Banister, C., Creek, K.E., Gallie, B.L.,
239 2007. Human retinoblastoma is not caused by known pRb-inactivating human DNA
240 tumor viruses. Int. J. Cancer 120, 1482–1490. <https://doi.org/10.1002/ijc.22516>
241 Hermonat, P.L., Kechelava, S., Lowery, C.L., Korourian, S., 1998. Trophoblasts are the
242 preferential target for human papilloma virus infection in spontaneously aborted
243 products of conception. Hum. Pathol. 29, 170–174. [https://doi.org/10.1016/s0046-8177\(98\)90228-3](https://doi.org/10.1016/s0046-8177(98)90228-3)
244 Javanmard, D., Moein, M., Esghaei, M., Naseripour, M., Monavari, S.H., Bokharaei-Salim, F.,
245 Sadeghipour, A., 2019. Molecular evidence of human papillomaviruses in the
246 retinoblastoma tumor. VirusDisease 30, 360–366. <https://doi.org/10.1007/s13337-019-00540-7>
247 Johnson, D.A., Zhang, J., Frase, S., Wilson, M., Rodriguez-Galindo, C., Dyer, M.A., 2007.
248 Neuronal Differentiation and Synaptogenesis in Retinoblastoma. Cancer Res. 67, 2701–
249 2711. <https://doi.org/10.1158/0008-5472.CAN-06-3754>
250 Knudson, A.G., 1971. Mutation and Cancer: Statistical Study of Retinoblastoma. Proc. Natl.
251 Acad. Sci. U. S. A. 68, 820–823.
252 Lee, S.M., Park, J.S., Norwitz, E.R., Koo, J.N., Oh, I.H., Park, J.W., Kim, S.M., Kim, Y.H., Park, C.-
253 W., Song, Y.S., 2013. Risk of vertical transmission of human papillomavirus throughout
254 pregnancy: a prospective study. PloS One 8, e66368.
255 <https://doi.org/10.1371/journal.pone.0066368>
256 McLaughlin-Drubin, M.E., Munger, K., 2009. Oncogenic Activities of Human Papillomaviruses.
257 Virus Res. 143, 195–208. <https://doi.org/10.1016/j.virusres.2009.06.008>

260 Mohan, A., Venkatesan, N., Kandalam, M., Pasricha, G., Acharya, P., Khetan, V., Gopal, L.,
261 Sharma, T., Biswas, J., Krishnakumar, S., 2009. Detection of human papillomavirus DNA
262 in retinoblastoma samples: a preliminary study. *J. Pediatr. Hematol. Oncol.* 31, 8–13.
263 <https://doi.org/10.1097/MPH.0b013e31818b373b>

264 Montoya-Fuentes, H., de la Paz Ramirez-Munoz, M., Villar-Calvo, V., Suarez-Rincon, A.E.,
265 Ornelas-Aguirre, J.M., Vazquez-Camacho, G., Orbach-Arbouys, S., Bravo-Cuellar, A.,
266 Sanchez-Corona, J., 2003. Identification of DNA sequences and viral proteins of 6
267 human papillomavirus types in retinoblastoma tissue. *Anticancer Res.* 23, 2853–2862.

268 Moody, C.A., Laimins, L.A., 2010. Human papillomavirus oncoproteins: pathways to
269 transformation. *Nat. Rev. Cancer* 10, 550–560. <https://doi.org/10.1038/nrc2886>

270 Orjuela, M., Castaneda, V.P., Ridaura, C., Lecona, E., Leal, C., Abramson, D.H., Orlow, I., Gerald,
271 W., Cordon-Cardo, C., 2000. Presence of human papilloma virus in tumor tissue from
272 children with retinoblastoma: an alternative mechanism for tumor development. *Clin.*
273 *Cancer Res. Off. J. Am. Assoc. Cancer Res.* 6, 4010–4016.

274 Palazzi, M.A., Yunes, J.A., Cardinali, I.A., Stangenhans, G.P., Brandalise, S.R., Ferreira, S.A.,
275 Sobrinho, J.S.P., Villa, L.L., 2003. Detection of oncogenic human papillomavirus in
276 sporadic retinoblastoma. *Acta Ophthalmol. Scand.* 81, 396–398.
277 <https://doi.org/10.1034/j.1600-0420.2003.00112.x>

278 Roman, A., Munger, K., 2013. The papillomavirus E7 proteins. *Virology*, Special Issue: The
279 Papillomavirus Episteme 445, 138–168. <https://doi.org/10.1016/j.virol.2013.04.013>

280 Ryoo, N.-K., Kim, J.-E., Choung, H.-K., Kim, N., Lee, M.-J., Khwarg, S.-I., 2013. Human Papilloma
281 Virus in Retinoblastoma Tissues from Korean Patients. *Korean J. Ophthalmol.* KJO 27,
282 368–371. <https://doi.org/10.3341/kjo.2013.27.5.368>

283 Sachdeva, U.M., O'Brien, J.M., 2012. Understanding pRb: toward the necessary development
284 of targeted treatments for retinoblastoma. *J. Clin. Invest.* 122, 425–434.
285 <https://doi.org/10.1172/JCI57114>

286 Shetty, O.A., Naresh, K.N., Banavali, S.D., Shet, T., Joshi, R., Qureshi, S., Mulherkar, R., Borges,
287 A., Desai, S.B., 2012. Evidence for the presence of high risk human papillomavirus in
288 retinoblastoma tissue from nonfamilial retinoblastoma in developing countries.
289 *Pediatr. Blood Cancer* 58, 185–190. <https://doi.org/10.1002/pbc.23346>

290 Shukla, S., Bharti, A.C., Mahata, S., Hussain, S., Kumar, R., Hedau, S., Das, B.C., 2009. Infection
291 of human papillomaviruses in cancers of different human organ sites. *Indian J. Med.*
292 *Res.* 130, 222–233.

293 Skoczyński, M., Goździcka-Józefiak, A., Kwaśniewska, A., 2014. Risk factors of the vertical
294 transmission of human papilloma virus in newborns from singleton pregnancy -
295 preliminary report. *J. Matern.-Fetal Neonatal Med. Off. J. Eur. Assoc. Perinat. Med.*
296 *Fed. Asia Ocean. Perinat. Soc. Int. Soc. Perinat. Obstet.* 27, 239–242.
297 <https://doi.org/10.3109/14767058.2013.807238>

298 Tseng, C.J., Lin, C.Y., Wang, R.L., Chen, L.J., Chang, Y.L., Hsieh, T.T., Pao, C.C., 1992. Possible
299 transplacental transmission of human papillomaviruses. *Am. J. Obstet. Gynecol.* 166,
300 35–40. [https://doi.org/10.1016/0002-9378\(92\)91825-u](https://doi.org/10.1016/0002-9378(92)91825-u)

301 Whyte, P., Buchkovich, K.J., Horowitz, J.M., Friend, S.H., Raybuck, M., Weinberg, R.A., Harlow,
302 E., 1988. Association between an oncogene and an anti-oncogene: the adenovirus E1A
303 proteins bind to the retinoblastoma gene product. *Nature* 334, 124–129.
304 <https://doi.org/10.1038/334124a0>

305 Xu, S., Liu, L., Lu, S., Ren, S., 1998. Clinical observation on vertical transmission of human
306 papillomavirus. *Chin. Med. Sci. J. Chung-Kuo Hsueh Ko Hsueh Tsa Chih* 13, 29–31.

307 zur Hausen, H., 2002. Papillomaviruses and cancer: from basic studies to clinical application.
308 *Nat. Rev. Cancer* 2, 342–350. <https://doi.org/10.1038/nrc798>

309

In situ hybridization

Retina

Immunohistochemistry

Figure 1. Comparative results of *in situ* hybridization and immunostaining for HPV in retinoblastoma, using the Green and DAB chromogens. A, C and E: hybridization in retinal areas, tumor and optic nerve, respectively. B, D and F: immunostaining in retinal areas (E6), tumor (E1+E4) and optic nerve (L1), respectively. Arrows in E and F.

Tumor cells

Optical nerve (endothelial cells and nerve bundles)

Immunohistochemical Detection of early and late Human Papillomaviruses (HPV) proteins in Retinoblastoma tissue

Daily et al., 2020.