Adaptive short term COVID-19 prediction for India

Shuvrangshu Jana^{a,*}, Debasish Ghose^a

^a*Department of Aerospace Engineering, IISc*

Abstract

In this paper, a data-driven adaptive model for infection of COVID-19 is formulated to predict the confirmed total cases and active cases of an area over 4 weeks. The parameter of the model is always updated based on daily observations. It is found that the short term prediction of up to 3-4 weeks can be possible with good accuracy. Detailed analysis of predicted value and the actual value of confirmed total cases and active cases for India from 1st June to 3rd July is provided. Prediction over 7, 14, 21, 28 days has the accuracy about $0.73\% \pm 1.97\%$, $1.92\% \pm 2.95\%$, $4.34\% \pm 3.91\%$, $6.40\% \pm 9.26\%$ of the actual value of confirmed total cases. Similarly, the 7, 14, 21, 28 days prediction has the accuracy about $1.24\% \pm 6.57\%$, $3.04\% \pm 10.00\%$, $6.33\% \pm 16.12\%$, $10.20\% \pm 24.14\%$ of the actual value of confirmed active cases.

Keywords: COVID-19, Prediction

1. Introduction

The accurate predictions of the spread of the novel coronavirus (COVID-19) are essential for planning and management of medical resources as well as lockdown strategy. A good prediction can avoid the shortage of critical medical resources, economical loss associated with unnecessary lockdowns. Mathematical
 5 modelling of infectious disease is generally performed by classifying the population into different compartments such as Susceptible, Infectious, Recovered, Deceased. Various classical compartmental model reported in literature such as

*Corresponding author

Email address: shuvra.ce@gmail.com (Shuvrangshu Jana)

SIR, SIS, SIRD, MSIR, SEIR, SEIS, MSEIR, MSEIRS. In order to model the
 10 spreading of corona virus, various modification of the SIR model is proposed
 in literature to include travel information [1], behavioural changes [2], various
 intervention measures such as quarantine, strict social distancing, tracing and
 isolation [3],[4], [5]. The model parameters are generally obtained for a specific
 15 area as there can be huge variation in these parameters based on the government
 policies, economical factors, adoption of social distancing measures. Prediction
 for the Spread of COVID-19 for a specific country is developed such for India
 [6, 7], China [8, 9, 10], South Korea-Italy [1], Iran [11],Korea [2], UK [12].

The estimation of spread of COVID-19 is made based on the assumption
 on rate of growth of total cumulative cases following a certain pattern such as
 20 gaussian [13], Verhulst equation [14], Lotka-Volterra dynamic model[15], Gom-
 pertz equation [16]. Estimation for Greece, Netherlands, Germany, Italy, Spain,
 France, United Kingdom and United States is developed using Gaussian fitting
 hypothesis based on the assumptions that evolutions of infected cases are of
 Gaussian in nature [13]. In [17], estimations for COVID-19 is performed from
 25 the fatality data for India under the assumption that the growth of infected
 cases is exponential. In [15], Lotka-Volterra dynamic model is used to repre-
 sent the growth rate and the model coefficients are derived using the Extended
 Kalman filter techniques. In [18], Kalman filter-based techniques are used for
 short term prediction model.

30 Time-invariant SIR model is not effective to predict the spread of corona
 virus[19], considering the fact the large variation in the transmission dynam-
 ics. Also, it is difficult to fit a single model in different areas. So, Prediction
 models are developed based on the time series analysis of the reported data,
 where the model parameters c are estimated based on the previous observations
 35 of the confirmed cases. Data-driven estimation methods using long short-term
 memory (LSTM) and curve fitting is used for prediction of COVID-19 cases for
 India in 30 days ahead [20]. Deceased cases of COVID-19 prediction for one
 month is developed using a hybrid model using discrete wavelet decomposition
 and autoregressive integrated moving average (ARIMA) models [21]. In [22],

40 Autoregressive Integrated moving average model is used to predict the daily number of COVID-9 cases up to 4 weeks. A mathematical model for prediction is developed using power series polynomials, where the coefficients of the polynomials are obtained using the least square approximations.

Generally, it is observed that it is difficult to predict in the long term due
45 to the various uncertainties involved in the government decision-making process, social distancing measured followed by the people, availability of medical resources. There is a time lag between the time of infection and the time of reporting of infection. So in short term prediction has less uncertainty compared to the long term. A good general prediction model should use information about
50 the local state as less as possible.

In this paper, a short term prediction model is proposed based on the concept of SIR model, where the parameters are updated continuously based on the observations using weighted least square techniques. It is observed that the spread of COVID-19, recovery and deceased rate can be approximated using
55 time-varying polynomials. The proposed model is validated by comparing the true and actual value of June for India over the past predictions on time different time windows such as 7 days, 14 days, 21 days, and 28 days. In case of 2 days time window, It is found that error in prediction is within $6.40\% \pm 9.26\%$ and $10.22\% \pm 24.14\%$ for the prediction of total cases and active cases.

60 The rest of the paper is described as follows. The basic model is described in Section 2. The model parameters are estimated in Section 3. The proposed model is validated with COVID-19 statistics of India in Section 5. Expected total and active cases for a month are predicted in Section 7. The prediction results are discussed and concluded in Discussions and Conclusions Sections
65 respectively.

2. Model formulation

The total number of people who have been infected with the virus can be classified as active, recovered or deceased cases. Few cases will be unreported

due to non-development of symptoms in the infected person. In Fig. 1, different categories of different cases are shown. We will consider the population as

Figure 1

Figure 2: Different compartment

70

susceptible (S), Infectious (I), and Removed (as shown in Fig. 2). Infectious cases are divided into two categories, active cases (A) and active unreported cases A_{ur} . Similarly, recovered and deceased cases are also classified as those who are reported and unreported cases. So, if N is the total population, then,

$$S + A + A_{ur} + R + R_{ur} + D + D_{ur} = N \quad (1)$$

75

where, R , R_{ur} are the reported recovered and unreported recovered cases respectively. Similarly, D , D_{ur} are the reported and unreported deceased cases

respectively. Also,

$$I = A + A_{ur} \quad (2)$$

As per standard SIR model, If β is the infection rate, then,

$$\frac{dS}{dt} = -\frac{\beta IS}{N} \quad (3)$$

Also, we can write,

$$\frac{dA}{dt} = -\frac{dS}{dt} - \gamma(t)A - \mu(t)A - \frac{dA_{ur}}{dt} - \gamma_{ur}(t)A_{ur} - \mu_{ur}(t)A_{ur} \quad (4)$$

80 where $\gamma(t)$ and $\mu(t)$ are the rate of recovered and death among the reported cases; and γ_{ur} and μ_{ur} are the rate of recovered and death cases among unreported cases.

$$\frac{dR}{dt} = \gamma(t)A \quad (5)$$

$$\frac{dD}{dt} = \mu(t)A \quad (6)$$

$$\frac{dR_{ur}}{dt} = \gamma_{ur}(t)A_{ur} \quad (7)$$

$$\frac{dD_{ur}}{dt} = \mu_{ur}(t)A_{ur} \quad (8)$$

3. Parameter estimation

85 The different parameter of the above model is estimated from the observed data and dynamically adjusted using the daily observations. The active cases, recovered and deceased from the reported cases are readily available; whereas, the similar statistics for the unreported cases need to be estimated from random antibody tests or serological tests. The confirmed active cases at each day are obtained by the following equation

$$A = T - R - D \quad (9)$$

90 where, T is the total number of reported infected person. Then we have,

$$\frac{dA}{dt} = \frac{dT}{dt} - \frac{dR}{dt} - \frac{dD}{dt} \quad (10)$$

From the previous reported daily statistics of daily new infected cases, recovered cases, and deceased cases; the rate of growth of the total number of cases, recovered cases and deceased cases are obtained.

$$\frac{dT}{dt} = f_T(t) \quad (11)$$

$$\frac{dD}{dt} = f_D(t), \quad \frac{dR}{dt} = f_R(t) \quad (12)$$

The rate of growth of total cases are obtained by taking derivative of the variable T . T can be represented by different functions. In the case of India, it is observed that T can be approximated using a fourth-order polynomial. Let consider T be,

$$T = a_T t^4 + b_T t^3 + c_T t^2 + d_T t + e_T \quad (13)$$

The coefficients (a_T, b_T, c_T, d_T, e_T) of the function T is obtained using the weighted least squares method by minimizing the error between the predicted value and the observed daily values. The weighted least squares method is used to provide more weightage on the recent values which in turn reflects the status of lockdown, social distancing measured followed by the citizens. The following cost function is used.

$$\min_{\zeta} \sum_{t=t_0}^{t=t_f} w_t (y_t - \sum_{j=1}^{j=n} X_{jt} \zeta_j)^2 \quad (14)$$

where y_t is the observed cumulative total cases, T is expressed as $T = X\zeta$, and ζ are the coefficients of the polynomial. The weights (w_t are selected from the previous observations as 1(last 7 days), 0.9 (last 7-14 days), 0.8 (last 14-21 days), 0.5 (last 21-28 days) and 0.2 for further previous days.

The weights need to be selected properly for an area considering the lockdown/unlock period. The optimum values of ζ are calculated from the equation 14 as follows.

$$\hat{\zeta} = (X^T W X)^{-1} X^T W y_t \quad (15)$$

where W is the diagonal matrix consist of weights (w_t). It is observed that reported recovered and deceased cases can be approximated using the fourth-order and second-order polynomial respectively.

$$R = a_R t^4 + b_R^t 3 + c_R t^2 + d_R t + e_R \quad (16)$$

$$D = a_D t^2 + b_D^t + c_D \quad (17)$$

4. Prediction

115 Prediction values are updated based on the current observations. The data of total cases recovered cases, and deceased cases from 14th March to 3rd July for India is considered in the analysis. The data is collected from the “<https://api.covid19india.org/>” website, where the daily COVID-19 updates of various states and central government’s of India are recorded. The variation of
120 total cases, cumulative recovered cases, cumulative deceased cases, and corresponding active cases on daily basis are plotted in Figure 3a, Figure 3b, Figure 3c, Figure 3d respectively. Based on the observations till 3rd July, the prediction for the next 14 days is shown in Figure 4a and Figure 4b. The variation of coefficients of the prediction curve of the total cases is tabulated in Table 1.

(a) Total cases

(b) Cumulative recovered cases

(c) Cumulative deceased cases

(d) Active cases

Figure 3: Total case, Recovered case, Deceased case, Active Case

(a) Total cases prediction

(b) Active cases prediction

Figure 4: Prediction of total cases and active cases

Table 1: Total cases prediction curve coefficients

Date of prediction	a_T	b_T	c_T	d_T	e_T
03.07.2020	0.0031	0.0442	8.85	-116.96	595.4
02.07.2020	0.003	0.0496	8.59	-112.71	580.6
01.07.2020	0.003	0.0548	8.38	-110.48	578.6
30.06.2020	0.0028	0.0889	6.34	-68.53	381.5
29.06.2020	0.0025	0.1615	1.86	26.12	-72.7
28.06.2020	0.002	0.2576	-4.13	153.64	-687.4
27.06.2020	0.0016	0.3317	-8.71	250.11	-1148.3
26.06.2020	0.0013	0.3811	-11.67	311.14	-1433.9
25.06.2020	0.0012	0.4033	-12.94	336.07	-1546.1
24.06.2020	0.0012	0.4038	-12.88	333.34	-1528.1
23.06.2020	0.012	0.3984	-12.47	323.59	-1478
22.06.2020	0.001	0.4402	-14.92	372.64	-1701.1
21.06.2020	0.0006	0.5134	-19.29	461.18	-2106.9
20.06.2020	0.0004	0.5559	-21.85	513.52	-2347.7
19.06.2020	0.0004	0.5502	-21.60	509.79	-2335.1
18.06.2020	0.0007	0.5007	-18.86	457.595	-2108.5
17.06.2020	0.0012	0.4045	-13.50	355.37	-1665
16.06.2020	0.0019	0.2981	-7.59	243.34	-1181.7
15.06.2020	0.0022	0.2338	-4.21	181.97	-925.9
14.06.2020	0.0024	0.2027	-2.61	153.62	-810.5
13.06.2020	0.0027	0.153	0.0002	106.73	-617.8
12.06.2020	0.0031	0.0952	3.0	53.13	-399.5
11.06.2020	0.0036	0.0046	7.72	-30.60	-59.9
10.06.2020	0.0042	-0.0945	12.85	-121.40	306
9.06.2020	0.0044	-0.12	14.08	-143.07	393.2
8.06.2020	0.0043	-0.11	13.43	-131.40	345.8
7.06.2020	0.0044	-0.116	13.92	-139.45	376.6
6.06.2020	0.0044	-0.119	14.06	-141.79	385.7
5.06.2020	0.0045	-0.13	14.67	-152.24	425.9
4.06.2020	0.0047	-0.16	16.17	-176.80	518.1
3.06.2020	0.005	-0.204	18.14	-208.85	637.4
2.06.2020	0.0047	-0.169	16.43	-180.86	533.3
1.06.2020	0.0044	-0.12	14.19	-144.43	399

125 5. Model Validation

5.1. Total case prediction validation

The difference between the predicted value and the actual value is compared to check the efficiency of the prediction algorithm. The predicted value on the 7th, 14th, 21th and 28th days back is considered for comparison. The predictions
130 from June 1st to July 3rd are considered for detailed analysis. The total case prediction over a 7 days time window and the corresponding actual value is shown in Table 2. In 26th June, the total cases prediction for 3rd July was 61130 and the actual value reported is 627065. So, the error in prediction is 2.54 % of the actual value of 627065. From Table 2, over the complete duration
135 of June 1st to July 3rd, the error between the actual and predicted value is $0.73\% \pm 1.97\%$ of the actual value.

The predicted and actual values over a 14 days prediction window from June 1st is tabulated in Table 3. As per Table 3 In case of prediction on 14 days duration, the error between the actual and the predicted value is found to
140 be $1.92\% \pm 2.95\%$ of the actual value. Similarly, prediction over 21 days and 28 days time window are tabulated in 4 and 5. The error between the predicted value and the actual value over 21 days and 28 days is $4.34\% \pm 3.91\%$ and $6.40\% \pm 9.26\%$ of the actual value respectively.

The error in prediction from mid-April to July 3rd is plotted in Figure 5a. In
145 Figure 5a, magenta points show the predicted value on 7 days back and the black points are the predicted value on that day. At each day, the difference between the black and magenta point is the difference in predicted and the actual value. Similar plots for the prediction window of 14 days, 21 days, and 28 days are plotted in the Figure 5b, Figure 5c, and Figure 5d respectively.

Table 2: Validation of prediction for 7 days window

Date during pre-diction	Date of prediction	Prediction	Actual	Error from actual(%)
26.06.2020	03.07.2020	611130	627065	-2.54
25.06.2020	02.07.2020	590940	605118	-2.27
24.06.2020	01.07.2020	571970	585689	-2.31
23.06.2020	30.06.2020	553680	567433	-2.26
22.06.2020	29.06.2020	534770	549094	-2.47
21.06.2020	28.06.2020	516140	529484	-2.40
20.06.2020	27.06.2020	499230	509342	-1.91
19.06.2020	26.06.2020	484050	491087	-1.30
18.06.2020	25.06.2020	470350	472882	-0.58
17.06.2020	24.06.2020	457990	456014	+0.40
16.06.2020	23.06.2020	445710	440358	+1.25
15.06.2020	22.06.2020	432190	426798	+1.62
14.06.2020	21.06.2020	418110	411647	+1.87
13.06.2020	20.06.2020	404670	395729	+2.44
12.06.2020	19.06.2020	391710	381091	+2.94
11.06.2020	18.06.2020	379170	367262	+3.33
10.06.2020	17.06.2020	367390	354154	+3.76
9.06.2020	16.06.2020	355690	343068	+3.67
8.06.2020	15.06.2020	342430	333036	+3.01
7.06.2020	14.06.2020	329090	321631	+2.55
6.06.2020	13.06.2020	316490	309592	+2.43
5.06.2020	12.06.2020	304000	298286	+2.17
4.06.2020	11.06.2020	292000	287151	+1.89
3.06.2020	10.06.2020	280810	275995	+1.53
2.06.2020	9.06.2020	270090	266014	+1.31
1.06.2020	8.06.2020	258580	257478	+ 0.77
31.05.2020	7.06.2020	247430	246596	+0.33
30.05.2020	6.06.2020	237260	236188	+0.25
29.05.2020	5.06.2020 ₁	228180	226716	+0.62
28.05.2020	4.06.2020	219640	216869	+0.13
27.05.2020	3.06.2020	211570	207180	+1.90
26.05.2020	2.06.2020	202590	198368	+1.95
25.05.2020	1.06.2020	192820	190645	+1.20

Table 3: Validation of prediction for 14 days window

Date during pre-diction	Date of prediction	Prediction	Actual	Error from actual(%)
19.06.2020	03.07.2020	603000	627065	-3.84
18.06.2020	02.07.2020	588280	605118	-2.78
17.06.2020	01.07.2020	576050	585689	-1.65
16.06.2020	30.06.2020	563900	567433	-0.62
15.06.2020	29.06.2020	549090	549094	-0.0007
14.06.2020	28.06.2020	532640	529484	+0.60
13.06.2020	27.06.2020	517310	509342	+1.56
12.06.2020	26.06.2020	502820	491087	+2.39
11.06.2020	25.06.2020	488880	472882	+3.38
10.06.2020	24.06.2020	476370	456014	+4.46
9.06.2020	23.06.2020	463910	440358	+5.35
8.06.2020	22.06.2020	448000	426798	+4.97
7.06.2020	21.06.2020	431400	411647	+4.80
6.06.2020	20.06.2020	416030	395729	+5.13
5.06.2020	19.06.2020	400600	381091	+5.12
4.06.2020	18.06.2020	385890	367262	+5.07
3.06.2020	17.06.2020	372490	354154	+5.18
2.06.2020	16.06.2020	359780	343068	+4.87
1.06.2020	15.06.2020	344840	333036	+3.54
31.05.2020	14.06.2020	330210	321631	+2.67
30.05.2020	13.06.2020	317490	309592	+2.55
29.05.2020	12.06.2020	306910	298286	+2.89
28.05.2020	11.06.2020	297380	287151	+3.56
27.05.2020	10.06.2020	288690	275995	+4.60
26.05.2020	9.06.2020	277750	266014	+4.41
25.05.2020	8.06.2020	264890	257478	+2.88
24.05.2020	7.06.2020	251330	246596	+1.92
23.05.2020	6.06.2020	238310	236188	+0.89
22.05.2020	5.06.2020 ₁₂	226270	226716	-0.20
21.05.2020	4.06.2020	214030	216869	-1.31
20.05.2020	3.06.2020	201940	207180	-2.53
19.05.2020	2.06.2020	192680	198368	-2.87
18.05.2020	1.06.2020	183700	190645	-3.64

Table 4: Validation of prediction for 21 days window

Date during pre-diction	Date of prediction	Prediction	Actual	Error from actual(%)
12.06.2020	03.07.2020	636020	627065	+1.43
11.06.2020	02.07.2020	621080	605118	+2.64
10.06.2020	01.07.2020	608540	585689	+3.90
9.06.2020	30.06.2020	596090	567433	+5.05
8.06.2020	29.06.2020	577340	549094	+5.14
7.06.2020	28.06.2020	556870	529484	+5.17
6.06.2020	27.06.2020	538410	509342	+5.71
5.06.2020	26.06.2020	519600	491087	+5.81
4.06.2020	25.06.2020	501850	472882	+6.13
3.06.2020	24.06.2020	486130	456014	+6.60
2.06.2020	23.06.2020	471440	440358	+7.06
1.06.2020	22.06.2020	452240	426798	+ 5.96
31.05.2020	21.06.2020	433210	411647	+ 5.24
30.05.2020	20.06.2020	417510	395729	+5.50
29.05.2020	19.06.2020	405600	381091	+ 6.43
28.05.2020	18.06.2020	395580	367262	+7.71
27.05.2020	17.06.2020	386990	354154	+9.27
26.05.2020	16.06.2020	374040	343068	+9.03
25.05.2020	15.06.2020	357380	333036	+7.31
24.05.2020	14.06.2020	338890	321631	+5.37
23.05.2020	13.06.2020	321070	309592	+3.71
22.05.2020	12.06.2020	304740	298286	+2.16
21.05.2020	11.06.2020	287530	287151	-0.13
20.05.2020	10.06.2020	270180	275995	-2.11
19.05.2020	9.06.2020	257870	266014	-3.06
18.05.2020	8.06.2020	245800	257478	-4.53
17.05.2020	7.06.2020	236170	246596	-4.23
16.05.2020	6.06.2020	234360	236188	-0.77
15.05.2020	5.06.2020 ₁₃	231570	226716	-2.14
14.05.2020	4.06.2020	228380	216869	+5.31
13.05.2020	3.06.2020	224320	207180	+8.27
12.05.2020	2.06.2020	218790	198368	+10.30
11.05.2020	1.06.2020	208670	190645	+9.45

Table 5: Validation of prediction for 28 days window

Date during pre-diction	Date of prediction	Prediction	Actual	Error from actual(%)
5.06.2020	03.07.2020	664400	627065	+5.95
4.06.2020	02.07.2020	643270	605118	+6.30
3.06.2020	01.07.2020	625210	585689	+6.75
2.06.2020	30.06.2020	608650	567433	+7.26
1.06.2020	29.06.2020	584200	549094	+6.39
31.05.2020	28.06.2020	559650	529484	+5.69
30.05.2020	27.06.2020	540520	509342	+6.12
29.05.2020	26.06.2020	527610	491087	+7.44
28.05.2020	25.06.2020	517820	472882	+9.50
27.05.2020	24.06.2020	510370	456014	+11.92
26.05.2020	23.06.2020	495480	440358	+12.52
25.06.2020	22.06.2020	474220	426798	+11.11
24.05.2020	21.06.2020	449360	411647	+9.16
23.05.2020	20.06.2020	425280	395729	+7.47
22.05.2020	19.06.2020	403440	381091	+5.86
21.05.2020	18.06.2020	379570	367262	+3.35
20.05.2020	17.06.2020	355050	354154	-0.25
19.05.2020	16.06.2020	338820	343068	-1.24
18.05.2020	15.06.2020	322730	332036	-3.09
17.05.2020	14.06.2020	310660	321631	-2.89
16.05.2020	13.06.2020	312340	309592	+0.88
15.05.2020	12.06.2020	312310	298286	+4.45
14.05.2020	11.06.2020	311560	287151	+8.19
13.05.2020	10.06.2020	309330	275995	+12.07
12.05.2020	9.06.2020	304560	266014	+14.49
11.05.2020	8.06.2020	291510	257478	+13.22
10.05.2020	7.06.2020	291040	246596	+18.02
9.05.2020	6.06.2020	287750	236188	+21.83
8.05.2020	5.06.2020 ₁₄	274110	226716	+20.90
7.05.2020	4.06.2020	250330	216869	+15.43
6.05.2020	3.06.2020	219070	207180	+5.74
5.05.2020	2.06.2020	174510	198368	-12.03
4.05.2020	1.06.2020	137740	190645	-27.75

Figure 5: Validation of prediction of total cases

5.2. Active case prediction validation

The prediction results in case of active cases are compared with the actual value of the active cases. The prediction values and the error in prediction of active cases over 7 days, 14 days, 21 days, 28 days time duration are tabulated in Table 6, Table 7, Table 8, and Table 9 respectively. The difference between the predicted active case and the actual active case is found to be $1.24\% \pm 6.57\%$, $3.04\% \pm 10.00\%$, $6.33\% \pm 16.12\%$, $10.20\% \pm 24.14\%$ for the prediction window of 7 days, 14 days, 21 days, 28 days respectively. The error in active case prediction from mid April to July 3rd over different time window is plotted in Figure 6a, Figure 6b, Figure 6c, and Figure 6d.

Table 6: Validation of active case prediction for 7 days window

Date during pre-diction	Date of prediction	Prediction	Actual	Error from actual(%)
26.06.2020	03.07.2020	218250	228936	-4.67
25.06.2020	02.07.2020	215180	227366	-5.36
24.06.2020	01.07.2020	211870	220439	-3.89
23.06.2020	30.06.2020	209110	215254	-2.85
22.06.2020	29.06.2020	205600	210829	-2.48
21.06.2020	28.06.2020	203200	203234	-0.02
20.06.2020	27.06.2020	200250	197735	+1.27
19.06.2020	26.06.2020	198350	190107	+4.34
18.06.2020	25.06.2020	196700	186286	+5.59
17.06.2020	24.06.2020	197200	182931	+7.80
16.06.2020	23.06.2020	197860	178205	+11.02
15.06.2020	22.06.2020	195120	175836	+10.97
14.06.2020	21.06.2020	190990	170186	+12.22
13.06.2020	20.06.2020	186400	168550	+10.59
12.06.2020	19.06.2020	181050	163305	+10.87
11.06.2020	18.06.2020	175630	160560	+9.39
10.06.2020	17.06.2020	170350	154683	+10.13
9.06.2020	16.06.2020	164770	152827	+7.81
8.06.2020	15.06.2020	157490	153830	+2.38
7.06.2020	14.06.2020	150030	150107	-0.05
6.06.2020	13.06.2020	142560	146469	-2.67
5.06.2020	12.06.2020	134830	142814	-5.59
4.06.2020	11.06.2020	126700	138117	-8.27
3.06.2020	10.06.2020	121200	133594	-9.28
2.06.2020	9.06.2020	117710	129519	-9.12
1.06.2020	8.06.2020	115620	126425	-8.55
31.05.2020	7.06.2020	118660	120995	-1.93
30.05.2020	6.06.2020	116160	116317	-0.13
29.05.2020	5.06.2020 ¹⁶	111900	111901	-0.0008
28.05.2020	4.06.2020	108110	106718	+1.30
27.05.2020	3.06.2020	102990	101077	+1.89
26.05.2020	2.06.2020	97190	97018	-0.18
25.05.2020	1.06.2020	91420	93378	-2.10

Table 7: Validation of active case prediction for 14 days window

Date during pre-diction	Date of prediction	Prediction	Actual	Error from actual(%)
19.06.2020	03.07.2020	231130	228936	+ 0.96
18.06.2020	02.07.2020	231470	227366	+1.81
17.06.2020	01.07.2020	236290	220439	+7.19
16.06.2020	30.06.2020	241850	215254	+12.36
15.06.2020	29.06.2020	240990	210829	+14.31
14.06.2020	28.06.2020	237250	203234	+16.74
13.06.2020	27.06.2020	232780	197735	+17.72
12.06.2020	26.06.2020	226850	190107	+19.33
11.06.2020	25.06.2020	220720	186286	+18.48
10.06.2020	24.06.2020	214960	182931	+17.51
9.06.2020	23.06.2020	208780	178205	+17.16
8.06.2020	22.06.2020	199000	175836	+13.17
7.06.2020	21.06.2020	188500	170186	+10.76
6.06.2020	20.06.2020	177690	168550	+5.42
5.06.2020	19.06.2020	166080	163305	-1.70
4.06.2020	18.06.2020	151960	160560	-5.36
3.06.2020	17.06.2020	143240	154683	-7.40
2.06.2020	16.06.2020	138830	152827	-9.16
1.06.2020	15.06.2020	137290	153830	-10.75
31.05.2020	14.06.2020	140460	150107	-6.43
30.05.2020	13.06.2020	150680	146469	+2.88
29.05.2020	12.06.2020	149620	142814	+4.77
28.05.2020	11.06.2020	144630	138117	+4.71
27.05.2020	10.06.2020	140540	133594	+5.20
26.05.2020	9.06.2020	133350	129519	+2.96
25.05.2020	8.06.2020	124460	126425	-1.55
24.05.2020	7.06.2020	115230	120995	-4.76
23.05.2020	6.06.2020	108420	116317	-6.79
22.05.2020	5.06.2020 ¹⁷	101750	111901	-9.07
21.05.2020	4.06.2020	95680	106718	-10.34
20.05.2020	3.06.2020	90980	101077	-9.99
19.05.2020	2.06.2020	89210	97018	-8.05
18.05.2020	1.06.2020	89190	93378	-4.48

Table 8: Validation of active case prediction for 21 days window

Date during pre-diction	Date of prediction	Prediction	Actual	Error from actual(%)
12.06.2020	03.07.2020	281980	228936	+23.17
11.06.2020	02.07.2020	275210	227366	+21.04
10.06.2020	01.07.2020	269210	220439	+22.12
9.06.2020	30.06.2020	262650	215254	+22.01
8.06.2020	29.06.2020	249590	210829	+18.39
7.06.2020	28.06.2020	234960	203234	+15.61
6.06.2020	27.06.2020	219520	197735	+11.01
5.06.2020	26.06.2020	202420	190107	+6.47
4.06.2020	25.06.2020	179170	186286	-3.82
3.06.2020	24.06.2020	165570	182931	-9.49
2.06.2020	23.06.2020	159800	178205	-10.33
1.06.2020	22.06.2020	159190	175836	-9.47
31.05.2020	21.06.2020	167410	170186	-1.63
30.05.2020	20.06.2020	189100	168550	+12.19
29.05.2020	19.06.2020	190740	163305	+16.80
28.05.2020	18.06.2020	185060	160560	+15.26
27.05.2020	17.06.2020	180950	154683	+16.98
26.05.2020	16.06.2020	170900	152827	+11.83
25.05.2020	15.06.2020	157520	153830	+2.40
24.05.2020	14.06.2020	143090	150107	-4.67
23.05.2020	13.06.2020	132980	146469	-9.21
22.05.2020	12.06.2020	122580	142814	-14.17
21.05.2020	11.06.2020	112990	138117	-18.19
20.05.2020	10.06.2020	106000	133594	-20.66
19.05.2020	9.06.2020	104500	129519	-19.32
18.05.2020	8.06.2020	106360	126425	-15.87
17.05.2020	7.06.2020	115190	120995	-4.80
16.05.2020	6.06.2020	122680	116317	+5.47
15.05.2020	5.06.2020 ¹⁸	124070	111901	+10.87
14.05.2020	4.06.2020	128170	106718	+20.10
13.05.2020	3.06.2020	131000	101077	+29.60
12.05.2020	2.06.2020	130930	97018	+34.95
11.05.2020	1.06.2020	125400	93378	+34.29

Table 9: Validation of active case prediction for 28 days window

Date during pre-diction	Date of prediction	Prediction	Actual	Error from actual(%)
5.06.2020	03.07.2020	244420	228936	+6.76
4.06.2020	02.07.2020	208060	227366	-8.49
3.06.2020	01.07.2020	187420	220439	-14.98
2.06.2020	30.06.2020	179740	215254	-16.50
1.06.2020	29.06.2020	180490	210829	-14.39
31.05.2020	28.06.2020	196170	203234	-3.47
30.05.2020	27.06.2020	234880	197735	+18.78
29.05.2020	26.06.2020	240880	190107	+26.71
28.05.2020	25.06.2020	234600	186286	+25.93
27.05.2020	24.06.2020	230870	182931	+26.20
26.05.2020	23.06.2020	217000	178205	+21.77
25.06.2020	22.06.2020	197330	175836	+12.22
24.05.2020	21.06.2020	175470	170186	+3.11
23.05.2020	20.06.2020	160680	168550	-4.67
22.05.2020	19.06.2020	144820	163305	-11.32
21.05.2020	18.06.2020	130030	160560	-19.01
20.05.2020	17.06.2020	119630	154683	-22.66
19.05.2020	16.06.2020	118580	152827	-22.41
18.05.2020	15.06.2020	123330	153830	-19.83
17.05.2020	14.06.2020	140410	150107	-6.46
16.05.2020	13.06.2020	155690	146469	+6.29
15.05.2020	12.06.2020	160390	142814	+12.31
14.05.2020	11.06.2020	170060	138117	+23.13
13.05.2020	10.06.2020	177650	133594	+32.98
12.05.2020	9.06.2020	180350	129519	+39.25
11.05.2020	8.06.2020	173260	126425	+37.05
10.05.2020	7.06.2020	176620	120995	+45.97
9.05.2020	6.06.2020	176260	116317	+51.54
8.05.2020	5.06.2020 ¹⁹	171530	111901	- 53.28
7.05.2020	4.06.2020	155630	106718	+45.83
6.05.2020	3.06.2020	135500	101077	+34.05
5.05.2020	2.06.2020	100131	97018	+3.21
4.05.2020	1.06.2020	69995	93378	-25.04

Figure 6: Validation of prediction of active cases

6. Discussions

The prediction is good in the smaller time window and error between the actual and predicted case grows with larger time window prediction. The prediction error from June 1st to 3rd July is tabulated in Table 10. The prediction is better in case of total cases compared to active cases. The reason behind the higher value of error bound could be the frequent change in discharge policy by the various state governments to adjust the growth of active patients in hospitals, which in turn affected the recovery rates. Also sometimes the deceased cases are adjusted in a single day after detailed accounting which caused a high

jump in the deceased curve. Discharge policy by the governments and reporting
of deceased cases have caused the large variation in the actual value of active
cases over daily-basis. However, the reporting of the total case is smooth (Figure
3a); therefore, the bound of prediction error is also less.

Table 10: **Summary of prediction error**

Parameter	7 days	14 days	21 days	28 days
Total cases	0.73 % $\pm$ 1.97 %	1.92 % $\pm$ 2.95 %	4.34 % $\pm$ 3.91 %	6.40 % $\pm$ 9.26 %
Active cases	1.24 % $\pm$ 6.57 %	3.04 % $\pm$ 10.00 %	6.33 % $\pm$ 16.12 %	10.20 % $\pm$ 24.14 %

7. Future prediction

Future predictions of total cases of India from the proposed algorithm over
different time windows are provided in Table 11, Table 12, Table 13, and Table
14. We will compare this table in future to further validate our algorithm.
Similar predictions for the active cases are tabulated in Table 15, Table 16,
Table 17, and Table 18. The prediction for next 28 days based on the upto date
observations are also included in Table 19 for reference.

Table 11: **Future Prediction of total cases for 7 days window**

Date during prediction	Date of prediction	Prediction	Actual	Error(%)
7.06.2020	14.07.2020	923271		
8.06.2020	15.07.2020	953582		
9.06.2020	16.07.2020	984856		
10.06.2020	17.07.2020	1017049		
11.07.2020	18.07.2020	1050612		
12.07.2020	19.07.2020	1085283		
13.07.2020	20.07.2020	1121190		

Table 12: Future prediction of total cases for 14 days window

Date during prediction	Date of prediction	Prediction	Actual	Error(%)
30.06.2020	14.07.2020	904382		
01.07.2020	15.07.2020	933350		
02.07.2020	16.07.2020	962644		
03.07.2020	17.07.2020	994203		
04.07.2020	18.07.2020	1028181		
05.07.2020	19.07.2020	1064689		
06.07.2020	20.07.2020	1102150		
07.07.2020	21.07.2020	1138688		
08.07.2020	22.07.2020	1175003		
09.07.2020	23.07.2020	1212579		
10.07.2020	24.07.2020	1251346		
11.07.2020	25.07.2020	1292120		
12.07.2020	26.07.2020	1334355		
13.07.2020	27.07.2020	1378048		

Table 13: Future prediction of total cases for 21 days window

Date during prediction	Date of prediction	Prediction	Actual	Error(%)
23.06.2020	14.07.2020	867245		
24.06.2020	15.07.2020	893187		
25.06.2020	16.07.2020	922109		
26.06.2020	17.07.2020	954784		
27.06.2020	18.07.2020	990776		
28.06.2020	19.07.2020	1030799		
29.06.2020	20.07.2020	1070675		
30.06.2020	21.07.2020	1107524		
01.07.2020	22.07.2020	1141728		
02.07.2020	23.07.2020	1176221		
03.07.2020	24.07.2020	1214099		
04.07.2020	25.07.2020	1255691		
05.07.2020	26.07.2020	1301005		
06.07.2020	27.07.2020	1347265		
07.07.2020	28.07.2020	1391397		
08.07.2020	29.07.2020	1434624		
09.07.2020	30.07.2020	1479500		
10.07.2020	31.07.2020	1525916		
11.07.2020	01.08.2020	1575199		
12.07.2020	02.08.2020	1626391		
13.07.2020	03.08.2020	1679281		

Table 14: Future prediction of total cases for 28 days window

Date during Prediction	Date of prediction	Prediction	Actual	Error(%)
16.06.2020	14.07.2020	875120		
17.06.2020	15.07.2020	883710		
18.06.2020	16.07.2020	898890		
19.06.2020	17.07.2020	920890		
20.06.2020	18.07.2020	950760		
21.06.2020	19.07.2020	987170		
22.06.2020	20.07.2020	1021400		
23.06.2020	21.07.2020	1049718		
24.06.2020	22.07.2020	1079705		
25.06.2020	23.07.2020	1114064		
26.06.2020	24.07.2020	1153961		
27.06.2020	25.07.2020	1198759		
28.06.2020	26.07.2020	1249335		
29.06.2020	27.07.2020	1298983		
30.06.2020	28.07.2020	1343492		
01.07.2020	29.07.2020	1383608		
02.07.2020	30.07.2020	1423955		
03.07.2020	31.07.2020	1469184		
04.07.2020	01.08.2020	1519869		
05.07.2020	02.08.2020	1575846		
06.07.2020	03.08.2020	1632683		
07.07.2020	04.08.2020	1685699		
08.07.2020	05.08.2020	1736842		
09.07.2020	06.08.2020	1790127		
10.07.2020	07.08.2020	1845392		
11.07.2020	08.08.2020	1904650		
12.07.2020	09.08.2020	1966379		
13.07.2020	10.08.2020	2030057		

Table 15: **Future prediction of active cases for 7 days window**

Date during prediction	Date of prediction	Prediction	Actual	Error(%)
7.06.2020	14.07.2020	305252		
8.06.2020	15.07.2020	314019		
9.06.2020	16.07.2020	322349		
10.06.2020	17.07.2020	330195		
11.07.2020	18.07.2020	338128		
12.07.2020	19.07.2020	346743		
13.07.2020	20.07.2020	357261		

Table 16: **Future prediction of active cases for 14 days window**

Date during prediction	Date of prediction	Prediction	Actual	Error(%)
30.06.2020	14.07.2020	284328		
01.07.2020	15.07.2020	293387		
02.07.2020	16.07.2020	302079		
03.07.2020	17.07.2020	307619		
04.07.2020	18.07.2020	316349		
05.07.2020	19.07.2020	327999		
06.07.2020	20.07.2020	340698		
07.07.2020	21.07.2020	352954		
08.07.2020	22.07.2020	363963		
09.07.2020	23.07.2020	373654		
10.07.2020	24.07.2020	382548		
11.07.2020	25.07.2020	391734		
12.07.2020	26.07.2020	402159		
13.07.2020	27.07.2020	415782		

Table 17: Future prediction of active cases for 21 days window

Date during prediction	Date of prediction	Prediction	Actual	Error(%)
23.06.2020	14.07.2020	265641		
24.06.2020	15.07.2020	267225		
25.06.2020	16.07.2020	267829		
26.06.2020	17.07.2020	271197		
27.06.2020	18.07.2020	280506		
28.06.2020	19.07.2020	292481		
29.06.2020	20.07.2020	307021		
30.06.2020	21.07.2020	318976		
01.07.2020	22.07.2020	330259		
02.07.2020	23.07.2020	340672		
03.07.2020	24.07.2020	346155		
04.07.2020	25.07.2020	356820		
05.07.2020	26.07.2020	372312		
06.07.2020	27.07.2020	389220		
07.07.2020	28.07.2020	405179		
08.07.2020	29.07.2020	419071		
09.07.2020	30.07.2020	430343		
10.07.2020	31.07.2020	440400		
11.07.2020	01.08.2020	451062		
12.07.2020	02.08.2020	463747		
13.07.2020	03.08.2020	481399		

Table 18: **Future prediction of active cases for 28 days window**

Date during prediction	Date of prediction	Prediction	Actual	Error(%)
16.06.2020	14.07.2020	327825		
17.06.2020	15.07.2020	307229		
18.06.2020	16.07.2020	299629		
19.06.2020	17.07.2020	294172		
20.06.2020	18.07.2020	294427		
21.06.2020	19.07.2020	292492		
22.06.2020	20.07.2020	293131		
23.06.2020	21.07.2020	289077		
24.06.2020	22.07.2020	289169		
25.06.2020	23.07.2020	287667		
26.06.2020	24.07.2020	290661		
27.06.2020	25.07.2020	302517		
28.06.2020	26.07.2020	318523		
29.06.2020	27.07.2020	338071		
30.06.2020	28.07.2020	353496		
01.07.2020	29.07.2020	367618		
02.07.2020	30.07.2020	380146		
03.07.2020	31.07.2020	385383		
04.07.2020	01.08.2020	398498		
05.07.2020	02.08.2020	419023		
06.07.2020	03.08.2020	441410		
07.07.2020	04.08.2020	462128		
08.07.2020	05.08.2020	479668		
09.07.2020	06.08.2020	492768		
10.07.2020	07.08.2020	504113		
11.07.2020	08.08.2020	516510		
12.07.2020	09.08.2020	531986		
13.07.2020	10.08.2020	554767		

Table 19: **Future prediction of active and total cases on 13.07.2020**

Date of prediction	Active case	Total case
14.07.2020	312279	932648
15.07.2020	319461	962181
16.07.2020	326767	992453
17.07.2020	334198	1023476
18.07.2020	341756	1055264
19.07.2020	349443	1087831
20.07.2020	357261	1121190
21.07.2020	365210	1155355
22.07.2020	373294	1190340
23.07.2020	381513	1226157
24.07.2020	389869	1262823
25.07.2020	398365	1300350
26.07.2020	407002	1338754
27.07.2020	415782	1378048
28.07.2020	424706	1418247
29.07.2020	433777	1459366
30.07.2020	442997	1501420
31.07.2020	452367	1544423
01.08.2020	461890	1588391
02.08.2020	471566	1633338
03.08.2020	481399	1679281
04.08.2020	491390	1726234
05.08.2020	501541	1774214
06.08.2020	511854	1823235
07.08.2020	522331	1873314
08.08.2020	532974	1924467
09.08.2020	543786	1976709
10.08.2020	554767	2030057

180 8. Conclusions

In this paper, a data-driven prediction model is proposed to predict the total cases and active cases of an area. The model can apply to an area. It is observed that the error between the actual value and predicted value is $4.34\% \pm 3.91\%$ and $6.33\% \pm 16.12\%$ for total cases and active cases respectively over the prediction window of 21 days. We have also provided the prediction for the next 28 days from today for further validation of the proposed algorithm. The short term prediction can be used in the allocation of scar medical resources among different units, optimal lockdown planning.

ACKNOWLEDGEMENTS

190 The authors would like to thank Mayur Shewale and GCDSL lab members for their suggestions in data processing and analysis of results.

Funding

Work is not funded by any funding agency.

Conflicts of interest

195 Authors have no conflict of interest.

References

- [1] C. Zhan, C. K. Tse, Z. Lai, T. Hao, J. Su, Prediction of covid-19 spreading profiles in south korea, italy and iran by data-driven coding, medRxiv.
- [2] S. Kim, Y. B. Seo, E. Jung, Prediction of covid-19 transmission dynamics using a mathematical model considering behavior changes in korea., Epidemiology and Health 42 (2020) 1–6.

- [3] M. Mandal, S. Jana, S. K. Nandi, A. Khatua, S. Adak, T. Kar, A model based study on the dynamics of covid-19: Prediction and control., *Chaos Solitons and Fractals* 136 (2020) 109889–109889.
- 205 [4] W. C. Roda, M. B. Varughese, D. Han, M. Y. Li, Why is it difficult to accurately predict the covid-19 epidemic?, *Infectious Disease Modelling* 5 (5) (2020) 271–281.
- [5] A. Rajesh, H. Pai, V. Roy, S. Samanta, S. Ghosh, Covid-19 prediction for india from the existing data and sir(d) model study, *medRxiv*.
- 210 [6] S. Mondal, S. Ghosh, Possibilities of exponential or sigmoid growth of covid19 data in different states of india, *medRxiv*.
- [7] J. Singh, P. K. Ahluwalia, A. Kumar, Mathematical model based covid-19 prediction in india and its different states, *medRxiv*.
- [8] Y. Zhang, C. You, Z. Cai, J. Sun, W. Hu, X.-H. Zhou, Prediction of the covid-19 outbreak based on a realistic stochastic model, *medRxiv*.
- 215 [9] A. J. Kucharski, T. W. Russell, C. Diamond, Y. Liu, J. Edmunds, S. Funk, R. M. Eggo, F. Sun, M. Jit, J. D. Munday, N. Davies, A. Gimma, K. van Zandvoort, H. Gibbs, J. Hellewell, C. I. Jarvis, S. Clifford, B. J. Quilty, N. I. Bosse, S. Abbott, P. Klepac, S. Flasche, Early dynamics of transmission and control of covid-19: a mathematical modelling study., *Lancet Infectious Diseases* 20 (5) (2020) 553–558.
- 220 [10] L. Peng, W. Yang, D. Zhang, C. Zhuge, L. Hong, Epidemic analysis of covid-19 in china by dynamical modeling., *arXiv preprint arXiv:2002.06563*.
- [11] B. Zareie, A. Roshani, M. A. Mansournia, M. A. Rasouli, G. Moradi, A model for covid-19 prediction in iran based on china parameters., *Archives of Iranian Medicine* 23 (4) (2020) 244–248.
- 225 [12] J. D. Annan, J. C. Hargreaves, Model calibration, nowcasting, and operational prediction of the covid-19 pandemic, *medRxiv*.

- [13] G. Barmparis, G. Tsironis, Estimating the infection horizon of covid-19 in
230 eight countries with a data-driven approach, *Chaos, Solitons and Fractals*
(2020) 109842.
- [14] S. Sanchez-Caballero, M. A. Selles, M. A. Peydro, E. Perez-Bernabeu, An
efficient covid-19 prediction model validated with the cases of china, italy
and spain: Total or partial lockdowns?, *Journal of Clinical Medicine* 9 (5)
235 (2020) 1547–1547.
- [15] A. B. Younes, Z. Hasan, Covid-19: Modeling, prediction, and control, *Ap-
plied Sciences* 10 (11) (2020) 3666.
- [16] M. C. Sabat, S. A. Muoz, E. lvarez Lacalle, D. L. Codina, P. J. C. Iglesias,
C. P. Soler, Empiric model for short-time prediction of covid-19 spreading,
240 *PLOS Computational Biology* (2020) 1–14.
- [17] S. Gupta, R. Shankar, Estimating the number of covid-19 infections in
indian hot-spots using fatality data, *arXiv preprint arXiv:2004.04025*.
- [18] D. K. K. Singh, S. Kumar, P. Dixit, D. M. K. Bajpai, Kalman filter based
short term prediction model for covid-19 spread, *medRxiv*.
- 245 [19] J.-P. Quadrat, 1-c nonlinear covid-19 epidemic model and application to
the epidemic prediction in france, *medRxiv*.
- [20] A. Tomar, N. Gupta, Prediction for the spread of covid-19 in india and
effectiveness of preventive measures, *Science of The Total Environment*
(2020) 138762.
- 250 [21] S. Singh, K. S. Parmar, J. Kumar, S. J. S. Makkhan, Development of
new hybrid model of discrete wavelet decomposition and autoregressive
integrated moving average (arima) models in application to one month
forecast the casualties cases of covid-19., *Chaos Solitons and Fractals* 135
(2020) 109866.

- 255 [22] S. I. Alzahrani, I. A. Aljamaan, E. A. Al-Fakih, Forecasting the spread of the covid-19 pandemic in saudi arabia using arima prediction model under current public health interventions., Journal of Infection and Public Health.