The adequacy of health system measures in reducing vulnerability to COVID-19 among the health care providers working in primary health care in Rajasthan, India: A Cross-sectional Study.

Arup Kumar Das^a, Ambey Kumar Srivastava^a, Saswata Ghosh^b, Ruchi Bhargava^a, Rajan Gupta^a and Rajesh Ranjan Singh^a

Abstract

Background

This paper examines the role of individual, facility and system level preparedness in reducing the physiological and psychological vulnerability among primary-level health care providers (HCPs) of COVID19 pandemic in Rajasthan, India.

Method and Material

Online and telephonic interviews are conducted among 274 HCPs working in 24 PHCs (17 rural and 7 urban), across 13 districts of Rajasthan. Five dimensions of vulnerability covering awareness, exposure to infection (daily contact; contact with high-risk individuals), physical and mental health conditions, while three aspects of preparedness – at individual (personal care) and facility (provider safety; management and supervision) level – are measured by employing factor analysis. Generalized ordered logit regression model is used to measure the effect of preparedness on COVID19 related vulnerability.

Result:

Among the 274 HCPs, majority of the staff are from rural PHCs (76 %), less than 35 years (87%), female (57%) and married (57 %). Almost half have high level exposure to COVID19, with mean contact rate is 90. Overall, 26% have comprehensive knowledge on COVID19, and 32% have any mental health issues. Although more than 70% of HCPs have reported more than one individual level preparedness, mental health measures adopted by the HCPs are comparably low. The facility level preparedness for enhancing safety are high such as social distance (79%) and maintaining record of each visitor (75%). However, management related measures adopted by the PHCs are perceived to be lower than the safety

^a Lords Education and Health Society (LEHS); Building No. 24 (3rd Floor), Okhla Industrial Area, Phase-III, New Delhi-110020, India

^b Institute of Development Studies Kolkata (IDSK), Kolkata, India

measures. The regression analyses suggest that safety related preparedness is significantly associated with reduction of vulnerability by 50%. The management-level preparedness has statistically no significant effect in explaining the variations in level of vulnerability.

Conclusion:

The facility-level safety measures, which lowers chances of acquiring infection has a positive effect on reducing vulnerability of COVID19. However, the HCPs do not have adequate preparedness at individual, facility management (PHC) and system level to reduce COVID19 vulnerability. Findings suggest that there is a need for a non-conventional approach of monitoring and supervision, in the absence of such measures there is a chance of moral injury that will make the HCPs at the primary level vulnerable to both physiologically and psychologically.

Introduction:

COVID-19 is possibly a novel zoonotic disease (SARS-COV-19), making the entire human population susceptible to infection in the absence of a vaccine [1]. The majority of the transmission happens through contact with droplets of the infected individuals, whether symptomatic or asymptomatic. The disease is considered to be contagious due to the high transmission rate including the asymptomatic phase [2]. There is also evidence of the airborne nature of the disease, particularly among HCPs working closely with COVID-19 infected persons [3]. Health care providers (HCPs) are at the forefront of the fight against the COVID-19 pandemic. Globally, HCPs are considered to be the most vulnerable group, not only in terms of infection and mortality, but also on account of hazards such as prolonged working hours, psychological distress, occupational burnout, stigma, violence [4], and possible transmission to family members [5]. Evidence from China and elsewhere suggests that HCPs face an enormous physiological and psychological vulnerability [6] and stress due to the new norms of wearing personal protective equipment (PPE), leading to hypoxia, hypoglycemia, or sudden cardiac arrest [7, 8, 9, 10]. A study conducted in China suggests that medical health workers are more vulnerable to psychosocial problems than their non-medical counterparts [11]. The study was focused on HCPs working in secondary or tertiary care, engaged directly in managing COVID-19 patients in ICUs and/or ventilation units.

The effects of this pandemic on health care providers working in primary level settings have not been explored in detail. This is of utmost importance in South Asian countries like India and in other lower-income and lower-middle-income countries (LMICs), where primary care is the first level of contact. Nevertheless, HCPs at all levels face common challenges [5], even if of different degrees of severity.

In India, recommendations have been made to engage community health workers, commonly referred as Frontline Health Workersⁱ (FLW), who are the foundation of primary health care [12], in activities related to the prevention of COVID-19. However, the vulnerability and preparedness in the wake of the pandemic faced by the primary health workers including FLWs have not been studied in detail. There is a growing realization that the PHCs have weak preparedness, leading to suboptimal patient safety and infection control measures in the context of COVID-19 [13].

Two important factors make India unique in the context of the current pandemic. First, India has the largest reliance on primary health care (PHCs and Sub-centers), especially in rural areas, which account for about 70% of the Indian population. Second, the pandemic is spreading from urban to rural areas due to the forced return migration after the implementation of the lockdown on 24th March 2020. As a result of this, it is highly likely that rural India would experience outbreaks soon, which will further burden the already overburdened primary health care system with questionable quality of care [14, 15].

The state of Rajasthan, located in the western parts of India, becomes a suitable case to understand the vulnerability and preparedness among primary healthcare providers as a consequence of the COVID-19 pandemic. As the state is 6th in terms of the number of COVID-19 cases (14,691) and the case fatality rate is 23.2 per thousand cases (as on June 21st, 2020) [16], within three months of the start of the pandemic. The state of Rajasthan accounts for 7% of the Indian population, 25% of the population lives in urban areas with a density of 200 population per square KMs and has female literacy of 52% as per the Census 2011. The state is one of the backward states, comes under Empowered Action Group States (EAG) statesⁱⁱ, and is lagging behind on several human development indicators compared to the national average (17). The dual burden of communicable and non-communicable diseases is also a major concern for

Rajasthan [18, 19]. Although there is no latest available data to understand the situation of the health system in Rajasthan the data available as of 2012-13 highlights major gap in human resource and infrastructure in primary health care facilities [20]. Though Rajasthan has made good progress in improving health outcomes in the recent past, such improvement stands far from achieving the Sustainable Development Goals (SDGs).

The Government of India, the state governments, and various other agencies have come out with many guidelines and have initiated a series of activities to prepare primary health care workers in the wake of the pandemic. These include raising awareness around COVID-19, building capacity, ensuring the availability of PPE, and motivating HCPs to prepare for the eventualities of the COVID-19 outbreak [21]. These policy measures are expected to have a positive effect on the individual, facility, and system-level preparedness so that the vulnerability of HCPs can be reduced.

While individual-level preparedness captures the individual measures adopted by the HCPs to maintain a balanced mental and physical wellbeing, facility-level preparedness captures an individual's perception about a facility's readiness in terms of management and safety aspects. Whereas, system-level preparedness is the translation of policy documents into capacity enhancement activities of the HCPs, in the form of training and orientations.

Against this backdrop, the main objectives of this paper are to understand:

- (1) the vulnerability of the health care workers and the preparedness measures adopted at the individual, facility, and system levels.
- (2) the effect of the preparedness on reducing vulnerability to the COVID-19 pandemic

Methods:

Data

A survey among HCPs was carried out between 29th April and 15th May 2020 in 24 PHCs (15 rural and 7 urban) managed by the Lords Education and Health Society (LEHS) through the public-private partnership (PPP) model. These PHCs are spread across 13 of the 33 districts of

Rajasthan that account for almost 43% of the total COVID-19 cases, as on June 21, 2020. Functionally, these PHCs are similar to the other government PHCs of Rajasthan, except that they have innovative components such as telemedicine consultants for general as well as specialized care, automatic medicine vending machines, and an additional layer of supervision. Out of the 284 staff working in these facilities, 272 (96.5%) participated in the study. We adopted a combination of the online google forms-based survey and a follow-up telephonic interview to collect the data. We have used a virtual meeting platform to train all interviewers., and randomly selected interviews were monitored for quality assurance. We also obtained ethical approval from the IRB Sigma ethical review board.

Outcome and explanatory Variables

Box 1 and Box 2 provide a detailed list of variables that were used in computing various dimensions of the outcome variable, the vulnerability; and the primary explanatory variables, which include preparedness at the individual, facility, and system-level respectively. Additionally, some background characteristics of HCPs such as respondent's age, sex, marital status, living arrangement, and years of experience were included as control variables to measure the adjusted effect of preparedness on vulnerability.

To achieve the study objectives, we explored various dimensions of vulnerabilities (Box 1). First, mixing with the high-risk population, often termed as crowding, is one of the major reasons for transmission [22]. This has been captured by using two variables, namely, the number of individuals the HCPs have contacted in the past 24 hours before the interview, and, among those contacts was there any high-risk population. The next dimension was the HCPs who have comprehensive knowledge of COVID-19. Having comprehensive knowledge helps a person with making better decisions and adopting healthy behaviors. Our definition of a person with comprehensive knowledge was when s/he had correct knowledge on four of the six knowledge items assessed during the survey. Finally, we have also included mental and physical health conditions as measures of vulnerability.

Box1: Description of the variables used for defining various dimensions of vulnerability

Dimensions	Descriptions
1	Individuals who met the 75 th quintile of the mixing distribution, which was 105 people in the past 24 hours.
2	staff who had contacted at least three high risk population group (which was 75 th quintile of the number of high-risk contacts). The High- risk population in this study is defined as those persons or patients who had influenza like symptoms, COVID19 positive person or person who had any direct contact with COVID19 positive cases, people who had any travel history of highly pandemic cities and countries, and persons who are in quarantine.
3	Individuals who had comprehensive knowledge of COVID19. The respondents were asked to list the persons who are most at the risk of acquiring infection and who are more likely to become critically ill or to die due to the infection. The respondents who were able to tell any four of the following six responses were considered to have comprehensive knowledge. Person who has high risk of acquiring infection: those who travelled to countries or regions which are affected by the pandemic, who has any family member visited countries or regions affected by the pandemic in recent past, migrant or mobile population, health workers who are providing services to COVID19 patients Person who are at risk of being critically ill: Elderly population and persons with any pre-existing chronic illness.
4	Individuals who had any pre-existing health condition that made them vulnerable to become critically ill due to the COVID19 infection. The pre-exist adverse health conditions included are of pregnancy, diabetic, hypertension and asthma.
5	Individuals who often experienced any specific mental health issue such as depression, irritation, insomnia, fear, anger, confusion and boredom.

The preparedness was measured at three levels: (a) at the individual level, it was measured in terms of the personal care measures adopted by a health care provider following the Government of India guidelines. The individual-level measures are the preventive physical and mental health practices adopted by HCPs in line with the Government of India guidelines (<https://www.mohfw.gov.in/pdf/PreventiveMeasures.pptx>); (b) at the facility level, it was measured in terms of the personal protective or safety measures and management measures put in place for a smooth functioning; (c) at the system level, it was measured through the number of orientations and training sessions conducted by the state health department based on the COVID-19 guidelines laid down by the central as well as the state government. To assess the coverage of the training programs conducted by the health department, we asked each respondent to report

the number of COVID-19 training programs they had undergone in the past 3-4 months and the topics covered in those training.

Box2: Variables included in computing individual and facility level preparedness

Individual level preparedness	Facility level preparedness (Providers' safety)
Do exercise every day	Arrange adequate number of PPE
Practice meditation	Attend patients showing ILI separately
Try to sleep well	Create separate waiting area for patients with respiratory symptoms
Do yoga every day	Advise patients not to come for routine visits to the OPD
Take more nutritious diet	Sanitize health facility everyday
Drink hot water	Ensure social distancing in patients
Take self-care and home remedies	Encourage patients to come with single/no attendant
Eat home prepared food	Disallow patients and attendants to enter the facility without masks
Take hydroxychloroquine tablets	Encourage sick employees to stay at home
Take green vegetables	Mop table/ wall with hypochlorite
	Sanitize hands of anybody entering facility
	Maintain record of every visitor to the facility
Facility level preparedness (management and supervision)	
Hold feedback sessions	
Conduct staff training	
Share tasks among staff	
Strengthened/developed a roster system	
Provide tips to avoid burnout	
Give rest breaks	
Provide food and water	
Plan every day for next day	
Keep motivating staff	
Keep reminding staff of safety	

Statistical analysis

To obtain a summary measure of vulnerability and preparedness, we used factor analysis, with principal factor, to compute the latent measures of vulnerability and preparedness (indices). The factor scores were categorized according to their tercile and classified as high, medium, and low. The first factor was chosen for constructing the indices, which explains maximum variations, with an eigenvalue greater than one. The factor scores of the first factor of vulnerability measures with an eigenvalue of 1.42 were used for computing the vulnerability index as it indicated that the factor had more information than the individual attributes. In the case of individual level preparedness index, the first factor explained 75% of the variation, and all coefficients were found to indicate a positive effect of each of the variables on the factor score. Similarly, the first factor for the two types of facility-level preparedness explained 92% and 98% of the total variation for safety and management indices respectively.

To understand the effect of various preparedness measures on vulnerability to COVID-19, we first used an ordered logit regression model. One of the main problems with such a model is the restriction of the parallel lines assumption, which is often violated. To examine the validity of such an assumption, we have used the brant test with the null hypothesis that the parallel lines assumption holds. The hypothesis was rejected with the brant test Chi2 value of 15.13, p=0.299. However, a detailed examination of the regression coefficients suggested that most of the coefficients varied greatly across regressions. Accordingly, we have used a generalized ordered logit model to allow some of the coefficients to vary, while others remained constant. We have STATA's user-written command, gologit2 to employ the model [23]. The model is explained with the following expression:

$$P(Y_i > j) = \frac{\exp(\alpha_j X_{1i} \beta_1 + X_{2i} \beta_2 + X_{2i} \beta_{3j})}{1 + \{\exp(\alpha_j X_{1i} \beta_1 + X_{2i} \beta_2 + X_{2i} \beta_{3j})\}}$$

Where $j=1, 2, \dots, M-1$

M is the number of categories in the ordinal variable (vulnerability Index) and β 's for X1 and X2 are the same for all values of J but β 's for X3 are free to differ. We have used the AIC and BIC criterion for comparing models. We have compared five variations of the model taking into

account one of the exposure variables at each time (model 1 to model 4) and finally, a full model (model 5) is created by including all exposure variables, after controlling for background characteristics of HCPs.

Results:

In this section, we elaborate on the variations in vulnerability and preparedness measures across the sub-groups of respondents. Following that, we assess the effect of preparedness on different levels of vulnerability.

Profile of the respondents

Table 1 provides the socio-demographic and professional details of the 272 staff interviewed in the study. Some of the key observations are that the majority of the staff were young, with 87% of them being less than 35 years old and 57% of them were married. The majority of the staff were working in rural areas (76%) and PHCs (77%). 92% of them were posted in a location that was outside of the district to which they belonged and 48% were living in a rented house. The majority of the staff were either ANM (39%) or Staff Nurse/GNM (20%). More than two-thirds of them (68%) had 2-5 years of work experience, while only 11% had more than 5 years of experience.

Table 1: Profile of the respondent staff working in 24 Primary Healthcare Centers (PHCs) Rajasthan, HCP survey, 2020		
	Percent (%)	Number (N)
Type of Facility (Rural/Urban)		
Rural	75.7	206
Urban	24.3	66
Facility type		
Sub-Center (SC)	22.8	62
Primary Health Centre (PHC)	77.2	210
Designation		
Medical Officer In-Charge (MOIC)	8.1	22
Pharmacist	8.1	22
Lady Health Visitor (LHV)	5.9	16
Staff Nurse (GNM)	19.5	53
Auxiliary Nurse Midwife (ANM)	38.6	105
Lab Technician (LT)	7.4	20

Multi-Purpose Worker (MPW)	12.5	34
Age category		
<=25 yrs.	23.9	65
25-30 yrs.	41.5	113
30-35 yrs.	21	57
35+	13.6	37
Sex		
Male	42.6	116
Female	57.4	156
Marital status		
Married	76.1	207
Unmarried	21.3	58
Separated/divorce/other	2.6	7
Residence (Within or outside districts)		
Outside district	91.9	250
Within district	8.1	22
Where do you live/ type of residence		
Staff quarter	23.5	64
On Rent	48.2	131
Own house	28.3	77
Years of experience		
< One yr.	21.3	58
2-5 yrs.	67.6	184
5+ yrs.	11	30
Any chronic illness		
No	94.5	257
Yes	5.5	15
Source: Authors' calculations from health care providers survey, 2020.		

Vulnerability

Table 2 describes variations in the level of vulnerability by selected background characteristics. Overall, 29% of staff belong to the highly vulnerable category with the highest among medical officers (50%) and the lowest among pharmacists (9%). Staff working in sub-centers (34%) were more likely to be highly vulnerable than those working in PHCs (26%). The proportion of staff who belong to the highly vulnerable category are those who belong to the age group of 35+ years (43%), female (30%), separated/divorced (43%), stay in staff quarters (41%). There were also substantial variations across facilities, with the percentage of staff that was highly vulnerable ranging from 9% to 66%.

Table 2: Percent distribution of respondents according to level of vulnerability by background characteristics, HCP survey, 2020					
Background Characteristics	Low	Medium	High	Total	N
Respondent classification (Rural/Urban)					
Rural	43.7	27.7	28.6	100	206
Urban	33.3	40.9	25.8	100	66
Facility type					
Sub-Center (SC)	43.5	22.6	33.9	100	62
Primary Health Centre (PHC)	40.5	33.3	26.2	100	210
Designation					
Medical Officer In-Charge (MOIC)	22.7	27.3	50	100	22
Pharmacist	63.6	27.3	9.1	100	22
Lady Health Visitor (LHV)	37.5	31.3	31.3	100	16
Staff Nurse (GNM)	45.3	32.1	22.6	100	53
Auxiliary Nurse Midwife (ANM)	34.3	32.4	33.3	100	105
Lab Technician (LT)	55	35.0	10.0	100	20
Multi-Purpose Worker (MPW)	47.1	26.5	26.5	100	34
Age					
<=25 yrs.	38.5	33.8	27.7	100	65
25-30 yrs.	44.2	31.9	23.9	100	113
30-35 yrs.	36.8	36.8	26.3	100	57
35+	43.2	13.5	43.2	100	37
Sex					
Male	47.4	28.4	24.1	100	116
Female	36.5	32.7	30.8	100	156
Marital status					
Married	43.5	27.5	29	100	207
Unmarried	36.2	41.4	22.4	100	58
Separated/divorce/other	14.3	42.9	42.9	100	7
Residence (Within or outside districts)					
Outside district	41.6	30.8	27.6	100	250
Within district	36.4	31.8	31.8	100	22
Where do you live/ type of residence					
Staff quarter	31.3	28.1	40.6	100	64
On rent	46.6	35.1	18.3	100	131
Own house	40.3	26.1	33.8	100	77
Years of experience					
< One yr.	37.9	37.9	24.1	100	58
2-5 yrs.	40.8	29.9	29.3	100	184
5+ yrs.	50	23.3	26.7	100	30
Total	41.2	30.9	27.9	100	272

N	112	84	76	272
Source: Authors' calculations from health care providers survey, 2020.				

Preparedness:

Variations at the individual, facility, and system-level preparedness have been presented in Table 3 and the salient findings are as follows.

Individual-level preparedness

It is evident from Table 3, that overall, 23% of the staff belongs to the category of highly prepared at the individual level. Individual-level preparedness was the highest among the medical officers (46%) and the lowest among the Multipurpose Health Workers (MPW) (9%). It is interesting to note that a higher proportion of staff with less than 5 years of experience was highly prepared (28%) than those who had more than 5 years of experience (13%). Likewise, a higher proportion of staff belonging to the sub-centers was highly prepared (29%) than those who were posted in PHCs. Female (28%), unmarried (31%), and staff from outside districts (24%), living in staff quarters (28%), among others, have a high level of individual preparedness than their counterparts.

Facility level preparedness

Both measures of facility-level preparedness vary across the profile of the respondents. A higher proportion of respondents from the Urban PHCs (56%) than the rural (26%), and PHCs (38%) than SCs (18%) are highly prepared in terms of management-related preparedness. A higher proportion of medical officers (59%) reported a high level of management preparedness than other staff. The safety-related preparedness did not show such substantial variations across the profile of the respondents (Table 3). We have also examined these measures of preparedness across facilities and found that the proportion of respondents who reported a high level of preparedness varies from 8% to 68% for management related preparedness, and from null to 64% for safety-related preparedness (not shown in the table). The correlation between the two measures of facility-level preparedness is not significant, the correlation coefficient being 0.43.

Table 3: Individual and facility level preparedness by background characteristics, HCP survey, 2020

Background Characteristics	Individual level preparedness			Facility level preparedness (Management)			Facility level preparedness (Safety)			System level	N
	Low	Medium	High	Low	Medium	High	Low	Medium	High	% trained	
Respondent classification (Rural/Urban)											
Rural	37.4	39.3	23.3	37.9	36.4	25.7	36.9	33.5	29.6	60.2	206
Urban	22.7	54.5	22.7	19.7	24.2	56.1	22.7	43.9	33.3	74.2	66
Facility type											
Sub-Center (SC)	32.3	38.7	29.0	43.5	38.7	17.7	50.0	21.0	29.0	71.0	62
Primary Health Centre (PHC)	34.3	44.3	21.4	30.5	31.9	37.6	28.6	40.5	31.0	61.4	210
Designation											
Medical Officer In-Charge (MOIC)	36.4	18.2	45.5	4.5	36.4	59.1	9.1	59.1	31.8	72.7	22
Pharmacist	50.0	50.0	0.0	54.5	31.8	13.6	45.5	27.3	27.3	36.4	22
Lady Health Visitor (LHV)	31.3	56.3	12.5	18.8	37.5	43.8	25.0	43.8	31.3	93.7	16
Staff Nurse (GNM)	34.0	37.7	28.3	30.2	35.8	34.0	20.8	47.2	32.1	60.4	53
Auxiliary Nurse Midwife (ANM)	26.7	45.7	27.6	40.0	27.6	32.4	41.9	29.5	28.6	77.1	105
Lab Technician (LT)	50.0	30.0	20.0	25.0	55.0	20.0	40.0	30.0	30.0	35.0	20
Multi-Purpose Worker (MPW)	35.3	55.9	8.8	35.3	32.4	32.4	35.3	29.4	35.3	41.2	34
Age											
<=25 yrs.	36.9	43.1	20.0	40.0	26.2	33.8	32.3	27.7	40.0	63.1	65
25-30 yrs.	25.7	46.0	28.3	28.3	43.4	28.3	33.6	42.5	23.9	62.8	113
30-35 yrs.	35.1	43.9	21.1	33.3	31.6	35.1	31.6	36.8	31.6	66.7	57
35+	51.4	32.4	16.2	37.8	18.9	43.2	37.8	29.7	32.4	62.2	37
Sex											
Male	43.1	39.7	17.2	26.7	37.9	35.3	27.6	38.8	33.6	46.6	116
Female	26.9	45.5	27.6	38.5	30.1	31.4	37.8	34.0	28.2	76.3	156
Marital status											
Married	33.3	45.4	21.3	34.3	33.8	31.9	34.8	36.2	29.0	60.9	207
Unmarried	37.9	31.0	31.0	34.5	29.3	36.2	29.3	34.5	36.2	72.4	58

Separated/divorce/other	14.3	71.4	14.3	0.0	57.1	42.9	28.6	42.9	28.6	71.4	7
Residence (Within or outside districts)											
Outside district	33.2	43.2	23.6	32.4	34.0	33.6	33.6	35.6	30.8	62.8	250
Within district	40.9	40.9	18.2	45.5	27.3	27.3	31.8	40.9	27.3	72.7	22
Where do you live/ type of residence											
Staff quarter	37.5	34.4	28.1	39.1	34.4	26.6	32.8	39.1	28.1	70.3	64
On rent	32.8	40.5	26.7	32.8	28.2	38.9	35.1	30.5	34.4	71.0	131
Own house	32.5	54.5	13.0	29.9	41.6	28.6	31.2	42.9	26.0	45.5	77
<i>Years of experience</i>											
< One yr.	24.1	48.3	27.6	34.5	31.0	34.5	25.9	37.9	36.2	65.5	58
2-5 yrs.	32.6	44.0	23.4	33.7	32.1	34.2	34.8	33.2	32.1	64.1	184
5+ yrs.	60.0	26.7	13.3	30.0	46.7	23.3	40.0	50.0	10.0	56.7	30
Total	33.8	43.0	23.2	33.5	33.5	33.1	33.5	36.0	30.5	63.6	272
N	92	117	63	91	91	90	91	98	83	272	

Source: Authors' calculations from health care providers survey, 2020.

System-level preparedness

We found that 63% of the staff had received some training or orientation on a COVID-19 topic, with the average being 1.18 training sessions per staff member. There were substantial variations in the proportion of staff that had received any training (Table 3). A higher proportion of urban staff (74%), sub-center staff (71%), Lady Health Visitors (LHV) (94%), and those who had less than one year of work experience (66%) had received at least one training on COVID-19 related issues. It is important to note that a large proportion of pharmacists (64%), lab-technicians (65%), and MPWs (59%) did not receive any training.

Role of preparedness on reducing vulnerability:

The descriptive analysis suggests that there was no significant association between individual preparedness and level of vulnerability. Preparedness of facilities in terms of management and supervision was also not associated with vulnerability. However, a significant association was observed between the preparedness of facilities in terms of safety measures and the level of vulnerability (Table 4). The proportion of staff who were highly vulnerable reduced monotonically from 35% to 21% as the safety-related preparedness increased from low to high ($\chi^2=9.67$, $p<0.05$). Similarly, as the number of training sessions increased from ‘no training’ to 3+ training sessions, the proportion of staff who were highly vulnerable reduced from 29% to 22%. However, the association was not statistically significant.

Table 4: Association between the various level of preparedness and vulnerability, HCP survey, 2020					
Preparedness	Low	Medium	High	Total	N
<i>Individual preparedness</i>					
Low	43.5	28.3	28.3	100	92
Medium	42.7	33.3	23.9	100	117
High	34.9	33.3	31.7	100	63
<i>Pearson $\chi^2(4) = 2.2576$ Pr = 0.688</i>					
<i>Facility preparedness (Safety-Infection prevention)</i>					
Low	35.2	29.7	35.2	100	91
Medium	36.7	37.8	25.5	100	98
High	53.0	26.5	20.5	100	83
<i>Pearson $\chi^2(4) = 9.6562$ Pr = 0.047</i>					
<i>Facility preparedness (management)</i>					
Low	39.6	35.2	25.3	100	91

Medium	45.1	30.8	24.2	100	91
High	38.9	28.9	32.2	100	90
<i>Pearson chi2(4) = 2.3636 Pr = 0.669</i>					
<i>No. of training attended on COVID-19</i>					
None	43.4	27.3	29.3	100	99
One	40.2	32.9	26.8	100	82
Two	34.5	38.2	27.3	100	55
Three+	47.2	30.6	22.2	100	36
<i>Pearson chi2(6) = 2.9362 Pr = 0.817</i>					
Total	41.2	31.6	27.2	100	272
N	112	86	74	272	
Source: Authors' calculations from health care providers survey, 2020.					

The adjusted odds ratios (AORs) of the generalized ordered logit regression – after adjusting for background characteristics including age, sex, marital status, designation, place of work (SC or PHC), facility location (rural or urban), years of experience, living arrangement, and whether they belonged to the same district where the facility was located or to some other district have been shown in Table 5. The results suggest that facility-level safety measures were highly significant in explaining the variations in the level of vulnerability. The comparison of the five models suggests that the M2 is the best fit with the lowest AIC and BIC values, 582.004 and 668.543, respectively. We have also observed from the model that facility-level safety preparedness has a positive and significant effect on reducing the COVID-19 vulnerability of HCPs working in the primary health setting. Those who belong to the medium and high level of preparedness, in terms of safety measures, are less likely to be highly vulnerable (AOR: 0.54; 95%CI: 0.30-0.97) and (AOR:0.36; 95%CI: 0.19-0.66) respectively, than the reference category. The second-best model, full model (M5), also substantiates the same findings. None of the other exposure variables are statistically significant. Among the background characteristics, the staff designation is found to be significant and it is found that medical officers are more likely to be highly vulnerable, followed by ANM and LHV with a predicted probability of being highly vulnerable as 0.45; 95%CI:0.24-0.73, 0.37; 95%CI:0.24-0.51 and 0.28; 95%CI: 0.08-0.48, respectively (Figure 1)

Figure 1: Predicted probability of low, medium and high level of vulnerability by staff designation

Source: Authors calculation from the health care providers survey

	M1			M2			M3			M4			M5		
Vulnerability index	AOR	[95% Conf.	Interv al]	AOR	[95% Conf.	Interv al]	AOR	[95% Conf.	Interv al]	AOR	[95% Conf.	Interv al]	AOR	[95% Conf.	Interv al]
Designation (ref.: Medical officer)	1.00			1.00			1.00			1.00			1.00		
pharmacist	0.12	0.03	0.42	0.09**	0.03	0.32	0.11**	0.03	0.40	0.10**	0.03	0.36	0.10**	0.02	0.37
LHV	0.37	0.08	1.74	0.38	0.08	1.74	0.37	0.08	1.67	0.39	0.09	1.77	0.42	0.09	2.03
Staff N (GNM)	0.30*	0.10	0.90	0.29*	0.10	0.86	0.29*	0.10	0.87	0.27*	0.09	0.81	0.29*	0.09	0.91
ANM	0.72	0.20	2.62	0.65	0.18	2.40	0.67	0.18	2.44	0.66	0.18	2.42	0.69	0.18	2.64
Lab Technician	0.18*	0.05	0.63	0.15**	0.04	0.53	0.18*	0.05	0.64	0.15**	0.04	0.54	0.14**	0.04	0.53
MPW	0.26*	0.08	0.83	0.20*	0.06	0.66	0.24	0.08	0.78	0.23*	0.07	0.75	0.20*	0.06	0.68
Preparedness- Individual Level (ref.: low)	1.00			1.00			1.00			1.00			1.00		
medium	1.00	0.56	1.81										0.96	0.52	1.77
high	1.11	0.61	2.01										1.03	0.56	1.90
Preparedness- Facility level safety (ref.: low)	1.00			1.00			1.00			1.00			1.00		
medium				0.54*	0.30	0.97							0.47*	0.25	0.89
high				0.36**	0.19	0.66							0.25**	0.12	0.53
Preparedness- Facility level management (ref.: low)							1.00						1.00		
medium							0.75	0.41	1.37				1.19	0.62	2.29
high							0.90	0.49	1.65				2.13	0.98	4.61
No. of training sessions (ref.: None)										1.00			1.00		
One													0.76	0.41	1.40

Two				0.79	0.43	1.45	0.65	0.31	1.34
Three+				0.75	0.38	1.51	0.50	0.21	1.20
Model comparison									
II (null)	-	-	-	-			-		
	294.7	294.7	294.7	294.7			294.7		
	32	32	32	32			32		
II(model)	-	-	-	-			-		
	272.5	267.0	272.1	270.9			263.8		
	05	02	29	99			23		
AIC	593.0	582.0	592.2	591.9			589.6		
	09	04	57	97			46		
BIC	679.5	668.5	678.7	682.1			701.4		
	49	43	96	42			25		

AOR: Adjusted odds ratio

*p < 0.05; **p < 0.01; ***p < 0.001.; CI: Confidence Intervals

Source: Authors' calculations from health care providers survey, 2020.

Discussion:

The present paper highlights the level of vulnerability of HCPs in primary health care vis-à-vis the strength of the support structure to protect their rights. It is emerging that HCPs need to be well-prepared to handle any eventualities which could emerge from the localized outbreak of COVID-19 in the PHC areas by adopting a rights-based approach, as highlighted by the World Health Organisation (WHO) [4, 24]. The facility-level measures directly reduce their chance of acquiring an infection while performing their roles. Inadequate supply of basic equipment in PHCs was a cause for concern across India and in other LMICs even before the COVID-19 pandemic [14].

However, the present study suggests that the HCPs do not have adequate preparedness at the individual level nor the health facility (management mechanisms), and system levels. In the absence of such a support structure, the HCPs are likely to experience more physical and psychological distress. One plausible reason why managerial mechanism at the facility level does not reduce vulnerability could be that there is a need for a non-conventional approach to increasing staff motivation. In the absence of such an approach, there is a possibility of moral injury [25] during the crisis. When a crisis begins, managers should be frank, check on the wellbeing of the staff, provide regular contact, and focus on active monitoring and supervision. By contrast, the data reveals that measures to reduce burnout and to provide personal care to the staff are substantially low. To safeguard the rights, roles, and responsibilities of the HCPs, including sustained social distancing and other safe working environment norms [4], there is a need for redesigning the functioning of the PHCs in terms of infrastructure and human resources and for leveraging innovative technologies like digital health and telemedicine, [26] non-touchable diagnostics, and automated medicine dispenser mechanism, etc. At present, India is facing challenges of efficient supply chain and inadequate stock of PPE [27].

Another important observation is the variations across the 24 facilities is substantial in terms of the vulnerability and preparedness indicators, indicating that there is a need for more localized and micro-planning for a better outcome. Considering the similar health system performance across the EAG states, the findings from the study plausibly can also be used for designing COVID-19 responses in other states.

Conclusion

As the post COVID-19 era will demand new norms in human society, the service delivery mechanisms will need to be intertwined in the same fashion. In this respect, the study identifies three major areas of change requirements. First, institutionalized systems and mechanisms for highly motivated staff through the alternative method of management and supervision, to prevent them from moral injury. Second, the learning acquired from digital health and telemedicine innovationsⁱⁱⁱ [28, 29] will have a role in maintaining social distance, efficient triaging, referral, and follow-up. Third, it is time to go beyond the dichotomous approach of horizontal and vertical integration and to learn from the experience of the diagonal approach [30] to the health system. All changes have to be customized taking into account concerns of health care providers to prepare the health system to be better equipped for any future crisis.

ⁱ The front-line health workers includes the Auxiliary Nurse Midwife (ANM) who are posted at the sub-health center level and responsible for 5000 population, while the Accredited Social Health Activist (ASHA) are village-level volunteers and interphase between community and health system. Usually, there is one ASHA per every 1000 population, as per the norm.

ⁱⁱ There are eight socioeconomically backward states of Bihar, Chhattisgarh, Jharkhand, Madhya Pradesh, Orissa, Rajasthan, Uttaranchal, and Uttar Pradesh, referred to as the Empowered Action Group (EAG) states, lag behind on many of the health and demographic indicators. The maternal mortality ratio is 186 against the India average of 122 and far below from states like Kerala (42) and Tamil Nadu (63). Similarly, the IMR is one of the highest in the country i.e., 38 per thousand live birth, while the same for India is 33 and few other better performing states such as Kerala (10) and Tamil Nadu (16) [Source: Authors compilation from various SRS bulletin]

ⁱⁱⁱ The digital health and telemedicine innovations is one of the successful initiatives in Rajasthan and Delhi implemented WISH [28]. The model is now scaled in Madhya Pradesh in 51 districts, where call centers were initially used for contact tracing, quarantine, and referral management. Similarly, E-Sanjeevani (<https://esanjeevaniopd.in/About>) has become the attraction of general out-patient care services. Such initiatives should be enhanced along with the enhancement of skills and competencies of HCPs in the primary health care sector.

References

- [1] Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. Journal of autoimmunity. 2020 Feb 26:102433. <https://doi.org/10.1016/j.jaut.2020.102433>

- [2] World Health Organization. Coronavirus disease 2019 (COVID-19): situation report, 73. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200402-sitrep-73-covid-19.pdf?sfvrsn=5ae25bc7_2 [Last accessed June 2020].
- [3] Bahl P, Doolan C, de Silva C, Chughtai AA, Bourouiba L, MacIntyre CR. Airborne or droplet precautions for health workers treating COVID-19? The Journal of infectious diseases. 2020 Apr 16. <https://doi.org/10.1093/infdis/jiaa189>
- [4] World Health Organization. Coronavirus disease (COVID-19) outbreak: rights, roles and responsibilities of health workers, including key considerations for occupational safety and health: interim guidance, 19 March 2020. World Health Organization. World Health Organization. <https://apps.who.int/iris/bitstream/handle/10665/331510/WHO-2019-nCov-HCWadvice-2020.2-eng.pdf?sequence=1&isAllowed=y> [Last accessed May 2020].
- [5] Adams JG, Walls RM. Supporting the health care workforce during the COVID-19 global epidemic. Jama. 2020 Apr 21;323(15):1439-40. <https://doi.org/10.1001/jama.2020.3972>
- [6] Li W, Zhang J, Xiao S, Sun L. Characteristics of health worker fatality in China during the outbreak of COVID-19 infection. J Infect. 2020 Jul;81(1):147-178. <https://doi.org/10.1016/j.jinf.2020.03.030>.
- [7] Wu P, Fang Y, Guan Z, Fan B, Kong J, Yao Z, Liu X, Fuller CJ, Susser E, Lu J, Hoven CW. The psychological impact of the SARS epidemic on hospital employees in China: exposure, risk perception, and altruistic acceptance of risk. The Canadian Journal of Psychiatry. 2009 May;54(5):302-11. <https://doi.org/10.1177/070674370905400504>
- [8] Zeng Y, Zhen Y. RETRACTED: Chinese medical staff request international medical assistance in fighting against COVID-19. The Lancet Global health. 2020 Feb 24. [https://doi.org/10.1016/S2214-109X\(20\)30065-6](https://doi.org/10.1016/S2214-109X(20)30065-6)
- [9] Calefi AS, de Queiroz Nunes CA, da Silva Fonseca JG, Quinteiro-Filho WM, Ferreira AJ, Palermo-Neto J. Heat stress reduces Eimeria spp. infection and interferes with C. perfringens infection via activation of the hypothalamic-pituitary-adrenal axis. Research in veterinary science. 2019 Apr 1;123:273-80. <https://doi.org/10.1016/j.rvsc.2019.01.026>
- [10] Scorza FA, Albuquerque RD, Arida RM, Albuquerque MD, Terra VC, Machado HR, Cysneiros RM, Scorza CA, Cavalheiro EA. What are the similarities between stress, sudden cardiac death in Gallus gallus and sudden unexpected death in people with epilepsy. Arquivos de neuro-psiquiatria. 2010 Oct;68(5):788-90. <https://doi.org/10.1590/s0004-282x2010000500022>
- [11] Zhang WR, Wang K, Yin L, Zhao WF, Xue Q, Peng M, Min BQ, Tian Q, Leng HX, Du JL, Chang H. Mental health and psychosocial problems of medical health workers during the

COVID-19 epidemic in China. *Psychotherapy and psychosomatics*. 2020 Apr 9:1-9.
<https://doi.org/10.1159/000507639>

[12] Haines A, de Barros EF, Berlin A, Heymann DL, Harris MJ. National UK programme of community health workers for COVID-19 response. *The Lancet*. 2020 Apr 11;395(10231):1173-5. [https://doi.org/10.1016/S0140-6736\(20\)30735-2](https://doi.org/10.1016/S0140-6736(20)30735-2)

[13] Garg S, Basu S, Rustagi R, Borle A. Primary Health Care Facility Preparedness for Outpatient Service Provision During the COVID-19 Pandemic in India: Cross-Sectional Study. *JMIR Public Health and Surveillance*. 2020;6(2):e19927. <https://doi.org/10.2196/19927>

[14] Khan PK, Banerjee K, Nandi S. Primary Healthcare Infrastructure and Reproductive Healthcare in Rural India: A District Level Analysis. In *The Demographic and Development Divide in India 2019* (pp. 417-466). Springer, Singapore. https://doi.org/10.1007/978-981-13-5820-3_8

[15] Mohan P, Kumar R. Strengthening primary care in rural India: Lessons from Indian and global evidence and experience. *J Family Med Prim Care*. 2019;8(7):2169-2172.
https://doi.org/10.4103/jfmpe.jfmpe_426_19

[16] Accessed from <https://www.covid19india.org/>, accessed on June 21st, 2020

[17] Accessed from https://censusindia.gov.in/vital_statistics/SRS_Bulletins/Bulletins.html, accessed on June 21st, 2020

[18] India State-Level Disease Burden Initiative Collaborators. Nations within a nation: variations in epidemiological transition across the states of India, 1990-2016 in the Global Burden of Disease Study [published correction appears in *Lancet*. 2017 Dec 2;390(10111):e49]. *Lancet*. 2017;390(10111):2437-2460. doi:10.1016/S0140-6736(17)32804-0

[19] Mohan P, Mohan SB, Dutta M. Communicable or noncommunicable diseases? Building strong primary health care systems to address double burden of disease in India. *J Family Med Prim Care*. 2019;8(2):326-329. doi:10.4103/jfmpe.jfmpe_67_19

[20] International Institute for Population Sciences (IIPS), 2014. *District Level Household and Facility Survey (DLHS-4), 2012-13:India. Rajasthan*: Mumbai: IIPS

[21] Accessed from <https://www.mohfw.gov.in/pdf/GuidelinesonpreventivemeasurestocontainspreadofCOVID19inworkplacesettings.pdf>, accessed on June 14th, 2020.

-
- [22] Rader B, Scarpino S, Nande A, Hill A, Reiner R, Pigott D, Gutierrez B, Shrestha M, Brownstein J, Castro M, Tian H. Crowding and the epidemic intensity of COVID-19 transmission. medRxiv 2020.04.15.20064980. <https://doi.org/10.1101/2020.04.15.20064980>
- [23] Richard Williams, Generalized ordered logit/partial proportional odds models for ordinal dependent variables, The Stata Journal (2006), Vol 6, Number 1, pp. 58–82, <https://doi.org/10.1177/1536867X0600600104>
- [24] World Health Organization. COVID-19: operational guidance for maintaining essential health services during an outbreak: interim guidance, 25 March 2020. World Health Organization; 2020. <https://apps.who.int/iris/handle/10665/331561> [Last accessed June 2020].
- [25] Greenberg N, Docherty M, Gnanapragasam S, Wessely S. Managing mental health challenges faced by healthcare workers during covid-19 pandemic. bmj. 2020 Mar 26;368. <https://doi.org/10.1136/bmj.m1211>
- [26] Nachiket Mor, Resources for Primary Care Providers to Meet Patients Needs During the COVID-19 Epidemic. The Banyan Academy of Leadership in Mental Health, Thiruvadanthai, Tamil Nadu, India, 2020, May 6. https://www.researchgate.net/publication/340095908_Resources_for_Primary_Care_Providers_to_Meet_Patient_Needs_During_the_COVID-19_Epidemic [Last accessed June 2020].
- [27] News article: Accessed from <https://www.forbesindia.com/article/coronavirus/indias-ppe-crisis-puts-workers-in-the-line-of-fire/59073/1>, accessed on May 23rd, 2020
- [28] News article: Accessed from <https://ehealth.eletsonline.com/2018/11/empowering-healthcare-through-technology-and-innovations/>, accessed on 23rd May 2020
- [29] News article: Accessed from <https://www.thehindubusinessline.com/specials/india-interior/doc-youre-wanted-online-for-consultation/article31121370.ece>, accessed on 29th May 2020
- [30] Frenk J, Gómez-Dantés O, Knaul F M. The Health Systems Agenda: Prospects for the Diagonal Approach. In The Handbook of Global Health Policy (eds G.W. Brown, G. Yamey and S. Wamala). 2014, pp. 425-439 <https://doi.org/10.1002/9781118509623.ch23>