The impact of climate temperature on counts, recovery, and death rates due to SARS-CoV-2 in South Africa.

Naven Chetty^{1*}, Bamise Adeleye² & Abiola Olawale Ilori³

^{1,2,3}School of Chemistry and Physics, University of KwaZulu–Natal, Private Bag X01, Scottsville, 3209, South Africa

*Corresponding author: chettyn3@ukzn.ac.za

ABSTRACT

The impact of climate temperature on the counts (number of positive COVID-19 cases reported), recovery, and death rates of COVID-19 cases in South Africa's nine provinces was investigated. The data for confirmed cases of COVID-19 were collected for March 25 and June 30, 2020 (14 weeks) from South Africa's Government COVID-19 online resource, while the daily provincial climate temperatures were collected from the website of the South African Weather Service. Our result indicates that a higher or lower climate temperature does not prevent or delay the spread and death rates but shows significant positive impacts on the recovery rates of COVID-19 patients. Thus, it indicates that the climate temperature is unlikely to impose a strict limit on the spread of COVID-19. There is no correlation between the cases and death rates, an indicator that no particular temperature range is closely associated with a faster or slower death rate of COVID-19 patients. As evidence from our study, a warm climate temperature can only increase the recovery rate of COVID-19 patients, ultimately impacting the death and active case rates and freeing up resources quicker to enable health facilities to deal with those patients' climbing rates who need treatment.

Keywords: climate temperature, COVID-19, rates, SARS-CoV-2, South Africa.

1. Introduction

The roles of climate temperature on transmitting the novel Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV-2) and patients' recovery and death rates seemed to have generated significant worldwide debates. Several kinds of research indicated the spreads of other seasonal flu virus outbreaks by reporting a substantial number of cases with colder climate conditions similar to the spreads of SARS-CoV-2 [1, 2]. In contrast, others found an inverse or no correlation between climate temperature variations and the Coronavirus [3–5].

SARS-CoV-2, popularly called Coronavirus (COVID-19), envelopes positive-stranded RNA viruses with a characteristic surface [6, 7]. It has been reported that the infection of the virus in humans causes severe gastroenteritis and respiratory tract diseases [6, 8]. These viruses' persistence was officially lost at around 56°C but somewhat differed depending on the infection [9]. In their study of SARS-CoV-2 aerosol and surface stability, Doremalen *et al.* [10] revealed that the virus could survive on hard surfaces for up to 72 hours at temperatures between 21°C and 23°C. Similarly, Zhou *et al.* [11] found the coronavirus to be most active between 9°C and 24°C, decreasing with higher temperatures. The fact that the virus causing the SARS-CoV-2 is new still limits information about how it impacted by variations in climatic temperature conditions. Peng *et al.* [2] studied the impacts of temperature and absolute humidity

on COVID-19 in 31 provincial regions of mainland China. They observed a decline in the transmission rate of the COVID-19 as temperature increases. A recent study examining the impact of temperature variation and humidity on COVID-19 mortality in Wuhan China found that mortality rates were lower for days with higher temperatures and humidity. The study also indicates high death rates on days with a broader maximum and minimum temperature range [12].

Therefore, this study aims to investigate the impact of climate temperature variation on the counts, recovery, and death rates of COVID-19 cases in all South Africa's provinces. The findings were compared with those of countries with comparable climate temperature values.

Fig 1: Map of South Africa showing the provinces [13].

2. Methods

2.1 Data Collection

The data for confirmed cases of COVID-19 were collected for March 25 and June 30 (14 weeks) for South African provinces, including daily counts, death, and recovery rates. The dates were grouped into two, wherein weeks 1-5 represent the periods of total lockdown to contain the spread of COVID-19 in South Africa. Weeks 6-14 are periods where the lockdown was eased to various levels 4 and 3. The daily information of COVID-19 count, death, and recovery was obtained from South Africa's Government COVID-19 online resource (<https://sacoronavirus.co.za>). Daily provincial climate temperatures were collected from the website of the South African Weather Service (<https://www.weathersa.co.za>). The provinces of South Africa are Eastern Cape, Western Cape, Northern Cape, Limpopo, Northwest, Mpumalanga, Free State, KwaZulu-Natal, Western Cape, and Gauteng. Weekly consideration was given to the daily climate temperature (average minimum and maximum). The recorded values were considered, respectively, to be in the ratio of death-to-count (D/C) and recovery-to-count (R/C).

2.2 Statistical Analysis

Descriptive statistics were performed for all the data collected for this study. The analyses were performed using the Person's bivariate correlation to analyze the association between climate temperature, death-to-count, and recovery-to-count ratios of COVID-19. The death and recovery rates were first accommodated to check the ratio of weekly COVID-19 counts to death/recovery rates. Then,

the climate temperature was introduced into the equation to verify the correlation between the parameters of interests. Equations 1 and 2 used are given as a pair of variables (C , D/C , and R/C) [14-16].

$$\rho_{(C, D/C)} = \text{cov}(C, D/C) \cdot (\sigma_C \cdot \sigma_{D/C})^{-1} \quad (1)$$

$$\rho_{(C, R/C)} = \text{cov}(C, R/C) \cdot (\sigma_C \cdot \sigma_{R/C})^{-1} \quad (2)$$

Where $\rho_{(C, D/C)}$ is the correlation between counts and death-to-count ratio; $\rho_{(C, R/C)}$ is the correlation between counts and recovery-to-count ratio; cov is the covariance of the count and the interested ratios; $\sigma_C, \sigma_{D/C}, \sigma_{R/C}$ are the standard deviation of $C, D/C$, and R/C respectively. The correlation curves between the parameters were fitted to get the coefficient of determination (R^2). The analysis was performed using OriginPro 8.5.0 SR1.

3. Results and Discussion

3.1 Description of COVID-19 counts, death and recovery rates within the study period

The average climate temperature within the study period for Eastern Cape Province ranges from 10.2 to 29.6 °C with 0.0295 (2.95 %) death-to-count ratio and 0.0207 (2.07 %) recovery-to-count ratio for the lockdown periods and average climate temperature of 6.8 to 29.4 °C with 0.0204 (2.04 %) death-to-count ratio and 0.0683 (6.83 %) recovery-to-count ratio for the eased-lockdown periods, respectively. The average climate temperature within the study period for Northern Cape Province ranges from 9.2 to 33.8 °C with 0.1000 (10 %) death-to-count ratio and 1.400 (140 %) recovery-to-count ratio for the lockdown periods and average climate temperature of 1.7 to 28.3 °C with 0.0000 (0 %) death-to-count ratio and 0.5732 (57.32 %) recovery-to-count ratio for the eased-lockdown periods, respectively. The average climate temperature within the study period for Limpopo Province ranges from 10.8 to 29.8 °C with 0.1833 (18.33 %) death-to-count ratio and 1.1667 (116.67 %) recovery-to-count ratio for the lockdown periods and average climate temperature of 4.4 to 25.2 °C with 0.0095 (0.95 %) death-to-count ratio and 0.4794 (47.94 %) recovery-to-count ratio for the eased-lockdown periods, respectively. The average climate temperature within the study period for Northwest Province ranges from 9.4 to 30 °C with 0.000 (0 %) death-to-count ratio and 0.8412 (84.12 %) recovery-to-count ratio for the lockdown periods and average climate temperature of 0.5 to 25.1 °C with 0.0073 (0.73 %) death-to-count ratio and 0.1889 (18.89 %) recovery-to-count ratio for the eased-lockdown periods, respectively. The average climate temperature within the study period for Mpumalanga Province ranges from 10.4 to 30 °C with 0.000 (0 %) death-to-count ratio and 0.8400 (84 %) recovery-to-count ratio for the lockdown periods and average climate temperature of 7.1 to 25.9 °C with 0.0033 (0.33 %) death-to-count ratio and 0.5681 (56.81 %) recovery-to-count ratio for the eased-lockdown periods, respectively. The average climate temperature within the study period for Free State Province ranges from 4.8 to 28.4 °C with 0.1071 (10.71 %) death-to-count ratio and 2.0232 (202.32 %) recovery-to-count ratio for the lockdown periods and average climate temperature of -2.7 to 24 °C with 0.0098 (0.98 %) death-to-count ratio and 0.3362 (33.62 %) recovery-to-count ratio for the eased-lockdown periods, respectively. The average climate temperature within the study period for KwaZulu-Natal Province ranges from 15.6 to 29.6 °C with 0.0388 (3.88 %) death-to-count ratio and 0.3342 (33.42 %) recovery-to-count ratio for the lockdown periods and average climate temperature of 10.4 to 28.7 °C with 0.0146 (1.46 %) death-to-count ratio and 0.5626 (56.26 %) recovery-to-count ratio for the eased-lockdown periods, respectively. The average climate temperature within the study period for Western Cape Province ranges from 7.6 to 28.6 °C with 0.0188 (1.88 %) death-to-count ratio and 0.2933 (29.33 %) recovery-to-count ratio for the lockdown periods and average climate temperature of 6.7 to 25 °C with 0.0274 (2.74 %) death-to-count ratio and 0.5967 (59.67 %) recovery-to-count ratio for the eased-lockdown periods, respectively.

recovery-to-count ratio for the eased-lockdown periods, respectively. The average climate temperature within the study period for Gauteng Province ranges from 8.8 to 24.6 °C with 0.0066 (0.66 %) death-to-count ratio and 0.8233 (82.33 %) recovery-to-count ratio for the lockdown periods and average climate temperature of 1.6 to 20.6 °C with 0.0099 (0.99 %) death-to-count ratio and 0.4077 (40.77 %) recovery-to-count ratio for the eased-lockdown periods, respectively.

The results of COVID-19 cases, death, recovery rates along with the death-to-count, recovery-to-count ratios, and average weekly climate temperature are presented in Table 1 (a-i).

Table 1 (a – i): COVID-19 counts, death, recovery rates along with the death-to-count, recovery-to-count ratios, and average weekly climate temperature.

(a) Eastern Cape

	Date	Counts (C)	Death (D)	Recovery (R)	D/C	R/C	Temp. (°C)
WK 1	March 25-31	12	0	0	0.0000	0.0000	12 – 30
WK 2	April 1-7	20	0	0	0.0000	0.0000	12 – 36
WK 3	April 8-14	161	0	3	0.0000	0.0186	10 – 25
WK 4	April 15-21	171	5	12	0.0292	0.0702	9 – 34
WK 5	April 22-28	271	32	4	0.1181	0.0148	8 – 23
	Total	635	37	19			
	Average	127	7.4	3.8	0.0295	0.0207	10.2 – 29.6
WK 6	April 29-30/May, 1-5	222	8	295	0.0360	1.3288	6 – 34
WK 7	May 6-12	666	6	178	0.0090	0.2673	11 – 33
WK 8	May 13-19	631	17	320	0.0269	0.5071	6 – 24
WK 9	20-26, May	729	29	652	0.0398	0.8944	7 – 32
WK 10	May 27-31/June, 1-2	1461	18	632	0.0123	0.4326	7 – 29
WK 11	June 3-9	2436	40	1418	0.0164	0.5821	9 – 30
WK 12	June 10-16	4259	90	1961	0.0211	0.4604	5 – 28
WK 13	June 17-23	7069	86	2784	0.0122	0.3938	5 – 26
WK 14	June 24-30	9578	91	5822	0.0095	0.6079	6 – 29
	Total	27051	385	14062			
	Average	3005.7	42.78	1562	0.0204	0.6083	6.8 – 29.4

(b) Northern Cape

	Date	Counts (C)	Death (D)	Recovery (R)	D/C	R/C	Temp. (°C)
WK 1	March 25-31	9	0	0	0.0000	0.0000	11 – 37
WK 2	April 1-7	2	0	0	0.0000	0.0000	10 – 33
WK 3	April 8-14	8	0	0	0.0000	0.0000	9 – 33
WK 4	April 15-21	2	1	6	0.5000	3.0000	8 – 35
WK 5	April 22-28	1	0	4	0.0000	4.0000	8 – 31
	Total	22	1	10			
	Average	4.4	0.2	2	0.1000	1.4000	9.2 – 33.8
WK 6	April 29-30/May, 1-5	9	0	3	0.0000	0.3333	6 – 30
WK 7	May 6-12	4	0	3	0.0000	0.7500	7 – 30
WK 8	May 13-19	7	0	14	0.0000	2.0000	4 – 31
WK 9	20-26, May	9	0	2	0.0000	0.2222	3 – 29
WK 10	May 27-31/June, 1-2	46	0	4	0.0000	0.0870	1 – 30
WK 11	June 3-9	29	0	8	0.0000	0.2759	0 – 32
WK 12	June 10-16	85	0	8	0.0000	0.0941	-8 – 24

WK 13	June 17-23	59	0	60	0.0000	1.0169	3 – 25
WK 14	June 24-30	195	0	74	0.0000	0.3795	0 – 24
Total		443	0	176			
Average		49	0	19.56	0.0000	0.5732	1.7 – 28.3

(c) Limpopo

	Date	Counts (C)	Death (D)	Recovery (R)	D/C	R/C	Temp. (°C)
WK 1	March 25-31	14	0	0	0.0000	0.0000	14 – 32
WK 2	April 1-7	5	0	0	0.0000	0.0000	12 – 31
WK 3	April 8-14	6	0	13	0.0000	2.1667	10 – 31
WK 4	April 15-21	3	2	8	0.6667	2.6667	9 – 28
WK 5	April 22-28	4	1	4	0.2500	1.0000	9 – 27
Total		32	3	25			
Average		6.4	0.6	5	0.1833	1.1667	10.8 – 29.8
WK 6	April 29-30/May, 1-5	9	0	6	0.0000	0.6667	9 – 25
WK 7	May 6-12	14	1	1	0.0714	0.0714	8 – 26
WK 8	May 13-19	40	0	11	0.0000	0.2750	7 – 26
WK 9	20-26, May	40	0	26	0.0000	0.6500	6 – 25
WK 10	May 27-31/June, 1-2	61	0	80	0.0000	1.3115	0 – 26
WK 11	June 3-9	54	0	13	0.0000	0.2407	5 – 26
WK 12	June 10-16	141	0	51	0.0000	0.3617	-1 – 26
WK 13	June 17-23	231	1	57	0.0043	0.2468	3 – 23
WK 14	June 24-30	509	5	250	0.0098	0.4912	3 – 24
Total		1099	7	495			
Average		122	0.78	55	0.0095	0.4794	4.4 – 25.2

(d) North West

	Date	Counts (C)	Death (D)	Recovery (R)	D/C	R/C	Temp. (°C)
WK 1	March 25-31	8	0	0	0.0000	0.0000	12 – 32
WK 2	April 1-7	3	0	0	0.0000	0.0000	9 – 29
WK 3	April 8-14	11	0	3	0.0000	0.2727	8 – 32
WK 4	April 15-21	3	0	10	0.0000	3.3333	8 – 29
WK 5	April 22-28	5	0	3	0.0000	0.6000	10 – 28
Total		30	0	16			
Average		6	0	3.2	0.0000	0.8412	9.4 – 30
WK 6	April 29-30/May, 1-5	7	0	4	0.0000	0.5714	6 – 27
WK 7	May 6-12	17	1	3	0.0588	0.1765	6 – 27
WK 8	May 13-19	21	0	5	0.0000	0.2381	5 – 26
WK 9	20-26, May	43	0	1	0.0000	0.0233	0 – 25
WK 10	May 27-31/June, 1-2	156	0	30	0.0000	0.1923	-3 – 27
WK 11	June 3-9	333	0	29	0.0000	0.0871	-1 – 24
WK 12	June 10-16	674	4	64	0.0059	0.0950	-8 – 25
WK 13	June 17-23	1173	0	104	0.0000	0.0887	1 – 21
WK 14	June 24-30	1733	2	395	0.0012	0.2279	-1 – 24
Total		4157	7	635			
Average		462	0.8	70.6	0.0073	0.1889	0.5 – 25.1

(e) Mpumalanga

	Date	Counts (C)	Death (D)	Recovery (R)	D/C	R/C	Temp. (°C)
--	------	---------------	-----------	-----------------	-----	-----	------------

WK 1	March 25-31	12	0	0	0.0000	0.0000	13 – 35
WK 2	April 1-7	6	0	0	0.0000	0.0000	0 – 33
WK 3	April 8-14	5	0	6	0.0000	1.2000	14 – 32
WK 4	April 15-21	4	0	8	0.0000	2.0000	12 – 24
WK 5	April 22-28	4	0	4	0.0000	1.0000	13 – 26
Total		31	0	18			
Average		6.2	0	3.6	0.0000	0.8400	10.4 – 30
WK 6	April 29-30/May, 1-5	34	0	4	0.0000	0.1176	11 – 27
WK 7	May 6-12	6	0	0	0.0000	0.0000	9 – 25
WK 8	May 13-19	13	0	35	0.0000	2.6923	10 – 27
WK 9	20-26, May	26	0	4	0.0000	0.1538	9 – 26
WK 10	May 27-31/June, 1-2	29	0	25	0.0000	0.8621	4 – 27
WK 11	June 3-9	58	1	25	0.0172	0.4310	7 – 23
WK 12	June 10-16	146	0	27	0.0000	0.1849	2 – 27
WK 13	June 17-23	319	1	43	0.0031	0.1348	6 – 24
WK 14	June 24-30	528	5	283	0.0095	0.5360	6 – 27
Total		1159	7	446			
Average		129	0.78	50	0.0033	0.5681	7.1 – 25.9

(f) Free State

	Date	Counts (C)	Death (D)	Recovery (R)	D/C	R/C	Temp. (°C)
WK 1	March 25-31	74	1	0	0.0135	0.0000	6 – 30
WK 2	April 1-7	15	0	0	0.0000	0.0000	5 – 27
WK 3	April 8-14	10	3	61	0.3000	6.1000	2 – 28
WK 4	April 15-21	9	2	13	0.2222	1.4444	4 – 29
WK 5	April 22-28	7	0	18	0.0000	2.5714	7 – 28
Total		115	6	92			
Average		23	1.2	18.4	0.1071	2.0232	4.8 – 28.4
WK 6	April 29-30/May, 1-5	15	1	4	0.0667	0.2667	2 – 25
WK 7	May 6-12	8	0	9	0.0000	1.1250	2 – 25
WK 8	May 13-19	33	0	13	0.0000	0.3939	-2 – 26
WK 9	20-26, May	40	0	5	0.0000	0.1250	1 – 24
WK 10	May 27-31/June, 1-2	93	2	2	0.0215	0.0215	-5 – 26
WK 11	June 3-9	92	0	39	0.0000	0.4239	-2 – 25
WK 12	June 10-16	182	0	58	0.0000	0.3187	-9 – 23
WK 13	June 17-23	236	0	42	0.0000	0.1780	-5 – 21
WK 14	June 24-30	700	0	121	0.0000	0.1729	-7 – 21
Total		1399	3	293			
Average		155	0.33	32.56	0.0098	0.3362	-2.7 – 24

(g) KwaZulu-Natal

	Date	Counts (C)	Death (D)	Recovery (R)	D/C	R/C	Temp. (°C)
WK 1	March 25-31	179	2	0	0.0112	0.0000	16 – 31
WK 2	April 1-7	78	6	0	0.0769	0.0000	15 – 29
WK 3	April 8-14	232	4	15	0.0172	0.0647	16 – 33
WK 4	April 15-21	182	11	136	0.0604	0.7473	15 – 28
WK 5	April 22-28	248	7	213	0.0282	0.8589	16 – 27
Total		919	30	364			
Average		184	6	72.8	0.0388	0.3342	15.6 – 29.6

WK 6	April 29-30/May, 1-5	223	6	51	0.0269	0.2287	13 – 30
WK 7	May 6-12	252	8	191	0.0317	0.7579	14 – 32
WK 8	May 13-19	222	2	212	0.0090	0.9550	12 – 26
WK 9	20-26, May	314	3	362	0.0096	1.1529	12 – 31
WK 10	May 27-31/June, 1-2	710	4	68	0.0056	0.0958	8 – 28
WK 11	June 3-9	618	9	320	0.0146	0.5178	11 – 28
WK 12	June 10-16	790	11	565	0.0139	0.7152	7 – 30
WK 13	June 17-23	1577	18	716	0.0114	0.4540	9 – 28
WK 14	June 24-30	4049	35	753	0.0086	0.1860	8 – 25
Total		8755	96	3238			
Average		973	11	360	0.0146	0.5626	10.4 – 28.7

(h) Western Cape

	Date	Counts (C)	Death (D)	Recovery (R)	D/C	R/C	Temp. (°C)
WK 1	March 25-31	325	1	0	0.0031	0.0000	8 – 24
WK 2	April 1-7	137	2	0	0.0146	0.0000	9 – 30
WK 3	April 8-14	181	5	152	0.0276	0.8398	4 – 30
WK 4	April 15-21	367	9	64	0.0245	0.1744	7 – 32
WK 5	April 22-28	860	21	389	0.0244	0.4523	10 – 27
Total		1870	38	605			
Average		374	7.6	121	0.0188	0.2933	7.6 – 28.6
WK 6	April 29-30/May, 1-5	1739	33	228	0.0190	0.1311	8 – 27
WK 7	May 6-12	2496	34	724	0.0136	0.2901	10 – 28
WK 8	May 13-19	4534	81	2114	0.0179	0.4663	6 – 23
WK 9	20-26, May	5190	171	3481	0.0329	0.6707	7 – 28
WK 10	May 27-31/June, 1-2	7754	225	4592	0.0290	0.5922	6 – 24
WK 11	June 3-9	11236	322	8355	0.0287	0.7436	6 – 25
WK 12	June 10-16	10538	296	10137	0.0281	0.9619	5 – 22
WK 13	June 17-23	8155	336	6998	0.0412	0.8581	6 – 27
WK 14	June 24-30	8969	323	5886	0.0360	0.6563	6 – 21
Total		60611	1821	42515			
Average		6734.6	202	4724	0.0274	0.5967	6.7 – 25

(i) Gauteng

	Date	Counts (C)	Death (D)	Recovery (R)	D/C	R/C	Temp. (°C)
WK 1	March 25-31	633	1	0	0.0016	0.0000	11 – 27
WK 2	April 1-7	80	0	0	0.0000	0.0000	9 – 24
WK 3	April 8-14	196	3	157	0.0153	0.8010	8 – 25
WK 4	April 15-21	290	3	388	0.0103	1.3379	7 – 24
WK 5	April 22-28	178	1	352	0.0056	1.9775	9 – 23
Total		1377	8	897			
Average		275.4	1.6	179.4	0.0066	0.8233	8.8 – 24.6
WK 6	April 29-30/May, 1-5	320	7	82	0.0219	0.2563	7 – 23
WK 7	May 6-12	317	7	268	0.0221	0.8454	5 – 22
WK 8	May 13-19	347	5	420	0.0144	1.2104	4 – 21
WK 9	20-26, May	682	4	293	0.0059	0.4296	1 – 21
WK 10	May 27-31/June, 1-2	1233	2	141	0.0016	0.1144	0 – 23
WK 11	June 3-9	2270	24	515	0.0106	0.2269	3 – 23
WK 12	June 10-16	6477	30	826	0.0046	0.1275	-4 – 17

WK 13	June 17-23	11018	35	2178	0.0032	0.1977	-2 – 17
WK 14	June 24-30	18840	94	4914	0.0050	0.2608	0 – 18
Total		41504	208	9637			
Average		4611.6	23.1	1071	0.0099	0.4077	1.6 – 20.6

As shown from the results in Table 1 (a – i), higher or lower climate temperatures cannot be reported to prevent or delay the transmissions and death rates of patients of COVID-19. Nonetheless, it can be concluded that a higher temperature of the climate raises recovery rates for COVID-19 patients. Figure 2 shows the average minimum and maximum climate temperatures and the death-to-count ratio for the lockdown period. In contrast, Figure 3 shows the average minimum and maximum climate temperatures and the COVID-19 recovery-to-count ratio for the lockdown period.

Fig. 2: Average minimum and maximum climate temperatures and death-to-count (D/C) ratio for the lockdown period.

Fig. 3: Average minimum and maximum climate temperatures and recovery-to-count (R/C) for the lockdown period.

Figure 4 shows the average minimum and maximum climate temperatures and the death-to-count ratio for the eased-lockdown period. In contrast, Figure 5 shows the average minimum and maximum climate temperatures and the COVID-19 recovery-to-count ratio for the eased-lockdown period.

Fig. 4: Average minimum and maximum climate temperatures and death-to-count (D/C) ratio for the eased-lockdown period.

Fig. 5: Average minimum and maximum climate temperatures and recovery-to-count (R/C) for the eased-lockdown period.

The results showed that higher climate temperatures aren't essential to avoid the COVID-19 from being spread. The present results conform to the reports that suggested that COVID-19 is unlike the seasonal flu, which does dissipate as the climate temperature rises [17]. Accordingly, the ratio of counts and death-to-count cannot be concluded to be influenced by variations in the climate temperatures within the study areas.

3.2 Correlation between climate temperature, death-to-count and recovery-to-death ratios of COVID-19 in the study areas

Table 2 showed the correlation values for climate temperatures, the death-to-count, and the recovery-to-count ratios of COVID-19 in South African provinces during the study period (March 25 to June 30, 2020). Figures 6 and 7 presented the correlation pattern of COVID-19 death and recovery rates and the daily minimum and maximum temperatures in the study period.

A positive association of $r = 0.27, 0.45$ for climate temperatures and death-to-count ratio and $r = 0.31, 0.38$ for climate temperatures and recovery-to-count ratio, respectively, was recorded for the Free State province. In contrast, a negative association of $r = -0.54, -0.10$ for climate temperatures and death-to-count ratio and $r = -0.71, -0.34$ for climate temperatures and recovery-to-count ratio, respectively, was recorded for the Western Cape Province.

Table 2: Correlation of climate temperature with death-to-count, and recovery-to-count rates of COVID-19 within the study period.

S/N	Study provinces	Corr. (Temp, D/C)		Corr. (Temp, R/C)	
		Min.	Max.	Min.	Max.
1	Eastern Cape	-0.25702	-0.36147	-0.62102	0.20832
2	Northern Cape	0.35047	0.20117	0.15736	0.24561

3	Western Cape	-0.54262	-0.10085	-0.71306	-0.33939
4	Limpopo	0.20555	0.10653	0.04314	0.18914
5	Gauteng	0.20852	0.12452	0.3466	0.14557
6	Mpumalanga	-0.17996	-0.3818	0.37143	-0.17437
7	Free State	0.2656	0.45127	0.3077	0.38355
8	KwaZulu-Natal	0.51814	0.05807	-0.05048	-0.11779
9	North West	0.05066	-0.0095	0.26437	0.22027

Generally, the correlation curve (Figure 6) tends not to show an association between climate temperature and death rates of COVID-19. The curve shows similar trends for both average minimum and maximum climate temperatures. However, the correlation curve (Figure 7) tends to show an association between climate temperature and recovery rates of COVID-19 patients with an almost perfect linear recovery correlation for the maximum temperature of 24°C and above. The curve shows a clear difference in recovery rates at the minimum and maximum climate temperature conditions.

Figure 6: correlation curve between climate temperature and death-to-count (D/C) ratio within the study period

Figure 7: correlation curve between climate temperature and recovery-to-count (R/C) ratio within the study period.

Table 3: Comparison of COVID-19 counts, death, recovery rates, and death-to-count, recovery-to-count ratios from the study provinces with values from other countries (as of June 30, 2020).

Country	Count (C)	Death (D)	Recovery (R)	D/C	R/C	References
Eastern Cape	27,686	422	14,081	0.0152	0.5086	
Northern Cape	465	1	186	0.0022	0.4000	
Limpopo	1,131	10	520	0.0088	0.4598	
North West	4,187	7	651	0.0017	0.1555	Present
Mpumalanga	1,190	7	464	0.0059	0.3899	Study
Free State	1,514	9	385	0.0059	0.2543	(South Africa)
KwaZulu-Natal	9,674	126	3,602	0.0130	0.3723	
Western Cape	62,481	1,859	43,120	0.0298	0.6901	
Gauteng	42,881	216	10,534	0.0050	0.2457	
Total				0.0212	0.5510	
Australia*	7,834	104	7,037	0.0133	0.8983	[18]
USA*	2,618,480	126,645	705,203	0.0484	0.2693	[19]
Argentina*	64,517	1,307	22,015	0.0203	0.3412	[20]
Russia**	647,849	9,320	412,650	0.0144	0.6370	[21]
Nigeria**	25,694	590	9,746	0.0230	0.3793	[22]
India**	566,840	16,893	334,822	0.0298	0.5907	[23]

*similar climate temperature as South Africa **higher climate temperature than South Africa.

The results show no significant difference in counts, death, and death-to-count rates for different climate temperatures compared with values from other countries with similar and higher climate temperatures, just like South Africa's. Higher recovery rates for countries with higher climate temperatures (Table 3) have also been reported. This study concludes that the virus can be spread at every climate temperature, with no significant difference in the count and death rates. However, higher recovery rates were recorded for countries with higher climate temperatures than South Africa, which coincides with the results from provinces with higher climate temperatures from the study areas having higher recovery rates of patients infected with COVID-19.

4. Conclusion

The study investigates the impact of climate temperature on the counts, recovery, and death rates of COVID-19 cases in all South Africa's provinces. The findings were compared with those of countries with comparable climate temperatures as South Africa. Our result indicates that a higher or lower climate temperature does not prevent or delay the spread and death rates but shows significant positive impacts on the recovery rates of COVID-19 patients. Warm climate temperatures seem not to restrict the spread of the COVID-19 as the count rate was substantial at every climate temperatures. Thus, it indicates that the climate temperature is unlikely to impose a strict limit on the spread of COVID-19. There is no correlation between the cases and death rates, an indicator that there is no particular temperature range of the climatic conditions closely associated with a faster or slower death rate of COVID-19 patients. However, other shortcomings in this study's process should not be ignored. Some other factors may have contributed to recovery rates, such as the South African government's timely intervention to announce a national lockdown at the early stage of the outbreak, the availability of intensive medical care, and social

distancing effects. Nevertheless, this study shows that a warm climate temperature can only help COVID-19 patients recover more quickly, thereby having huge impacts on the death and active case rates.

References

1. Melanie Bannister-Tyrrell, Anne Meyer, Celine Faverjon, Angus Cameron (2020). Preliminary evidence that higher temperatures are associated with a lower incidence of COVID-19, for cases reported globally up to 29th February 2020. medRxiv 2020.03.18.20036731. <https://doi.org/10.1101/2020.03.18.20036731>
2. Peng Shi, Yinqiao Dong, Huanchang Yan, Xiaoyang Li, Chenkai Zhao, Wei Liu, Miao He, Shixi ng Tang, Shuhua Xi (2019). *The impact of temperature and absolute humidity on the coronavirus disease (COVID-19) outbreak - evidence from China*. medRxiv 2020.03.22.20038919. <https://doi.org/10.1101/2020.03.22.20038919>
3. Jin Bu, Dong-Dong Peng, Hui Xiao, Qian Yue, Yan Han, Yu Lin, Gang Hu, Jing Chen (2020). *Analysis of meteorological conditions and prediction of the epidemic trend of 2019-nCoV infection*. medRxiv 2020.02.13.20022715. <https://doi.org/10.1101/2020.02.13.20022715>
4. Sajadi, Mohammad M. and Habibzadeh, Parham and Vintzileos, Augustin and Shokouhi, Shervin and Miralles-Wilhelm, Fernando, and Amoroso, Anthony (2020). *Temperature, Humidity, and Latitude Analysis to Predict Potential Spread and Seasonality for COVID-19*. Available at <https://dx.doi.org/10.2139/ssrn.3550308>
5. Wei Luo, Maimuna S. Majumder, Diamo Liu, Canelle Poirier, Kenneth D Mandl, Marc Lipsitch, Mauricio Santillana (2020). *The role of absolute humidity on transmission rates of the COVID-19 outbreak*. medRxiv 2020.02.12.20022467. <https://doi.org/10.1101/2020.02.12.20022467>
6. van der Hoek, L., P. Krzysztof, and B. Berkhout. (2006). *Human coronavirus NL63, a new respiratory virus*. FEMS Microbiol. Rev. 30:760-773. <https://doi.org/10.1111/j.1574-6976.2006.00032.x>
7. Zhao, Q., S. Li, F. Xue, Y. Zou, C. Chen, M. Bartlam, and Z. Rao. (2008). *Structure of the main protease from a global infectious human coronavirus, HCoV-HKU1*. J. Virol. 82:8647-8655. <https://jvi.asm.org/content/82/17/8647>
8. Gaunt ER, Hardie A, Claas EC, Simmonds P, Templeton KE. (2010). *Epidemiology and clinical presentations of the four human coronaviruses 229E, HKU1, NL63, and OC43 detected over three years using a novel multiplex real-time PCR method*. J Clin Microbiol.; 48(8):2940-2947. <https://jcm.asm.org/content/48/8/2940>
9. K.H. Chan, J.S. Peiris, SY. Lam, L.L. Poon, K.Y. Yuen, W.H. Seto (2011). *The Effects of Temperature and Relative Humidity on the Viability of the SARS Coronavirus*. Adv Virol, p. 734690. <https://doi.org/10.1155/2011/734690>
10. N. van Doremalen, T. Bushmaker, D.H. Morris, M.G. Holbrook, A. Gamble, B.N. Williamson, A. Tamin, J.L. Harcourt, N.J. Thornburg, S.I. Gerber *et al.* (2020). *Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1*. Engl J Med; 382:1564-1567. <https://www.nejm.org/doi/full/10.1056/nejmc2004973>
11. Zhou P, Yang XL, Wang XG, *et al.* (2020). *A pneumonia outbreak associated with a new coronavirus of probable bat origin*. Nature 579, 270–273. <https://doi.org/10.1038/s41586-020-2012-7>
12. Yueling Ma, Yadong Zhao, Jiangtao Liu, Xiaotao He, Bo Wang, Shihua Fu, Jun Yan, Jingping Ni u, Bin Luo (2020). *Effects of temperature variation and humidity on the mortality of COVID-19 in Wuhan*. Science of the Total Environment, Volume 724, 138226. <https://doi.org/10.1016/j.scitotenv.2020.138226>
13. The map of South Africa's provinces. Available at <https://www.southafrica.to/provinces/provinces.php>

14. Fulekar, M.H. (2009). *Bioinformatics in Life and Environmental Sciences*, Springer (pp. 110) <https://stanford.idm.oclc.org/login?url=http://link.springer.com/10.1007/978-1-4020-8880-3>
15. Buda, A., Jarynowski, A. (2010). *Life Time of Correlations and its Applications*. Wydawnictwo Niezależne, Vol. 1, pp. 5–21. <https://arxiv.org/abs/1105.6272>
16. Huber, P.J. (2011). *Robust Statistics*. In: Lovric M. (eds) International Encyclopedia of Statistical Science. Springer, Berlin, Heidelberg. <https://doi.org/10.1007/978-3-642-04898-2>
17. Kay Lazar and Andrew Ryan (2020). *No, warm weather will not kill the Coronavirus*. <https://www.bostonglobe.com/2020/04/24/nation/no-warm-weather-will-not-kill-coronavirus/>
18. Department of Health of Australian Government (2020). *Coronavirus (COVID-19) situation in Australia*. <https://www.health.gov.au/resources/publications/coronavirus-covid-19-at-a-glance-30-june-2020>
19. Coronavirus USA news updates for Tuesday 30 June 2020 https://en.as.com/en/2020/06/30/latest_news/1593468452_648953.html
20. COVID-19 cases in Argentina. <https://www.argentina.gob.ar/coronavirus/informe-diario>
21. Cumulative number of coronavirus (COVID-19) cases, active cases, recoveries, and deaths in Russia <https://www.statista.com/statistics/1107929/cumulative-coronavirus-cases-in-russia/>
22. COVID-19 Update in Nigeria. <https://nairametrics.com/2020/06/27/covid-19-update-in-nigeria/>
23. Johns Hopkins University (2020). *Coronavirus India news summary as at Tuesday 16 June* https://en.as.com/en/2020/06/30/latest_news/1593474002_618328.html

Author contributions statement

NC conceived the study, supervised and guided the process of the research. BA and AOI performed data collection and analysis. All authors wrote and reviewed the manuscript.

Competing Interests

The authors declare no competing interests.