Autoimmune Encephalitis Presenting with Malignant Catatonia in a 40-Year-Old Male Patient with Covid-19

Jan Mulder, PhD^{1*} and Amalia Feresiadou MD^{2*}, David Fällmar MD PhD³, Robert Frithiof, MD PhD⁴, Johan Virhammar MD PhD², Annica Rasmusson PhD⁵, Elham Rostami MD PhD⁶, Eva Kumlien MD PhD², Janet L. Cunningham MD PhD^{5§}

*Equal contribution

§Corresponding author

¹Department of Neuroscience, Karolinska Institute, Stockholm, Sweden

²Department of Neuroscience, Neurology, Uppsala University, Uppsala, Sweden

³Department of Surgical Sciences, Radiology, Uppsala University, Uppsala, Sweden

⁴Department of Surgical Sciences, Anaesthesia and Intensive Care, Uppsala University, Uppsala, Sweden

⁵Department of Neuroscience, Psychiatry, Uppsala University, Uppsala, Sweden

⁶Department of Neuroscience, Neurosurgery, Uppsala University, Uppsala, Sweden

SUMMARY

Acute malignant catatonia with autonomic instability developed in a previously healthy man with PCR-verified SARS-CoV-2. CT and MRI were normal, EEG showed slowing and cerebrospinal fluid showed a subtle indication of inflammation. There were no signs of pathology in other organs. ^{18}F -FDG-PET conveyed high bilateral uptake in the striatum. While commercial tests were negative, immunohistochemical staining of mouse brain revealed anti-neuronal IgG antibodies against neuronal targets in the hippocampus, thalamus, striatum and cortex. Early treatment with plasmapheresis and corticosteroid reversed disease progression and may have prevented large-scale neurological damage. We are not aware of other types of encephalitis with such distinct pyramidal tract symptoms and raise the possibility that this may be a novel form of autoimmune encephalitis induced by infection with SARS-CoV-2.

1 INTRODUCTION

2 A 40-year-old man showed acute debut of agitation, grimacing, repetitive speech and movements
3 (verbigitation and stereotypies) that started earlier the same day. His Glasgow coma scale score
4 was 14. Because his behaviour was bizarre, disorganised, hyperkinetic and uncooperative, a
5 decision was made for involuntary commitment. He had been experiencing high levels of stress
6 in recent months as his business was struggling. Twenty-two days before he developed Covid-19
7 related respiratory symptoms, dyspnoea and fatigue, which did not require hospital care and he
8 had tested positive for SARS-CoV-2 RNA in a nasopharyngeal swab (day 14; Figure 1A).
9 Anosmia and ageusia were not present. Before admission, he had suffered from a headache. On
10 admission (day 22), he no longer had respiratory symptoms but did have a fever (38.4 °C). He
11 made no eye contact, reflexes were normal and Babinski's sign was absent. Brain CT, metabolic
12 parameters and blood tests were unremarkable. The patient was lightly sedated, with midazolam
13 followed with dexmedetomidine. Neuroleptics were not used. Treatment with antibiotics and
14 acyclovir was initiated until the tests excluded bacterial infection and herpes encephalitis.

15
16 Lumbar puncture showed high red blood cells, 19000 cells $\times 10^6/L$, secondary to traumatic
17 lumbar puncture. Cerebral spinal fluid (CSF) cell count indicated pleocytosis with 23
18 mononuclear and 8 polymorphonuclear cells $\times 10^6/L$. Signs of blood-brain barrier disruption with
19 elevated albumin levels in CSF of 838 mg/L (reference <400 mg/L) and the
20 CSF/serum quotient was 15.6 (>6.8). Interleukin 6 (IL6) in CSF was high: 102.1 pg/mL (<5
21 pg/mL). IL6 in plasma, taken a day later, was 30 pg/ml (<7.0). Neurofilament (NFL), glial
22 fibrillary acidic protein (GFAP) and tau protein in CSF were normal. PCR tests were negative in
23 CSF for SARS-CoV-2. Antibodies against anti-neuronal antibodies were negative using a
24 commercial assay (Euroimmun, Germany). Brain MRI was normal.

25

26 A few hours later, the patient's state deteriorated, temperature rose to 39°C and he became
 27 mutistic, showing signs of autonomic instability with recurrent episodes of fluctuating heart rate
 28 and arterial blood pressure. He also exhibited periods of oxygen desaturation (Figure 1B). When
 29 hypertension ensued, it was difficult to treat, despite high doses of clonidine and labetalol. Plasma
 30 lactate varied between 0.6 and 8 mmol/L but myoglobin but CKMB remained normal. His
 31 pupil size, reaction to light and the oculo-cephalic reflex was normal. Slow, horizontal roving eye
 32 moments were noted. He displayed decorticate posturing, increased tonus, resisted movement of
 33 arms and jaw but had normal tonus in the legs. Hyperreflexia was present with bilateral foot
 34 clonus and Babinski's sign but no neck stiffness. Anaesthesia was induced to facilitate
 35 endotracheal intubation. The patient was thereafter continuously sedated with propofol and
 36 clonidine. D-Dimer (1.2mg/L) was slightly elevated but there were no signs of thromboembolic
 37 events. Respiration and cardiovascular function remained stable. Continuous EEG monitoring
 38 showed non-specific slowing with left hemisphere predominance without epileptiform activity.
 39 An episode of asystole with spontaneous recovery, episodes of bradycardia of 27 bpm and
 40 repeated p-waves without QRS waves were interpreted as atrioventricular block III and the
 41 patient was monitored for 3 days with telemetry.

42

43 RESULTS

44 Although the diagnosis remained uncertain, post-infectious autoimmune encephalitis was
 45 suspected. Plasmapheresis was therefore initiated and repeated three times over 4 days. On day
 46 26 and after two courses of plasmapheresis, the patient was extubated and was autonomically
 47 stable. Eye movement was normalised, hyperreflexia was less prominent and bilateral Babinski's
 48 sign persisted. This case shared features with a recently published steroid-responsive case¹ and
 49 treatment was therefore initiated with 1g methylprednisolone per day and discontinued after 3
 50 days because of insomnia and elevated mood.

51

52 On day 28, the patient showed a dramatic improvement. He was awake, oriented, communicative
53 but had no memories from the past days. He had visual hallucinations and cognitive symptoms
54 that improved over the coming weeks. The patient's EEG was normal.

55

56 A second lumbar puncture, performed on day 29, showed pleocytosis, 10 mononuclear cells and
57 1 polymorphonuclear cell $\times 10^6/L$, elevated IgG levels and IgG index, as well as two oligoclonal
58 bands in CSF not represented in serum, indicating intrathecal production of antibodies. PCR
59 tests were still negative in CSF for SARS-CoV-2. IL6 in CSF was normalised at 4.8 pg/ml. GFAP
60 and Tau remained normal but NFL was increased to 1030 ng/L (<890). A second MRI and a
61 standard neurological exam on day 31 were normal, and no abnormal eye movements. The
62 hallucinations were less frequent. He described increased emotional lability and mental fatigue
63 with disturbed short-term memory and decision making. He also found it challenging to
64 recognise the voices and faces of acquaintances. Serology taken on day 33 was strongly positive
65 for IgG against SARS-CoV-2. ^{18}F -FDG PET scan, day 35 (13 days after treatment initiation),
66 showed high bilateral uptake in the striatum (caudate nucleus and putamen) when compared to
67 the cortex (Figure 1C).

68

69 EVIDENCE OF AUTOIMMUNITY

70 Using immunohistochemistry we detected IgG autoantibodies against mouse brain neuronal
71 proteins in serum and CSF at admission (Figure 1D-W). Immunoreactivity was observed in all
72 brain regions investigated (cerebral cortex, hippocampal formation, thalamus, caudate-putamen)
73 with strongest signal observed in the nuclear and perinuclear compartments without reactivity in
74 the nucleoli. Neuronal labelling intensity was strongest in the CA3 in the hippocampal formation,
75 layer V in the somatosensory cortex and the paraventricular and reticular nucleus in the thalamus.

76 A subset of ependymal cells located in the ventricle wall and choroid plexus revealed strong
77 immunoreactivity of the (peri)nuclear compartment and cytoplasm. Immunoreactivity of neuropil
78 was most intense in the caudate-putamen, revealing neuronal processes and spine-like structures.
79 Post-treatment IgG immunoreactivity in the (peri)nuclear compartment and neuropil was notably
80 reduced, reaching the levels of reference CSF and serum.

81

82 DISCUSSION

83 This report is a case of suspected autoimmune encephalitis in a patient with Covid-19 who
84 presented with excited catatonia progressing to malignant catatonia with autonomic instability
85 Signs of autoimmune encephalitis were present but this case did not initially meet the proposed
86 criteria.² Standard radiological findings were normal and the discrete pleocytosis and elevated
87 protein in CSF may have easily been ignored as non-specific. Cases are reported with rapid onset
88 of encephalitis with diffuse corticospinal tract signs in conjunction with Covid-19.^{3,4} Neurological
89 symptoms may also occur as an early Covid-19 manifestation in patients who later develop
90 typical symptoms.⁴ Neurological symptoms in Covid-19 patients include one case of catatonia⁵ as
91 well as mutism.¹ Autonomic dysfunction after Covid-19 has been reported in a patient with
92 psoriasis.⁶ Several forms of autoimmunity after Covid-19 are emerging and this coronavirus may
93 join other neurotropic viruses as a risk factor for autoimmune encephalitis.⁷

94

95 Malignant catatonia is a more severe form of the catatonia spectrum but perhaps less recognised
96 than the typical form with immobility, waxy flexibility and stupor.^{8,9} A recent review identified
97 124 cases of catatonia in conjunction with infections, of which 38% were viral.¹⁰ Catatonia may
98 occur in patients with no psychiatric history and is co-occurring with several autoimmune
99 conditions, especially NMDAr encephalitis.¹⁰ The staining, clinical presentation and radiology are
100 consistent with a synaptic target highly present in the striatum, where binding leads to increased

excitation. Possible mechanisms include disinhibition or activation of excitatory receptors as well as disturbance of excitatory neurotransmitter uptake/clearance or interference with ion homeostasis.

In patients with acute respiratory distress syndrome related to Covid-19, agitation diffuse corticospinal tract signs along with enhanced reflexes and ankle clonus are common findings seen in 67% of cases³. In the same study, dysexecutive symptoms were observed in fewer cases (37%).³ The suggested causes are critical illness-related encephalopathy, elevated cytokines, or the effect of withdrawal of medication. Autoimmunity in these cases, though speculated, has not yet been reported.

The present case was not classed as a critical illness and IL6 was only moderately elevated. Normal MRI and NFL levels in CSF did not support neuronal injury. The ¹⁸F-FDG PET findings in the striatum are consistent with increased metabolism in local neurons, glial cells, or both. Striatal hypermetabolism has been previously reported for autoimmune encephalitis caused by antibodies against NMDAr and VGKC encephalitis.¹¹⁻¹³

The swift introduction of treatment with plasmapheresis and high dose steroids may have prevented the development of more severe neurological damage.. The treatment reduced autoreactive IgG in both serum and CSF but may also have other actions.¹⁴

We conclude that malignant catatonia with potentially life-threatening autonomic instability can occur in patients with Covid-19 and is likely a form of para-infectious autoimmune encephalitis. This condition responded with dramatic improvement and there was no evidence of structural brain damage. We are not aware of other types of encephalitis with such distinct pyramidal tract

126 symptoms and raise the possibility that this may be a novel form of autoimmune encephalitis
127 induced by infection with SARS-CoV-2.

REFERENCES

1. Pilotto A, Odolini S, Masciocchi S, et al. Steroid-Responsive Encephalitis in Coronavirus Disease 2019. *Ann Neurol* 2020.
2. Pollak TA, Lennox BR, Muller S, et al. Autoimmune psychosis: an international consensus on an approach to the diagnosis and management of psychosis of suspected autoimmune origin. *Lancet Psychiatry* 2020;7:93-108.
3. Helms J, Kremer S, Merdji H, et al. Neurologic Features in Severe SARS-CoV-2 Infection. *N Engl J Med* 2020;382:2268-70.
4. Mao L, Jin H, Wang M, et al. Neurologic Manifestations of Hospitalized Patients With Coronavirus Disease 2019 in Wuhan, China. *JAMA Neurol* 2020.
5. Rogers JP, Chesney E, Oliver D, et al. Psychiatric and neuropsychiatric presentations associated with severe coronavirus infections: a systematic review and meta-analysis with comparison to the COVID-19 pandemic. *Lancet Psychiatry* 2020;7:611-27.
6. Logmin K, Karam M, Schichel T, Harmel J, Wojtecki L. Non-epileptic seizures in autonomic dysfunction as the initial symptom of COVID-19. *J Neurol* 2020.
7. Ehrenfeld M, Tincani A, Andreoli L, et al. Covid-19 and autoimmunity. *Autoimmun Rev* 2020:102597.
8. Saddawi-Konefka D, Berg SM, Nejad SH, Bittner EA. Catatonia in the ICU: an important and underdiagnosed cause of altered mental status. a case series and review of the literature*. *Crit Care Med* 2014;42:e234-41.
9. Wilson JE, Niu K, Nicolson SE, Levine SZ, Heckers S. The diagnostic criteria and structure of catatonia. *Schizophr Res* 2015;164:256-62.
10. Rogers JP, Pollak TA, Blackman G, David AS. Catatonia and the immune system: a review. *Lancet Psychiatry* 2019;6:620-30.
11. Kremer S, Lersy F, de Seze J, et al. Brain MRI Findings in Severe COVID-19: A Retrospective Observational Study. *Radiology* 2020:202222.
12. Kandemirli SG, Dogan L, Sarikaya ZT, et al. Brain MRI Findings in Patients in the Intensive Care Unit with COVID-19 Infection. *Radiology* 2020:201697.
13. Moloney P, Boylan R, Elamin M, O'Riordan S, Killeen R, McGuigan C. Semi-quantitative analysis of cerebral FDG-PET reveals striatal hypermetabolism and normal cortical metabolism in a case of VGKCC limbic encephalitis. *Neuroradiol J* 2017;30:160-3.
14. Reeves HM, Winters JL. The mechanisms of action of plasma exchange. *Br J Haematol* 2014;164:342-51.
15. Virhammar J, Kumlien E, Fallmar D, et al. Acute necrotizing encephalopathy with SARS-CoV-2 RNA confirmed in cerebrospinal fluid. *Neurology* 2020.
16. Baumgartner A, Rauer S, Mader I, Meyer PT. Cerebral FDG-PET and MRI findings in autoimmune limbic encephalitis: correlation with autoantibody types. *J Neurol* 2013;260:2744-53.

AUTHOR CONTRIBUTIONS: AF and JLC collected the clinical data and developed the theoretical framework; JM performed the immunohistochemistry experiments; DF analysed the radiological data; EK interpreted the neurophysiology results; EA and AR aided in sample acquirement; JLC designed the study and drafted the manuscript; AR, JM, JV and JLC designed the figure. All authors helped interpret the results and worked on the manuscript. All authors discussed the results and commented on the manuscript.

ETHICS: This study was approved by the Swedish Ethical Board. The patient has approved the publication of his case and signed an informed consent form. All animal experiments conformed to the European Communities Council Directive (86/609/EEC) and in accordance with Swedish laws and regulations, and all experiments were approved by the local ethical committee (Stockholms Norra Djurförsöksetiska Nämnd N183/14).

Figure 1: Panel A shows a timeline of evaluations, treatments and progression of symptoms. PLEX, plasmapheresis; ICU, intensive care unit; LP, lumbar puncture; MRI, magnetic resonance imaging, PET- positron emission tomography.

Panel B shows an original 12-hour chart recording taken before plasmapheresis (PLEX) (Day 23) with peripheral oxygen saturation (% SpO₂, blue), arterial blood pressure (BP, red) and heart rate (HR, green). The vertical bars represent hours. The patient was initially sedated with an intravenous dexmedetomidine infusion. A hypoxic event of unknown origin was noted (arrow 1). Blood pressure and heart rate were generally variable with episodes of hypotension associated with hyperlactatemia (arrow 2) and hypertension (arrow 3). Hypotension was treated with intravenous crystalloid fluid and noradrenalin. Induction of anaesthesia and endotracheal intubation (arrow 3) was followed by hypertension (arrow 4) that persisted after treatment with bolus doses of propofol, morphine, clonidine and labetalol. After endotracheal intubation, this treatment regimen was changed to propofol, clonidine and morphine.

Panel C shows a brain PET scan 60 minutes after injection of 209 MBq of ¹⁸F-FDG 13 days after plasmapheresis treatment. The uptake in the caudate nucleus and putamen is bilaterally more prominent than the uptake in the cortex. This observation could represent a global cortical decrease in metabolism or symmetrical hypermetabolism of the striatum. On visual inspection, there were no regions with focal cortical anomalies.

Panel D shows profiling of IgG immunoreactivity on mouse brain tissue. CSF (diluted 1:4) collected at the first lumbar puncture revealed a moderate to strong immunoreactivity in neuron-like cells in all investigated brain regions. In the hippocampal formation the strongest immunoreactivity was observed in the CA3 pyramidal neurons (E) labelling the nucleus, soma and proximal dendrites in the stratum lucidum (arrowheads in F). In other regions of the hippocampal formation IgG immunoreactivity was found in a small number of neuron-like cells in the pyramidal layer, the oriens layer (G) and the polymorph layer of the dentate gyrus (E). A

similar neuronal staining pattern was observed in the somatosensory cortex (H), with the strongest labelling of pyramidal neurons in layer V and the thalamus (I) and the strongest labelling of neurons in the paraventricular nucleus and reticular nucleus compared to dorsolateral and ventrolateral thalamic nuclei (J and K). A subpopulation of ependymal cells lining the dorsal 3rd ventricle and choroid plexus (L) reveals nuclear and cytoplasm staining. The highest level of immunoreactivity was found in neurons and neuropil in the caudate-putamen (M). Note the dotted spine-like staining pattern (arrows in M). CSF (N) and serum (S) from before treatment compared to after treatment (O and T) revealed less of the same pattern of immunoreactivity with a remarkably reduced intensity when compared. Reference was CSF (diluted 1:4; P and Q) and serum (diluted 1:1,000 U and V) from two age-matched men with Bipolar I disorder in manic phase. Quantification of immunoreactivity before and after treatment is shown for CSF respective serum (R and W). Abbreviations: CA(1-3), cornu ammonis (1-3); CPU, caudate-putamen; CTX, cerebral cortex; ic, internal capsule; DG, dentate gyrus; DL, dorsolateral thalamus; DM, dorsomedial thalamus; fi, fimbria of the hippocampus; HIFO, hippocampal formation; LV, lateral ventricle; or, oriens layer; PV, paraventricular thalamic nucleus; pyr, pyramidal layer; rad, radiatum layer; Rt, reticular nucleus; Slu, stratum lucidum; THAL, thalamus; VL, ventrolateral thalamus.

A

B

C PET Day 35

D

Hippocampal formation

Cerebral cortex

Thalamus

Ch. plexus

Caudate putamen

admission

post-treatment

reference #1

reference #2

relative mean intensity