A Comprehensive Analysis of COVID-19 Transmission and Fatality Rates at the County level in the United States considering Socio-Demographics, Health Indicators, Mobility Trends and Health Care Infrastructure Attributes

Tanmoy Bhowmik*

Post-Doctoral Scholar

Department of Civil, Environmental & Construction Engineering, University of Central Florida

Tel: 407-927-6574; Fax: 1-407-823-3315; Email: tanmoy78@knights.ucf.edu

ORCID number: 0000-0002-0258-1692

Sudipta Dey Tirtha

Graduate Student

Department of Civil, Environmental & Construction Engineering, University of Central Florida

Tel: 407-543-7521; Email: sudiptadeytirtha2018@knights.ucf.edu

ORCID number: 0000-0002-6228-0904

Naveen Chandra Iraganaboina

Graduate Student

Department of Civil, Environmental & Construction Engineering, University of Central Florida

Tel: 321-295-2134; Email: naveen.chandra@knights.ucf.edu

ORCID number: 0000-0002-1425-6897

Naveen Eluru

Professor

Department of Civil, Environmental & Construction Engineering, University of Central Florida

Tel: 407-823-4815; Fax: 407-823-3315; Email: naveen.eluru@ucf.edu

ORCID number: 0000-0003-1221-4113

*Corresponding author

SUMMARY

Background: Several research efforts have evaluated the impact of various factors including a) socio-demographics, (b) health indicators, (c) mobility trends, and (d) health care infrastructure attributes on COVID-19 transmission and mortality rate. However, earlier research focused only on a subset of variable groups (predominantly one or two) that can contribute to the COVID-19 transmission/mortality rate. The current study effort is designed to remedy this by analyzing COVID-19 transmission/mortality rates considering a comprehensive set of factors in a unified framework.

Method: We study two per capita dependent variables: (1) daily COVID-19 transmission rates and (2) total COVID-19 mortality rates. The first variable is modeled using a linear mixed model while the later dimension is analyzed using a linear regression approach. The model results are augmented with a sensitivity analysis to predict the impact of mobility restrictions at a county level.

Findings: Several county level factors including proportion of African-Americans, income inequality, health indicators associated with Asthma, Cancer, HIV and heart disease, percentage of stay at home individuals, testing infrastructure and Intensive Care Unit capacity impact transmission and/or mortality rates. From the policy analysis, we find that enforcing a stay at home order that can ensure a 50% stay at home rate can result in a potential reduction of about 30% in daily cases.

Interpretation: The model framework developed can be employed by government agencies to evaluate the influence of reduced mobility on transmission rates at a county level while accommodating for various county specific factors. Based on our policy analysis, the study findings support a county level stay at home order for regions currently experiencing a surge in transmission. The model framework can also be employed to identify vulnerable counties that need to be prioritized based on health indicators for current support and/or preferential vaccination plans (when available).

Funding: None.

Keywords: COVID-19, transmission rate, mortality rate, linear mixed model, policy analysis, vulnerable counties

RESEARCH IN CONTEXT

Evidence before this study

We conducted an exhaustive review of studies examining the factors affecting COVID-19 transmission and mortality rates at an aggregate spatial location such as national, regional, state, county, city and zip code levels. The review considered articles published in peer-reviewed journals (via PubMed and Web of Science) and working articles uploaded in preprint platforms (such as medRxiv). A majority of these studies focused on a small number of counties (up to 100 counties) and considered COVID-19 data only up to the month of April. While these studies are informative, cases in the US grew substantially in recent months. Further, earlier studies have considered factors selectively from the four variable groups - socio-demographics, health indicators, mobility trends, and health care infrastructure attributes. The exclusion of variables from these groups is likely to yield incorrect/biased estimates for the factors considered.

Added value of this study

The proposed study enhances the coverage of COVID-19 data in our analysis. Spatially, we consider 1258 counties encompassing 87% of the total population and 96% of the total confirmed COVID-19 cases. Temporally, we consider data from March 25th to July 3rd, 2020. The model system developed comprehensively examines factors affecting COVID-19 from all four categories of variables described above. The county level daily transmission data has multiple observations for each county. To accommodate for these repeated measures, we employ a linear mixed modeling framework for model estimation. The model estimation results are augmented with policy scenarios imposing hypothetical mobility restrictions.

Implications of all the available evidence

The proposed framework and the results can allow policy makers to (a) evaluate the influence of population behavior factors such as mobility trends on virus transmission (while accounting for other county level factors), (b) identify priority locations for health infrastructure support as the pandemic evolves, and (c) prioritize vulnerable counties across the country for vaccination (when available).

INTRODUCTION

Coronavirus disease 2019 (COVID-19) pandemic, as of July 19th, has spread to 188 countries with a reported 14.4 million cases and 603 thousand fatalities¹. The pandemic has affected the mental and physical health of people across the world significantly taxing the social, health and economic systems². Among the various countries affected, United States has reported the highest number of confirmed cases (3.8 million) and deaths (140 thousand) in the world³. In this context, it is important that we clearly understand the factors affecting COVID-19 transmission and mortality rate to prescribe policy actions grounded in empirical evidence to slow the spread of the transmission and/or prepare action plans for potential vaccination programs in the near future. Towards contributing to these objectives, the current study develops a comprehensive framework for examining COVID-19 transmission and fatality rates in the United States using COVID-19 data at a county level encompassing about 87% of the US population. The study effort is designed with the objective of including a universal set of factors affecting COVID-19 in the analysis of transmission and mortality rates. We employ an exhaustive set of county level characteristics including (a) socio-demographics, (b) health indicators, (c) mobility trends, and (d) health care infrastructure attributes. We recognize that analysis of COVID-19 data with a subset of factors, as has been the case with earlier work, is likely to yield incorrect/biased estimates for the factors considered. The framework proposed for understanding and quantifying the influence of these factors can allow policy makers to (a) evaluate the influence of population behavior factors such as mobility trends on virus transmission (while accounting for other county level factors), (b) identify priority locations for health infrastructure support as the pandemic evolves, and (c) prioritize vulnerable counties across the country for vaccination (when available).

In recent months, a number of research efforts have examined COVID-19 data in several countries to identify the factors influencing COVID-19 transmission and mortality. Given the focus of our current study, we restrict our review to studies that explore COVID-19 transmission and mortality rate at an aggregated spatial scale. To elaborate, these studies explored COVID-19 transmission and mortality rates at the national⁴⁻⁶, regional⁷, state⁸, county⁹⁻¹⁴, city¹⁵ and zip code levels¹⁶. A majority of these studies considered transmission rate as the response variable (transmission rate per capita). The main approach employed to identify the factors affecting the response variables is the linear regression approach. In their analysis, researchers employed a host of independent variables from four variable categories: socio-demographics, health indicators, mobility trends and health care infrastructure attributes. For socio- demographics, studies found income, race and age distribution have a positive association with the COVID-19 transmission^{11,16-18}. Regarding health indicators, earlier research found that smokers, obese and individuals with existing health conditions are more likely to be severely affected by COVID-19¹¹. In terms of mobility trends, studies showed that staying at home and effective mobility restriction measures significantly lower the COVID-19 transmission rate^{4,7,9,10,14} while increased mobility resulted in increased COVID-19 transmission^{12,19}. Finally, among health care infrastructure attributes, testing rate is linked with reduced risk of COVID-19 transmission^{4,5}. While earlier research efforts have considered the factors from all variable categories, it is important to recognize that each individual study focused only on a subset of variable groups (predominantly one or two) and have not controlled explicitly for other variable groups that can contribute to the COVID-19 transmission/mortality rate.

The current study builds on earlier literature examining the factors affecting COVID-19 transmission and mortality rate and contributes along the following directions. *First*, we extensively enhance the spatial and temporal coverage of COVID-19 data in our analysis. *Spatially*, earlier research on COVID-19 aggregate data has focused on a small number of counties (up to 100 counties). In our study, we consider all counties with total number of cases greater than 100 on July 3rd. The 1258 counties selected encompass 87% of the total population and 96% of the total confirmed COVID-19 cases. *Temporally*, earlier research has only considered data up to the month of April. While these studies are informative, cases in the US grew substantially in the recent months. Hence, in our study we have considered data from March 25th to July 3rd, 2020. The longer period of data (101 days) also enables us to study/test for the evolution of variable effects over time. *Second*, earlier research studies have considered factors from one or two of the categories of variables identified above. Further, studies that tested health indicators employed one or two measures selectively. In our analysis, we conduct a comprehensive examination of factors affecting COVID-19 from all four categories of variables including (a) socio-demographics: distribution by age, gender, race, income, location (urban or rural), education status, income inequality and employment, (b) health indicators: percentage of population suffering from cancer, cardiovascular disease, hepatitis, Chronic Obstructive Pulmonary Disease (COPD); diabetes, obesity, Human Immunodeficiency Virus (HIV), heart disease, kidney disease, asthma; drinking and smoking habits, (c) mobility trends: daily average exposure, social distancing matrices, percentage of people staying at home, and (d) health care infrastructure attributes: hospitals per capita, ICU beds per capita, COVID-19 testing measures. *Finally*, the research study employs a robust modeling framework in terms of model structure and dependent variable representation. A linear mixed model system that addresses the limitations of the traditional linear regression framework for handling repeated measures is employed. For dependent variable, alternative functional forms of COVID-19 transmission – natural logarithm of daily cases per 100 thousand people and natural logarithm of 7-day moving average of cases per 100 thousand people - are considered in model estimation. The overall approach allows us to robustly quantify the impact of factors affecting COVID-19 transmission.

METHODS

Data Collection

Independent variables: Table 1 summarizes sample characteristics of the explanatory variables with the definition considered for final model estimation, the data source, and sample characteristics (minimum, maximum and mean values). For the sake of brevity, in the ensuing discussion we only present details of two groups of independent variables: health indicators and mobility trends. Using health indicator data, we ranked the 1,258 counties in a descending order of health metric and provided it in Figure 1. Further, we compute the average values for different health indicators across the healthiest and unhealthiest 10 counties to highlight the change in health conditions across the two groups. The values clearly emphasize the vulnerability of the unhealthiest counties relative to the healthiest counties. For instance, number of HIV patients in the healthy counties are 75·83 while in the unhealthiest counties, it is almost 430% higher (407).

<i>Health Status</i>	<i>% Diabetic</i>	<i>% Hepatitis C</i>	<i>% Asthma</i>	<i>Cancer rate (100K people)</i>	<i>HIV rate (100K people)</i>	<i>Cardio Rate (1k people)</i>
Healthiest 10 Counties	8.09	0.21	8.39	393.26	75.83	30.92
Unhealthiest 10 Counties	12.60	5.01	11.79	468.61	407.00	72.23

Figure 1 Ranking of Counties Based on Health Indicators

To incorporate mobility trends, we considered two variables: daily average exposure²⁰ and social distancing metric (from SafeGraph¹) to serve as surrogate measures for the mobility patterns (see Supplementary Material for detail). Figure 2 provides a summary of both these measures at a state level from January 22nd to July 3rd. From the figure, we can clearly see the reduction in average daily exposure in March as many states and local jurisdictions imposed lockdowns. By late April, exposure activity started to increase again across all the states while still being lower than the levels for February. In terms of the staying at home measure, as expected, we find an exactly opposite trend.

Figure 2 Average Daily Exposure and Percentage of People Staying at Home

Dependent variables: We analyze two county level dependent variables: (1) COVID-19 daily transmission rate per 100K population and (2) COVID-19 mortality rates per 100K population. For the transmission rate analysis, we tested two alternative functional forms: daily cases per 100 thousand people and 7-day moving average of cases per 100 thousand people. The moving average data is likely to be less volatile and serves as a stability test for the daily cases model. The reader would note that we used a natural logarithmic transformation for all the dependent variables. The COVID-19 dataset from Center for Systems Science and Engineering (CSSE) Coronavirus Resource Center at Johns Hopkins University²¹ provides information on the daily confirmed

¹ SafeGraph is a data company that aggregates anonymized location data from numerous applications in order to provide insights about physical places. To enhance privacy, SafeGraph excludes census block group information if fewer than five devices visited an establishment in a month from a given census block group

COVID-19 cases, number of people recovered (when available) and the number of deaths from COVID-19 starting from January 22nd to the current date across 3,142 counties in the United States. In our research, we confined our analysis to the cases between March 25th to July 3rd resulting in 101 days of data. Further, we focus on counties that have at least 100 cases by July 3rd and have available information on the mobility trends. With this requirement, a total of 1,258 counties are included in the analysis providing a coverage of 87% of the total population in the United States. For mortality rate, we considered the fatalities within the same time frame across all the 1,258 counties as the transmission rate variable. The summary statistics of the dependent variable are presented in bottom row panel of Table 1.

Data Analysis

For the analysis of daily COVID-19 transmission rate, we have repeated measures of the variable (101 repetitions for each county). The traditional linear regression model is not appropriate to study data with multiple repeated observations²². Hence, we employ a linear mixed modeling approach that builds on the linear regression model while incorporating the influence of repeated observations from the same county. A brief description of the linear mixed model is provided in the Supplementary Material. For modeling the COVID 19 mortality rate, we rely on simple linear regression approach as the dependent variable here is the total number of COVID-19 deaths per 100K population at a county level. Both models are estimated in SPSS.

Role of the Funding Source

There was no funding source for this study.

RESULTS

COVID-19 Transmission Rate Model Results

The estimation results for the linear mixed model are presented in Table 2².

Socio-demographics: In terms of female population, we find that higher proportion of females in the population has a positive impact on transmission rate. The result is in contrast to earlier studies that show women are less likely to be affected by COVID-19 transmission relative to men¹⁶. Among age and racial distribution proportions, we found that increased percentage of younger individuals (<18 years) and African-Americans is associated with more transmission (see earlier work for similar findings^{11,18}). It has been suggested that African-Americans in general reside in densely populated low income neighborhoods with lower access to amenities and are employed in industries that requires more public exposure¹⁷. Educational attainment in a county also plays an important role in influencing the COVID-19 transmission. The counties with higher share of individuals with less than high school education are likely to report increased incidence of COVID-19. In terms of income, we find that higher median income counties have a higher incidence of COVID-19. The effect of income might appear counter-intuitive at first glance. However, it is possible that higher income individuals are more likely to get tested (even in the absence of symptoms) due to higher health insurance affordability. With respect to employment rate, counties with higher employment rate reflect more exposure and have a positive association with

² As discussed earlier, we also developed the same mixed linear model to estimate the 7-day moving average of COVID-19 cases per capita and find similar results as in the daily COVID-19 transmission model (results are available upon request from the authors). This further reinforces the stability of the transmission model.

transmission. The percentage of people living in rural area offers a negative association with the daily COVID-19 incidence. This is intuitive as rural areas are sparsely populated and hence have more opportunity for social distancing thus lowering transmission rates.

Health indicators: With respect to health indicators, we tried several variables in the transmission rate model. Of these, two variables - number of people suffering from HIV and hepatitis C in a county offered significant impacts. We observe that counties with higher percentage of HIV and hepatitis C patients have an increased incidence of COVID-19 transmission. Individuals with these diseases have weaker immune systems and hence are more susceptible to COVID-19 transmission.

Mobility Trends: In terms of mobility trends, we tested two measures: daily average exposure and percentage of people staying at home. In considering these variables in the model, we recognize that exposure will have a lagged effect on transmission i.e. exposure to virus today is likely to manifest as a case in the next 5 to 14 days. In our analysis, we tested several lag combinations and selected the 10 day lag exposure as it offered the best fit. The exposure variable offers interesting results. Until April 25th, exposure variable does not have any impact on transmission. This trend strongly coincides with the lower exposure trends (see Figure 2). After April 25th, increased exposure is associated with higher transmission rates 10 days into the future (see Hamada and colleagues¹⁹ for similar findings). For the second measure, staying at home with 14 days lag, we find that daily transmission rates are affected as expected^{4,10}. The reader would note that the two measures considered were not found to be correlated and thus were simultaneously considered in the model.

Health Care Infrastructure Attributes: From Table 2, we find that counties with more hospitals per capita are more likely to report higher COVID-19 transmission rate. This result perhaps accounts for the higher availability of COVID-19 testing. The last set of variables within this category corresponds to COVID-19 testing effects. Again, we select a 5 day lag as testing results are likely to be reported in 3-5 days. The coefficient of this variable is positive as expected and highly significant⁴. However, after May 10th, the effect has a lower magnitude, which suggests that compared to the previous time period (before May 10th), higher testing rate will increase the daily COVID-19 transmission at a marginally lower rate.

Temporal factors: With data available for 101 days, we can evaluate the effect of the transmission rate in previous days on the current day. As expected, we find a positive association between the daily COVID-19 transmission rate and the number of cases 7 and 14 days prior. The result suggests higher transmission rate in previous time periods (7 and 14 days earlier) is likely to result in increased transmission. However, the effect is higher for the 7 day lagged variable, as evidenced by the higher magnitude associated with the corresponding time period in Table 2. Further, the 7 day lagged transmission rate after June 21st implies a higher impact perhaps explaining the sudden surge in COVID-19 cases in recent weeks. Finally, the weekend variable highlights that the COVID-19 transmission rate is lower during weekends possibly because of reduced testing rate on weekends²³.

Correlation: As indicated earlier, we developed the mixed linear model for estimating the daily COVID-19 transmission rate per 100,000 people while incorporating the dependencies across each county for various repetition levels (such as week and month). We found that the model

accommodating weekly correlations provided the best result in terms of statistical data fit and variable interpretation. The final set of variables in Table 2 corresponds to the correlation parameter across every 7 days within a county.

COVID-19 Mortality Rate

The coefficients in Table 3 represent the effect of different independent variables on COVID-19 mortality rate at a county level.

Socio-demographics: A higher percentage of older people in a county leads to an increased COVID-19 mortality rate (see ^{14,18} for similar findings). Further, consistent with previous research¹⁷, the current analysis also found that the percentage of African-Americans is positively associated with COVID-19 mortality rate. The variable specific to education attainment indicates that the likelihood of COVID-19 mortality increases with increasing share of people with less than high school education in a county. We find that counties with higher income inequality are more likely to experience higher number of COVID-19 deaths per capita relative to the counties with lower income disparities²⁴. Finally, higher employment rate has a positive association with COVID-19 mortality rate.

Health Indicators: Several variables significantly influence the COVID-19 mortality rate in a county. For instance, in comparison to other counties, counties with higher number of HIV, cancer, asthma and cardiovascular patients are more likely to have higher number of COVID-19 deaths. This is expected²⁵⁻²⁸ as people with such conditions usually have weaker immune system which makes them vulnerable to the disease.

Health Care Infrastructure Attributes: The number of ICU beds per capita at a county is found to have a negative impact on COVID-19 mortality rate suggesting a reduced death rate with higher number of ICU bed per person in a county. The result is intuitive as more ICU bed per capita indicates the county is well equipped to handle higher patient demand and treatment is accessible to more COVID-19 patients.

Policy Implications

To illustrate the applicability of the proposed COVID-19 transmission model, we conduct a scenario analysis exercise by imposing mobility restrictions. While earlier researchers explored the influence of mobility measures, these models did not account for county level factors such as socio-demographics, health indicators and hospital infrastructure attributes. In our framework, the sensitivity analysis is conducted while controlling for these factors. The hypothetical restrictions on mobility are considered through the following changes to two variables:

- (1) county level average daily exposure reduced by 10%, 25% and 50%
- (2) county level percentage of stay at home population increased to 40%, 50% and 60%.

The changes to the independent variables were used to predict the dependent variable. Subsequently, the variable was converted to the daily cases per 100 thousand people. The results from this exercise are presented in Table 4. We present the average change in cases for all counties (1,258), and for the 25 counties with the highest overall transmission rates. From Table 4, two important observations can be made. First, changes to average daily exposure and stay at home population influence COVID-19 transmission significantly. In fact, by increasing stay at home population share to 50% the model predicts a reduction of the number of cases by about 30%.

Second, the benefit from mobility restrictions is slightly higher for the 25 counties with higher overall cases. The two observations provide evidence that issuing lockdown orders in counties with a recent surge is a potential mitigation measure to curb future transmission.

The COVID-19 total mortality rate model can be employed to identify vulnerable counties that need to be prioritized for vaccination programs (when available). While prioritizing the counties based on mortality rate might be a potential approach, it might not always be feasible. To elaborate, vaccination programs have to be planned well in advance (say 2 months) of the vaccine availability. As total mortality rates for 2 months into the future are unavailable, we need a model to predict total mortality into the future. The estimated mortality rate model provides a framework for such analysis. To be sure, it would be prudent to update the proposed model with the latest data to develop a more accurate prediction system.

DISCUSSION

The current study develops a comprehensive framework for examining COVID-19 transmission and fatality rates in the United States at a county level including an exhaustive set of independent variables: socio-demographics, health indicators, mobility trends and health care infrastructure attributes. In our analysis, we consider all counties with total number of cases greater than 100 on July 3rd and analyze daily cases data from March 25th to July 3rd, 2020. The COVID-19 transmission rate is modeled at a daily basis using a linear mixed method while the total mortality rate is analyzed adopting a linear regression approach.

Several county level factors including proportion of African-Americans, income inequality, health indicators associated with Asthma, Cancer, HIV and heart disease, percentage of stay at home individuals, testing infrastructure and Intensive Care Unit capacity impact transmission and/or mortality rates. The results clearly support our hypothesis of considering a universal set of factors in analyzing the COVID-19 data. Further we conducted policy scenario analysis to evaluate the influence of social distancing on the COVID-19 transmission rate. The results highlight the effectiveness of social distancing in mitigating the virus transmission. In fact, we found that by increasing stay at home population share to 50% the model predicts a reduction of the number of cases by about 30%. The finding provides evidence that issuing lockdown orders in counties with a recent surge is a potential mitigation measure to curb future transmission. To be sure, the study is not without limitations. The study is focused on county level analysis and is intended to reflect associations as opposed to causation. For such causation based analysis, data from individuals would be more suitable. While exposure data were reasonably addressed, data was not available for mask wearing behavior across all counties. Finally, the data on transmission and mortality are updated for few counties to correct for errors or omissions. These were carefully considered in our data preparation. However, it is possible that further updates might be made after we finished our analysis.

Contributors

NE conceptualized the study. TB and NE finalized the study design. TB, SD and NC conducted the literature review. TB, SD and NC collected the data. TB, SD, NC, and NE analyzed and interpreted the model results. TB, NC and SD prepared the figures. TB, SD, NC and NE drafted the main manuscript. All authors reviewed the results and approved the final version of the manuscript.

Declaration of Interests

We declare no competing interests.

Acknowledgement

The authors would like to gratefully acknowledge SafeGraph COVID-19 Data Consortium, County Health Ranking and Road Maps, Centers for Disease Control System for providing access to the data at county level for United States.

REFERENCES

1. Worldometer. Coronavirus Cases [Internet]. [cited 2020 Jul 12]. p. 1–22. Available from: <https://www.worldometers.info/coronavirus/>
2. The Global Economic Outlook During the COVID-19 Pandemic: A Changed World [Internet]. [cited 2020 Jul 12]. Available from: <https://www.worldbank.org/en/news/feature/2020/06/08/the-global-economic-outlook-during-the-covid-19-pandemic-a-changed-world>
3. Cases in the U.S. | CDC [Internet]. [cited 2020 Jul 12]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/cases-in-us.html>
4. Berger DW, Herkenhoff KF, Mongey S. An seir infectious disease model with testing and conditional quarantine. National Bureau of Economic Research. 2020.
5. Omori R, Mizumoto K, Chowell G. Changes in testing rates could mask the novel coronavirus disease (COVID-19) growth rate. International Journal of Infectious Diseases. 2020.
6. Yuan X, Xu J, Hussain S, Wang H, Gao N, Zhang L. Trends and Prediction in Daily New Cases and Deaths of COVID-19 in the United States: An Internet Search-Interest Based Model. Exploratory research and hypothesis in medicine. 2020;5(2):1.
7. Courtemanche CJ, Garuccio J, Le A, Pinkston JC, Yelowitz A. Did Social-Distancing Measures in Kentucky Help to Flatten the COVID-19 Curve? 2020.
8. Hu B, Gong J. Support vector machine based classification analysis of SARS spatial distribution. In: 2010 Sixth international conference on natural computation; 2010 (Vol. 2, pp. 924-927). IEEE.
9. Engle S, Stromme J, Zhou A. Staying at home: mobility effects of covid-19. Available at SSRN. 2020.
10. Bilgin NM. Tracking COVID-19 Spread in Italy with Mobility Data. Available at SSRN. 2020.
11. Mollalo A, Vahedi B, Rivera KM. GIS-based spatial modeling of COVID-19 incidence rate in the continental United States. Science of The Total Environment. 2020.
12. Sharkey P, Wood G. The Causal Effect of Social Distancing on the Spread of SARS-CoV-2. 2020.
13. Wu X, Nethery RC, Sabath BM, Braun D, Dominici F. Exposure to air pollution and COVID-19 mortality in the United States. medRxiv. 2020.
14. Wieland T. Flatten the Curve! Modeling SARS-CoV-2/COVID-19 Growth in Germany on the County Level. medRxiv. 2020.
15. Xie J, Zhu Y. Association between ambient temperature and COVID-19 infection in 122 cities from China. Science of The Total Environment. 2020.
16. Borjas GJ. Demographic determinants of testing incidence and COVID-19 infections in New York City neighborhoods. National Bureau of Economic Research. 2020.

17. Backer A. Why COVID-19 May Be Disproportionately Killing African Americans: Black Overrepresentation among COVID-19 Mortality Increases with Lower Irradiance, Where Ethnicity Is More Predictive of COVID-19 Infection and Mortality Than Median Income. SSRN Electron J. 2020.
18. Xie Z, Li D. Health and Demographic Impact on COVID-19 Infection and Mortality in US Counties. Medrxiv. 2020.
19. Badr HS, Du H, Marshall M, Dong E, Squire MM, Gardner LM. Association between mobility patterns and COVID-19 transmission in the USA: a mathematical modelling study. The Lancet Infectious Diseases. 2020.
20. Couture V, Dingel JI, Green A, Handbury J, Williams KR. Measuring Movement and Social Contact with Smartphone Data: A Real-Time Application to COVID-19. University of Chicago Booth School of Business Working Paper. 2020.
21. COVID-19 Map - Johns Hopkins Coronavirus Resource Center [Internet]. [cited 2020 Jul 11]. Available from: <https://coronavirus.jhu.edu/map.html>
22. Faghih-Imani A, Eluru N, El-Geneidy AM, Rabbat M, Haq U. How land-use and urban form impact bicycle flows: evidence from the bicycle-sharing system (BIXI) in Montreal. Journal of Transport Geography. 2014;41:306-14.
23. Reduced testing suggested as reason for weekend drop in confirmed COVID-19 deaths | Michigan Radio [Internet]. [cited 2020 Jul 11]. Available from: <https://www.michiganradio.org/post/reduced-testing-suggested-reason-weekend-drop-confirmed-covid-19-deaths>
24. Income and wealth inequality in the U.S. has fueled COVID-19 deaths - MarketWatch [Internet]. [cited 2020 Jul 11]. Available from: <https://www.marketwatch.com/story/income-and-wealth-inequality-in-the-us-has-fueled-covid-19-deaths-2020-06-29>
25. Mehra MR, Desai SS, Kuy S, Henry TD, Patel AN. Cardiovascular disease, drug therapy, and mortality in COVID-19. New England Journal of Medicine. 2020.
26. Bansal M. Cardiovascular disease and COVID-19. Diabetes & Metabolic Syndrome: Clinical Research & Reviews. 2020.
27. Common Questions About the New Coronavirus Outbreak [Internet]. [cited 2020 Jul 11]. Available from: <https://www.cancer.org/latest-news/common-questions-about-the-new-coronavirus-outbreak.html>
28. Zhang L, Zhu F, Xie L, Wang C, Wang J, Chen R, Jia P, Guan HQ, Peng L, Chen Y, Peng P. Clinical characteristics of COVID-19-infected cancer patients: a retrospective case study in three hospitals within Wuhan, China. Annals of Oncology. 2020.
29. The COVID-19 tracking project [Internet]. [cited 2020 June 12]. Available from: <https://covidtracking.com/data/>.

Table 1 Descriptive Statistics of the Dependent and Independent Variables

Variables	Source	Mean	Min/Max	Sample Size
Independent Variables				
<i>Demographic Characteristics</i>				
Percentage of population aged 18 years and lower	ACS ^a	22.705	7.155/33.882	1258
Percentage of population aged 65 years and over	ACS	16.644	7.545/56.944	1258
Percentage of African American	ACS	13.133	0.113/80.507	1258
Percentage of Hispanic	ACS	11.331	0.623/96.323	1258
Percentage of Female	ACS	50.507	37.041/54.495	1258
Ln (Median income)	ACS	10.903	10.150/11.820	1258
Percentage of people less than high school education	ACS	14.003	3.127/47.053	1258
Employment rate per capita	ACS	0.446	0.195/0.634	1258
Income inequality ratio (80 th percentile/20 th percentile)	CHRR ^b	4.574	3.150/9.148	1258
<i>Health Indicators</i>				
Ln (HIV Prevalence Rate per 100K people)	CHRR	5.043	0.723/7.859	1258
Hepatitis B Cases per 100K people in 2017	CDC ^c	1.250	0.000/11.700	1258
Hepatitis C Cases per 100K people in 2017	CDC	0.980	0.000/5.600	1258
Asthma % for >= 18 years	CDC	9.328	7.400/12.300	1258
COPD % for >= 18 years	CDC	6.692	3.300/13.700	1258
Reported cancer case per 100K people	CDC	455.918	241.000/576.400	1258
Percentage of diabetic	CHRR	11.353	3.300/20.400	1258
Percentage of obesity among adults	CHRR	31.653	13.600/46.700	1258
Cardiovascular Disease Hospitalization Rate per 1,000 Medicare Beneficiaries	CDC	63.609	0.300/110.500	1258
<i>Mobility Trends</i>				
Ln (Daily Average Exposure), 10 days lag				
From April 25th	CEI ^d	3.325	0.102/0.644	127058
% People staying at home				
14 days lag	Safegraph	4.128	0.847/7.049	127058
<i>Healthcare Related Attributes</i>				
Hospitals per 100K people	CHRR	1.967	0.000/15.644	1258
Number of ICU beds per capita	CHRR	20.111	0.000/171.850	1258
Ln (No of tests with 5 days lag)	CTP ^e	11.178	4.320/15.190	5151
<i>Temporal Factors</i>				
Day is weekend	--	0.277	0.000/1.000	127058
Dependent Variables				
Ln (Daily COVID-19 transmission rate per 100K people)	CSSE ^f	1.455	0.000/7.670	127058
Ln (Total COVID-19 mortality rate per 100K people)	CSSE	2.722	0.000/7.120	1258

^a = American Community Survey

^b = County Health Rankings & Roadmaps

^c = Central for Disease Control System

^d = COVID Exposure Indices²⁰

^e = COVID-19 Tracking Project²⁹

^f = Center for Systems Science and Engineering Coronavirus Resource Center at Johns Hopkins University²¹

Table 2 Estimation Results for Daily COVID-19 Transmission Rate per 100K Population

Variables	Estimates	t-statistic	p-value
Constant	-6.071	-16.215	0.000
Demographics			
% of Female population	0.024	7.496	0.000
% Young population (<=18 years)	0.005	2.524	0.012
% of Black population	0.007	15.061	0.000
% of People less than high school education	0.025	18.454	0.000
Ln (median income)	0.396	12.650	0.000
Employment rate per capita	1.409	9.640	0.000
Ln (% of People living in rural areas)	-0.309	-9.249	0.000
Health Indicators			
Ln (HIV rate per 100K People)	0.052	6.306	0.000
Hepatitis C rate per 100K People	0.027	5.152	0.000
Mobility Trends			
Ln (Daily Average Exposure), 10 days lag			
From April 25th	0.027	9.618	0.000
% People staying at home			
14 days lag	-1.027	-7.690	0.000
Health Care Infrastructure Attributes			
Hospital per 100K People	0.028	8.958	0.000
Ln (Testing), 5 days lag			
March 25 th to May 10th	0.020	10.481	0.000
After May 10th	0.016	7.155	0.000
Temporal Factors			
Temporal Lagged Variables			
7 days lag (March 25 th to June 21 st)	0.205	68.500	0.000
7 days lag (After June 21 st)	0.340	60.026	0.000
14 days lag	0.148	50.225	0.000
Day is Weekend	-0.033	-5.925	0.000
Correlation			
σ	0.973	194.881	0.000
ρ	0.960	305.240	0.000
Φ	0.321	83.977	0.000

Table 3 Estimation Results for COVID-19 Mortality Rate per 100K Population

Variables	Estimates	t-statistic	p-value
Constant	-6.952	-7.715	0.000
Demographics			
Older people % (>65 years old)	0.068	6.649	0.000
Black people%	0.018	5.749	0.000
% of People less than high school education	0.050	6.153	0.000
Employment rate per capita	8.369	7.853	0.000
Income inequality ratio	0.274	4.795	0.000
Ln (% of People living in rural areas)	-0.772	-3.173	0.002
Health Indicators			
Ln (HIV rate per 100K people)	0.082	1.557	0.120
Cancer rate per 100K people	0.003	3.397	0.001
% People having Asthma	0.091	2.720	0.007
Cardiovascular disease per 1K people	0.007	2.304	0.021
Health Care Infrastructure Attributes			
ICU beds per capita	-0.009	-4.794	0.000

Table 4 Policy Scenario Analysis of Social Distancing in COVID-19 Transmission Rate per 100K Population

Hypothetical Scenarios	1,258 Counties	Worst 25 Counties
1: daily average exposure reduced by 10%	-0.225	-0.240
2: daily average exposure reduced by 25%	-0.612	-0.653
3: daily average exposure reduced by 50%	-1.465	-1.565
4: 40% people stay at home	-22.949	-23.206
5: 50% people stay at home	-30.469	-30.701
6: 60% people stay at home	-37.256	-37.465