Hypertension among people living with HIV/AIDS in Cameroon: A cross-sectional analysis from Central Africa International Epidemiology Databases to Evaluate AIDS.

Authors

Anastase Dzudie^{§1,2,3} MD PhD FESC, Don Hoover⁴ PhD, Hae-Young Kim⁵ DrPH, Rogers Ajeh¹ PhD, Adebola Adedimeji⁶ PhD MPH MBA, Qiuhu Shi⁵ PhD, Walter Pefura Yone^{2,7} MD PhD, Denis Nsame Nforiwe⁸ MD, Kinge Thompson Njie⁹ MD, Andre Pascal Kengne¹ MD PhD, Vanes Peter Ebasone¹ MD, Blaise Barche¹ MD, Zoung-Kany Bissek Anne Cecile^{2,10} MD, Denis Nash¹¹ PhD, Marcel Yotebieng⁶ MD PhD, Kathryn Anastos⁶ MD

Affiliations

- 1) Clinical Research Education, Networking and Consultancy
- 2) Faculty of Medicine and Biomedical Sciences, University of Yaounde I
- 3) Service of Internal Medicine, Douala General Hospital, Douala, Cameroon
- 4) Department of Statistics and Institute for Health, Health Care Policy and Aging Research, Rutgers the State University of New Jersey, NJ, USA
- 5) Department of Public Health, New York Medical College, Valhalla, NY, USA
- 6) Albert Einstein College of Medicine, Bronx, NY, USA
- 7) Jamot Hospital, Yaounde, Cameroon
- 8) Limbe Regional Hospital, Limbe, Cameroon
- 9) Bamenda Regional Hospital, Bamenda, Cameroon
- 10) Institute for Division of operational health research (DROS), Ministry of Public Health (MSP), Yaoundé, Cameroon
- 11) Implementation Science in Population Health, Graduate School of Public Health and Health Policy, City University of New York (CUNY), USA

Key words - HIV infection, Hypertension, Non communicable diseases, Africa

§ Corresponding author: Professor Anastase Dzudie, *Service of Internal Medicine and Cardiology, Douala General Hospital, Cameroon* Po box 4856, Douala, Cameroon, Email: aitdzudie@yahoo.com

Abstract

Background: Antiretroviral therapy (ART) success has led people to live longer with HIV/AIDS (PLWH) and thus be exposed to increasing risk of cardiovascular diseases (CVD). Hypertension (HTN), the biggest contributor to CVD burden, is a growing concern among PLWH. The current report describes the prevalence and predictors of HTN among PLWH in care in Cameroon.

Methods: This cross-sectional study included all PLWH aged 20 years and above who received care between 2016 and 2019 at one of the three Central Africa International Epidemiology Databases to Evaluate AIDS (CA-IeDEA) sites in Cameroon (Bamenda, Limbe, and Yaoundé). HTN was defined as blood pressure (BP) $\geq 140/90$ mm Hg or self-reported use of antihypertensive medication. Logistic regressions models examined the relationship between HTN and clinical characteristics, and HIV-related factors.

Results

Among 9,839 eligible PLWH, 66.2% were female and 25.0% had prevalent HTN [age-standardized prevalence 23.9% (95% CI: 22.2–25.6)], among whom 28 (1.1%) were on BP lowering treatment, and 6 of those (21.4%) were at target BP levels. Median age (47.4 vs. 40.5 years), self-reported duration of HIV infection (5.1 vs 2.8 years), duration of ART exposure (4.7 vs 2.3 years), and CD4 count (408 vs 359 cell/mm³) were higher in hypertensives than non-hypertensives (all $p < 0.001$). Age and body mass index (BMI) were independently associated with higher prevalent HTN risk. PLWH starting ART had a 30% lower risk of prevalent HTN, but this advantage disappeared after a cumulative 2-year exposure to ART. There was no significant association between other HIV predictive characteristics and HTN.

Conclusion

About a quarter of these Cameroonian PLWH had HTN, driven among others by age and adiposity. Appropriate integration of HIV and NCDs services is needed to improve early detection, treatment and control of common comorbid NCD risk factors like hypertension and safeguard cardiovascular health in PLWH.

INTRODUCTION

The advent and widespread use of antiretroviral therapy (ART) has successfully achieved viral suppression in 76% of sub-Saharan Africans living with HIV/AIDS [1]. Since 2005, there has also been a decline in AIDS-related deaths by 35% globally and by 39% in sub-Saharan Africa (SSA). This decline is partly attributed to new recommendations which entails providing antiretroviral therapy (ART) for all people living with HIV (PLWH) at diagnosis irrespective of CD4 counts, otherwise known as “*Treat All*” [2]. These improvements have narrowed the gap in life expectancy between persons with and without HIV [3][4]. However, increases in life expectancy among PLWH are accompanied by increased occurrence of non-communicable diseases (NCDs), particularly cardiovascular disease (CVD) [5][6].

Hypertension (HTN) remains the leading risk factor for CVD which is the main cause of death globally, accounting for 10.4 million deaths annually [7][8]. In 2015, the World Health Organization (WHO) indicated that the number of people with HTN in the general population was highest in sub-Sahara Africa (SSA), at about 46% of adults aged 25 years and older, compared to 35 to 40% elsewhere [8]. In PLWH, previous studies in Africa and elsewhere showed a high prevalence of HTN [6,9] with some studies estimating that one-quarter of PLWH have HTN [10].

In SSA, reported HTN prevalence in PLWH ranged from 12.5% to 28.5%, including in Cameroon where two single center cross-sectional studies of small sample size estimated that 25% and 28% of PLWH were hypertensive in 2015 and 2016 [11–13]. In a systematic narrative review, Nguyen et al estimated that HTN prevalence in PLWH from low- and middle-income countries ranged from 8.7 to 45.9% [14], suggesting that HTN burden in PLWH populations has not been well-established or otherwise may vary greatly. Accurate estimates of HTN burden in SSA PLWH is needed to guide policies that will facilitate optimization of both HIV and HTN in HIV clinics as recommended by the Pan African Society of cardiology [15].

Cameroon bears the largest burden of HIV/AIDS in Central Africa and one of the highest in SSA with an estimated number of PLWH around 540,000 in 2018 [16]. According to the Cameroon

Population-Based HIV Impact Assessment study, 52% of this HIV-infected population was receiving ART and had access to healthcare services in 2017[17]. However, HIV care is generally provided in dedicated HIV clinics while patients with NCDs receive care in general medical services, with little or no overlap in management of HIV and NCDs, particularly HTN, limiting the assessment of the true burden of NCDs in PLWH. Similarly, because of the lack of integration of NCDs and HIV care services, data are limited on access to HTN treatment and outcomes among PLWH in SSA in general and in Cameroon in particular. Such information would help in allocation of resources, and development of cost-effective therapeutic strategies and programs for integrating HIV and HTN management. We therefore sought to examine the prevalence and predictors of HTN among PLWH attending three large clinics in Cameroon.

METHODS

Study design and setting

This cross-sectional study used data from the International Epidemiology Databases to Evaluate AIDS (IeDEA) in Cameroon. Cameroon, Burundi, the Democratic Republic of Congo, the Republic of Congo and Rwanda, comprise central Africa-IeDEA (CA-IeDEA) contributing secondary data from medical records to the regional data center, one of the seven that compose the global IeDEA consortium [18]. Each regional data center collaborates with clinical sites to identify and define key variables, harmonize and effectively analyze the data, and generate large datasets. In Cameroon, three clinical sites participate in the CA-IeDEA: the Limbe Regional Hospital (LRH), the Bamenda Regional Hospital (BRH), and the Yaoundé Jamot Hospital (YJH). At these public health sites, ART and care delivery is supported by the Cameroon National AIDS Control Committee.

Population, enrolment into IeDEA

In Cameroon, the National Ethical Committee requires a written consent from participants before their data are extracted from medical records into the IeDEA database. Starting in January 2016, all PLWH in care at the HIV clinic in each of the three sites were approached during a routine clinic visit and invited to participate in IeDEA and consented. During the consenting visit

(enrolment into leDEA), additional demographic and clinical variables were collected to supplement routine data, including blood pressure (BP), weight and height.

For this study, we included all non-pregnant (at enrolment) HIV-positive participants aged 20 years or older enrolled into leDEA from January 2016 through December 2019. We excluded patients with missing systolic and diastolic BP measurements within the first six months of enrolment into leDEA.

Patient data are routinely collected by research assistants on case report forms. Data are then captured from paper forms into REDCap[19,20], a free, secure and flexible web-based clinical research data capture platform. These data included demographic characteristics, clinical conditions including weight, height, and systolic and diastolic blood pressure (BP), HIV diagnosis, duration and disease control, laboratory test results, medication use (antiretroviral (ARV) and non-ARV). During the study period, the standard first-line regimen for PLWH in Cameroon was a combination of three drugs including two nucleoside or nucleotide reverse transcriptase inhibitors (NRTI) like zidovudine (AZT) or lamivudine (3TC) or Tenofovir (TDF) and a non-nucleoside reverse transcriptase inhibitor (NNRTI) like efavirenz (EFV) or nevirapine (NVP) as either AZT/3TC/EFV, AZT/3TC/NVP, TDF/3TC/EFV or TDF/3TC/NVP. Second-line regimens during this period included protease inhibitor- (PI) and Integrase Strand Transfer Inhibitor (INSTI) based therapy.

Blood pressure measurement

Blood pressure measurements were obtained during the enrolment visit by trained healthcare professionals, using validated automated BP monitors and following a standardized protocol for office BP measurement. Briefly, patients were seated comfortably in a quiet office for 5 min. BP measurements were done on the right arm using an OMRON M3 HEM-7131-E device (Omron Healthcare Co Ltd, Kyoto, Japan) and appropriate cuff size. Three BP measurements were recorded 1–2 min apart, and an average of the two last readings used in the analysis.

Definition of outcomes

Our primary outcome was HTN at enrolment in leDEA. HTN was defined following the European Society of Cardiology(ESC)/European Society of HTN (ESH) guidelines for HTN[21] as Systolic BP (SBP) ≥ 140 mmHg and/or diastolic BP (DBP) ≥ 90 mmHg and/or current use of antihypertensive medication from medical records. HTN was further graded following the ESC/ESH guidelines of HTN as optimal for SBP < 120 mmHg and DBP < 80 mmHg, normal for SBP 120-129 mmHg and/or DBP 80-84 mmHg, high normal for SBP 130-139 mmHg and/or DBP 85-89 mmHg, grade 1 HTN for SBP 140-159 mmHg and/or DBP 90-99 mmHg, grade 2 for SBP 160-179 mmHg and/or DBP 100-109 mmHg, grade 3 for SBP ≥ 180 mmHg and/or DBP ≥ 110 mmHg and isolated systemic hypertension for SBP ≥ 140 mmHg and DBP < 90 mmHg [21]. Secondary outcomes included the proportion of all hypertensive participants receiving treatment for HTN at the time of enrolment into leDEA and controlled HTN defined as BP $< 140/90$ mmHg in those on antihypertensive treatment.

Potential predictors of hypertension

Values of all predictors investigated were collected at time of enrolment. Candidate predictors included age (categorized into 18-29, 30-49, ≥ 50 years), sex (male vs female), marital status (single, married, live with a partner, separated or divorced and widowed), education level (none, primary, secondary up to high school, high school or more), employment status (employed and unemployed), smoking status (never, current, former) and alcohol consumption (never, monthly or less, 2- 3 standard drinks per months, $\geq$ one per standard drink week). A standard alcoholic drink was considered as equivalent to 1 can (340mL) of beer, 1 glass (125mL) wine, or 1 shot (25mL) of spirits.

The following clinical parameters were also considered as candidate predictors: body mass index (BMI, kg/m^2 , categorized as underweight ≤ 18 , normal 18-24.9, mildly overweight 25-29.9, overweight 30-34.9, or obese ≥ 35 kg/m^2); blood glucose categorized based on fasting plasma glucose (FPG) as normal (FPG < 100 mg/dL), impaired fasting glucose (FPG between 100 and 125 mg/dL) and diabetes (FPG ≥ 126 mg/d or ongoing antidiabetic medications). HIV disease categories were based on the WHO classification [22]. ART exposure was categorized as having

received ART prior to the date of enrollment in study. Estimated duration on ART among users was the time from the first prescription of ART to the date of enrolment BP measurement. Diagnosed duration of HIV infection was estimated time from first recorded or self-reported HIV positive test to the date of enrollment into CA-leDEA project.

Statistical analysis

The crude prevalence of HTN was calculated among study participants with available BP measurement, while the proportion receiving treatment was calculated among all hypertensive participants and the proportion with controlled calculated among those receiving treatment. Age-standardized sex-specific prevalence was calculated using the age distribution of PLWH in the 2018 distribution of the Cameroon HIV population [23]. Medians and quartiles (Q1 = 25th percentile, Q3 = 75th percentile) and proportions were used to summarize continuous and categorical variables, respectively. Chi-square (χ^2) tests and equivalents compared proportions, Student t-test, Analysis of the variance (ANOVA) or rank tests as appropriate compared continuous variables. Logistic regression models estimated the odds ratios (OR) and 95% confidence intervals (95% CI) as measure of the association between candidate predictors and HTN. Only variables found to be associated with the outcome in crude analysis ($p < 0.05$) were included in the multivariable logistic models, but duration of HIV infection was excluded due to collinearity with age and duration of ART use. The limited number of hypertensive participants on antihypertensive treatment precluded similar investigations for the two secondary outcomes. Data were analysed using SAS Version 9.4 (Cary, NC). A p -value < 0.05 was considered statistically significant.

Results

Cohort description

Of 9,988 PLWH eligible for study inclusion, 149 (1.5%) were excluded due to missing data on BP measurement (Figure 1). The remaining 9,839 participants were included in this analysis. Table 1 depicts the distribution of participants' characteristics overall and by HTN status. In all, 6,502 (66.2%) were women, the median age was 42 years (Q1, Q3: 35-50), 3,225 (32.9%) were single;

only 1,720 (17.5%) had high school training or more and 6,245 (64.8%) were unemployed. Median time from HIV diagnosis to enrollment into leDEA was 3.4 years (Q1, Q3: 0.2-8.1), 6,645 (67.5%) participants were on ART at enrollment with 6,312 (95.4%) on NNRTI-, 405 (6.1%) on PI- and only 4 (0.1%) on INSTI-based regimen. Among those on ART, median duration of use was 2.9 years (Q1, Q3: 0.0-7.6). Poor general health status described by WHO clinical stage III and IV was noted in 1,249 (29.9%) and 419 (10.0%) respectively. Median CD4 count was 373 cells/mm³ (Q1, Q3: 184-573), median systolic and diastolic BP were 122 (Q1, Q3: 107-133) and 76 (Q1, Q3: 69-86) mmHg, respectively, median BMI was 24.1 (Q1, Q3: 21.4-27.7) kg/m² and median hemoglobin 11.2 (Q1, Q3: 9.8-12.6) g/dl.

Table 1. Demographic and clinical characteristics of people leaving with HIV/AIDS at enrolment with and without hypertension

Characteristics	Total n=9839	Hypertension n=2450	No hypertension n=7389	p- value
Age (years)				
Median (25th-75 th percentiles)	42.0 (35.0,50.0)	47.4 (40.1,55.0)	40.5 (34.0- 48.0)	<0.001
(Min, Max)	18.1 -105.8	19.0-87.1	18.1-105.8	
Age category, n (%)				
18-29	1146 (11.16)	106 (4.3)	1040 (14.1)	<0.001
30-39	3053 (31.0)	505 (20.6)	2548 (34.5)	
40-49	3220 (32.7)	860 (35.1)	2360 (31.9)	
≥50	2420 (24.6)	979 (40.0)	1441(19.5)	
Gender (Female), n (%)	6502 (66.2)	1555 (63.5)	4947 (67.1)	0.001
Marital Status, n (%)				
Single	3225 (32.9)	615 (25.2)	2610 (35.4)	<0.001
Married	3287 (33.5)	916 (37.5)	2371(32.2)	
Live with a partner	872 (8.9)	148 (6.1)	724 (9.8)	
Separated/Divorced	695 (7.1)	193 (7.9)	502 (6.8)	
Widowed	1734(17.7)	572(23.4)	1162(15.8)	
Level of education, n (%)				
Never went to school	959 (9.8)	271(11.1)	688 (9.3)	0.0021
Primary	4256 (43.4)	1059 (43.4)	3197(43.4)	
Secondary	2875 (29.3)	655 (26.9)	2220 (30.1)	
High school or more	1720 (17.5)	454 (18.6)	1266 (17.2)	
Employment status, n (%)				
Employed	3542 (35.2)	856 (35.2)	2686 (35.5)	0.228
Unemployed	6245(64.8)	1578(64.8)	4668(63.5)	
Smoking status, n (%)				
Never smoked	8237 (84.0)	2037 (83.5)	6200 (84.1)	0.003

Characteristics	Total n=9839	Hypertension n=2450	No hypertension n=7389	p- value
Current smoker	325 (3.3)	60 (2.5)	265 (3.6)	
Former smoker	1244 (12.7)	341(14.0)	903 (12.3)	
Alcohol consumption, n (%)				
Never	3937 (40.2)	959 (39.3)	2978 (40.5)	0.708
Monthly or less	2988 (30.5)	760 (31.2)	2228 (30.3)	
2-3 times per month	1170 (11.9)	299 (12.3)	871 (11.8)	
Once a week or more	1706 (17.4)	421 (17.3)	1285 (17.5)	
History of diabetes, n (%)				
No	9704 (98.6)	2383 (97.3)	7321 (99.1)	<0.001
Yes	135 (1.4)	67 (2.7)	68 (0.9)	
Estimated duration of HIV infection (years)				
Median (25th-75 th percentiles)	3.4 (0.2, 8.1)	5.1(1.1, 9.6)	2.8 (0.1, 7.4)	<0.001
(Min, Max)	(0.0,25.8)	(0.0, 25.4)	(0.0, 25.8)	
Any ART use, n (%)				
No ART	3194 (32.5)	956 (39.0)	2238 (30.3)	<0.001
Yes	6645 (67.5)	1494 (61.0)	5151 (69.7)	
PI among ART users, n (%)				
No	6203(93.9)	1392(93.7)	4817(93.9)	0.799
Yes	405(6.1)	93(6.3)	312(6.1)	
NRTI among ART users, n (%)				
Yes	6614(100.0)	1485(100.0)	5129(100.0)	
NNRTI among ART users, n (%)				
No	302(4.6)	60(4.0)	242(4.7)	0.271
Yes	6312(95.4)	1425(96.0)	4887(95.3)	

Characteristics	Total n=9839	Hypertension n=2450	No hypertension n=7389	p- value
INSTI among ART users, n (%)				
No	6610(99.9)	1483(99.9)	5127(100.0)	0.186
Yes	4(0.1)	2(0.1)	2(0.0)	
ART duration (years)				
Median (25th-75 th percentiles)	2.9(0.0, 7.6)	4.7(0.7, 9.2)	2.3(0.0, 6.9)	<0.001
(Min, Max)	(0.0, 18.5)	(0.0, 17.6)	(0.0, 18.5)	
CD4 cell count				
Median (IQR)	373 (184, 573)	409 (23, 598)	359 (171, 558)	<0.001
(Min, Max)	(0, 1825)	(0, 1825)	(0, 1753)	
Bp lowering medication, n (%)				
No	9811(99.7)	2422 (98.9)	7389 (100.0)	
Yes	28 (0.3)	28 (1.1)		
Systolic blood pressure (mmHg)				
Median (25th-75 th percentiles)	119.0 (107.0, 134.0)	147.0 (139.0, 160.0)	113.0 (103.0, 123.0)	<0.001
(Min, Max)	(47.0, 276)	(93.0, 276)	47.0, 139.0)	
Diastolic blood pressure (mmHg)				
Median (25th-75 th percentiles)	77.0(69.0, 86.0)	94.0(90.0, 101.0)	73.0(66.0, 80.0)	<0.001
(Min, Max)	(20.0, 190.0)	(38.0, 190.0)	(20.0, 89.0)	
Body mass index (kg/m²)				
Median (25th-75 th percentiles)	24.1(21.4, 27.7)	25.9(22.8, 30.1)	23.5(21.0, 27.0)	<0.001
(Min, Max)	(10.4, 87.2)	(10.9, 87.2)	(10.4, 73.5)	

Characteristics	Total n=9839	Hypertension n=2450	No hypertension n=7389	p- value
Body mass index categories				
(kg/m²), n (%)				
<18	431(4.6)	47(2.0)	384(5.5)	<0.001
18-24.9	4943(53.2)	976(41.7)	3967(57.1)	
25-29.9	2474(26.6)	720(30.8)	1754(25.2)	
30-35	1038(11.2)	398(17.0)	640(9.2)	
>35	401(4.3)	199(8.5)	202(2.9)	
Fasting blood sugar (mg/dl)				
Median (25th-75 th percentiles)	88.0(79.0, 97.0)	90.0(80.0, 99.0)	87.0(78.0, 96.0)	<0.001
(Min, Max)	(41.0, 860.0)	(47.6, 519.0)	(41.0, 860.0)	
Glucose categories, n (%)				
Normal fasting glucose	3228(32.8)	784(32.0)	2444(33.1)	<0.001
Impaired fasting glucose	666(6.8)	200(8.2)	466(6.3)	
Diabetes / provisional diabetes	264(2.7)	102(4.2)	162(2.2)	
No glucose measure	5681(57.7)	1364(55.7)	4317(58.4)	

ART: antiretroviral therapy, BP: blood pressure, INSTI: integrase strand transfer inhibitor, NRTI: nucleoside reverse-transcriptase inhibitor, NNRTI: non-nucleoside reverse-transcriptase inhibitor.

Prevalence of HTN

A quarter of our cohort (n = 2,450, 25.0%) had hypertension HTN at the time of enrollment into leDEA (figure 1). The sex-specific prevalence of HTN was 26.6% (95% CI: 25.1-28.1) in men and 23.7% (95% CI: 22.7-24.7) in women for crude prevalence (Figure 2) (P < 0.001); and 23.9% (95% CI: 22.2-25.6) and 20.6% (95% CI: 19.5-21.7) after age-standardization. Overall, 1,328 (13.5%) PLWH had high normal BP, 966 (9.8%) had mild HTN, and 511 (5.2%) grade 2 and 373 (3.8%) grade 3 (Figure 2).

Figure 2. Grade of Blood pressure by 2018 ESC/ESH Guidelines for the overall group (left), female (middle) and male (right).

Table 1 compares a comparison of demographic and clinical characteristics between hypertensive and non-hypertensive PLWH. HTN prevalence increased with increasing age, starting at 9.2% among the <18 year old rising to 40.5% among those aged ≥ 50 years ($p < 0.001$) (Table 1). Median diagnosed duration of HIV infection [5.1(1.1, 9.6) vs 2.8(0.1, 7.4) years, $p < 0.001$] and duration on ART [4.7(0.0, 7.6) vs 2.3(0.0, 6.9), $p < 0.001$] was longer in participants with HTN than those without HTN. Hypertension was lower among those who did than did not initiate ART (61.0 Vs. 69.7%, $p < 0.001$). However, among those who initiated ART duration on ART was longer for those with than without HTN [4.7(0.0, 7.6) vs 2.3(0.0, 6.9), $p < 0.001$]. Based on Loess plots and analyses using indicator variables among ART users once duration reached 2 years, prevalent HTN did not increase further with duration on ART after adjusting for age;

meanwhile participants with HTN exhibited higher CD4 count than those without HTN [409 cells/mm³ (231 598) vs 359 (171, 558), $p < 0.001$], Table 1.

Figure 1: flow chart of the study participants.

BP: blood pressure, HTN: hypertension

Predictors of hypertension

Table 2 presents OR and 95%CI from crude and multivariable logistic regression models. In multivariable models including age, sex, marital status, BMI, blood glucose level, ART use, ART duration up to 2 years and age, factors significantly associated with HTN were age, sex, BMI, ART use and duration. A 5-year increase in age was independently associated with 28% higher odds of prevalent HTN (adjusted odds ratio (aOR)=1.28, 95%CI: 1.25-1.32). The adjusted OR of prevalent HTN comparing participants aged ≥ 50 years versus those aged 18-29 years was 4.80 (95%CI: 3.78-6.10). Compared to participants with BMI < 18 kg/m², the adjusted odds of prevalent HTN was 1.87 (95%CI: 1.36-2.57), 2.97 (95%CI: 2.15-4.10), 4.37 (95%CI: 3.11-6.13), and 7.32 (95%CI: 5.04- 10.64) respectively among in those with BMI between 18 to 24.9, 25 to 29.9, 30 to 35, and > 35 kg/m². Among those with this measure, higher blood glucose was also independently associated with prevalent HTN with an aOR=1.47 (95% CI:1.10-1.95) for those with diabetes compared to those with normal blood glucose.

Table 2. Predictors of prevalent hypertension

Variable	Unadjusted OR** (95% CI)	Adjusted OR (95% CI)
Socio-demographic factors		
Age Group		
18-29	Ref	Ref
30-39	1.94 (1.56, 2.42)	1.60 (1.27, 2.02)
40-49	3.57 (2.88, 4.43)	2.65 (2.10, 3.34)
50+	6.66 (5.37, 8.26)	4.80 (3.78, 6.10)
Gender		
Women	Ref	Ref
Men	1.17 (1.06, 1.29)	1.38 (1.23, 1.54)
Marital status		
Single	Ref	Ref
Married	1.64 (1.46, 1.84)	1.01 (0.89, 1.16)
Live with a partner	0.87 (0.71, 1.06)	0.87 (0.71, 1.07)
Separated/Divorced	1.63 (1.35, 1.97)	0.98 (0.79, 1.20)
Widowed	2.09 (1.83, 2.39)	1.12 (0.96, 1.32)
Education		
Never went to school	Ref	
Primary	0.84 (0.72, 0.98)	
Secondary	0.75 (0.63, 0.88)	
High School or more	0.91 (0.76, 1.09)	
Employment		
Unemployed	Ref	
Employed	1.06 (0.96, 1.17)	
Cardiovascular risk factors		
Smoking status		
Never smoked	Ref	
Current smoker	0.69 (0.52, 0.92)	
Former smoker	1.15 (1.00, 1.31)	
Alcohol consumption		
Never	Ref	
Monthly or less	1.06 (0.95, 1.18)	
2-3 times per month	1.07 (0.92, 1.24)	
Once a week or more	1.02 (0.89, 1.16)	
Diabetes		
No	Ref	
Yes	3.03 (2.15, 4.26)	
Systolic BP (mmHg)	6.85 (6.26, 7.49)	
Diastolic BP (mmHg)	14.22 (12.61, 16.03)	

Variable	Unadjusted OR** (95% CI)	Adjusted OR (95% CI)
BMI (kg/m²)	1.09 (1.08, 1.10)	
BMI (kg/m²) categories		
< 18	Ref	Ref
18 to 24.9	2.01 (1.47, 2.74)	1.87 (1.36, 2.57)
25 to 29.9	3.35 (2.45, 4.59)	2.97 (2.15, 4.10)
30 to 35	5.08 (3.66, 7.05)	4.37 (3.11, 6.13)
> 35	8.05 (5.61, 11.54)	7.32 (5.04, 10.64)
Fasting blood sugar (mg/dl)	1.02 (1.00, 1.04)	
Glucose categories		
Normal fasting glucose	Ref	Ref
Impaired fasting glucose	1.34 (1.11, 1.61)	1.23 (1.00, 1.50)
Diabetes / provisional diabetes	1.96 (1.51, 2.55)	1.47 (1.10, 1.95)
No glucose measure	0.98 (0.89, 1.09)	1.09 (0.97, 1.23)
HIV-related factors		
Estimated duration of HIV infection (per year of use)	1.07 (1.06, 1.08)	1.15 (1.07-1.24)
Any ART use		
No	Ref	Ref
Yes	0.68 (0.62, 0.75)	0.70 (0.62, 0.80)
PI among ART users		
No	Ref	
Yes	1.03 (0.81, 1.31)	
NNRTI among ART users		
No	Ref	
Yes	1.18 (0.88, 1.57)	
INSTI among ART users		
No	Ref	
Yes	3.46 (0.49, 24.56)	
ART Duration up to 2 years (capped at 2 years, non-ART use considered as 0 years)	1.10 (1.05, 1.16)	1.15 (1.07, 1.24)
CD4 count	1.06 (1.03, 1.09)	
Log10-transformed Viral Load	0.87 (0.77, 0.99)	

ART: antiretroviral therapy, BMI: Body Mass Index, BP: blood pressure, NRTI: nucleoside reverse-transcriptase inhibitor, NNRTI: non-nucleoside reverse-transcriptase inhibitor.

Participants on ART at enrollment independently had 30% lower odds of prevalent HTN compared to those not on ART (aOR: 0.70; 95% CI: 0.62-0.80), but there was no significant association between ART regimen (NNRTI or PI) and prevalent HTN (Table 2). Among those using ART, being on ART for up to 2 years was independently associated with higher prevalent HTN (aOR=1.15/year of ART use, 95% CI:1.07-1.24). Further duration on ART beyond 2 years did not increase the risk of hypertension beyond the fact that those using ART longer tended to be older based on Loess plots (Figure 3). There was a very slow rise in HTN prevalence after 2 years (Figure 3).

Figure 3. Loess for prevalent hypertension and duration of Antiretroviral therapy (ART) use.

Treatment and control of HTN

Among participants with HTN, 2422(98.9%) were not receiving any antihypertensive medication; and of the 28 (1.1%) receiving treatment, 6 (21.4%) had a controlled BP (figure 1).

Discussion

Our study represents the first multicenter and the largest study of HTN in PLWH in west and central Africa and one of the largest in SSA. We found a high crude prevalence of HTN (25%) in PLWH receiving routine care at three HIV clinics across three major cities in the Cameroon, coupled with unacceptably low treatment and control rates. This high prevalence was largely driven by conventional risk factors such as age, sex, and excess weight, and to some extent associated with HIV-related factors such as ART use and duration on ART.

Our findings extend to the population of PLWH in Cameroon and SSA at large, previous observations on the high burden of HTN in the general population across Africa. For instance, we have previously demonstrated in several studies [24–26] that despite being the main cause of stroke, heart and renal disease in Africa, HTN while largely prevalent is still poorly diagnosed, poorly treated and controlled across the continent. Our study clearly highlights the missed opportunity of optimally co-addressing comorbid HTN in PLWH who are already in regular contacts with the health system and at risk of HTN-mediated organ damage.

Prevalence of hypertension

Over the last decade, three systematic reviews have exclusively focused on the burden of HTN in PLWH. In 2015, Nguyen et al reported prevalence rates ranging from 8% to 45.9% across 13 studies in Africa, of which none were from Cameroon [14]. In a systematic review and metanalysis of 49 studies published during 2011–2016 (10 from Africa including two from Cameroon) with 63,554 participants, Xu et al. reported a global HTN prevalence of 25.2% (95% CI: 21.2% -29.6%) in PLWH [10]. The third and most recent review including 194 studies (396,776 PLWH from 61 countries) reported a 23.6% (95% CI: 21.6-25.5) global prevalence of HTN with little regional variation, 28.1% (95% CI: 24.5-31.9) in Western and Central Europe and North America, with 23.5% (16.6-31.0) in West and Central Africa[27].

Prevalence ranged from 19% to 44.4% across 3 previous Cameroon studies[13][28][29]. It should be noted that these studies all had small samples size ranging from 108 [28] to 311 [29] PLWH and were restricted to single HIV clinics. These shortcomings of Cameroon studies also occurred in studies from other SSA countries and to a certain extent from other regions around the world. For instance, the pooled sample size of PLWH of 17 studies conducted in PLWH in West and Central Africa between 2011-2017 was around 9,010 non randomly selected participants[27].

Apart from the high prevalence, other similarities of HTN burden between our sample of PLWH and the general population lie in the very poor detection, treatment and control rates, as previously described by Mutemwa et al [30] and confirmed by our findings. In 2018, 59.2% of the 8.9 (95% CI: 8.3-9.6) million PLWH with HTN globally were living in Sub-Saharan Africa [27]. In the absence of an integrated HTN and HIV care and no universal health coverage in many SSA countries including Cameroon, PLWH receive ART for free but do not get any BP check and when eventually checked, they do not seek treatment for HTN because of the relatively high cost and poor availability of the medications[31]. This exposes the growing PLWH population to HTN-mediated organ damage and raises strong concerns about the integration of HTN management in HIV clinics including identification of the main drivers of HTN that might be controlled in PLWH.

Predictors of hypertension

Our findings are generally consistent with previous reports emphasizing the key role of conventional factors in predicting HTN. We found that male sex, higher age and higher BMI increased the odds of prevalent HTN. This is similar to what has been reported in several previous publications both in the general population [26] and in PLWH [32]. With regard to HIV related factors, being on ART at enrolment was associated with 30% lower odds of having HTN, but estimated ART duration up to 2 years was associated with an increased prevalent HTN for both sexes. But this study did not show any significant association between ART regimen (NNRTI, NRTI or PI) and prevalent HTN. The role of HIV-specific risk factors contributing to the development of HTN in PLWH is poorly understood and potentially includes ART exposure and related changes in body composition, immunodeficiency, immune activation and inflammation, as well as effects

from antiretroviral therapy itself. Some studies, mostly cross-sectional in design, described higher prevalence rates of HTN in PLWH on ART compared to those not on treatment [29] while others did not find any significant association between ART use and HTN [33][30]. In their metaanalysis and opposite of what we observed, Xu et al[10] described that HTN was 2.7 times more prevalent among ART-exposed PLWH compared with ART naïve. But falling somewhat between, the Strategic Timing of Antiretroviral Treatment clinical trial recently demonstrated that over a follow-up period of 48 months, there were no significant differences in incident HTN between PLWH in the immediate ART group versus deferred treatment group, nor when comparing treated versus untreated HIV infection[34].

We postulate that the protective association of any ART use in our study (aOR = 0.70) may be due to the fact that people initiating ART had more advanced HIV disease i.e. prior to the “treat all” era, had an indication for ART. As proven elsewhere, more advanced HIV disease (WHO stage III/IV, low CD4, and low Hb) is protective against prevalent HTN while low BMI (<18 kg/m²) is protective against incident hypertension [32]. Indeed, before starting ART, PLWH in SSA usually suffer poor nutrition with severe weight loss, which is associated with a reduction in the BP and may be protective against or otherwise mask diagnosis of HTN [31]. Excess weight stands as one of the main risk factors for development of HTN and weight reduction is key in HTN prevention and control strategies [35]. After starting ART, the initial weight gain and immune reconstitution leading to improved overall health status are likely to explain the increased HTN risk among PLWH, especially after sustained exposure to ART exposure.

Our findings have important implications for policy, practice and research. In the face of this double burden of HIV infection and HTN, integrating HTN’s detection, treatment and control into successful vertical programs for diseases such as HIV/AIDS and tuberculosis as recommended by the Pan African Society of Cardiology[15] is urgently needed to curb the burden. Secondly, healthcare providers should pay more attention to NCD and particularly HTN to prevent the occurrence of cardiovascular events in PLWH. And last but not least, longitudinal studies are needed to understand the pathophysiology of HTN in HIV infection as well as interventional

studies to direct innovative and efficient strategies to support prevention and treatment of HTN and HTN-mediated organ damage in PLWH.

Potential limitations and strengths of our study

Our results should be interpreted considering some limitations. First, there were high levels of missing data for some parameters that could impact on HTN status like WHO stage, creatinine, and hemoglobin. As proven elsewhere, more advanced disease (WHO stage III/IV, low CD4, and low Hemoglobin) is protective against prevalent HTN [32]. Second, we did not have access to HIV-negative populations for comparison. Adding this to the cross-sectional nature of the study, we are unable to draw any causal inference between the diagnosis of HTN and HIV. Considering our recruitment at three large HIV clinics, and targeting nearly 2% of the entire HIV population in the country, these are major strengths and vehicles for accuracy and generalizability. Indeed, in several previous published studies of the burden of HTN in Cameroon, we and others have demonstrated that HTN occurs around the fifth decade in the general population and affects 30% of Cameroonians older than 21 years of age [24,26,37], roughly only 5% higher than the estimated 25% we reported in younger population of PLWH and consistent with reports from other robust studies [32] and a contemporary systematic review metanalysis [27]. Finally, we may have underestimated the proportion of participants with HTN receiving treatment, if that treatment was occurring in another location.

Conclusions

In conclusion, our study found that about one of every four PLWH in Cameroon treated at three major clinics has HTN, which is largely under-recognized and under-treated, exposing this specific population group to HTN-mediated organ damages. Drivers of HTN in PLWH include both conventional risk factors like older age, male sex and excess weight as well as specific HIV-related factors including ART exposure and estimated duration of ART exposure. These findings should motivate the integration of HTN services at HIV clinics to stop the growing burden of NCDs among PLWH.

Author contributions

AD, AA, MY, DN, and KA conceived the study and acquired the funding.

AD, RA, WPY, DNN, KTN, VPE, ZKBAC operationalized and supervised data collection

AD, DH, HYK, QS, KA, MY, DN and APK analyzed the data and drafted the manuscript.

All co-authors substantially revised the manuscript and approved the submission.

Acknowledgments list

Nimbona Pélagie, Association Nationale de Soutien aux Séropositifs et Malade du Sida (ANSS), Burundi; Patrick Gateretse, Jeanine Munezero, Valentin Nitereka, Théodore Niyongabo, Christelle Twizere, Centre National de Référence en Matière de VIH/SIDA, Burundi; Hélène Bukuru, Thierry Nahimana, Centre de Prise en Charge Ambulatoire et Multidisciplinaire des PVVIH/SIDA du Centre Hospitalo-Universitaire de Kamenge (CPAMP-CHUK), Burundi; Elysée Baransaka, Patrice Barasukana, Eugene Kabanda, Martin Manirakiza, François Ndikumwenayo, CHUK/Burundi National University, Burundi; Jérémie Biziragusenyuka, Ange Marie Michelline Munezero, Centre de Prise en Charge Ambulatoire et Multidisciplinaire des PVVIH/SIDA de l'Hôpital Prince Régent Charles (CPAMP-HPRC), Burundi; Tabeyang Mbuh, Kinge Thompson Njie, Edmond Tchasse, Kien-Atsu Tsi, Bamenda Hospital, Cameroon; Rogers Ajeh, Mark Benwi, Marc Lionel Ngamani, Victorine Nkome, Grace Toutou, Clinical Research Education and Consultancy (CRENC), Cameroon; Anastase Dzudie, CRENC and Douala General Hospital, Cameroon; Akindéh Mbuh, CRENC and University of Yaoundé, Cameroon; Djenabou Amadou, Amadou Dodo Balkissou, Eric Ngassam, Eric Walter Pefura Yone, Jamot Hospital, Cameroon; Alice Ndelle Ewanoge, Norbert Fuhngwa, Ernestine Kendowo, Chris Moki, Denis Nsame Nforniwe, Limbe Regional Hospital, Cameroon; Catherine Akele, Akili Clever, Faustin Kitetele, Patricia Lelo, Martine Tabala, Kalembелеmbе Pediatric Hospital, Democratic Republic of Congo; Cherubin Ekembe, Didine Kaba, Kinshasa School of Public Health, Democratic Republic of Congo; Merlin Diafouka, Martin Herbas Ekat, Dominique Mahambou Nsonde, CTA Brazzaville, Republic of Congo; Adolphe Mafoua, Massamba Ndala Christ, CTA Pointe-Noire, Republic of Congo; Jules Igrimbabazi, Nicole Ayinkamiye, Bethsaida Health Center, Rwanda; Providance Uwineza, Emmanuel Ndamijimana, Busanza Health Center, Rwanda; Emmanuel Habarurema, Marie Luise Nyiraneza, Gahanga Health Center, Rwanda; Marie Louise Nyiransabimana, Liliane Tuyisenge,

Gikondo Health Center, Rwanda; Christian Shyaka, Catherine Kankindi, Kabuga Health Center, Rwanda; Bonheur Uwakijijwe, Marie Grace Ingabire, Kicukiro Health Center, Rwanda; Jules Ndumuhire, Marie Goretti Nyirabahutu, Masaka Health Center, Rwanda; Fred Muyango, Jean Christophe Bihibindi, Nyagasambu Health Center, Rwanda; Oliver Uwamahoro, Yvette Ndoli, Nyarugunga Health Center, Rwanda; Sabin Nsanzimana, Placidie Mugwaneza, Eric Remera, Esperance Umumararungu, Gallican Nshogoza Rwibasira, Dominique Savio Habimana, Rwanda Biomedical Center, Rwanda; Josephine Gasana, Faustin Kanyabwisha, Gallican Kubwimana, Benjamin Muhoza, Athanase Munyaneza, Gad Murenzi, Francoise Musabyimana, Francine Umwiza, Charles Ingabire, Patrick Tuyisenge, Alex M. Butera, Jules Kabahizi, Ephrem Rurangwa, Rwanda Military Hospital, Rwanda; Rosine Feza, Eugenie Mukashyaka, Shyorongi Health Center, Rwanda; Chantal Benekigeri, Jacqueline Musaninyange, WE-ACTx Health Center, Rwanda.

Coordinating and Data Centers:

Adebola Adedimeji, Kathryn Anastos, Madeline Dilorenzo, Lynn Murchison, Jonathan Ross, Marcel Yotebieng, Albert Einstein College of Medicine, USA; Diane Addison, Ellen Brazier, Heidi Jones, Elizabeth Kelvin, Sarah Kulkarni, Denis Nash, Matthew Romo, Olga Tymejczyk, Institute for Implementation Science in Population Health, Graduate School of Public Health and Health Policy, City University of New York (CUNY), USA; Batya Elul, Columbia University, USA; Xiatao Cai, Allan Dong, Don Hoover, Hae-Young Kim, Chunshan Li, Qihu Shi, Data Solutions, USA; Kathryn Lancaster, The Ohio State University, USA; Mark Kuniholm, University at Albany, State University of New York, USA; Andrew Edmonds, Angela Parcesepe, Jess Edwards, University of North Carolina at Chapel Hill, USA; Olivia Keiser, University of Geneva; Stephany Duda; Vanderbilt University School of Medicine, USA; April Kimmel, Virginia Commonwealth University School of Medicine, USA.

Supporting institutions

Professor Zoung-Kany Bisseck Anne Cecile, Division of operational health research (DROS), Ministry of Public Health (MSP), Cameroon, Dr Elat Mfetam Jean Bosco and Dr Bonono Leonard, National AIDS Control Committee (NACC), MSP, Cameroon Ministry of Public Health, Albert Einstein College of Medicine, USA.

Participants to the study

All PLWH for attending the clinics for their continued trust in our health care system

Funding acknowledgement

Research reported in this publication was supported by the National Institutes of Health's National Institute of Allergy and Infectious Diseases (NIAID), the *Eunice Kennedy Shriver* National Institute of Child Health & Human Development (NICHD), the National Cancer Institute (NCI), the National Institute on Drug Abuse (NIDA), the National Heart, Lung, and Blood Institute (NHLBI), the National Institute on Alcohol Abuse and Alcoholism (NIAAA), the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), the Fogarty International Center (FIC), the National Library of Medicine (NLM), and the Office of the Director (OD) under Award Number U01AI096299 (Central Africa-IeDEA). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. UNIADS. UNAIDS Data 2017. Program HIV/AIDS [Internet]. 2017;1–248. Available from: http://www.unaids.org/sites/default/files/media_asset/20170720_Data_book_2017_en.pdf
2. WHO. WHO | Treatment and care. WHO [Internet]. World Health Organization; 2020 [cited 2020 Jun 30]; Available from: <http://www.who.int/hiv/topics/treatment/en/>
3. Wandeler G, Johnson LF, Egger M. Trends in life expectancy of HIV-positive adults on antiretroviral therapy across the globe. *Curr Opin HIV AIDS* [Internet]. 2016;11:492–500. Available from: <https://journals.lww.com/01222929-201609000-00007>
4. Wang H, Naghavi M, Allen C, Barber RM, Carter A, Casey DC, et al. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. Lancet Publishing Group; 2016;388:1459–544.
5. Hadigan C, Paules CI, Fauci AS. Association Between Human Immunodeficiency Virus Infection and Cardiovascular Diseases. *JAMA Cardiol* [Internet]. 2017;2:123. Available from: <http://cardiology.jamanetwork.com/article.aspx?doi=10.1001/jamacardio.2016.5177>
6. Chow DC, Souza SA, Chen R, Richmond-Crum SM, Grandinetti A, Shikuma C. Elevated Blood Pressure in HIV-Infected Individuals Receiving Highly Active Antiretroviral Therapy.

- HIV Clin Trials [Internet]. 2003;4:411–6. Available from:
<http://www.tandfonline.com/doi/full/10.1310/5E7Q-PGWB-16UE-J48U>
7. Beaney T, Schutte AE, Tomaszewski M, Ariti C, Burrell LM, Castillo RR, et al. May Measurement Month 2017: an analysis of blood pressure screening results worldwide. *Lancet Glob Heal*. 2018;6.
8. WHO | A global brief on hypertension. WHO. World Health Organization; 2013;
9. Okello S, Kanyesigye M, Muyindike WR, Annex BH, Hunt PW, Haneuse S, et al. Incidence and predictors of hypertension in adults with HIV-initiating antiretroviral therapy in south-western Uganda. *J Hypertens* [Internet]. 2015;33:2039–45. Available from:
<http://journals.lww.com/00004872-201510000-00012>
10. Xu Y, Chen X, Wang K. Global prevalence of hypertension among people living with HIV: a systematic review and meta-analysis. *J Am Soc Hypertens* [Internet]. 2017;11:530–40. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1933171117301900>
11. Njelekela M, Muhihi A, Aveika A, Spiegelman D, Hawkins C, Armstrong C, et al. Prevalence of Hypertension and Its Associated Risk Factors among 34,111 HAART Naïve HIV-Infected Adults in Dar es Salaam, Tanzania. *Int J Hypertens* [Internet]. 2016;2016:1–9. Available from: <https://www.hindawi.com/journals/ijhy/2016/5958382/>
12. Kalyesubula R, Kayongo A, Semitala FC, Muhanguzi A, Katantazi N, Ayers D, et al. Trends and level of control of hypertension among adults attending an ambulatory HIV clinic in Kampala, Uganda: a retrospective study. *BMJ Glob Heal* [Internet]. 2016;1:e000055. Available from: <http://gh.bmj.com/lookup/doi/10.1136/bmjgh-2016-000055>
13. Nsagha DS, Assob JCN, Njunda AL, Tanue EA, Kibu OD, Ayima CW, et al. Risk Factors of Cardiovascular Diseases in HIV/AIDS Patients on HAART. *Open AIDS J* [Internet]. 2015;9:51–9. Available from: <https://openaidsjournal.com/VOLUME/9/PAGE/51/>
14. Nguyen KA, Peer N, Mills EJ, Kengne AP. Burden, Determinants, and Pharmacological Management of Hypertension in HIV-Positive Patients and Populations: A Systematic Narrative Review. *AIDS Rev. Spain*; 2015;17:83–95.
15. Dzudie A, Rayner B, Ojji D, Schutte AE, Twagirumukiza M, Damasceno A, et al. Roadmap to achieve 25% hypertension control in Africa by 2025. *Cardiovasc J Afr*. 2017;28:262–73.
16. Cameroon | UNAIDS [Internet]. [cited 2020 Jun 30]. Available from:
<https://www.unaids.org/en/regionscountries/countries/cameroon>

17. Cameroon Population-Based HIV Impact Assessment, CAMPHIA 2017 | Fast-Track Cities.
18. IeDEA International epidemiology Databases to Evaluate AIDS.
19. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)-A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* [Internet]. NIH Public Access; 2009 [cited 2020 Jun 30];42:377–81. Available from: [/pmc/articles/PMC2700030/?report=abstract](https://pubmed.ncbi.nlm.nih.gov/31438304/)
20. Ngamani L, Ajeh R, Mbuh A, Dzudie A, Duda SN. Implementation of a redcap-based research data collection system in Cameroon. *Stud Health Technol Inform* [Internet]. IOS Press; 2019 [cited 2020 Jun 30]. p. 1708–9. Available from: <https://pubmed.ncbi.nlm.nih.gov/31438304/>
21. Williams B, Masera G. 2018 ESC / ESH Guidelines for the management of arterial hypertension The Task Force for the management of arterial hypertension of the [Internet]. *Eur. Heart J*. 2018 [cited 2019 Nov 17]. Available from: www.wjpr.net
22. Organization WH. WHO case definitions of hiv for surveillance and revised clinical staging and immunological classification of hiv-related disease in adults and children hiv/aids Programme. 2007.
23. PEPFAR. Cameroon Country Operational Plan (COP/ROP) 2019 Strategic Direction Summary. 2019;80.
24. Dzudie A, Djomou A, Ba H, Njume E, Ndom MS, Mfekeu LK, et al. MMM17-Cameroon, analysis and opportunities—Sub-Saharan Africa. *Eur Hear J Suppl*. 2019;21:D31–3.
25. Katte J-C, Dzudie A, Sobngwi E, Mbong EN, Fetse GT, Kouam CK, et al. Coincidence of diabetes mellitus and hypertension in a semi-urban Cameroonian population: A cross-sectional study. *BMC Public Health*. 2014;14.
26. Dzudie A, Kengne AP, Muna WFT, Ba H, Menanga A, Kouam Kouam C, et al. Prevalence, awareness, treatment and control of hypertension in a self-selected sub-Saharan African urban population: a cross-sectional study. *BMJ Open* [Internet]. 2012;2:e001217. Available from: <http://bmjopen.bmj.com/lookup/doi/10.1136/bmjopen-2012-001217>
27. Bigna JJ, Ndoadoumgue AL, Nansseu JR, Tochie JN, Nyaga UF, Nkeck JR, et al. Global burden of hypertension among people living with HIV in the era of increased life expectancy. *J Hypertens* [Internet]. 2020;Publish Ah. Available from: <https://journals.lww.com/10.1097/HJH.0000000000002446>

28. Ngatchou W, Lemougoum D, Nga E, Yiagnigni, Tiogou, Kouanfack, et al. Increased burden and severity of metabolic syndrome and arterial stiffness in treatment naïve HIV+ patients from Cameroon. *Vasc Health Risk Manag* [Internet]. 2013;509. Available from: <http://www.dovepress.com/increased-burden-and-severity-of-metabolic-syndrome-and-arterial-stiff-peer-reviewed-article-VHRM>
29. Dimala CA, Atashili J, Mbuagbaw JC, Wilfred A, Monekosso GL. Prevalence of Hypertension in HIV/AIDS Patients on Highly Active Antiretroviral Therapy (HAART) Compared with HAART-Naïve Patients at the Limbe Regional Hospital, Cameroon. Burdmann EA, editor. *PLoS One* [Internet]. 2016;11:e0148100. Available from: <https://dx.plos.org/10.1371/journal.pone.0148100>
30. Mutemwa M, Peer N, de Villiers A, Mukasa B, Matsha TE, Mills EJ, et al. Prevalence, detection, treatment, and control of hypertension in human immunodeficiency virus (HIV)-infected patients attending HIV clinics in the Western Cape Province, South Africa. *Medicine (Baltimore)* [Internet]. 2018;97:e12121. Available from: <http://journals.lww.com/00005792-201808310-00077>
31. Dzudie A, Njume E, Abanda M, Aminde L, Hamadou B, Dzekem B, et al. Availability, cost and affordability of essential cardiovascular disease medicines in the south west region of Cameroon: Preliminary findings from the Cameroon science for disease study. *PLoS One*. 2020;15:e0229307.
32. Brennan AT, Jamieson L, Crowther NJ, Fox MP, George JA, Berry KM, et al. Prevalence, incidence, predictors, treatment, and control of hypertension among HIV-positive adults on antiretroviral treatment in public sector treatment programs in South Africa. Graham SM, editor. *PLoS One* [Internet]. 2018;13:e0204020. Available from: <https://dx.plos.org/10.1371/journal.pone.0204020>
33. Jericó C, Knobel H, Montero M, Sorli ML, Guelar A, Gimeno JL, et al. Hypertension in HIV-infected patients: prevalence and related factors. *Am J Hypertens*. United States; 2005;18:1396–401.
34. Baker J V., Sharma S, Achhra AC, Bernardino JI, Bogner JR, Duprez D, et al. Changes in Cardiovascular Disease Risk Factors With Immediate Versus Deferred Antiretroviral Therapy Initiation Among HIV-Positive Participants in the START (Strategic Timing of Antiretroviral Treatment) Trial. *J Am Heart Assoc*. 2017;6.

35. Wang L, Heizhati M, Zhang D, Chang G, Yao X, Hong J, et al. Excess weight loss is a vital strategy for controlling hypertension among multi-ethnic population in northwest China: A cross-sectional analysis. *Med (United States)*. Lippincott Williams and Wilkins; 2019;98.
36. Guehi C, Badjé A, Gabillard D, Ouattara E, Koulé SO, Moh R, et al. High prevalence of being Overweight and Obese HIV-infected persons, before and after 24months on early ART in the ANRS 12136 Temprano Trial. *AIDS Res Ther*. BioMed Central Ltd.; 2016;13.
37. Kingue S, Ngoe CN, Menanga AP, Jingi AM, Noubiap JJN, Fesuh B, et al. Prevalence and Risk Factors of Hypertension in Urban Areas of Cameroon: A Nationwide Population-Based Cross-Sectional Study. *J Clin Hypertens*. 2015;17:819–24.